

Advanced paternal age at birth: phenotypic and etiologic associations with eating pathology in offspring

S. E. Racine¹, K. M. Culbert², S. A. Burt¹ and K. L. Klump^{1*}

¹Department of Psychology, Michigan State University, East Lansing, MI, USA

²Department of Psychiatry and Behavioral Neuroscience, University of Chicago, Chicago, IL, USA

Background. Advanced paternal age at birth has been linked to several psychiatric disorders in offspring (e.g. schizophrenia) and genetic mechanisms are thought to underlie these associations. This study is the first to investigate whether advanced paternal age at birth is associated with eating disorder risk using a twin study design capable of examining both phenotypic and genetic associations.

Method. In a large, population-based sample of female twins aged 8–17 years in mid-puberty or beyond ($n=1722$), we investigated whether advanced paternal age was positively associated with disordered eating symptoms and an eating disorder history [i.e. anorexia nervosa (AN), bulimia nervosa (BN) or binge eating disorder (BED)] in offspring. Biometric twin models examined whether genetic and/or environmental factors underlie paternal age effects for disordered eating symptoms.

Results. Advanced paternal age was positively associated with disordered eating symptoms and an eating disorder history, where the highest level of pathology was observed in offspring born to fathers ≥ 40 years old. The results were not accounted for by maternal age at birth, body mass index (BMI), socio-economic status (SES), fertility treatment or parental psychiatric history. Twin models indicated decreased genetic, and increased environmental, effects on disordered eating with advanced paternal age.

Conclusions. Advanced paternal age increased risk for the full spectrum of eating pathology, independent of several important covariates. However, contrary to leading hypotheses, environmental rather than genetic factors accounted for paternal age–disordered eating associations. These data highlight the need to explore novel (potentially environmental) mechanisms underlying the effects of advanced paternal age on offspring eating disorder risk.

Received 4 February 2013; Revised 17 May 2013; Accepted 18 May 2013; First published online 24 June 2013

Key words: Advanced paternal age, disordered eating, eating disorders, environmental, genetic, twin study.

Introduction

Advanced paternal age at birth is a risk factor for psychiatric disorders in offspring, including schizophrenia, autism and bipolar disorder (Malaspina *et al.* 2001; Reichenberg *et al.* 2006; Frans *et al.* 2008). These relationships have been confirmed in meta-analytic studies (Wohl & Gorwood, 2007; Hultman *et al.* 2011; Miller *et al.* 2011) and are independent of important confounds [i.e. maternal age at birth, socio-economic status (SES) and parental psychiatric history] that increase psychiatric risk and are associated with later

entry into parenthood (Byrne *et al.* 2003; Croen *et al.* 2007; Menezes *et al.* 2010). Recent data also link paternal age to obesity (Eriksen *et al.* 2012). Despite these robust effects, advanced paternal age has never been examined as a risk factor for eating disorders in offspring. Examining this possibility could provide new insights into the complex etiology of eating disorders.

Leading theories propose that *de novo* genetic mutations are primary mechanisms underlying paternal age–psychiatric risk associations (Malaspina *et al.* 2002). The increased probability of DNA copy error problems with each sperm replication (~610 replications by age 40) and the accumulation of mutations in the germline of older fathers are thought to lead to disease phenotypes in offspring (Crow, 2000). Genetic studies indicate that offspring *de novo* mutation rates are influenced by paternal age at conception (Kong *et al.* 2012) and *de novo* mutations are implicated in

* Address for correspondence: K. L. Klump, Ph.D., Department of Psychology, Michigan State University, 316 Physics Road-107B, East Lansing, MI 48824-1116, USA.

(Email: klump@msu.edu)

Parts of this paper were presented at the Eating Disorders Research Society meeting, Edinburgh, Scotland, UK, 22–24 September 2011.

schizophrenia and autism (Awadalla *et al.* 2010; Xu *et al.* 2011; Iossifov *et al.* 2012). However, to our knowledge, no study has directly confirmed that *de novo* mutations underlie paternal age-offspring psychiatric risk associations.

Twin studies can be used to indirectly examine the role of *de novo* mutations in paternal age-psychiatric disorder relationships. Because monozygotic (MZ) twins share all of their genes and dizygotic (DZ) twins share, on average, half of their segregating genetic material, MZ twins would be expected to share all *de novo* mutations whereas DZ twins would be very unlikely to share genetic mutations given their rarity (Zhao *et al.* 2007; Liu *et al.* 2010; Ronald & Hoekstra, 2011). If *de novo* mutations underlie paternal age-psychiatric risk associations, larger differences between MZ and DZ twin correlations should be observed with advancing paternal age because DZ co-twins should become less similar as mutation rates increase.

One previous study examined these processes by investigating twin similarity for autism across paternal age (Lundström *et al.* 2010). Importantly, differences in MZ/DZ concordance for autism decreased with advanced paternal age, implicating environmental rather than genetic processes. Although the results were limited by the small number of cases in each age category ($n=3-44$), they highlight the need to test hypotheses regarding mechanisms for paternal age effects. Twin registries are excellent low-cost resources for such investigations, as they can indirectly examine whether *de novo* mutations or other competing processes are most likely to underlie paternal age-psychiatric risk associations.

The current study investigated phenotypic and etiologic associations between advanced paternal age and disordered eating symptoms (e.g. weight/shape concerns) in a large, population-based sample of female twins. These symptoms are core features of eating disorders and prospectively predict the development of diagnoses (Jacobi *et al.* 2004); thus, our results should inform risk models of eating disorders. Exploratory analyses examining associations between advanced paternal age and an eating disorder history [anorexia nervosa (AN), bulimia nervosa (BN), binge eating disorder (BED)] were also conducted to investigate whether the results generalize across categorical and continuous eating disorder dimensions (APA, 2000; Insel *et al.* 2010). Several covariates were included [e.g. maternal age, body mass index (BMI) percentile, SES, fertility treatment, parental psychiatric history] to ensure that associations were not due to confounding factors. Finally, we compared twin similarity for disordered eating across paternal age to explore the mechanisms (i.e. genetic/environmental) underlying paternal age-disordered eating associations.

Method

Participants

Participants were 1722 female twins [488 (28%) MZ, 611 (35%) same-sex DZ and 623 (36%) opposite-sex DZ] between the ages of 8 and 17 years (mean \pm s.d. = 13.86 ± 2.39 years) from the ongoing Michigan Twins Project (MTP), a study conducted within the Michigan State University Twin Registry (MSUTR; Klump & Burt, 2006; Burt & Klump, 2013). Twins were recruited using birth records and driver's license databases (see Klump & Burt, 2006). MTP recruitment began in 2008, and response rates ($\sim 56\%$) are similar to those for other population-based twin registries (Kendler *et al.* 1992; Lichtenstein, 2002).

Because increases in genetic influences on disordered eating are observed during puberty in females (Klump *et al.* 2007; Culbert *et al.* 2009), we only included twins who were in mid-puberty or beyond (i.e. score ≥ 2.5 on the Pubertal Development Scale; Petersen *et al.* 1988). The twins spanned a range of racial/ethnic backgrounds (e.g. Caucasian, Black/African American, Asian/Pacific Rim) but, consistent with the region (see www.michigan.gov), most (87%) were Caucasian.

Measures

All data came from the MTP questionnaire completed by the twins' mother or father (93% biological mothers, 6% biological fathers, 1% step/adoptive parents).

Zygosity

Similar to other twin registries (Kendler *et al.* 1992; Lichtenstein, 2002), zygosity was determined using five physical similarity items. These items have demonstrated more than 95% accuracy when compared to genotyping (Lykken *et al.* 1990).

Parental age at birth

Paternal and maternal ages at offspring birth were calculated using birth dates. Average ages at birth (paternal mean \pm s.d. = 32.70 ± 5.92 years, range = 15–55; maternal mean \pm s.d. = 30.49 ± 4.77 years, range = 17–48) were similar to those from other studies examining parental age effects (e.g. paternal mean \pm s.d. = 31.5 ± 6.8 years, range = 13–70; maternal mean \pm s.d. = 28.8 ± 5.9 years, range = 12–53) (Croen *et al.* 2007).

Disordered eating symptoms

Disordered eating was assessed using nine items from the Eating Disorder Examination Questionnaire (EDE-Q; Fairburn & Beglin, 1994) (see Table 1). The EDE-Q assesses core eating disorder symptoms

Table 1. Disordered eating items and frequency of item endorsement

Disordered eating items	Item endorsement frequency (% of twins)		
	0 (Not true)	1 (Somewhat true)	2 (Certainly true)
1. Feels fat	70.7	21.4	7.9
2. Definite fear of gaining weight or of becoming fat	66.3	25.3	8.5
3. Has a strong desire to lose weight	77.1	17.1	5.9
4. Diets to control weight	86.2	11.2	2.6
5. Very dissatisfied with his/her body weight or shape	65.1	24.9	10.0
6. Body weight and/or shape influences how thinks about (judges) him/herself as a person	69.7	25.1	5.2
7. Feels guilty about what he/she eats because of the effect on body weight and/or shape	80.1	16.6	3.3
8. Eats lots of food and can't stop	89.5	8.7	1.9
9. Has vomited to control his/her weight	99.4	0.20	0.30

including weight/shape concerns, dietary restraint, binge eating and compensatory behaviors. EDE-Q items were included if they: (1) represented key attitudinal and behavioral symptoms; (2) exhibited significant correlations with the EDE-Q Global Score (mean $r=0.76$); and (3) were developmentally appropriate for pre- and early-adolescent participants. Importantly, these symptoms are well-established risk factors for eating disorders (Jacobi *et al.* 2004).

EDE-Q items were modified for use in this large-scale, mail-in twin registry. Parent, rather than child, reports were collected to keep the MTP questionnaire brief and easy to complete. This was necessary for recruiting as many families as possible into the registry (which serves as a participant bank; see Burt & Klump 2013). Although parent reports of disordered eating are less commonly used than parent reports of other psychiatric symptoms, data from another ongoing MSUTR study (Klump *et al.* 2010) show that the parent-child correlation for the MTP disordered eating items ($r=0.52$) is similar or better than parent-child correlations for externalizing/internalizing symptoms (r 's= 0.14 – 0.48) (Kolko & Kazdin, 1993; Youngstrom *et al.* 2000). EDE-Q parent reports also demonstrated expected correlations with external correlates in the Klump *et al.* (2010) sample (i.e. BMI, $r=0.58$; depressive symptoms, $r=0.35$).

Parents rated their children's disordered eating based on what they 'generally are like' using a three-point scale ('not true', 'sometimes true', 'certainly true') rather than the original EDE-Q format (i.e. number of days a symptom was present in the past month). This was necessary to match other MTP items and to index trait rather than state levels of disordered eating. Our data suggest that this rating format does not

substantially affect the validity of the EDE-Q. In a subsample of twins whose mothers rated disordered eating using the MTP EDE-Q and the original EDE-Q, ratings were highly correlated ($r=0.59$) despite occurring an average of 1.5 years apart (s.d.= 0.77 , range= 0.23 – 2.90 years).

Internal consistency for the MTP items was excellent ($\alpha=0.86$). There was significant variability in disordered eating, with the rate at which most symptoms were 'sometimes' or 'certainly' true ranging from 10–35% (see Table 1). Levels of disordered eating were similar to those from other MSUTR studies examining a similar age range but using the original EDE-Q (item endorsement= 7 – 42%) (Klump *et al.* 2010, 2013). As would be expected in a population-based, adolescent sample (Wade *et al.* 2008), attitudinal symptoms (e.g. fear of fatness) were the most highly endorsed, although behaviors (e.g. dieting, binge eating) also showed sufficient variation. Correlations with BMI were in the low-to-moderate range (range= 0.009 – 0.42 ; mean $r=0.25$; average percentage variance shared = 6%), suggesting that our measure taps disordered eating symptoms that are not merely a reflection of weight status/obesity.

Eating disorder diagnosis

Lifetime histories of eating disorders (AN, BN or BED) were assessed through parent report. Of the 1668 twins with available data, 11 (1%) had a history of an eating disorder [6/11 (55%) AN, 4/11 (36%) BED, 1/11 (9%) AN and BN], of whom seven (64%) received previous treatment and four (36%) did not. These percentages are similar to the prevalence of eating disorders in female children/adolescents (Merikangas *et al.* 2010;

Swanson *et al.* 2011). As expected, twins with an eating disorder history had substantially higher MTP disordered eating scores (mean \pm s.d.=7.81 \pm 5.98) than those with no history (mean \pm s.d.=2.31 \pm 3.16, $p=0.01$, Cohen's $d=1.15$). Disordered eating scores did not differ meaningfully between patients with a history of AN (or AN/BN) (mean \pm s.d.=8.14 \pm 7.40) versus BED (mean \pm s.d.=7.25 \pm 2.98).

Covariates

Several covariates were examined to ensure that associations between paternal age and eating pathology were not accounted for by confounding factors. Covariates included: twin age, ethnicity and BMI percentile (i.e. BMI adjusted for age); SES; parental fertility treatment and psychiatric history. Selected covariates have been associated with disordered eating and/or later paternal age at birth (Hare & Moran, 1979; Lilenfeld *et al.* 1998; O'Dea & Caputi, 2001; Croll *et al.* 2002; Doornbos *et al.* 2007; Sobotka, 2010; Eriksen *et al.* 2012). Parent reports of twin height and weight, which correlate highly with laboratory measurements (r 's=0.79–0.81; Huybrechts *et al.* 2011), were used to calculate BMI. BMI values were transformed to percentiles (see www.cdc.gov) to capture age-related variation in our child/adolescent sample. Family yearly income and parental education (i.e. highest level of education achieved by mother or father) were used to assess SES. A history of depression, anxiety disorder (i.e. panic, obsessive-compulsive, post-traumatic stress, separation anxiety) or eating disorder (i.e. AN, BN, BED) in the biological mother and/or father was assessed using a family history checklist coded 'yes' if at least one parent suffered from one or more disorder or 'no' if neither parent reported a history of the disorders.

Statistical analyses

Phenotypic associations

Paternal age at birth was examined as a predictor of disordered eating symptoms and history of an eating disorder. Paternal age was modeled continuously as well as categorically (i.e. <25, 25–29, 30–34, 35–39, ≥ 40 years) to increase statistical power and identify particularly high-risk age thresholds (e.g. paternal age >35 years; see Wohl & Gorwood, 2007).

Generalized linear mixed models (GLMMs) were used as they could account for the non-independence of twin data [i.e. by nesting a level 1 variable (individual twin) within a level 2 unit (family)] and could examine both continuous and categorical outcomes. GLMM linear regressions (normal distribution with identity link) and binary logistic regressions (binomial distribution with logit link) were used to examine

disordered eating and eating disorder history respectively. We standardized all variables prior to analysis to interpret unstandardized coefficients as standardized coefficients.

GLMMs were run twice: (1) controlling only for maternal age at birth and (2) controlling for all covariates. This two-step process allowed us to identify the effects of covariates on paternal age–eating pathology associations while always controlling for maternal age because maternal age correlated highly with paternal age ($r=0.72$)[†] and is associated with other psychiatric disorders (Croen *et al.* 2007; Menezes *et al.* 2010). Controlling for maternal age also ensured that our independent variable (i.e. paternal age at birth) and dependent variables (i.e. parent-reported eating pathology) were not confounded, as 93% of MTP questionnaires were completed by biological mothers.²

Genetic associations

We investigated genetic influences on paternal age effects by comparing MZ and DZ twin similarity for disordered eating as a function of paternal age at birth. Given that very few participants had an eating disorder history, analyses focused on disordered eating symptoms. Furthermore, although same-sex and opposite-sex twins were included in phenotypic analyses, twin analyses included same-sex twins only, as male co-twins of opposite-sex females were not examined because of low eating disorder prevalence in males (Swanson *et al.* 2011).

Twin intraclass correlations were calculated in the full sample and each paternal age category to provide an initial indication of genetic/environmental effects on disordered eating across paternal age. Additive genetic effects (A; genetic influences that add across genes) are indicated if the MZ twin correlation is approximately twice the DZ twin correlation whereas non-additive genetic effects (D; interaction of genetic effects at same locus) are suggested if MZ correlations are more than double DZ correlations. *De novo* mutations are considered a non-additive genetic process in twin studies, as DZ twins would share far less than half of their mutations because of their rarity (Liu *et al.* 2010). Therefore, if *de novo* mutations underlie paternal age–disordered eating relationships, we would expect larger MZ/DZ differences with advanced paternal age.

Twin studies also decompose variance into two forms of environmental effects. Shared environmental influences (C; factors that make co-twins similar to one another) are inferred if MZ and DZ twin

[†] The notes appear after the main text.

correlations are approximately equal whereas non-shared environmental influences (E; factors that make co-twins different from one another, including measurement error) are implied if the MZ correlation is less than 1.0. If differences between MZ/DZ correlations decrease with paternal age, shared environmental processes would be implicated, as co-twins would increase in similarity regardless of genetic sharing. Finally, decreases in the MZ correlation would indicate non-shared environmental influences on paternal age–disordered eating effects.

We then used twin moderation models (see Purcell, 2002) to statistically quantify the degree to which genetic/environmental influences on disordered eating vary by paternal age (the moderator). ACE and ADE models were examined to test for differences in both additive and non-additive genetic effects. Model fitting was conducted using full information maximum-likelihood raw data techniques with Mx statistical software (Neale, 1997).

Following previous recommendations (Purcell, 2002), we fit ‘full’ ACE and ADE moderator models that included linear and non-linear moderators to directly test whether genetic/environmental estimates vary linearly or non-linearly with paternal age. We compared full models to two more restrictive models: (1) linear moderation models that dropped nonlinear moderation coefficients and (2) no moderation models that estimated only genetic/environmental paths. To minimize the number of models fit, we did not drop individual moderation coefficients one by one. Instead, we determined whether each etiologic influence varied by paternal age by examining whether confidence intervals for moderator estimates overlapped with zero.

The best-fitting model was determined using both the difference in minus twice the log likelihood ($\Delta -2\ln L$) and Akaike’s Information Criteria (AIC). $\Delta -2\ln L$ was used to compare full and nested moderation models, and AIC was used to compare the un-nested ACE and ADE models. Statistically significant differences in $-2\ln L$ suggest that dropping moderator coefficients results in a significantly worse model fit whereas lower AIC values indicate a better model fit.

Although definition variables can be used to account for covariate effects in twin models (Neale *et al.* 2006), it was not possible to simultaneously include all eight covariates as definition variables in one model, as this would reduce the statistical power to detect differences in etiologic effects across paternal age (Agrawal *et al.* 2010). Consequently, we ran individual models that included each covariate as a definition variable one by one to examine individual covariate effects. Because findings were unchanged in each run of the model, we focused our results on models that included

maternal age at birth as the definition variable to mirror our first set of phenotypic analyses described earlier.

Results

Phenotypic associations

Consistent with hypotheses, paternal age was positively associated with disordered eating symptoms in female offspring, even after controlling for several important covariates (e.g. maternal age, BMI percentile; see Table 2). Similar to other disorders (Byrne *et al.* 2003), a threshold effect was detected, such that mean levels of disordered eating were significantly higher in offspring of fathers aged ≥ 40 years compared to offspring of younger fathers (see Table 2). Effect sizes indicate that the phenotypic effect of paternal age on disordered eating is small to medium in magnitude (β 's=0.08–0.09, d 's=0.18–0.41).³

Exploratory analyses revealed that paternal age at birth, measured continuously, also predicts a lifetime eating disorder history. Small sample sizes prevented examining paternal age as a categorical predictor of eating disorder diagnosis; however, there were a disproportionate number of cases in older age categories (see Table 2), and this effect was not driven by a specific diagnosis (i.e. AN and BED were both present in the older age groups).

Genetic associations

Similar to previous research in pubertal females (Klump *et al.* 2007; Culbert *et al.* 2009), higher MZ than DZ correlations and an MZ correlation less than 1.0 indicated the presence of genetic and non-shared environmental influences on disordered eating in the full sample. However, differences in MZ/DZ correlations across paternal age suggested significant heterogeneity in etiologic effects (see Table 3). Before age 35, the MZ correlation was twice the DZ correlation, suggesting additive genetic effects but little to no shared environmental or non-additive genetic influences. By contrast, after age 35, the DZ correlation increased and was comparable to most of the MZ correlations, indicating a complete lack of additive or non-additive genetic effects but significant shared environmental influences. The lower MZ correlation in the oldest paternal age group could reflect greater non-shared environmental effects, although conclusions are difficult to draw given the group’s small sample size ($n=16$ pairs) and the fact that this correlation does not differ significantly from MZ correlations in most younger paternal age groups (as evidenced by overlapping confidence intervals; see Table 3). Overall, twin correlations suggest increasing shared environmental influences on disordered

Table 2. Paternal age at birth as a predictor of disordered eating symptoms and lifetime eating disorder history in offspring

Paternal age (years)	Maternal age only model					Full covariate model			
	Coefficient (s.e.)	<i>t</i> (df)	<i>p</i>	<i>d</i>	Coefficient (s.e.)	<i>t</i> (df)	<i>p</i>	<i>d</i>	
Disordered eating symptoms									
	<i>n</i> (%)								
Continuous	1722	0.09 (0.04)	2.37 (1719)	0.02	–	0.08 (0.03)	2.44 (1709)	0.01	–
Categorical									
<25	128 (7.4)	–0.02 (0.10)	–2.08 (1716)	0.04	0.28	–0.06 (0.10)	–2.53 (1706)	0.01	0.30
25–29	439 (25.5)	–0.15 (0.05)	–4.14 (1716)	<0.001	0.41	–0.10 (0.06)	–3.98 (1706)	<0.001	0.31
30–34	589 (34.2)	0.01 (0.04)	–3.10 (1716)	0.002	0.27	0.04 (0.06)	–2.89 (1706)	0.004	0.18
35–39	383 (22.2)	0.04 (0.06)	–2.80 (1716)	0.005	0.32	0.05 (0.07)	–2.85 (1706)	0.004	0.18
≥40	183 (10.6)	0.29 (0.08)	–	–	–	0.28 (0.09)	–	–	–
Lifetime eating disorder (ED) history									
	ED	No ED							
	<i>n</i> (%)	<i>n</i> (%)							
Continuous	11	1657	0.64 (0.32)	2.01 (1666)	0.04	–	0.67 (0.31)	2.14 (1655)	0.03
Categorical									
<25	1 (9.1)	122 (7.4)	–	–	–	–	–	–	–
25–29	0 (0)	422 (25.5)	–	–	–	–	–	–	–
30–34	5 (45.5)	566 (34.2)	–	–	–	–	–	–	–
35–39	2 (18.2)	369 (22.3)	–	–	–	–	–	–	–
≥40	3 (27.3)	178 (10.7)	–	–	–	–	–	–	–

s.e., Standard error; df, degrees of freedom.

Significant *p* values are indicated in bold font.

The ‘maternal age only’ model controlled for maternal age at birth only. The ‘full covariate’ model controlled for maternal age at birth, child age, child ethnicity, child body mass index (BMI) percentile, family education and income, fertility treatment, and parental psychiatric history. Coefficient=standardized β in continuous paternal age models; standardized mean in categorical paternal age models. The ≥ 40 years group was coded as the reference group for *t* test comparisons in categorical paternal age models.

Table 3. Twin correlations for disordered eating by paternal age category

	MZ (95% CI) [<i>n</i>]	DZ (95% CI) [<i>n</i>]	<i>Z</i>	<i>p</i>
Full sample	0.69 (0.63 to 0.75) [227]	0.43 (0.35 to 0.50) [269]	4.28	<0.001
Paternal age (years)				
<25	0.73 (0.50 to 0.87) [26]	0.32 (–0.03 to 0.66) [17]	1.76	0.04
25–29	0.67 (0.53 to 0.79) [67]	0.36 (0.21 to 0.50) [77]	2.54	0.005
30–34	0.77 (0.67 to 0.84) [72]	0.33 (0.18 to 0.46) [88]	4.18	<0.001
35–39	0.65 (0.50 to 0.79) [46]	0.57 (0.41 to 0.70) [54]	0.62	0.24
≥40	0.30 (–0.01 to 0.56) [16]	0.64 (0.46 to 0.78) [33]	–1.35	0.99

MZ, Monozygotic; DZ, dizygotic same-sex twins; CI, confidence interval; *n*, number of twin pairs; *Z*, test of equality examining whether MZ correlation is higher than DZ twin correlations; *p*, one-tailed significance value for *Z* test of equality.

Maternal age at birth was regressed from each twin’s disordered eating score, and standardized residual scores were used to calculate twin correlations.

eating after age 35, arguing against increases in genetic or *de novo* mutation effects with advanced paternal age.

Because twin correlations suggested no differences in etiologic effects until age 35, we created two paternal age groups for analyses (<35 versus ≥ 35 years) to

maximize power and capture key etiologic differences across paternal age.⁴ Non-linear moderation cannot be examined with only two moderator levels, so we focused on linear and no moderation models. ACE models fit better than ADE models (i.e. lower AIC), which is not surprising given that non-additive genetic

effects were non-significant (i.e. 95% confidence intervals included 0).

Within the ACE models, a statistically significant $\Delta-2\ln L$ indicated that the linear moderation model fit better than the no moderation model and that genetic/environmental influences on disordered eating varied by paternal age. Following previous recommendations (Purcell, 2002), we report unstandardized path and moderator estimates in Table 4 and Fig. 1 so as to depict absolute changes in genetic/environmental influences rather than changes in proportions of total variance. Nonetheless, to compare genetic/environmental estimates to previous twin studies of disordered eating, we also report standardized estimates (calculated by squaring the path coefficients in the <35 age group, and squaring the sum of the moderator estimates plus the path coefficients in the ≥ 35 age group).

As shown in Table 4, a, c and e moderator estimates were statistically significant, suggesting that all three differ across paternal age. The most substantial change occurs for additive genetic and shared environmental influences (see Fig. 1). Before age 35, additive genetic effects contributed primarily to disordered eating ($A=75\%$), with the remaining variance due to the non-shared environment ($E=25\%$). By contrast, after age 35, the variance in disordered eating was entirely due to shared and non-shared environmental factors ($C=59\%$, $E=41\%$).⁵

Discussion

Using a large, population-based sample of female twins, we have demonstrated for the first time that advanced paternal age at birth is a significant risk factor for offspring eating pathology. This phenotypic effect was robust, in that it was present for eating disorder symptoms and diagnoses, and effects were independent of several important covariates (e.g. maternal age, BMI percentile, SES and parental psychiatric history). We also examined differences in etiologic influences across paternal age to investigate whether genetic factors (more specifically, *de novo* mutations) might underlie paternal age-disordered eating relationships. However, the results strongly suggested that genetic contributions to disordered eating decreased, rather than increased, with paternal age.

These findings contribute to a growing body of literature implicating advanced paternal age in offspring psychiatric risk, as indicated for schizophrenia, autism and bipolar disorder (Malaspina *et al.* 2001; Reichenberg *et al.* 2006; Frans *et al.* 2008). A relationship with disordered eating is novel and suggests that eating disorders may share some, perhaps more general, etiologic risk factors with these disorders. This is important given that sociocultural influences

(e.g. pressures for thinness) have led to views that eating disorders are etiologically distinct from 'neuropsychiatric' illnesses (Klump *et al.* 2009). Instead, eating disorders may be uniquely associated with both the internalizing and neuropsychiatric spectrums of psychopathology. This possibility is being explored by researchers investigating social/cognitive similarities between AN and autism (Zucker *et al.* 2007) and also other neuropsychiatric disorders (Steinglass *et al.* 2007).

De novo mutations have been proposed as a potential biological mechanism underlying paternal age-psychiatric risk associations, although our results argue against *de novo* mutation effects for paternal age-disordered eating relationships. We observed a sharp decrease in the magnitude of genetic influences on disordered eating in the paternal age ≥ 35 group, as DZ twins (who would not share *de novo* mutations) became more rather than less similar with paternal age. Our results converge with those of an autism twin study (Lundström *et al.* 2010), suggesting that *de novo* mutations may not account for paternal age effects. However, if only the frequency (rather than the location) of mutations confers risk, paternal age-related mutations could theoretically increase the similarity of DZ twins born to older fathers, as DZ twins may have a similar mutation burden but, unlike MZ twins, would not share specific mutations. We presumed that sharing specific mutations, rather than frequency alone, would influence twin similarity for eating pathology, consistent with the prediction that *de novo* mutations increase heritability estimates in twin studies (Liu *et al.* 2010; Ronald & Hoekstra, 2011). Nonetheless, we cannot definitively rule out the importance of *de novo* mutations. Future twin and molecular genetic studies are needed to further examine the mutation hypothesis for paternal age-psychiatric disorder relationships.

Another leading mechanism proposed to underlie paternal age-psychiatric disorder associations is disruptions in epigenetic processes (i.e. activation-inactivation of genes and/or changes to chromatin structure) that occur with advanced paternal age (Perrin *et al.* 2007). Although epigenetic mechanisms have been implicated in eating disorders and other psychiatric illnesses (e.g. Schanen, 2006; Frieling *et al.* 2010), epigenetic disruptions that accumulate in the male germline and are passed to twin offspring should increase MZ/DZ twin differences. Indeed, MZ twins, formed from one sperm, should share these epigenetic errors at a higher rate than DZ twins who develop from two different sperm. Thus, similar to studies showing greater epigenetic similarity for MZ *versus* DZ twins (Kaminsky *et al.* 2009), paternal age-related epigenetic errors would be expected to increase genetic influences on disordered eating. We instead observed

Table 4. Unstandardized parameter estimates and test statistics for the comparison of linear moderation and no moderation models

Model	Parameter estimates			Test statistics					
	A	C/D	E	-2lnL	df	$\Delta - 2\ln L$	Δdf	<i>p</i>	AIC
ACE									
Linear moderation ^a				2594.36	982	–	–	–	630.36
Path estimates	0.87 (0.75 to 0.95)	0 (–0.42 to 0.42)	0.50 (0.44 to 0.56)						
Linear moderators	–0.87 (–1.47 to –0.26)	0.77 (0.27 to 1.21)	0.14 (0.02 to 0.23)						
No moderation				2607.80	985	13.44	3	0.004	637.80
Path estimates	0.75 (0.60 to 0.88)	0.37 (–0.57 to 0.57)	0.54 (0.50 to 0.60)						
ADE									
Linear moderation				2606.36	982	–	–	–	642.36
Path estimates	0.79 (0.30 to 0.94)	–0.37 (–0.82 to 0.82)	0.50 (0.45 to 0.56)						
Linear moderators	0 (–0.24 to 0.49)	0.74 (–1.04 to 1.04)	0.09 (–0.01 to 0.20)						
No moderation				2609.70	985	3.34	3	0.34	639.76
Path estimates	0.84 (0.69 to 0.90)	0 (–0.47 to 0.47)	0.53 (0.49 to 0.58)						

A, Additive genetic; C, shared environmental; D, non-additive genetic; E, non-shared environmental; –2lnL, minus twice the log likelihood; df, degrees of freedom; AIC, Akaike's Information Criteria.

Linear moderation: genetic, shared environmental, and non-shared environmental parameter estimates vary linearly by paternal age at birth.

No moderation: genetic, shared environmental, and non-shared environmental parameter estimates do not vary by paternal age at birth.

^a Best-fitting model.

Significant paths and moderator estimates are indicated by confidence intervals that do not overlap with zero.

Significant *p* values are indicated in bold font.




Fig. 1. Unstandardized additive genetic (a), shared environmental (c) and non-shared environmental (e) variance contributions to disordered eating by paternal age at birth.

an increase in DZ twin similarity, and thus shared environmental effects with paternal age, arguing against epigenetic mechanisms. Of course, replication, particularly with molecular genetic designs, is needed.

Several covariates examined (e.g. maternal age, SES) could contribute to paternal age effects by decreasing genetic and increasing shared environmental influences on disordered eating. However, these factors did not account for paternal age–disordered eating associations and are unlikely to be strong mechanistic candidates. Additional unexamined factors that could account for paternal age effects and increased shared environmental influences include birth/obstetric complications and paternal weight concerns/behaviors. First, older paternal age has been associated with several birth/obstetric complications in offspring (e.g. pre-term birth, pre-eclampsia, cesarean section) (Harlap *et al.* 2002; Tang *et al.* 2006; Sartorius & Nieschlag, 2010; Shah, 2010), and birth/obstetric complications are linked to risk for eating disorders and other psychiatric illnesses (Cannon *et al.* 2002; Favaro *et al.* 2006). These paternal age and birth/obstetric complication associations are present after controlling for maternal age (Harlap *et al.* 2002; Tang *et al.* 2006), suggesting that they might account for the unique effects of paternal age on offspring psychiatric risk. However, maternal age has been shown to be a stronger predictor of birth/obstetric complications than paternal age (Harlap *et al.* 2002; Shah, 2010), and the lack of association between maternal age and disordered eating in our data makes it less likely that birth/obstetric complications could entirely account for our findings.

Second, longitudinal research suggests that weight concerns and dieting increase with age in men, such that these attitudes/behaviors are higher in older adulthood than earlier adulthood (Keel *et al.* 2007). In older

fathers, weight concerns and dieting may be highest when children are being raised and could be transmitted environmentally to offspring in the form of paternal expectations and/or criticism. Importantly, although older women continue to report significant weight/shape concerns and eating disorder behaviors in mid-life (Gagne *et al.* 2012), longitudinal data suggest that women experience a relative decrease in weight concerns/behaviors with advancing age (Keel *et al.* 2007). Sex differences in these longitudinal trajectories could potentially explain differential relationships between paternal and maternal ages at birth and offspring eating pathology. Clearly, these possibilities are speculative, particularly given that paternal weight concerns/behaviors may also have genetic (and/or gene–environment interaction) effects on offspring disordered eating. Further research is necessary to examine whether these types of mechanisms contribute environmentally to paternal age–eating pathology associations.

Despite notable strengths of our study (e.g. a large, population-based twin sample, examination of confounding factors), our study was not without limitations. Given that the maximum age of our participants was 17 years, we were unable to capture the entire period of risk for eating disorders, which can extend up until at least age 25 (Lewinsohn *et al.* 2000). However, disordered eating symptoms are present as early as childhood and increase across the pubertal period in females (Maloney *et al.* 1989; Killen *et al.* 1992). Therefore, meaningful variance in disordered eating was probably captured in our sample of twins, who are in mid-puberty and beyond. Nonetheless, future research should examine young adult samples to ensure results generalize to other periods of risk.

Eating disorder symptoms and diagnoses, along with covariates, were assessed through parent report. Parent reports allowed us to collect data on a large sample of twins, which was necessary for twin models investigating mechanisms underlying paternal age effects. However, parent reports may miss significant eating disorder symptoms/diagnoses that would be captured by self-reports, and limiting assessment of all covariates to a single parent report could introduce error due to shared method variance. Fortunately, our identified eating disorder cases performed as expected on measures of disordered eating and external correlates, and initial MSUTR data suggest that parent–child concordance for disordered eating is better than that observed for other phenotypes routinely assessed by parent report (see Method). Future studies should nevertheless confirm that advanced paternal age predicts eating pathology when examining self-reported symptoms/diagnoses using questionnaire and interview data. Using multiple methods to assess covariates

(e.g. laboratory height/weight measures) could also help to minimize potential influences of shared method variance.

Finally, given the small number of twins with eating disorder histories, phenotypic links between paternal age and eating disorder diagnoses could only be examined cursorily, and paternal age–diagnosis relationships were not investigated using twin models. Although the symptoms we examined are risk factors for eating disorders (Jacobi *et al.* 2004), additional studies with larger samples are needed to investigate phenotypic and etiologic relationships between paternal age and the broad eating disorder category, along with the specific diagnoses of AN, BN and BED. Such analyses could increase our understanding of the role of paternal age in eating disorder risk and identify specific mechanisms contributing to differential symptom presentations.

Notes

- ¹ We tested for multi-collinearity due to the high correlation between paternal and maternal age at birth, but tolerance and the variance inflation factor (VIF) were well within the acceptable range (tolerance=0.50, threshold <0.10; VIF=2.01, threshold >10).
- ² The results remained unchanged when excluding participants whose biological father completed the questionnaire (6% of the sample; data not shown).
- ³ Advanced maternal age at birth was not associated with offspring eating pathology. Maternal age exhibited a modest, negative association with disordered eating ($b=-0.06$, $p=0.03$) but this association became non-significant after controlling for covariates ($b=-0.03$, $p=0.15$). Furthermore, maternal age at birth did not significantly predict history of an eating disorder diagnosis ($b=0.21$, $p=0.48$).
- ⁴ We also tested the five paternal age categories to ensure that dichotomizing paternal age did not influence results. The results were identical to those presented in the text (data not shown), where genetic effects decreased and shared environmental effects increased with older paternal age.
- ⁵ We examined whether the same pattern of results (i.e. decrease in additive genetic effects and increase in shared environmental effects with paternal age) was observed when twins in the paternal ≥ 40 years age group were excluded from analyses, given the low MZ correlation in this age group. The results were identical, in that the ACE linear moderation model was best fitting, and both additive genetic and shared environmental moderation coefficients were significant and in the same direction as in the original models.

Acknowledgments

Data collection was supported by grants from Michigan State University (K. L. Klump and

S. A. Burt). Data analysis was supported by grants from the Canadian Institutes of Health Research (MDR-96630; S. E. Racine) and the National Institute of Mental Health (5 T32 MH082761 and 1 F31 MH0844701; K. M. Culbert; 1 R01 MH0820-54 and 1 R01 MH092377-01; K. L. Klump and S. A. Burt). The content is solely the responsibility of the authors and does not necessarily represent the official views of Michigan State University, the Canadian Institutes of Health Research or the National Institute of Mental Health.

Declaration of Interest

None.

References

- Agrawal A, Balasubramanian S, Smith EK, Madden PA, Bucholz KK, Heath AC, Lynskey MT (2010). Peer substance involvement modifies genetic influences on regular substance involvement in young women. *Addiction* **105**, 1844–1853.
- APA (2000). *Diagnostic and Statistical Manual of Mental Disorders, Fourth Edition, Text Revision*. American Psychiatric Association: Washington, DC.
- Awadalla P, Gauthier J, Myers RA, Casals F, Hamdan FF, Griffing AR, Côté M, Henrion E, Spiegelman D, Tarabeux J, Piton A, Yang Y, Boyko A, Bustamante C, Xiong L, Rapoport JL, Addington AM, DeLisi JLE, Krebs M, Joobor R, Millet B, Fombonne E, Motttron L, Zilversmit M, Keebler J, Daoud H, Marineau C, Roy-Gagnon M, Dube M, Eyre-Walker A, Drapeau P, Stone EA, Lafreniere RG, Rouleau GA (2010). Direct measure of the de novo mutation rate in autism and schizophrenia cohorts. *American Journal of Human Genetics* **87**, 316–324.
- Burt SA, Klump KL (2013). The Michigan State University Twin Registry: an update. *Twin Research and Human Genetics* **16**, 344–350.
- Byrne M, Agerbo E, Ewald H, Eaton WW, Mortensen PB (2003). Parental age and risk of schizophrenia: a case-control study. *Archives of General Psychiatry* **60**, 673–678.
- Cannon M, Jones PB, Murray RM (2002). Obstetric complications and schizophrenia: historical and meta-analytic review. *American Journal of Psychiatry* **159**, 1080–1092.
- Croen LA, Najjar DV, Fireman B, Grether JK (2007). Maternal and paternal age and risk of autism spectrum disorders. *Archives of Pediatrics and Adolescent Medicine* **161**, 334–340.
- Croll J, Neumark-Sztainer D, Story M, Ireland M (2002). Prevalence and risk and protective factors related to disordered eating behaviors among adolescents: relationship to gender and ethnicity. *Journal of Adolescent Health* **31**, 166–175.

- Crow JF (2000). The origins, patterns and implications of human spontaneous mutation. *Nature Reviews Genetics* **1**, 40–47.
- Culbert KM, Burt SA, McGue M, Iacono WG, Klump KL (2009). Puberty and the genetic diathesis of disordered eating attitudes and behaviors. *Journal of Abnormal Psychology* **118**, 788–796.
- Doornbos ME, Maas SM, McDonnell J, Vermeiden JPW, Hennekam RCM (2007). Infertility, assisted reproduction technologies and imprinting disturbances: a Dutch study. *Human Reproduction* **22**, 2476–2480.
- Eriksen W, Sundet JM, Tambs K (2012). Paternal age at birth and the risk of obesity in young adulthood: a register-based birth cohort study of Norwegian males. *American Journal of Human Biology* **25**, 29–34.
- Fairburn CG, Beglin SJ (1994). Assessment of eating disorders: interview or self-report questionnaire? *International Journal of Eating Disorders* **16**, 363–370.
- Favaro A, Tenconi E, Santonastaso P (2006). Perinatal factors and the risk of developing anorexia nervosa and bulimia nervosa. *Archives of General Psychiatry* **63**, 82–88.
- Frans EM, Sandin S, Reichenberg A, Lichtenstein P, Langstrom N, Hultman CM (2008). Advancing paternal age and bipolar disorder. *Archives of General Psychiatry* **65**, 1034–1040.
- Frieling H, Römer KD, Scholz S, Mittelbach F, Wilhelm J, De Zwaan M, Jacoby GE, Kornhuber J, Hillemecher T, Bleich S (2010). Epigenetic dysregulation of dopaminergic genes in eating disorders. *International Journal of Eating Disorders* **43**, 577–583.
- Gagne DA, Von Holle A, Brownley KA, Runfola CD, Hofmeier S, Branch KE, Bulik CM (2012). Eating disorder symptoms and weight and shape concerns in a large web-based convenience sample of women ages 50 and above: results of the gender and body image (GABI) study. *International Journal of Eating Disorders* **45**, 832–844.
- Hare E, Moran P (1979). Raised parental age in psychiatric patients: evidence for the constitutional hypothesis. *British Journal of Psychiatry* **134**, 169–177.
- Harlap S, Paltiel O, Deutsch L, Knaanie A, Masalha S, Tiram E, Caplan LS, Malaspina D, Friedlander Y (2002). Paternal age and preeclampsia. *Epidemiology* **13**, 660–667.
- Hultman C, Sandin S, Levine S, Lichtenstein P, Reichenberg A (2011). Advancing paternal age and risk of autism: new evidence from a population-based study and a meta-analysis of epidemiological studies. *Molecular Psychiatry* **16**, 1203–1212.
- Huybrechts I, Himes JH, Ottevaere C, De Vriendt T, De Keyzer W, Cox B, Van Trimont I, De Bacquer D, De Henauw S (2011). Validity of parent-reported weight and height of preschool children measured at home or estimated without home measurement: a validation study. *BMC Pediatrics* **11**, 63.
- Insel TR, Cuthbert BN, Garvey MA, Heinssen RK, Pine DS, Quinn KJ, Sanislow CA, Wang PS (2010). Research domain criteria (RDoC): toward a new classification framework for research on mental disorders. *American Journal of Psychiatry* **167**, 748–751.
- Iossifov I, Ronemus M, Levy D, Wang Z, Hakker I, Rosenbaum J, Yamrom B, Lee Y, Narzisi G, Leotta A, Kendall J, Grabowska E, Ma B, Marks S, Rodgers L, Stepansky A, Troge J, Andrews P, Bekritsky M, Pradhan K, Ghiban E, Kramer M, Parla J, Demeter R, Fulton LL, Fulton RS, Magrini VJ, Ye K, Darnell JC, Darnell RB, Mardis ER, Wilson RK, Schatz MC, McCombie WR, Wigler M (2012). De novo gene disruptions in children on the autistic spectrum. *Neuron* **74**, 285–299.
- Jacobi C, Hayward C, de Zwaan M, Kraemer HC, Agras WS (2004). Coming to terms with risk factors for eating disorders: application of risk terminology and suggestions for a general taxonomy. *Psychological Bulletin* **130**, 19–65.
- Kaminsky ZA, Tang T, Wang SC, Ptak C, Oh GHT, Wong AHC, Feldcamp LA, Virtanen C, Halfvarson J, Tysk C, McRae AF, Visscher PM, Montgomery GW, Gottesman II, Martin NG, Petronis A (2009). DNA methylation profiles in monozygotic and dizygotic twins. *Nature Genetics* **41**, 240–245.
- Keel PK, Baxter MG, Heatherton TF, Joiner TE (2007). A 20-year longitudinal study of body weight, dieting, and eating disorder symptoms. *Journal of Abnormal Psychology* **116**, 422–432.
- Kendler KS, Heath AC, Neale MC, Kessler RC, Eaves LJ (1992). A population-based twin study of alcoholism in women. *Journal of the American Medical Association* **268**, 1877–1882.
- Killen J, Hayward C, Litt I, Hammer L, Wilson D, Miner B, Taylor C, Varady A, Shisslak C (1992). Is puberty a risk factor for eating disorders? *Archives of Pediatrics and Adolescent Medicine* **146**, 323–325.
- Klump KL, Bulik CM, Kaye WH, Treasure J, Tyson E (2009). Academy for eating disorders position paper: eating disorders are serious mental illnesses. *International Journal of Eating Disorders* **42**, 97–103.
- Klump KL, Burt SA (2006). The Michigan State University Twin Registry (MSUTR): genetic, environmental and neurobiological influences on behavior across development. *Twin Research and Human Genetics* **9**, 971–977.
- Klump KL, Keel PK, Racine SE, Burt SA, Neale M, Sisk C, Boker S, Hu JY (2013). The interactive effects of estrogen and progesterone on changes in emotional eating across the menstrual cycle. *Journal of Abnormal Psychology* **122**, 131–137.
- Klump KL, Keel PK, Sisk C, Burt SA (2010). Preliminary evidence that estradiol moderates genetic influences on disordered eating attitudes and behaviors during puberty. *Psychological Medicine* **40**, 1745–1753.
- Klump KL, Perkins PS, Alexandra Burt S, McGue M, Iacono WG (2007). Puberty moderates genetic influences on disordered eating. *Psychological Medicine* **37**, 627–634.
- Kolko DJ, Kazdin AE (1993). Emotional/behavioral problems in clinic and nonclinic children: correspondence among child, parent and teacher reports. *Journal of Child Psychology and Psychiatry* **34**, 991–1006.

- Kong A, Frigge ML, Masson G, Besenbacher S, Sulem P, Magnusson G, Gudjonsson SA, Sigurdsson A, Jonasdottir A, Jonasdottir A, Wong WSW, Sigurdsson G, Walters GB, Steinberg S, Helgason H, Thorleifsson G, Gudbjartsson DF, Helgason A, Magnusson OT, Thorsteinsdottir U, Stefansson K (2012). Rate of de novo mutations and the importance of father's age to disease risk. *Nature* **488**, 471–475.
- Lewinsohn PM, Striegel-Moore RH, Seeley JR (2000). Epidemiology and natural course of eating disorders in young women from adolescence to young adulthood. *Journal of the American Academy of Child and Adolescent Psychiatry* **39**, 1284–1292.
- Lichtenstein P (2002). The Swedish Twin Registry: a unique resource for clinical, epidemiological and genetic studies. *Journal of Internal Medicine* **252**, 184–205.
- Lilienfeld LR, Kaye WH, Greeno CG, Merikangas KR, Plotnicov K, Pollice C, Rao R, Strober M, Bulik CM, Nagy L (1998). A controlled family study of anorexia nervosa and bulimia nervosa: psychiatric disorders in first-degree relatives and effects of proband comorbidity. *Archives of General Psychiatry* **55**, 603–610.
- Liu K, Zerubavel N, Bearman P (2010). Social demographic change and autism. *Demography* **47**, 327–343.
- Lundström S, Haworth C, Carlström E, Gillberg C, Mill J, Råstam M, Hultman CM, Ronald A, Anckarsäter H, Plomin R, Lichtenstein P, Reichenberg A (2010). Trajectories leading to autism spectrum disorders are affected by paternal age: findings from two nationally representative twin studies. *Journal of Child Psychology and Psychiatry* **51**, 850–856.
- Lykken D, Bouchard T, McGue M, Tellegen A (1990). The Minnesota Twin Family Registry. *Acta Geneticae Medicae et Gemellologiae* **39**, 35–70.
- Malaspina D, Corcoran C, Fahim C, Berman A, Harkave-Friedman J, Yale S, Goetz D, Goetz R, Harlap S, Gorman J (2002). Paternal age and sporadic schizophrenia: evidence for de novo mutations. *American Journal of Medical Genetics* **114**, 299–303.
- Malaspina D, Harlap S, Fennig S, Heiman D, Nahon D, Feldman D, Susser ES (2001). Advancing paternal age and the risk of schizophrenia. *Archives of General Psychiatry* **58**, 361–367.
- Maloney MJ, McGuire J, Daniels SR, Specker B (1989). Dieting behavior and eating attitudes in children. *Pediatrics* **84**, 482–489.
- Menezes P, Lewis G, Rasmussen F, Zammit S, Sipos A, Harrison G, Tynelius P, Gunnell D (2010). Paternal and maternal ages at conception and risk of bipolar affective disorder in their offspring. *Psychological Medicine* **40**, 477–485.
- Merikangas KR, He JP, Brody D, Fisher PW, Bourdon K, Koretz DS (2010). Prevalence and treatment of mental disorders among US children in the 2001–2004 NHANES. *Pediatrics* **125**, 75–81.
- Miller B, Messias E, Miettunen J, Alaräsänen A, Järvelin MR, Koponen H, Räsänen P, Isohanni M, Kirkpatrick B (2011). Meta-analysis of paternal age and schizophrenia risk in male versus female offspring. *Schizophrenia Bulletin* **37**, 1039–1047.
- Neale MC (1997). *Mx: Statistical Modeling*, 4th edn. Department of Psychiatry, Medical College of Virginia: Richmond, VA.
- Neale MC, Aggen SH, Maes HH, Kubarych TS, Schmitt JE (2006). Methodological issues in the assessment of substance use phenotypes. *Addictive Behaviors* **31**, 1010–1034.
- O'Dea JA, Caputi P (2001). Association between socioeconomic status, weight, age and gender, and the body image and weight control practices of 6- to 19-year-old children and adolescents. *Health Education Research* **16**, 521–532.
- Perrin MC, Brown AS, Malaspina D (2007). Aberrant epigenetic regulation could explain the relationship of paternal age to schizophrenia. *Schizophrenia Bulletin* **33**, 1270–1273.
- Petersen AC, Crockett L, Richards M, Boxer A (1988). A self-report measure of pubertal status: reliability, validity, and initial norms. *Journal of Youth and Adolescence* **17**, 117–133.
- Purcell S (2002). Variance components models for gene-environment interaction in twin analysis. *Twin Research* **5**, 554–571.
- Reichenberg A, Gross R, Weiser M, Bresnahan M, Silverman J, Harlap S, Rabinowitz J, Shulman C, Malaspina D, Lubin G, Knobler HY, Davidson M, Susser E (2006). Advancing paternal age and autism. *Archives of General Psychiatry* **63**, 1026–1032.
- Ronald A, Hoekstra RA (2011). Autism spectrum disorders and autistic traits: a decade of new twin studies. *American Journal of Medical Genetics. Part B, Neuropsychiatric Genetics* **156**, 255–274.
- Sartorius GA, Nieschlag E (2010). Paternal age and reproduction. *Human Reproduction Update* **16**, 65–79.
- Schanen NC (2006). Epigenetics of autism spectrum disorders. *Human Molecular Genetics* **15**, R138–R150.
- Shah PS (2010). Paternal factors and low birthweight, preterm, and small for gestational age births: a systematic review. *American Journal of Obstetrics and Gynecology* **202**, 103–123.
- Sobotka T (2010). Shifting parenthood to advanced reproductive ages: trends, causes and consequences. In *A Young Generation Under Pressure?* (ed. J. C. Tremmel), pp. 129–154. Springer: New York.
- Steinglass JE, Eisen JL, Attia E, Mayer L, Walsh BT (2007). Is anorexia nervosa a delusional disorder? An assessment of eating beliefs in anorexia nervosa. *Journal of Psychiatric Practice* **13**, 65–71.
- Swanson SA, Crow SJ, Le Grange D, Swendsen J, Merikangas KR (2011). Prevalence and correlates of eating disorders in adolescents: results from the national comorbidity survey replication adolescent supplement. *Archives of General Psychiatry* **68**, 714–723.
- Tang CH, Wu MP, Liu JT, Lin HC, Hsu CC (2006). Delayed parenthood and the risk of cesarean delivery – is paternal age an independent risk factor? *Birth* **33**, 18–26.

- Wade TD, Byrne S, Bryant-Waugh R** (2008). The Eating Disorder Examination: norms and construct validity with young and middle adolescent girls. *International Journal of Eating Disorders* **41**, 551–558.
- Wohl M, Gorwood P** (2007). Paternal ages below or above 35 years old are associated with a different risk of schizophrenia in the offspring. *European Psychiatry* **22**, 22–26.
- Xu B, Roos JL, Dexheimer P, Boone B, Plummer B, Levy S, Gogos JA, Karayiorgou M** (2011). Exome sequencing supports a de novo mutational paradigm for schizophrenia. *Nature Genetics* **43**, 864–868.
- Youngstrom E, Loeber R, Stouthamer-Loeber M** (2000). Patterns and correlates of agreement between parent, teacher, and male adolescent ratings of externalizing and internalizing problems. *Journal of Consulting and Clinical Psychology* **68**, 1038–1050.
- Zhao X, Leotta A, Kustanovich V, Lajonchere C, Geschwind DH, Law K, Law P, Qiu S, Lord C, Sebat J, Ye K, Wigler M** (2007). A unified genetic theory for sporadic and inherited autism. *Proceedings of the National Academy of Sciences USA* **104**, 12831–12836.
- Zucker NL, Losh M, Bulik CM, LaBar KS, Piven J, Pelphrey KA** (2007). Anorexia nervosa and autism spectrum disorders: guided investigation of social cognitive endophenotypes. *Psychological Bulletin* **133**, 976–1006.