

# Effects of different sources and levels of tannins on live performance and antioxidant response of Ossimi lambs

Tamer M. M. Hassan<sup>1</sup> , Omar A. Ahmed-Farid<sup>2</sup> and Fathy A. I. Abdel-Fattah<sup>3</sup>

## Animal Research Paper

**Cite this article:** Hassan TMM, Ahmed-Farid OA, Abdel-Fattah FAI (2020). Effects of different sources and levels of tannins on live performance and antioxidant response of Ossimi lambs. *The Journal of Agricultural Science* **158**, 339–348. <https://doi.org/10.1017/S0021859620000684>

Received: 26 February 2018

Revised: 3 July 2020

Accepted: 21 July 2020

First published online: 20 August 2020

### Key words:

Antioxidant response; lambs; mango leaves; pomegranate peels; tannins

### Author for correspondence:

Tamer M. M. Hassan,

E-mail: [tamer.mohamed@fagr.bu.edu.eg](mailto:tamer.mohamed@fagr.bu.edu.eg)

<sup>1</sup>Department of Animal Production, Faculty of Agriculture, Benha University, Qalyubia 13736, Egypt; <sup>2</sup>Physiology Department, National Organization for Drug Control and Research (NODCAR), Giza 12553, Egypt and <sup>3</sup>Department of Nutrition and Clinical Nutrition, Faculty of Veterinary Medicine, Benha University, Qalyubia 13736, Egypt

### Abstract

Pomegranate peels (PP) and mango leaves (ML) were analysed for nutrients and tannin contents. In an *in vitro* test, ten diets were prepared; six contained 2, 4 and 6% of PP or ML, three diets supplemented with mixed levels of PP and ML (1 + 1%, 2 + 2% and 3 + 3%) and control diet free of them. Gas was measured after 3, 6, 12, 24, 48 and 72 h of incubation. Methane and rumen parameters were estimated. In an *in vivo* experiment, 40 Ossimi lambs were divided into four groups; the first was control, other groups (T1, T2 and T3) fed diets containing 6% PP, 6% ML or mix levels (3% PP + 3%ML), respectively, for 2 months. Results showed that PP and ML were rich in tannins. In the *in vitro* test, a maximum reduction in gas, methane and NH<sub>3</sub>-N was in 6% PP, 6% ML and mixed levels (3% PP + 3% ML). In the *in vivo* experiment, there were no differences in growth and digestibility of DM and nutrients between treatment groups. Only a lowered DM intake and protein digestibility in lambs fed 6% PP. Gas and methane emission was decreased significantly in lambs fed 6% PP, compared to other groups. TVFAs and NH<sub>3</sub>-N were decreased for treatment groups. Also, all treatments did not show any pathological changes in liver function or on oxidative stress markers. In conclusion, PP and ML can be used in sheep diets at inclusion levels of 6% and mixture without detrimental effects on general health of Ossimi lambs.

### Introduction

Livestock, especially ruminants like sheep, contribute to the accumulation of methane emission in the atmosphere due to methanogenesis by archaeometanogen in the rumen. Such emission does not only affect the global warming, but it also represents energy loss from the animals, in which the lost can be between 80 and 140 g/kg from the total of digestible energy. Recently, nutritional strategies aimed to mitigate methane emission based on natural substances are preferred over the synthetic ones (Yogianto *et al.*, 2014). Production of volatile fatty acids in the rumen affects the process of methanogenesis depending on their metabolism and microbial activity. Fermentation of crude fibre in the rumen produces acetate and butyrate, results in increasing the formation of methane. On the other hand, fermentation of soluble carbohydrates produced propionate which is accompanied by a decrease in methane production (Bhatta *et al.*, 2014). Assuming 9 kg of ruminal CH<sub>4</sub> emission per sheep annually (Mbanzamihiho *et al.*, 2002) with a world population of more than 1172 million sheep in the year 2013 (FAOSTAT, 2015), sheep accounts for considerable CH<sub>4</sub> emission from enteric fermentation. Enteric CH<sub>4</sub> emission by ruminants also represents a loss of 50–150 g/kg of feed gross energy intakes (Singh *et al.*, 2005). Currently, there is increasing interest in the use of plants and plant by-products and organic farming to reduce enteric ruminal methane emissions and ammonia production. Secondary metabolites (tannins, antioxidants and flavonoids) in pomegranate (*Punica granatum L.*) peels (PP) and mango (*Mangifera indica*) leaves (ML) are produced by plants in their intermediary metabolism. Tannins are polyphenolic and important plant secondary metabolites in PP and ML and have an effect on ruminal methane production when included in the diet in significant proportions. Dietary tannins in livestock feeds have both beneficial and detrimental effects on animal performance and health. The valuable effects may comprise better performance and production. The adverse effects have been related to a reduction in palatability and digestibility of diet and growth inhibition of rumen microbes. The positive or negative effects depend on the source and concentration of tannins in the diet (Jayanegara *et al.*, 2012). Pomegranate components have attracted attention for their apparent immunomodulatory activity, antibacterial activity, antiatherosclerotic and antioxidative capacities (Sadq *et al.*, 2016). ML has various biomedical applications including antioxidative, anti-inflammation, anti-allergic, anti-cancer, hepatoprotective and immunomodulatory activities. It is a rich source of various biologically active compounds (Garrido *et al.*, 2004). To the best authors knowledge, a little reports cleared the effects of PP and ML on gas,

**Table 1.** Proximate analysis and total tannins in PP and ML (g/kg on dry matter basis)<sup>a</sup>

Items	PP	ML
Dry matter	800.1	925
Organic matter	942.1	967.4
Crude protein	33	49.6
Ether extract	58	31
Nitrogen-free extract	542.3	676.8
Crude fibre	224.7	200.2
NDF	189	168.3
ADF	118.5	105.5
Ash	142	42.4
ME (MJ/kg)	8.8 <sup>b</sup>	6.3 <sup>c</sup>
Total tannins (g/kg)	68.2	27.2

PP, pomegranate peel; ML, mango leaves; NDF, neutral detergent fibre; ADF, acid detergent fibre; ME, metabolizable energy.

<sup>a</sup>All analyses were carried out in triplicate.

<sup>b</sup>Calculated according to Mirzaei-Aghsaghalii *et al.* (2011).

<sup>c</sup>Calculated according to Cheema *et al.* (2014).

methane production and antioxidant response of Ossimi lambs was found. Therefore, the main aim of this study was to reduce gas production, methane emission and enhancing the immune response of Ossimi sheep by using tannins in PP, ML and its mixture through an *in vitro* and *in vivo* assay.

## Materials and methods

### Pomegranate peel and mango leaf preparation

Pomegranate peel was obtained manually by separation from fruits, while ML was collected from mango trees, rinsed with distilled water and cut into small pieces. Five hundred grams of fresh ML and PP were dried in air circulatory tray drier at 60°C for 6 h (Singh and Sethi, 2003) to obtain approximately 900 g/kg dry matter (DM), then ground to pass a 1 mm sieve for further chemical analysis as shown in Table 1.

### In vitro experiment

A test diet (substrate) was shown in Table 2. The commercial concentrate mixture was composed of cotton seed cake 350 g/kg, wheat bran 330 g/kg, yellow maize 220 g/kg, rice bran 40 g/kg, molasses 30 g/kg and calcium carbonate 30 g/kg. A total of ten treatments were prepared with different replacement levels of PP and ML. The test diets 1, 2 and 3 were prepared by replacing 20, 40 and 60 g/kg of PP with 20, 40 and 60 g/kg of wheat straw based on weight. Diets 4, 5 and 6 contained ML (20, 40 and 60 g/kg). An equal combination of PP and ML (10 + 10, 20 + 20 and 30 + 30 g/kg) was mixed to constitute test diets 7, 8 and 9, in addition to a control diet with zero PP or ML levels.

Rumen content was collected individually from ten mature cannulated Ossimi lambs before feeding in the morning. The animals were fed twice a day on the aforementioned concentrate mixture with alfalfa hay and wheat straw. After filtration using four layers of cheese-cloth, the pH-value of rumen fluid was measured immediately using a pH meter (Genway, Model 3520, USA). Five millilitres of rumen fluid was added to 1 ml of protein

precipitant, and then centrifuged at 3000 rpm (EBA 21, Hettich, Germany) for 15 min. Short-chain fatty acids (SCFAs) were determined according to Hoeltershinken *et al.* (1997). Concentration of ammonia nitrogen (NH<sub>3</sub>-N) was determined according to AOAC (1995). For rumen bacteria and protozoa counts, the filtered rumen liquor was stained with methyl-green formalin saline solution as described by Ogimoto and Imai (1981), then stoked in a dark place until examination. After gentle mixing of fixed samples, one drop was poured on haemocytometer slide, covered with a cover slip and examined under a light microscope. Samples were used after 72 h of incubation.

The *in vitro* gas test was performed according to Menke and Steingass (1988) to determine the optimum level of PP and ML that minimize CH<sub>4</sub> production. Fifty millilitres of filtered rumen fluid was flushed with CO<sub>2</sub>, then added (1:2, v/v) to buffered mineral solution and incubated in a shaker water bath at 39°C with a continuous flow of CO<sub>2</sub>. All laboratory handling of rumen fluid was carried out under a continuous flow of CO<sub>2</sub>. Approximately 200 mg of test diets (substrate) was placed into calibrated glass syringes of 100 mL capacity. Thirty syringes were divided into ten groups; each group composed of five replicate, three syringes each. Thirty millilitres of rumen liquor–buffer solution was injected into the warmed up syringes and incubated at 39°C. The volume of gas production was recorded from the calibrated scale on the syringe. Readings of total gas output were taken after 3, 6, 12, 24, 48 and 72 h of incubation according to Menke and Steingass (1988) recommendation.

CH<sub>4</sub> emission was determined at the end of incubation to get the full of methane production in samples by gas chromatography (GC-Agilent 6890) with flame ionization detector according to the method of Soliva and Hess (2007).

### In vivo experiment

#### Animals and experimental design

Forty Ossimi lambs at 6 months of age with an average weight of 29.25 ± 1.39 kg were obtained from experimental farm of the Faculty of Agriculture, Benha University, Egypt. Animals were divided into four groups (ten lambs each) and housed individually in clean and hygienic pens (0.97 m × 2.82 m). Lambs were subjected to the routine vaccination against infectious diseases and deworming programmes just before the onset of the experiment. The four experimental groups were fed diets formulated to meet the requirements of finishing lambs (Table 3) according to NRC of sheep (1985). The control diet was free of PP or ML; other test diets were supplemented with 60 g/kg of PP (T1), 60 g/kg ML (T2), while the diet T3 contained a mixture of 30 g/kg PP plus 30 g/kg ML. These replacement levels were selected according to the results of the *in vitro* test. The experimental animals were fed twice daily in equal portions at 09:00 and 17:00 h with an additional allowance of 100 g/kg more than the requirements for the period of 2 months. Feed residues were daily collected and weighed to determine the dry matter intake (DMI). Fresh and clean water was supplied at all times. Body weight changes of individual lambs were recorded at weekly intervals in the morning before feeding and feed conversion ratio (FCR) was calculated for each animal at the end of the experimental period.

#### Gas production and rumen metabolites

The rumen liquor was collected from three lambs in each group after 4 h of morning feed at the monthly interval by oral-gastric tube. The pH-value of rumen fluid, SCFAs, NH<sub>3</sub>-N and counts

**Table 2.** Ingredients and chemical composition of *in vitro* experiment diets

Items	PP			ML			MIX			
	Control	Diet 1	Diet 2	Diet 3	Diet 4	Diet 5	Diet 6	Diet 7	Diet 8	Diet 9
PP	0	20	40	60	0	0	0	10	20	30
ML	0	0	0	0	20	40	60	10	20	30
Concentrate mixture <sup>a</sup>	550	550	550	550	550	550	550	550	550	550
Alfalfa hay	262	262	262	262	262	262	262	262	262	262
Wheat straw	175	155	135	115	155	135	115	155	135	115
Common salt	5	5	5	5	5	5	5	5	5	5
Vitamin and minerals premix <sup>b</sup>	3	3	3	3	3	3	3	3	3	3
Ammonium chloride	5	5	5	5	5	5	5	5	5	5
Total	1000	1000	1000	1000	1000	1000	1000	1000	1000	1000
Calculated analysis (g/kg on dry matter basis) <sup>c</sup>										
ME (MJ/kg)	10.4	10.5	10.6	10.7	10.2	10.2	10.2	10.2	10.3	10.4
Crude protein	145	145	145	145	145	145	145	144	145	145
Crude fibre	215	202	202	203	200	202	202	201	203	203
NDF	414	389	383	379	381	389	394	384	389	387
ADF	259	231	234	235	239	240	243	231	236	240
Calcium	8.9	8.7	8.5	8.8	9.8	10.0	10.2	8.5	8.8	9
Phosphorus	5.8	5.8	6	6.2	5.7	5.8	5.9	5.6	5.8	6
Total tannins (g/kg diet)	--	1.3	2.7	4	0.5	1	1.63	0.9	1.9	2.8

PP, pomegranate peel; ML, mango leaves; ME, metabolizable energy; CP, crude protein; CF, crude fibre; NDF, neutral detergent fibre; ADF, acid detergent fibre.

<sup>a</sup>Concentrate mixture composition: cotton seed cake 350 g/kg, wheat bran 330 g/kg, yellow maize 220 g/kg, rice bran 40 g/kg, molasses 30 g/kg and calcium carbonate 30 g/kg.

<sup>b</sup>Purchased from Misr Feed Additives for animal nutrition, Egypt. Each 3 kg contain: Vitamin A = 12 000 000 IU, D<sub>3</sub> = 2 500 000 IU, E = 15 000 mg, Zinc = 60 000 mg, Manganese = 70 000 mg, Iron = 60 000 mg, Copper = 30 000 mg, Iodine = 5000 mg, Selenium = 300 mg, Cobalt = 1000 mg, Cobalt = 1000 mg.

<sup>c</sup>Calculated according to feed composition table, NRC for sheep, (1985).

of rumen microbes were measured. Total gas production was estimated after 3, 6, 12, 24, 48 and 72 h incubation periods according to Menke and Steingass (1988) recommendation, as previously described in the *in vitro* experiment.

### Digestibility trial

After a preliminary period of 3 weeks, a digestibility trial was carried out for each group to determine the digestion of diet DM and nutrients. A stainless steel wire mesh was placed on the floor of pens to separate the faecal matter from urine, which was passed into the drainage. Feed consumption was recorded daily and faecal matter was collected in morning for successive 7 days and weighed. Faecal samples (100 g/kg) were taken daily and dried out at 60°C for 48 h, ground and stored at -20°C till analysis. The nutrient digestible component (NDC) of diet was calculated for DM, organic matter (OM), crude protein (CP), neutral detergent fibre (NDF), acid detergent fibre (ADF), ether extract (EE)

and nitrogen-free extract (NFE) from each dietary treatment using the equation proposed by McDonald *et al.* (2002).

$$\text{NDC} = \frac{\text{total nutrient consumed} - \text{total nutrient excreted}}{\text{total nutrient consumed}} \times 100$$

### Blood metabolites

Blood samples from lambs were withdrawn in plain tubes from the jugular vein at two intervals; after 1 and 2 months of the experiment. To obtain serum, blood samples were centrifuged at 3000 rpm (Sigma 3-18K) for 15 min. Serum alanine aminotransferase (ALT) and aspartate aminotransferase (AST) were estimated according to Reitman and Frankels (1957). Total protein (TP) and albumin were measured according to Bradford (1976) and Gustafsson (1976), respectively. Globulin is calculated by

**Table 3.** Ingredients and chemical composition of *in vivo* experiment diets

Items	Control	T1	T2	T3
PP	0	60	0	30
ML	0	0	60	30
Concentrate mixture <sup>a</sup>	550	550	550	550
Alfalfa hay	262	262	257	257
Wheat straw	175	115	120	120
Common salt	5	5	5	5
Vitamin and minerals premix <sup>b</sup>	3	3	3	3
Ammonium chloride	5	5	5	5
Total	1000	1000	1000	1000
Calculated analysis (g/kg on dry matter basis) <sup>c</sup>				
ME (MJ/kg)	10.4	10.7	10.2	10.4
Crude protein	145	145	145	145
Crude fibre	215	203	202	203
NDF	414	379	394	387
ADF	259	235	243	240
Calcium	8.9	8.8	10.2	9
Phosphorus	5.8	6.2	5.9	6
Total tannins (g/kg diet)	--	4	1.6	2.8

T1, T2 and T3: diets containing 6% PP, 6% ML or mix levels (3% PP + 3%ML), respectively; PP, pomegranate peel; ML, mango leaves; ME, metabolizable energy; CP, crude protein; CF, crude fibre; NDF, neutral detergent fibre; ADF, acid detergent fibre.

<sup>a</sup>Concentrate mixture composition: cotton seed cake 350 g/kg, wheat bran 330 g/kg yellow maize 220 g/kg, rice bran 40 g/kg, molasses 30 g/kg and calcium carbonate 30 g/kg.

<sup>b</sup>Purchased from Misr Feed Additives for animal nutrition, Egypt. Each 3 kg contain: Vitamin A = 12 000 000 IU, D<sub>3</sub> = 2 500 000 IU, E = 15 000 mg, Zinc = 60 000 mg, Manganese = 70 000 mg, Iron = 60 000 mg, Copper = 30 000 mg, Iodine = 5000 mg, Selenium = 300 mg, Cobalt = 1000 mg, Cobalt = 1000 mg.

<sup>c</sup>Calculated according to feed composition table, NRC for sheep, (1985).

subtracting the measured albumin from the measured TP. Serum malondialdehyde (MDA) level was measured according to the method of Karatepe (2004). Serum ATP and serotonin (5HT) was assayed by high-performance liquid chromatography (HPLC) (Agilent series 1200, USA) according to the method of Teerlink *et al.* (1993) and Arafa *et al.* (2010), respectively. Serum reduced glutathione (GSH), oxidized glutathione (GSSG) and nitric oxide (NO, as nitrite and nitrate) were measured by HPLC according to Jayatilleke and Shaw (1993). Serum superoxide dismutase (SOD) activity was assayed at 420 nm on a UV-Vis Shimadzu spectrophotometer (2450) measured by Marklund and Marklund (1974). Catalase activity (CAT) was measured by a spectrophotometric method based on the decomposition of H<sub>2</sub>O<sub>2</sub> (Aebi, 1984).

### Chemical analyses

Samples of dried PP, ML, experimental diets and faecal matter were analysed according to AOAC (1995). DM was measured using a hot air circulation oven (Heraeus Ut20, Germany) at 105°C for 4 h (method no. 930.15), ash was measured using Burnout furnace Ney Vulcan D-550, USA (method no. 942.05). CP was determined using Kjeltac<sup>®</sup> system 2100, FOSS-Sweden

(method no. 984.13), and EE with Soxtec<sup>®</sup> system 2045, FOSS-Sweden (method no. 920.39). NDF and ADF were estimated using Fibretherm<sup>®</sup> FT12 according to the method of Van Soest *et al.* (1991). OM was calculated from the difference between DM and ash. For determination of total tannins, about 120 mL of ethanol (800 g/kg) was added to 50 g of dried PP or ML powder and heated for 5 min, then cooled, washed and filtered using filter paper. To the filtrate, 100 mL of ethanol was further added and the mixture was heated for 10 min followed by filtration. The liquid filtrate was partially evaporated at a temperature of 50°C for 30 min. Total tannins were determined by HPLC method (Agilent 1200 series, quaternary pump and UV detector) according to Durgawale *et al.* (2016).

### Statistical analysis

Data were analysed using the Statistical Analysis System (SAS, 2004) Computer Program, using the General Linear Model (GLM) procedure for analysing the data with equal subclass numbers. The statistical model was applied as follows:

*In vitro* experiment (Model 1)

$$Y_{ij} = \mu + T_i + L_j + (TL)_{ij} + e_{ij}$$

Where:  $Y_{ij}$  is the observation of measured traits;  $\mu$  is the overall mean;  $T_i$  is the fixed effect of  $i^{\text{th}}$  treatment in the diet ( $i = \text{PP, ML and Mix}$ );  $L_j$  is the fixed effect of  $j^{\text{th}}$  levels of treatment in the diet ( $j = 20, 40, 60 \text{ g/kg}$ );  $(TL)_{ij}$  is the fixed effect of interaction between treatment and levels;  $e_{ij}$  is a random error associated with the individual observation.

*In vivo* experiment (Model 2)

$$Y_{ij} = \mu + T_i + e_{ij}$$

Where:  $Y_{ij}$  is the observation of measured traits;  $\mu$  is the overall mean;  $T_i$  is the fixed effect of  $i^{\text{th}}$  treatment group ( $i = \text{control, PP, ML and Mix}$ ); and  $e_{ij}$  is a random error associated with the individual observation.

Blood metabolites (Model 3)

$$Y_{ij} = \mu + T_i + S_j + (TS)_{ij} + e_{ij}$$

Where:  $Y_{ij}$  is the observation of measured traits;  $\mu$  is the overall mean;  $T_i$  is the fixed effect of  $i^{\text{th}}$  treatment in the diet ( $i = \text{PP, ML and Mix}$ );  $S_j$  is the fixed effect of  $j^{\text{th}}$  sample month ( $j = \text{first month and second month}$ );  $(TS)_{ij}$  is the fixed effect of interaction between treatment and sample month;  $e_{ij}$  is a random error associated with the individual observation.

The significant differences between means of treatments were measured according to Duncan (1955).

## Results

### Tannins in PP and ML and their proximate composition

Data of proximate analysis and tannins content of PP and ML are summarized in Table 1. PP in general contained less CP and NFE than ML, while the contents of EE, CF, NDF, ADF and ash were higher than that of ML. Overall, total tannins concentration in PP and ML was high, PP contained more tannins than ML.

**Table 4.** Total gas, methane production after incubation periods and fermentation metabolites of the *in vitro* experiment

Items	Treatment (T)					Levels % (L)				P value		
	Control	PP	ML	MIX	SEM*	2	4	6	SEM*	T	L	T × L
Total gas production (ml/200 mg substrate) after incubation (hours)												
3	9.9	8.7	8.9	8.4	0.22	9.1	8.5	8.4	0.22	0.370	0.110	0.290
6	23.7	18.2	21.8	21.3	0.50	20.8	20.6	19.8	0.50	0.001	0.360	0.100
12	69	38	54	55	2.0	52	48	47	2.0	0.001	0.260	0.010
24	84	68	70	79	1.2	76	71	71	1.2	0.001	0.010	0.020
48	86.9	76.5	78.3	82.9	0.94	80.1	79.4	78.2	0.94	<0.001	0.370	0.200
72	98	86	93	92	1.6	92	91	88	1.6	0.003	0.230	0.580
Methane production	21.0	18.4	20.8	20.1	0.17	20.7	19.7	19.3	0.17	0.001	0.001	0.330
Fermentation metabolites after 72 h of incubation												
SCFAs (mmol/L) <sup>a</sup>	84	80	78	81	1.6	80	79	81	1.6	0.550	0.660	0.690
NH <sub>3</sub> -N (mg/dL)	24.6	18.3	19.2	20.0	0.40	22.1	19.5	19.0	0.40	<0.001	0.003	0.310
Bacteria count (10 × 6)/mL	9.3	8.6	7.8	7.4	0.33	7.6	8.2	7.9	0.33	0.070	0.480	0.020
Protozoa count (10 × 3)/mL	6.0	5.0	4.8	4.9	0.20	5.0	4.5	5.2	0.20	0.660	0.060	0.380
pH-value	6.4	6.5	6.5	6.4	0.11	6.3	6.7	6.4	0.11	0.920	0.060	0.370

SEM, standard error of mean; PP, pomegranate peel; ML, mango leaf; SEM, standard error of mean; SCFAs, short-chain fatty acids; NH<sub>3</sub>-N, ammoniacal nitrogen.

All data are the results of a measurement performed in triplicate.

<sup>a</sup>Short-chain fatty acids.

### The *in vitro* experiment

The mean values of total gas and methane production were presented in Table 4. The lowest CH<sub>4</sub> values ( $P < 0.05$ ) were for test diets contained 60 g/kg of PP, 60 g/kg of ML and the mix diet of 30 g/kg PP plus 30 g/kg ML. With these levels, there was insignificant ( $P > 0.05$ ) reduction in SCFAs, NH<sub>3</sub>-N, rumen bacteria and protozoa counts were detected. pH values were not affected by PP or ML dietary replacement.

### *In vivo* experiment

Results in Table 5 show the effect of replacement of PP, ML or their mixture on DMI, body weight gain (BWG), FCR and apparent digestibility of DM, OM and nutrients. Lambs fed on a diet containing 60 g/kg PP (T1) had significantly lower DMI ( $P < 0.05$ ) than that of the control group while the total BWG was comparable to other groups. FCR mean values of animals fed 60 g/kg PP were better than that of all test groups. There were no significant differences between means of the NDC of diet DM and nutrients between experimental groups except for CP. There was significant ( $P < 0.05$ ) reduction in CP digestibility in lambs fed 60 g/kg PP comparatively with other groups.

Total gas production with the results of rumen parameters is demonstrated in Table 6. Result values of total gas production were significantly lower ( $P < 0.05$ ) in lambs fed ration contained 60 g/kg PP than that of the control group. Rumen parameters (pH value, SCFAs, bacteria and protozoa count) were not affected by the replacement of PP or ML in the diet. NH<sub>3</sub>-N values were reduced ( $P < 0.05$ ) in animals fed PP or ML when compared with the control group.

### Blood metabolites

Data in Table 7 present the effect of PP, ML and their combinations on lamb's liver function (ALT, AST) and protein production (TP, albumin and globulin) during 2 months experimental period. Inclusion of PP, ML or their combination in the diet for lambs did not show any pathological changes in liver function and protein production that is markedly by normal level of serum ALT, AST, TP, albumin and globulin. Replacement of PP, ML or their combination showed significant ( $P < 0.05$ ) decrease in MDA, NO and GSSG values and increase GSH, endogenous antioxidant enzymes (CAT, SOD), cell energy (ATP) and neurotransmitter stimulate growth factor (5-HT) in comparison with the control group. Blood sample month (first or second month) had no significant effect on blood metabolites means except for GSH, CAT and ATP level was higher in the second month than the first month of the feeding period. Interaction between treatment and sample month effects was not significant.

### Discussion

The obtained chemical composition values of PP were in line with previous studies (Mirzaei-Aghsaghali *et al.*, 2011; Shabtay *et al.*, 2008). For ML, the nutritive contents were in the range of another study (Cheema *et al.*, 2014). Differences in the chemical composition of PP and ML may be attributed to the different geographical distribution of plant species, climate, maturity and growing conditions (Cheema *et al.*, 2014). Despite the high fibre and low protein contents in PP and ML, they can provide a good source of nutrients for feeding ruminants. Total tannins in PP and ML were found to be 68.25 and 27.21 mg/g DM, respectively. Tannins were in this study more abundant in PP than ML (27.21

**Table 5.** Live performance and nutrient digestible component of experimental lambs the *in vivo* experiment

Item	Control	T1	T2	T3	SEM*	P value
Total DMI, kg	87	80	85	84	1.8	0.040
Daily DMI, kg/day	1.5	1.3	1.4	1.4	0.03	0.040
Total intake of tannins g/kg DMI	0	327	139	241	8.0	0.055
Daily intake of tannins, g/day	0	5.3	2.2	4.0	0.03	0.041
Initial weight, kg	30	29	29	29	1.4	0.980
Final weight, kg	42	42	41	41	1.4	0.920
Total BWG, kg	11.7	12.4	11.1	11.9	0.45	0.290
Average daily BWG, g/day	198	206	185	199	7.0	0.220
FCR	7.7	6.6	7.8	7.1	0.40	0.150
Nutrient digestible component (g/kg)						
Dry matter	674.0	679.0	654.0	659.0	0.64	0.070
Organic matter	679	696	676	671	1.1	0.430
Crude protein	600.0	549.0	584.0	586.0	0.98	0.030
NDF	710.0	719.0	696.0	705.0	0.82	0.320
ADF	670	675	653	663	1.6	0.790
Ether extract	616.0	626.0	601.0	605.0	0.80	0.160
NFE	715.0	724.0	705.0	713.0	0.73	0.400

T1, T2 and T3: diets containing 6% PP, 6% ML or mix levels (3% PP + 3%ML), respectively; SEM, standard error of mean; DMI, dry matter intake; BWG, body weight gain; FCR, feed conversion ratio; NDF, neutral detergent fibre; ADF, acid detergent fibre; NFE, nitrogen-free extract.

**Table 6.** Total gas production and rumen metabolites of lambs fed the experimental diet *in vivo* experiment

Item	Control	T1	T2	T3	SEM*	P value
Total gas production (ml) after incubation (hours)						
3	9.4	8.4	8.9	8.7	0.20	0.010
6	22.5	18.2	21.8	21.3	0.48	<0.001
12	54	38	55	54	2.1	<0.001
24	79	68	71	70	1.6	<0.001
48	82.9	76.5	79.4	78.3	0.97	<0.003
72	95	86	94	92	1.6	<0.007
Rumen metabolites at the end of <i>in vivo</i> experiment						
SCFAs (mmol/L) <sup>a</sup>	81	80	78	81	1.5	0.590
NH <sub>3</sub> -N (mg/dL)	23.4	19.6	20.9	20.4	0.50	0.020
Bacteria count (10 <sup>6</sup> )/mL	8.6	7.4	7.8	7.5	0.35	0.100
Protozoa count (10 <sup>3</sup> )/mL	5.1	4.7	5	4.9	0.22	0.700
pH	6.3	6.5	6.5	6.4	0.11	0.780

T1, T2 and T3: diets containing 6% PP, 6% ML or mix levels (3% PP + 3%ML), respectively; SEM, standard error of mean; SCFAs, short-chain fatty acids; NH<sub>3</sub>-N, ammoniacal nitrogen. <sup>a</sup>Short-chain fatty acids.

mg/g DM) that was near the level (21.3 g/kg DM) determined by Cheema *et al.* (2014).

In this study, the *in vitro* test for methane measurement was conducted to select the optimum level which induces maximum reduction of total gas and CH<sub>4</sub> concentration without any detrimental effect on rumen health. With the gradual increase of PP or ML levels, there was a gradual suppression of total gas and

CH<sub>4</sub> production. From three levels of PP, ML or a combination of them, it was found that the maximum dietary incorporation of 60 g/kg PP or ML and the combined level (30 g/kg PP + 30 g/kg ML) reduced total gas and CH<sub>4</sub> emission effectively in comparison with the control (zero PP or ML) or other test levels (20, 40 and 60 g/kg of PP and ML). Results in this study were in the line of previous *in vitro* study (Bhatta *et al.*, 2014) who

**Table 7.** Effects of PP, ML and MIX on lambs liver function, oxidative stress markers, endogenous antioxidant enzymes, cell energy and neurotransmitters in serum

Items	Treatment (T)				SEM*	Sample month (S)			P value		
	Control	PP	ML	MIX		First	Second	SEM*	T	S	T × S
ALT, U/L	22	17	25	20	1.6	20	20	1.1	0.001	0.795	0.999
AST, U/L	31	28	33	26	2.1	27	28	1.5	<0.001	0.408	0.998
Total protein, g/dL	7.1	7.3	7.1	7.2	0.06	7.3	7.2	0.04	0.468	0.574	0.233
Albumin, g/dL	4.1	4.1	4.0	4.1	0.04	4.1	4.2	0.03	0.459	0.610	0.710
Globulin, g/dL	3.0	3.1	3.1	3	0.08	3.1	3.1	0.05	0.574	0.712	0.625
MDA, nmol/mL	1.8	1.2	1.5	1.3	0.09	1.3	1.5	0.06	<0.001	0.210	0.832
GSH, $\mu$ mol/mL	47	65	50	53	3.2	51	61	2.3	0.018	0.002	0.462
GSSG, $\mu$ mol/mL	1.1	0.6	1.0	0.9	0.06	0.9	0.9	0.04	<0.001	0.330	0.930
NO, $\mu$ mol/mL	33	23	30	28	2.0	24	29	1.5	<0.001	0.138	0.905
SOD, U/mL	72	90	76	79	5.2	81	83	3.7	0.012	0.490	0.996
CAT, $\mu$ moles/min/mL	40	52	44	47	2.4	43	52	1.7	0.004	0.017	0.997
ATP, $\mu$ mol/mL	18.5	25.8	20.7	23.7	0.83	21.7	25.2	0.59	0.001	0.029	0.991
5HT, $\mu$ g/mL	0.6	0.9	0.8	0.7	0.04	0.8	0.9	0.03	0.003	0.062	0.995

All data are the results of a measurement performed in triplicate.

PP, pomegranate peel; ML, mango leaf; SEM, standard error of mean; ALT, alanine transaminase; AST, aspartate transaminase; MDA, malondialdehyde; GSH, reduced glutathione; GSSG, oxidized glutathione; NO, nitric oxide; SOD, superoxide dismutase; CAT, catalase; ATP, adenosine triphosphate; 5HT, 5 hydroxytryptamine (serotonin).

indicated that total gas and CH<sub>4</sub> concentration decreases as the level of tannins and period of incubation increases. Tannins can reduce CH<sub>4</sub> emission through a reduction in fibre digestion which decreases H<sub>2</sub> production, inhibition of the growth of methanogens (Tavendale *et al.*, 2005). The slight decrease in SCFAs concentrations in this study may be attributed to the slight depression in fermentative activity from fibre degradation which could be a sequel of depressed methanogenesis (Jayanegara *et al.*, 2012). It was observed that the addition of condensed tannin decreased SCFAs levels in an *in vitro* fermentation study of Bhatta *et al.* (2013). In all levels of PP or ML, a decrease in NH<sub>3</sub>-N was observed. This reduction in NH<sub>3</sub>-N may be attributed to binding of tannins with proteins; forming undegradable tannin–protein complexes by proteases (Makkar, 2003).

With increasing PP or ML levels, the count of bacteria and protozoa was slightly reduced. In this trend, Manasri *et al.* (2012) have proven that plants containing tannins have antimicrobial properties. This inhibitory effect could be attributed to the affinity of tannins to form complexes with the cell wall and membrane of bacteria causing morphological changes of the cell wall and the extracellular enzymes secreted (Jones *et al.*, 1994). Ruminal pH values varied from 6.4 to 6.9, which were within the optimum ranges (6–7) for maintaining a normal cellulolytic organism. The effects of tannins on ruminal pH were variably reported among studies; either no effect (Dschaak *et al.*, 2011) or increase (Jami *et al.*, 2012).

Regarding lambs performance, the total DMI of animals fed 60 g/kg PP (T1) was the lowest followed by T3, T2 and the control group, respectively. Similar results obtained by Archimede *et al.* (2016) who indicated that DMI was reduced with tannin-rich plants in the diet. This finding agrees with a previous study (Ngwa *et al.*, 2003) who observed that tannins usually decreased palatability and preference of plants by cattle, sheep and goats. The reduction in palatability was referred to: firstly, a reaction between tannins and salivary mucoproteins, or through a direct

negative impact on the taste receptors, annoying an astringent sensation (Frutos *et al.*, 2004). Secondly, impaired digestion of DM in the rumen prolongs the emptying of the digestive tract, providing signals to the nerve centres. The third mechanism is based on the identification of negative post-prandial actions mediated by microbial fermentation following the ingestion of tannins (Waghorn, 1996). There are some exceptions to tannin effect on DMI, as in some cases, there was an increase (Puchala *et al.*, 2005) or insignificant reduction in voluntary feed intake in sheep (McSweeney *et al.*, 1988). Variations in DMI due to the presence of tannins in the diet could be elucidated through its concentration in the diet; low tannin content exhibits a positive effect on DMI (Aerts *et al.*, 1999), while higher levels in the diet reduced dramatically DMI (Frutos *et al.*, 2004).

Although the final BWG of all animals in this study showed no differences between treatments, there was insignificant ( $P > 0.05$ ) improvement of FCR for T1 (60 g/kg PP) and T3 (30 g/kg PP + 30 g/kg ML). These beneficial FCR values were due to lower DMI and efficient utilization of nutrients for groups T1 and T3 compared to other treatments. In this line, the final body weight did not have any change when lambs fed PP at levels (20, 40 and 60 g/kg PP). DM and nutrients retention in all treatments showed no change relative to the control group. Only CP digestibility was reduced ( $P < 0.05$ ) for lambs fed 60 g/kg PP (T1) comparatively with other treatment groups. This finding was attributed to the formation of tannin–protein complex which reduces plant protein degradation to NH<sub>3</sub> in the rumen. Low pH-value in abomasum promotes disassociation of tannin–protein complexes and enhances the proteins flow for digestion in the small intestine (Frutos *et al.*, 2004). However, the reduction in apparent digestibility of CP in this study indicates that the tannin–protein complexes did not undergo complete disassociation in the abomasum, thereby lowering the digestion of CP in the total tract. Similar results observed by Beauchemin *et al.* (2007) who concluded that tannin extract had no effect on DM, energy or fibre (ADF

and NDF) digestibility, but apparent digestibility of CP decreased linearly. Also Archimede *et al.* (2016) revealed that using tannin-rich plants in sheep diet did not affect apparent OM digestibility. In the study of McNabb *et al.* (1996), up to 45 g/kg condensed tannins in the diet improved production efficiency in ruminants by binding and protection of dietary proteins from microbial degradation in the rumen and consequently increasing the flow of essential amino acids and non-ammonia nitrogen for intestinal absorption. Therefore, improvement of FCR with less DMI in this study could be attributed to the increment of intestinal supply of metabolizable proteins (Waghorn, 1996), carbohydrates (Makkar *et al.*, 1995) and fats (Patra and Saxena, 2011) for maximum absorption. With 60 g/kg PP, ML or their mix (30 g/kg PP + 30 g/kg ML), there was no adverse effect either on DM or nutrients utilization by lambs. This favourable effect might be the result of added valuable nutrients in PP and ML as a feed ingredient in the diet. In this line, findings of Jami *et al.* (2012) proved that tannins in PP had no large negative effects on digestibility of nutrients which was reflected on improved performance of dairy cows. Shabay *et al.* (2008) found that the tannin-rich PP intake up to 200 g/kg of the total DM intake does not possess deleterious or positive effects on fattening ration intake of feedlot calves.

Pomegranate peel or ML replacement in the diet reduced CH<sub>4</sub> and total gas production linearly in comparison with the control group similarly as described in the *in vitro* experiment. Due to the binding of tannins to dietary protein, and also to a reduction in the activity of a large proportion of microflora in the rumen as well as binding to carbohydrates, a reduction in total gas emission is resulted (Sadq *et al.*, 2016). PP replacement in the diet of sheep at 60 g/kg induced maximum reduction of CH<sub>4</sub> emissions (166.2 g/kg) compared with the control. Similar results were obtained by Hess *et al.* (2006) in sheep fed diet with added tannins where CH<sub>4</sub> emissions were reduced by 130 g/kg without major losses in feeding value of the diet. Animut *et al.* (2008) reported that the goat's diet containing tannins decreased CH<sub>4</sub> emissions effectively. Most feasible conditions responsible for reduced CH<sub>4</sub> production are changes in the number and/or activity of bacteria other than ones highly fibrolytic, including methanogens (Field *et al.*, 1989) and protozoa (Schönhusen *et al.*, 2003). Tannins reduce CH<sub>4</sub> emission through several ways; (1) reduction in fibre digestion which decreases H<sub>2</sub> production by prevention or at least interfering with the attachment of rumen microorganisms to plant cell walls for further degradation (McAllister *et al.*, 1994). (2) Direct inhibitory effect of methanogens and declined digestibility of proteins and carbohydrates (Jayanegara *et al.*, 2012), and (3) reduction of protozoa population where part of the methanogens are associated symbiotically on the surface or inside the micro-fauna (Vogels *et al.*, 1980). Also TVFAs level was reduced ( $P < 0.05$ ) by PP or ML supply in the diet, while no change in SCFAs concentration between experimental groups. This reduction in TVFAs might be due to lowered fibre degradation in the rumen by adhering of tannins to plant cell wall hindering microbial attack. According to Bhatta *et al.* (2014), tannins in PP or ML exhibited effectively CH<sub>4</sub> reduction but did not cause a real inhibition of the TVFAs concentration that would indicate that CH<sub>4</sub> suppression was primarily due to antimethanogenic activity rather than lowered fibre digestibility. In this trend, no significant reduction in SCFAs concentration in this study was observed. The authors indicated that feeding tannin sometimes results in a decrease in ammonia concentration, enhancing the efficient use of TVFAs for microbial protein synthesis. In contrast, the results of other studies indicated no difference in TVFAs level

between groups (Min *et al.*, 1998; Puchala *et al.*, 2005). As a consequence of reduced CP digestibility, NH<sub>3</sub>-N was reduced ( $P < 0.05$ ) in all treatment groups compared with control. Min *et al.* (1998) noted that ewes fed forage containing condensed tannin had lower ruminal ammonia than ewes grazing tannin-free forage. The undegradable tannin-protein complex by microflora decreases the level of ruminal NH<sub>3</sub>-N (Makkar, 2003). Tannins can bind to proteins and carbohydrates, leading also to a reduction in ruminal gas production and methane emission (Makkar *et al.*, 1995). Due to a combination of these activities, tannins can be associated with improvements in animal growth and productivity and consequently minimization of effects on the environment (Silanikova *et al.*, 1993).

Dietary supply of PP or ML did not affect the counts of bacteria or protozoa as well as ruminal pH-value in this study. This result is supported by the meta-analysis study (Jayanegara *et al.*, 2012) in which tannins did not significantly reduce log protozoa population in both *in vitro* and *in vivo* experiments. Pilajun and Wanapat (2013) also reported that diet supplementation with tannins-rich plants did not affect the microbial population in swamp buffalo. In other studies, plants containing tannins and saponins had antimicrobial properties (Manasri *et al.*, 2012). Variations in results of this work with those obtained by other studies could be referred to the different sources of diets, dose of tannins, animal type, feeding programme and environmental conditions (Anantasook *et al.*, 2013).

Data did not show any pathological response with PP, ML and their combined replacement compared with the control group. ML was found to contain various biomedical substances of antioxidative, anti-inflammation, anti-allergic, anti-cancer, hepatoprotective and immunomodulatory activities (Garrido *et al.*, 2004). Also the presence of photochemical constituents especially flavonoids and tannins in PP set animal at normal homeostasis markedly by normal liver function (ALT, AST, TP, albumin and globulin) and normal cell structure markedly by normal level of MDA, NO, GSSG and GSH. These results are in agreement with those obtained by Abdel-Moneim *et al.* (2011) who found that PP and ML did not show any pathological alteration in liver and kidney functions. Recent studies showed the beneficial effects of ML on the liver through mitigating serum enzymes and the decreasing liver ALT and AST (Ojekale *et al.*, 2007).

Data of SOD and CAT revealed improvement in the endogenous defence mechanism in animal fed diet supplemented with PP or ML owing to their antioxidant activities. Similarly, studies of Parmar and Kar (2008) demonstrated that PP decreased lipid peroxidation in cardiac, hepatic and renal tissues. ML extract contains many polyphenols and mangiferin which is one of the biologically active cytoprotective compounds prevents mitochondrial stress and iron-induced oxidative damage in the liver (Kawpoomhae *et al.*, 2010). In this study, a considerable increase in the ATP level in lambs fed diets T1, T2 and T3. This indicates that tannins are able to protect the mitochondria from the damaging effects by scavenging the reactive oxygen species. These findings were similar to those obtained by Gulcin *et al.* (2010) who reported that tannic acid is the effective natural antioxidant component for food preservation.

Serotonin (5-HT) is a tryptamine, monoamine neurotransmitter synthesized in serotonergic neurons and in the gastrointestinal tract. According to the fundamental activity of 5-HT, its influence in the secretion of pituitary growth hormone is by inhibiting the production of hypothalamic somatostatin (Lesch, 2001). Phytochemical constituent of PP and ML showed an increased



level of 5-HT that may be due to the synergetic effect of flavonoid mixture in the plant. GSH, CAT and ATP level was higher in the second month of feeding period than the first month that may be due to increasing of metabolism activity.

## Conclusion

Replacement of 60 g/kg PP, 60 g/kg ML or their mix (30 g/kg PP + 30 g/kg ML) in sheep diet reduced total gas emission. Also, it enhanced the growth performance, digestibility and rumen functions, as well as it decreased the oxidative stress markers, and improvement in the antioxidant defence mechanisms of fattening Ossimi lambs.

**Acknowledgements.** The authors are grateful to Professor Fawzy Abo-Donia, Professor of Animal Nutrition, and Dr Mahmoud El-Attrouy for their assistance in carrying out the experiment and the statistical analysis. Thanks are also due to the central laboratory of Animal Production Department and Agricultural Research Centre, Sheep Farm, Faculty of Agriculture at Moshtohor, Benha University, Egypt.

**Financial support.** This work was supported by the central laboratory of Animal Production Department and Agricultural Research Centre, Sheep Farm, Faculty of Agriculture at Moshtohor, Benha University, Egypt.

**Conflict of interest.** The authors declare no conflict of interests.

**Ethical standards.** Humane animal care and handling procedures were conducted in accordance with the Animal Care Committee of the Department of Animal Production, Faculty of Agriculture at Moshtohor, Benha University and with the instructions from the Ministry of Agriculture in Egypt.

## References

- Abdel-Moneim AE, Dkhil MA and Al-Quraishy S (2011) Studies on the effect of pomegranate (*Punica granatum*) juice and peel on liver and kidney in adult male rats. *Journal of Medicinal Plants Research* **5**, 5083–5088.
- Aebi H (1984) Catalase *in vitro*. *Methods in Enzymology* **105**, 121–126.
- Aerts RJ, Barry TN and McNabb WC (1999) Polyphenols and agriculture. Beneficial effects of proanthocyanidins in forages. *Agriculture Ecosystems and Environment* **75**, 1–12.
- Anantasook N, Wanapat M, Cherdthong A and Gunun P (2013) Effect of plants containing secondary compounds with palm oil on feed intake, digestibility, microbial protein synthesis and microbial population in dairy cows. *Asian Australian Journal of Animal Science* **26**, 820–826.
- Animut G, Puchala R, Goetsch AL, Patra AK, Sahlu T, Varel VH and Wells J (2008) Methane emission by goats consuming diets with different levels of condensed tannins from lespedeza. *Animal Feed Science and Technology* **144**, 212–227.
- Arafa NMS, Salem SM and Ahmed-Farid OA (2010) Influence of Echinacea extract pre- or postnatal supplementation on immune and oxidative status of growing rabbits. *Italian Journal of Animal Science* **9**, 338–343.
- Archimede H, Rira M, Barde DJ, Labirin F, Marie-Magdeleine C, Calif B, Periacarpin F, Fleury J, Rochette Y, Morgavi DP and Doreau M (2016) Potential of tannin-rich plants, *Leucaena leucocephala*, *Glyricidia sepium* and *Manihot esculenta*, to reduce enteric methane emissions in sheep. *Journal of Animal Physiology and Animal Nutrition* **100**, 1149–1158.
- Association of Official Analytical Chemists (AOAC) (1995) *Analysis of the Association of Official Analytical Chemists*, 16th Edn edn. Arlington, VA, USA.
- Beauchemin KA, McGinn SM, Martinez TF and McAllister TA (2007) Use of condensed tannin extract from quebracho trees to reduce methane emissions from cattle. *Journal of Animal Science* **85**, 1990–1996.
- Bhatta R, Baruah L, Saravanan M, Suresh KP and Sampath KT (2013) Effect of medicinal and aromatic plants on rumen fermentation, protozoa population and methanogenesis *in vitro*. *Journal of Animal Physiology and Animal Nutrition* **97**, 446–456.
- Bhatta R, Saravanan M, Baruah L and Prasad CS (2014) Effects of graded levels of tannin-containing tropical tree leaves on *in vitro* rumen fermentation, total protozoa and methane production. *Journal of Applied Microbiology* **118**, 557–564.
- Bradford MM (1976) A rapid and sensitive method for the quantitation of microgram quantities of protein utilizing the principle of proteindye binding. *Analytical Biochemistry* **72**, 248–254.
- Cheema UB, Sultan JI, Javaid A, Mustafa MI and Younas M (2014) Screening of fodder tree leaves by chemical composition, mineral profile, anti-nutritional factors and *in sacco* digestion kinetics. *Scholarly Journal of Agricultural Science* **4**, 558–564.
- Dschaak CM, Williams CM, Holt MS, Eun JS, Young AJ and Min BR (2011) Effects of supplementing condensed tannin extract on intake, digestion, ruminal fermentation, and milk production of lactating dairy cows. *Journal of Dairy Science* **94**, 2508–2519.
- Duncan DB (1955) Multiple range and multiple F tests. *Biometrics* **11**, 1–42.
- Durgawale TP, Durgawale PP and Khanwelkar CC (2016) Quantitative estimation of tannins by HPLC. *Der Pharmacia Lettre* **8**, 123–126.
- Field JA, Kortekaas S and Lettinga G (1989) The tannin theory of methanogenic toxicity. *Biological Wastes* **29**, 241–262.
- Food and Agriculture Organization (FAO), *Statistical pocketbook world food and agriculture* (2015) <http://www.fao.org/3/a-i4691e.pdf>.
- Frutos P, Raso M, Hervás G, Mantecón ÁR, Pérez V and Giráldez F (2004) Is there any detrimental effect when a chestnut hydrolysable tannin extract is included in the diet of finishing lambs? *Animal Research* **53**, 127–136.
- Garrido G, Gonzalez D, Lemus Y, Garcia D, Lodeiro L, Quintero G, Delporte C, Nunez-Selles AJ and Delgado R (2004) *In vivo* and *in vitro* anti-inflammatory activity of *Mangifera indica* L. extract (VIMANG). *Pharmacological Research* **50**, 143–149.
- Gulcin I, Huyut Z, Elmastas M and Aboul-Enein HY (2010) Radical scavenging and antioxidant activity of tannic acid. *Arabian Journal of Chemistry* **3**, 43–53.
- Gustafsson J (1976) Improved specificity of serum albumin determination and estimation of 'acute phase reactants' by use of the bromocresol green reaction. *Clinical Chemistry* **22**, 616–622.
- Hess HD, Tiemann TT, Noto F, Carulla JE and Kreuzer M (2006) Strategic use of tannins as means to limit methane emission from ruminant livestock. *International Congress Series* **1293**, 164–167.
- Hoeltershinken M, Pitt U, Tammen FC, Hoffman P and Scholz H (1997) Influence of mouldy grass on fermentation and thiamine metabolism in bovine rumen fluid (*In vitro*). *Duetsche Tierarztliche Wochenschrift* **104**, 17–22.
- Jami SEA, Nikbatchat M, Yosef E, Miron J and Mizrahi I (2012) Effects of adding a concentrated pomegranate-residue extract to the ration of lactating cows on *in vivo* digestibility and profile of rumen bacterial population. *Journal of Dairy Science* **95**, 1–10.
- Jayanegara A, Leiber F and Kreuzer M (2012) Meta-analysis of the relationship between dietary tannin level and methane formation in ruminants from *in vivo* and *in vitro* experiments. *Journal of Animal Physiology and Animal Nutrition* **96**, 365–375.
- Jayatilleke E and Shaw S (1993) A high-performance liquid chromatographic assay for reduced and oxidized glutathione in biological samples. *Analytical Biochemistry* **214**, 452–457.
- Jones GA, McAllister TA, Cheng KJ and Muir AD (1994) Effect of sainfoin (*Onobrychis viciifolia* Scop) on growth and proteolysis by 4 strains of rumen bacteria. Resistance of *Prevotella* (Bacteroides) *ruminicola* B14. *Applied Environmental Microbiology* **60**, 1374–1378.
- Karatepe M (2004) Simultaneous determination of ascorbic acid and free malondialdehyde in human serum by HPLC-UV. *Chromatographic Line* **12**, 362–365.
- Kawpoomhae K, Sukma M, Ngawhirunpat T, Opanasopit P and Sripattanaporn A (2010) Antioxidant and neuroprotective effects of standardized extracts of *Mangifera indica* leaf. *Thai Journal of Pharmaceutical Science* **34**, 32–43.
- Lesch KP (2001) Variation of serotonergic gene expression. Neurodevelopment and the complexity of response to psychopharmacologic drugs. *European Neuropsychopharmacology* **11**, 457–474.
- Makkar HPS (2003) Effects and fate of tannins in ruminant animals, adaptation to tannins, and strategies to overcome detrimental effects of feeding tannin-rich feeds. *Small Ruminant Research* **49**, 241–256.

- Makkar HPS, Blümmel M and Becker K** (1995) In vitro effects and interactions between tannins, saponins and fate of tannins in the rumen. *Journal of the Science of Food and Agriculture* **69**, 481–493.
- Manasri N, Wanapat M and Navanukraw C** (2012) Improving rumen fermentation and feed digestibility in cattle by mangosteen peel and garlic pellet supplementation. *Livestock Science* **148**, 291–295.
- Marklund S and Marklund G** (1974) Involvement of the superoxide anion radical in the autoxidation of pyrogallol and a convenient assay for superoxide dismutase. *European Journal of Biochemistry* **47**, 469–474.
- Mbanzamihigo L, Fievez V, da Costa Gomez C, Piattoni F, Carlier L and Demeyer D** (2002) Methane emission from the rumen of sheep fed a mixed grass-clover pasture at two fertilisation rates in early and late season. *Canadian Journal of Animal Science* **82**, 69–77.
- McAllister TA, Bae HD, Jones GA and Cheng KJ** (1994) Microbial attachment and feed digestion in the rumen. *Journal of Animal Science* **72**, 3004–3018.
- McDonald P, Edward RA, Greenhalgh JFD and Morgan GA** (2002) *Animal Nutrition*, 6th Edn. Edinburgh, Great Britain: Pearson Educational Limited.
- McNabb WC, Waghorn GC, Peters JS and Barry TN** (1996) The effect of condensed tannins in *Lotus pedunculatus* on the solubilisation and degradation of ribulose-1,5-bisphosphate carboxylase (EC 4.1.1.39, Rubisco) protein in the rumen and the sites of rubisco digestion. *British Journal of Nutrition* **76**, 535–549.
- McSweeney CS, Kennedy PM and John A** (1988) Effect of ingestion of hydrolysable tannins in *Terminalia oblongata* on digestion in sheep fed *Stylosanthes hamata*. *Australian Journal of Agricultural Research* **39**, 235–244.
- Menke KH and Steingass H** (1988) Estimation of the energetic feed value obtained from chemical analysis and in vitro gas production using rumen fluid. *Animal Research and Development* **28**, 7–55.
- Min BR, Barry TN, McNabb WC and Kemp PD** (1998) Effect of condensed tannins on the production of wool and on its processing characteristics in sheep grazing *Lotus corniculatus*. *Australian Journal of Agricultural Research* **49**, 597–605.
- Mirzaei-Aghsaghali A, Maheri-Sis N, Mansouri H, Razeghi ME, Mirza-Aghazadeh A, Cheraghi H and Aghajanzadeh-Golshani A** (2011) Evaluating potential nutritive value of pomegranate processing by-products for ruminants using in vitro gas production technique. *ARPJ Journal of Agricultural and Biological Science* **6**, 45–51.
- Ngwa AT, Nsahlai IV and Bonsi ML** (2003) Feed intake and dietary preferences of sheep and goats offered hay and legume-tree pods in South Africa. *Agroforestry Systems* **57**, 29–37.
- N.R.C** (1985) Nutrient requirements of sheep, sixth revised edition. Subcommittee on sheep nutrition, committee of animal nutrition, board on agriculture, National Research Council.
- Ogimoto KH and Imai S** (1981) *Atlas of Rumen Microbiology*. Tokyo: Japan Scientific Societies Press.
- Ojekale AB, Ojiako OA, Saibu GM, Lala A and Oludude OA** (2007) Long term effects of aqueous stem bark extract of *Cissus populnea* (Guill. and Per.) on some biochemical parameters in normal rabbits. *African Journal of Biotechnology* **6**, 247–251.
- Parmar HS and Kar A** (2008) Medicinal values of fruit peels from *Citrus sinensis*, *Punica granatum*, and *Musa paradisiaca* with respect to alterations in tissue lipid peroxidation and serum concentration of glucose, insulin, and thyroid hormones. *Journal of Medicinal Food* **11**, 376–381.
- Patra AK and Saxena J** (2011) Exploitation of dietary tannins to improve rumen metabolism and ruminant nutrition. *Journal of the Science of Food and Agriculture* **91**, 24–37.
- Pilajun R and Wanapat M** (2013) Microbial population in the rumen of swamp buffalo (*Bubalus bubalis*) as influenced by coconut oil and mangosteen peel supplementation. *Journal of Animal Physiology and Animal Nutrition* **97**, 439–445.
- Puchala R, Min BR, Goetsch AL and Ahlu T** (2005) The effect of a condensed tannin-containing forage on methane emission by goats. *Journal of Animal Science* **83**, 182–186.
- Puchala R, Min BR, Goetsch AL and Ahlu T** (2005) The effect of a condensed tannin-containing forage on methane emission by goats. *Journal of Animal Science* **83**, 182–186.
- Reitman S and Frankel S** (1957) Colorimetric determination of AST and ALT activity. *American Journal of Clinical Pathology* **28**, 56–63.
- Sadq SM, Ramzi DO, Hamasalim HJ and Ahmed KA** (2016) Growth performance and digestibility in Karadi lambs receiving different levels of pomegranate peels. *Open Journal of Animal Sciences* **6**, 16–23.
- S.A.S** (2004) SAS Procedure Guide 'version 9 Ed'. SAS Institute Inc., Cary, NS. USA.
- Schönhusen U, Zitnan R, Kuhla S, Jentsch W, Derno M and Voigt J** (2003) Effects of protozoa on methane production in rumen and hindgut of calves around time of weaning. *Arch Tierernähr* **57**, 279–295.
- Shabtay A, Eitam H, Tadmor Y, Orlov A, Meir A, Weinberg P, Weinberg ZG, Chen Y, Brosh A, Izhaki I and Kerem ZJ** (2008) Nutritive and anti-oxidative potential of fresh and stored pomegranate industrial byproduct as a novel beef cattle feed. *Journal of Agricultural and Food Chemistry* **56**, 10063–10070.
- Silanikove N, Tagari H and Shkolnik A** (1993) Comparison of rate passage, fermentation rate and efficiency of digestion of high fiber diet in desert black Bedouin goats as compared to Swiss Saanen goats. *Small Ruminant Research* **12**, 45–60.
- Singh D and Sethi V** (2003) Screening of pomegranate genotypes for the preparation of quality grade Anardana. *Journal of Food Science and Technology* **40**, 236–238.
- Singh GP, Nagpal AK and Sainj N** (2005) Methane production in relation to productivity of livestock and environment. A review. *Indian Journal of Animal Science* **75**, 143–148.
- Soliva CR and Hess HD** (2007) Measuring methane emission of ruminants by in vitro and in vivo techniques. In Makkar HPS and Vercoe PE (eds). *Measuring Methane Production from Ruminants*. Dordrecht, The Netherlands: Springer, pp. 15–31.
- Tavendale MH, Meagher LP, Pacheco D, Walker N, Attwood GT and Sivakumaran S** (2005) Methane production from *In vitro* Rumen incubations with *Lotus pedunculatus* and *Medicago sativa*, and effects of extractable condensed tannin fractions on methanogenesis. *Animal Feed Science and Technology* **123–124**, 403–419.
- Teerlink T, Marcel H, Joli B and John G** (1993) Simultaneous determination of certain compounds and adenine nucleotides, in mitochondrial tissue rat by high performance liquid chromatography. *Analytical Biochemistry* **214**, 278–283.
- Van Soest PJ, Robertson JB and Lewis BA** (1991) Methods for dietary fiber, neutral detergent fiber and non-starch polysaccharides in relation to animal nutrition. *Journal of Dairy Science* **74**, 3583–3597.
- Vogels GD, Hoppe WF and Stumm CK** (1980) Association of methanogenic bacteria with rumen ciliates. *Applied and Environmental Microbiology* **40**, 608–612.
- Waghorn G** (1996) Condensed tannins and nutrient absorption from the small intestine. In Rode LM (ed) Proc of the 1996 Canadian Society of Animal Science Annual Meeting, Lethbridge, Canada, 175–194.
- Yogianto D, Sudarman A, Wina E and Jayanegara A** (2014) Supplementation effects of tannins and saponin extracts to diets with different forage to concentrate ratio on *In vitro* rumen fermentation and methanogenesis. *Journal of the Indonesian Tropical Animal Agriculture* **39**, 144–151.