

## Standard Paper

# *Neoplaca mirabilis*, a new genus and a new epigaeic species containing naphthopyrans from the family *Teloschistaceae*

Ivan V. Frolov<sup>1</sup> , Ilya A. Prokopiev<sup>2</sup> and Liudmila A. Konoreva<sup>3,4,5</sup> 

<sup>1</sup>Institute Botanic Garden, Ural Branch of the Russian Academy of Sciences (RAS), 620144 Yekaterinburg, Russia; <sup>2</sup>Laboratory of Analytical Phytochemistry, Komarov Botanical Institute RAS, 197376 St Petersburg, Russia; <sup>3</sup>Laboratory of Lichenology and Bryology, Komarov Botanical Institute RAS, 197376 St Petersburg, Russia; <sup>4</sup>Botanical Garden-Institute, Far Eastern Branch RAS, Vladivostok, Russia and <sup>5</sup>Polar-Alpine Botanical Garden and Institute N. A. Avrorin, a separate branch of the Federal Research Center 'Kola Science Center' RAS, Russia

### Abstract

The production of anthraquinones is a major characteristic of most species in the *Teloschistaceae*. Other secondary metabolites are quite rare in this family, but some species are known to produce depsides, depsidones, xanthonones and usnic acid. A new monotypic genus, *Neoplaca*, with a new species *N. mirabilis*, is described from the subfamily *Caloplocoideae* of the family *Teloschistaceae*, lacking anthraquinones but containing the naphthopyrans simonyellin and consimonyellin. This is the first time this class of organic compounds has been found in the family *Teloschistaceae* and the second in the order *Teloschistales*, where simonyellin has been detected in *Brigantiaaceae*. Simonyellin and consimonyellin have also previously been reported in the family *Roccellaceae*. *Neoplaca mirabilis* is currently known from the two nearby localities in Yakutia, Russia, where it is common and grows on base-rich soil on exposed south-facing siliceous outcrops. The thallus consists of scattered whitish to greyish, or rarely with pale yellow tinge, squamules 1–4.5 mm diam. and 0.3–1 mm thick with citrine to orange-yellow blastidia produced from their margin; apothecia and pycnidia are unknown. In addition to naphthopyrans, *N. mirabilis* contains an unidentified brown pigment similar in some features to melanin. The new species is also interesting in that the pigments are apparently located inside the cells of the cortex, not on their surface, where anthraquinones are found in *Teloschistaceae*.

**Keywords:** consimonyellin; lichen; melanin-like pigment; simonyellin; Yakutia

(Accepted 17 July 2023)

### Introduction

The production of yellow-orange-red anthraquinone pigments in the superficial tissues is a major characteristic of most species in the family *Teloschistaceae* and in many taxa these are the only lichen compounds present (Santesson 1970). Our knowledge of these anthraquinones is increasing (Elix *et al.* 2000; Søchting *et al.* 2014). Depending on the combination of the biosynthetically related anthraquinones and their proportions, several chemosyndromes have been distinguished (Søchting 1997, 2001). Chemosyndrome is a fairly constant and specific character, and most species have only one syndrome.

Secondary metabolites other than anthraquinones are quite rare in *Teloschistaceae*. Some species, in addition to anthraquinones, produce depsides (atranorin, gyrophoric acid and lecanoric acid; see Joshi *et al.* 2010; Arup *et al.* 2013; Vondrák *et al.* 2020; Zhang *et al.* 2019), depsidones (caloploicin, vicanicin and isofulgidin; see Søchting & Frödén 2002; Søchting & Figueras 2007; Bungartz *et al.* 2020), xanthonones (lichexanthone; see Bungartz *et al.* 2020), and usnic acid (Arup *et al.* 2013). Together with anthraquinones, many species also contain dark pigments of an unknown chemical structure, as for example, *Cinereorufa*-green and *Sedifolia*-grey,

which are insoluble in acetone and therefore unextractable by common methods (Wetmore 1996; Meyer & Printzen 2000; Vondrák *et al.* 2012, 2020; Frolov *et al.* 2021). In a small number of lineages of *Teloschistaceae*, anthraquinones are occasionally or entirely absent and replaced by other compounds (Vondrák *et al.* 2012), such as depsides and depsidones in *Olegblumia demissa* (Flot. ex Körb.) S. Y. Kondr. *et al.* and *Sucioplaca diplacia* (Ach.) Bungartz *et al.* (Søchting & Figueras 2007; Bungartz *et al.* 2020) or *Sedifolia*-grey in *Pyrenodesmia* (Frolov *et al.* 2021).

During three field trips in Yakutia (East Siberia, Russia), we collected a peculiar sterile epigaeic lichen crust lacking anthraquinones. Although collected only from the two nearby localities, it was common there. Subsequent analyses demonstrated that it belonged to the family *Teloschistaceae* and contained a remarkable chemistry (naphthopyrans) that is uncommon in lichens and previously unknown in *Teloschistaceae*, as well as an unidentified brown pigment resembling melanin. The lichen was not identified as any known taxon and due to its remarkable chemistry and morphology, and considering the current status of the taxonomy of *Teloschistaceae*, it is described here as a new species in a new genus.

### Material and Methods

#### Sampling and phenotype evaluation

Several thalli were collected from the two localities in the Republic of Sakha (Yakutia) in Russia by L. Konoreva in 2016 and by

**Corresponding author:** Ivan V. Frolov; Email: [ivfrolov@gmail.com](mailto:ivfrolov@gmail.com)

**Cite this article:** Frolov IV, Prokopiev IA and Konoreva LA (2023) *Neoplaca mirabilis*, a new genus and a new epigaeic species containing naphthopyrans from the family *Teloschistaceae*. *Lichenologist* 55, 443–450. <https://doi.org/10.1017/S0024282923000567>



I. Prokopiev and I. Frolov in 2021 and 2023. The specimens are deposited in LE, C, PRA and the personal herbarium of I. Frolov. Specimens from 2016 and 2021 were subjected to morphological, biochemical and phylogenetic analyses.

Measurements of morphological characters follow Vondrák *et al.* (2013) and microscopic observations are based on hand-cut sections mounted in water without chemical treatment. Measurements are accurate to 0.5  $\mu\text{m}$  for cells and 5–10  $\mu\text{m}$  for larger structures. Measurements of particular characters are given as  $\bar{x}_1 - \bar{x}_2 - \bar{x}_3$  ( $n$ ), where  $\bar{x}_1$  is the minimum value,  $\bar{x}_2$  is the arithmetic mean of all measurements,  $\bar{x}_3$  is the maximum value, and  $n$  is the total number of measurements. Morphological terminology follows Vondrák *et al.* (2013) and the LIAS glossary (available at <https://glossary.lias.net/wiki/>).

The following were used for spot tests: 10% aqueous KOH solution (K), aqueous solution of calcium hypochlorite (C) and 5% alcoholic p-phenylenediamine solution (P).

### Identification of secondary metabolites (TLC, UV/Vis spectrometric and HPLC-UV-ESI-QTOF/MS analyses)

For thin-layer chromatography (TLC) and high-performance liquid chromatography with UV detection coupled with electrospray-ionization quadrupole time-of-flight mass spectrometry (HPLC-UV-ESI-QTOF/MS), air-dried samples (4 mg) of the lichen were ground to a powder. Secondary substances were extracted from each sample with 0.3 ml of acetone. Extraction was carried out with constant stirring for 12 h at 20–25 °C. The obtained extracts were centrifuged for 10 min at 6000 g and kept at 4 °C until analysis.

TLC was performed on silica gel 60 plates (Merck, Darmstadt, Germany) using solvent system C (toluene:acetic acid = 170:30 v/v) according to Huneck & Yoshimura (1996). Authentic atranorin and norstictic acid were used as controls. After chromatographic development, the plates were examined under UV light (254 and 366 nm), then sprayed with a 10% sulphuric acid solution and heated at 100 °C for 15 min. Finally, the plates were cooled to room temperature and studied in daylight.

For UV/Vis spectrometric analysis, spots from the TLC plate (before a 10% sulphuric acid treatment) were scraped off with a scalpel. The resulting silica, containing the investigated substances, was extracted with 0.5 ml of acetonitrile. UV/Vis absorption spectra were recorded on a Beckman DU 800 spectrometer in the range 200–800 nm.

HPLC-UV-ESI-QTOF/MS analyses were performed with an Agilent 1290 Series chromatograph with UV detection. For chromatographic separation, a Thermo Hypersil-Keystone C18 column (150  $\times$  2.1 mm  $\times$  5  $\mu\text{m}$ ) was used. The mobile phase consisted of (A) water:acetonitrile:formic acid (95:5:0.1 v/v), and (B) acetonitrile:water:formic acid (90:10:0.1 v/v). Analyses were performed at 30 °C and a flow rate of 0.3 ml min<sup>-1</sup> in the gradient elution mode, and the percentage of B was programmed as follows: 5% (2 min) – 50% (5 min) – 70% (15 min) – 100% (25 min) – 100% (35 min). The volume of the injected sample was 5  $\mu\text{l}$ . Spectra of eluting substances were recorded in UV at 250 nm. After separation, the samples were also analyzed with a quadrupole time-of-flight mass spectrometer (6538 Series, Agilent, USA). Ionization was achieved by electrospray in the negative mode. Voltage on the capillary was 2.5 kV, capillary temperature 350 °C, atomizing gas pressure 45 psi, desiccant gas (nitrogen) temperature 225 °C, and drying gas flow rate 5 l min<sup>-1</sup>. Mass spectra were recorded in the range 100–1000 m/z. For

confirmation purposes, a targeted MS/MS analysis was performed. The precursor ions were filtered by the quadrupole. The collision energy of 30 eV was defined for MS/MS experiments. The resulting chromatograms were processed with MassHunter WorkStation v. B.07.00 software (Agilent, USA). In order to identify lichen substances, the obtained MS/MS spectra were compared with spectra from the MassBank of North America (MoNA).

### DNA extraction, amplification and sequencing

DNA was extracted with a CTAB-based protocol (Aras & Cansaran 2006). Amplifications were made of the internal transcribed spacer regions (nrITS), the large subunit (nrLSU) of the nuclear ribosomal RNA genes, and the small subunit of the mitochondrial ribosomal RNA gene (mtSSU). Primers for PCR amplification were ITS1F (Gardes & Bruns 1993) and ITS4 (White *et al.* 1990) for ITS, AL1R (Döring *et al.* 2000) and LR5 (Vilgalys & Hester 1990) for nrLSU, and mrSSU1 (Zoller *et al.* 1999) and mrSSU7 (Zhou & Stanosz 2001) for mtSSU. The PCR settings followed Ekman (2001). Sequences obtained were uploaded onto the NCBI database (GenBank) and Accession numbers are provided in the species protologue.

### Alignments and phylogenetic analyses

Newly obtained sequences were edited in FinchTV v. 1.4.0 (Geospiza Inc., Seattle, USA; <http://www.geospiza.com>) and BioEdit v. 7.2.5 (Hall 1999). Together with sequences downloaded from GenBank, datasets were aligned online using MAFFT v. 7 (Katoh & Standley 2013; available at <http://mafft.cbrc.jp/alignment/server/>), with the L-INS-i method (Katoh *et al.* 2005) selected automatically by the program. To exclude ambiguously aligned positions, alignments were subsequently analyzed by the gappout algorithm as implemented in the TrimAl software package (Capella-Gutierrez *et al.* 2009). Phylogenetic reconstructions of the concatenated dataset, as well as of the three single loci, were carried out using Bayesian inference in MrBayes v. 3.2.6 (Ronquist & Huelsenbeck 2003) and the analyses were run on the CIPRES Web Portal (<http://www.phylo.org/portal2/>). In the concatenated dataset, three partitions corresponding to three DNA loci were analyzed separately. Models of nucleotide substitutions for each partition (or single locus) were selected using the corrected Akaike information criterion implemented in jModelTest v. 0.1.1 (Posada 2008). The SYM + I + G model was selected for nrITS, the GTR + I + G for nrLSU, and the HKY + I + G for mtSSU. MrBayes analyses were performed using two independent runs with four MCMC chains (three cold and one heated) in each run. Trees were sampled every 500th generation. The analyses were stopped when the average standard deviation of split frequencies between the simultaneous runs dropped below 0.01 (140 000 generations for the concatenated dataset, 1 165 000 for nrITS, 1 690 000 for nrLSU, and 1 065 000 for mtSSU). The first 25% of trees was discarded as burn-in, and the remaining trees were used for construction of a 50% majority-rule consensus tree. Accession numbers of sequences downloaded from GenBank and used in the analyses are provided in Supplementary Material Table S1 (available online).

## Results and Discussion

### Phylogeny and taxonomic position of the new species

Performing an online NCBI BLAST search with sequences of all three loci of the new species demonstrated that it belongs in the

subfamily *Caloplacoideae* of the family *Teloschistaceae*, but without a distinct generic affiliation, with *c.* 87% identity of nrITS to species of *Caloplaca* s. str., *Lendemeriella* and *Pyrenodesmia*. For a more accurate determination of the taxonomic position of the new species, it was included in the combined analysis of the nrITS, nrLSU and mtSSU dataset, together with the main genera of the subfamily *Caloplacoideae* and a few genera of the subfamilies *Teloschistoideae* and *Xanthorioideae*. The combined alignment included 75 terminal species and a total of 2273 positions before trimming and 2051 positions thereafter. The phylogeny was rooted with taxa of the families *Brigantiaeaceae* and *Physciaceae* following Arup *et al.* (2013). The three single loci were also analyzed separately.


According to our phylogenetic analysis of the combined alignment, the two specimens of the new species are nested within a highly supported clade encompassing the genera *Kuettlingeria*, *Lendemeriella* and *Pyrenodesmia*, and the species '*Caloplaca*' *conversa* auct. and '*C.*' *obscura* (Fig. 1). The new species, however, does not belong to any of these genera forming a sister lineage to the specimen labelled as '*C. conversa*' in GenBank, but here it is given as '*C. conversa* auct.' since '*C.*' *conversa sensu* Krempelhuber (1861) is not related to this specimen and currently belongs in the genus *Pisutiella*. Analyses of the single loci generated the same position for the new species (not supported in nrITS and mtSSU; Supplementary Material Figs S1–S3, available online).

As a result, the new species could not be assigned to any currently established genus of *Teloschistaceae*. Due to the outstanding features of the new species, as well as the distinct position in the *Teloschistaceae* phylogeny, a new genus to encompass it is proposed.


### Secondary metabolites: HPLC-UV-ESI-QTOF/MS analysis and naphthopyrans

HPLC-UV analysis of the samples of thalli of the new species revealed two compounds with  $R_t = 13.2$  min–major and  $R_t = 13.9$  min–trace (Fig. 2). The obtained mass spectra of the substances revealed molecular ions  $[M-H]^-$  with  $m/z$  273.0403 (for  $R_t = 13.2$ ) and 305.0659 (for  $R_t = 13.9$ ) corresponding to the molecular formulas  $C_{14}H_{10}O_6$  and  $C_{15}H_{14}O_7$  respectively. Analysis of the MS/MS fragments showed that the compound  $C_{14}H_{10}O_6$  was simonyellin while  $C_{15}H_{14}O_7$  was a methoxylated derivative of simonyellin, namely consimonyellin (Fig. 3). Both chemicals belong in the class of organic compounds known as naphthopyrans. The obtained ESI-MS/MS spectra of simonyellin and consimonyellin are provided in Supplementary Material Fig. S4 (available online).


To explore the cause of the yellow colour of blastidia in the new species, these vegetative propagules were analyzed separately. After the twice-repeated extraction of separated blastidia in 0.1 ml of acetone, their colour changed from yellow to grey-green. In the



**Figure 1.** Phylogeny of the subfamily *Caloplacoideae* (*Teloschistaceae*) based on the combined Bayesian analysis of nrITS, nrLSU and mtSSU data. Genera are collapsed into single terminals. Numbers at branches represent posterior probability (PP) values  $\geq 0.95$ . Numbers in parentheses correspond to the number of species of a genus used in the analysis. I = *Teloschistaceae*; II = *Caloplacoideae*; III = *Xanthorioideae*; IV = *Teloschistoideae*.



**Figure 2.** HPLC-UV chromatograms (250 nm) of acetone extracts of *Neoplaca mirabilis*. 1 = simonyellin; 2 = consimonyellin.




**Figure 3.** Structural formulas of the *Neoplaca mirabilis* secondary metabolites.

yellow-dyed extract, naphthopyrans in the same ratio as in the whole thallus were detected by HPLC-UV. Since consimonyellin is a colourless substance (Huneck 2001), and simonyellin is a yellow pigment (Elix 2014), it can be concluded that the yellow colour of blastidia is caused by the latter compound.

This is the first time naphthopyrans have been detected in the family *Teloschistaceae* and the second in the order *Teloschistales*; here, simonyellin was previously known in the family *Brigantiaaceae*, and more particularly in the species *Brigantiaea leucoxantha* and *B. tricolor* (Elix 2008). Naphthopyran derivatives are not widespread lichen substances. Besides *Brigantiaaceae*, simonyellin was known only in the family *Roccellaceae*, in the genera *Cresponea* and *Simonyella*, and in the genus *Bactrospora* that has an uncertain position in the *Arthoniales* (Elix et al. 1995, 2011; Kalb 2004; Berger & Aptroot 2008; Kantvilas 2020), whereas consimonyellin has only been detected in the genus *Cresponea* (Elix et al. 2011; Kantvilas 2020). However, phenalenone, the precursor of naphthopyrans, is a well-known metabolite of non-lichenized fungi (Cooke & Edwards 1981). Takenaka et al. (2010) demonstrated that the isolated mycobiont of *Lecanora leprosa* was able to biosynthesize the naphthopyran-lecanopyrone, which was not found in the lichenized state of the mycobiont. It can be assumed that the ability of lichens to biosynthesize naphthopyrans represents the expression of the biosynthetic gene clusters of free-living fungi normally suppressed in the lichenized condition in lichens.

### Secondary metabolites: TLC and UV/Vis spectrometric analyses and an unknown brown pigment

TLC analysis (solvent system C; Fig. 4) revealed a yellow-coloured compound with  $R_f=0.6$  corresponding to simonyellin and a brown pigment with  $R_f=0.2$ , which was not detected by



**Figure 4.** TLC analysis in solvent C (toluene:acetic acid = 170:30 v/v). Controls = C; Nm = *Neoplaca mirabilis*. 1 = atranorin; 2 = norstictic acid; 3 = simonyellin; 4 = brown pigment. In daylight after spraying with 10% sulphuric acid and drying at 100 °C for 10 min. In colour online.

HPLC-UV. UV/Vis spectrum of acetonitrile solution of simonyellin shows absorption maxima at wavelengths 241, 264, 274, 307, 326 and 365 nm. The obtained UV/Vis spectrum of simonyellin is provided in Supplementary Material Fig. S5 (available online).

The brown pigment was readily soluble in 10% KOH and sparingly in water, methanol, ethanol and acetone. This pigment discoloured in reaction with oxidizing agents (0.5 M KMnO<sub>4</sub>, 30% H<sub>2</sub>O<sub>2</sub>) and produced a brown precipitate with 1% FeCl<sub>3</sub>. These characteristics resemble those displayed by the pigment melanin (Khabibrakhmanova *et al.* 2022); however, the solubility of our pigment in the organic solvents is not typical for melanin. Due to the scarcity of the available material, it was not possible to isolate enough brown pigment to obtain the infrared spectra that could confirm its assignment to melanin.

## Taxonomy

### *Neoplaca* I. V. Frolov, Prokopiev & Konoreva gen. nov.

Mycobank No.: MB 849087

Thallus consists of squamules lacking anthraquinones and containing naphthopyrans.

Type species: *Neoplaca mirabilis* I. V. Frolov, Prokopiev & Konoreva.

*Thallus* whitish, greyish to yellowish, squamulose with citrine to orange-yellow blastidia on margin.

*Apothecia* and *pycnidia* unknown.

**Chemistry.** Thallus contains simonyellin as a major compound, traces of consimonyellin, and an unknown brown pigment. Pigments are apparently located inside the cells of the cortex.

**Etymology.** The genus name is a combination of the syllables ‘-placa’, hinting at *Caloplaca* s. lat., and ‘neo-’, indicating the innovative chemical and morphological characters of the new species of the genus.

**Distribution.** Currently the genus is known only from the two nearby localities in Yakutia, East Siberia.

**Notes.** *Neoplaca mirabilis* is the only species included in the genus, and despite its remarkable morphological and chemical features uncharacteristic of the *Teloschistaceae*, the new genus has just a common position among the other genera of the subfamily *Caloplacoideae* (Fig. 1). Due to the absence of anthraquinones, apothecia and pycnidia (see protologue), the taxonomic position of the new species within *Teloschistaceae* is based solely on molecular data.

### *Neoplaca mirabilis* I. V. Frolov, Prokopiev & Konoreva sp. nov.

Mycobank No.: MB 848237

Thallus epigeaic, consisting of scattered whitish to greyish, or rarely with pale yellow tinge, squamules 1–4.5 mm diam., 0.3–1 mm thick. Blastidia common, produced from the margin of squamules, citrine to orange-yellow, highly contrasting with the colour of the upper surface. Apothecia and pycnidia are unknown. Contains simonyellin, consimonyellin and an unknown brown pigment. Anthraquinones are absent.

Type: Russia, Republic of Sakha (Yakutia), Tomponsky District, 145 km NE of Khandyga, right bank of the Vostochnaya Khandyga River, along the stream crossing the Kolyma Highway at the 585th km, 580 m alt., 63°07'17.2"N, 138°14'33.8"E, soil on exposed south-facing siliceous outcrops on the left bank of the stream, 12 August 2021, I. Frolov 3706 & I. Prokopiev (LE L-21846—holotype; C, PRA—iso-types). GenBank Accession numbers of the holotype sequences: OQ721918 (nrITS), OQ721950 (nrLSU), OQ724518 (mtSSU).

(Fig. 5A–D)


*Thallus* epigeaic, consisting of scattered whitish to greyish, or with pale yellow tinge, squamules, starting as tiny granules c. 0.1 mm diam., developing into small convex areoles/squamules 0.5–1 mm diam. and then into mature squamules 1–4.5 mm diam., sometimes with obscurely lobed margin, usually with blastidia and surface with wide, convoluted folds, in some places resembling the appearance of the brain in mammals; sometimes the squamules overlap each other like roof tiles or merge to form effigurate thalli c. 1 cm diam. (Fig. 5B); mature squamules are often surrounded by numerous tiny primordia, apparently germinating from blastidia (Fig. 5C), probably indicating the considerable ability of the species to reproduce vegetatively. *Squamules* in areas attached to substratum 0.7–1 mm thick, and those in detached parts thinner, up to 0.3–0.4 mm. *Cortex* paraplectenchymatous, 33–40 µm thick, consisting of two clearly distinguishable layers (Fig. 5D), a lower layer, c. 20 µm thick, consisting of colourless cells, and an upper layer, 13–20 µm thick, composed of cells with brown protoplasts and transparent cell walls (Fig. 5D). *Cortex* covered with uneven layer of white pruina, 12.5–45 µm thick, consisting of colourless crystals insoluble in KOH of various shapes and sizes up to 20 µm; cells of cortex isodiametric, 5–5.9–7 µm diam. (*n* = 10); on the lower side of squamules, cortex is absent. *Algal layer* 55–83 µm thick; algal cells globose, 9–12.2–15 µm diam. (*n* = 10). *Medulla* 0.2–0.8 mm thick, white, lax, composed of loosely interwoven, irregularly arranged hyphae up to 5 µm thick, with walls up to 2 µm thick and narrow lumina 1–2 µm. *Blastidia* common, produced from margin of squamules, citrine to orange-yellow, highly contrasting with the colour of the upper surface, rarely whitish, 58–89–125 µm diam. (*n* = 10), covered with a thin layer of brown pigment and tiny hook-shaped colourless hairs; hyphal sheath of blastidia 35–38 µm thick, consisting of isodiametric cells 5–7 µm diam.

*Apothecia* and *pycnidia* not observed.

**Chemistry.** Simonyellin (detected by both TLC and HPLC, major), consimonyellin (HPLC, trace), and an unknown brown pigment (TLC). Anthraquinones not detected. Squamules and blastidia K+ orange-brown, C–, P+ yellow (fleeting reaction). The reaction of the thallus with K does not correspond to simonyellin, which should be K– (Elix 2014), but to the brown pigment. Its spot on the TLC plate demonstrates the same reaction with K.

Pigments of the new species are apparently located inside the cells of the cortex (Fig. 5D), whereas anthraquinones and, for example, Sedifolia-grey are located on the cell surface. After adding K to the cortex cross-section, the brown grains in the protoplasts dissolve, and the cell walls and protoplasts become uniformly coloured yellow or orange.

**Etymology.** The epithet reflects the highly unusual chemistry and habit of the new species for the *Teloschistaceae*.




**Figure 5.** A–D, *Neoplaca mirabilis* (holotype). A, squamule with folded surface and blastidia. B, effigurate thallus formed by merged squamules. C, mature squamule surrounded by numerous primordia. D, cross-section of the cortex with the cell protoplasts coloured by a brown pigment. Scales: A–C = 2 mm; D = 10  $\mu$ m. In colour online.

**Similar taxa.** *Neoplaca mirabilis* is an unusual species owing to its large, whitish pruinose squamules with contrasting citrine to orange-yellow blastidia. Combined with its outstanding chemistry, the new species cannot really be confused with any other known taxon. Nevertheless, it may resemble the sterile epigeaic thalli of squamulose *Xanthocarpia tominii* (Savicz) Frödén *et al.* which, however, has bright, orange to ochre-yellowish, much smaller squamules up to 1.5 mm diam., soredia not contrasting with the thallus, and a purple reaction in K (contains anthraquinones). When squamules of the new species merge to form effigurate thalli, *N. mirabilis* may resemble *Calogaya decipiens* (Arnold) Arup *et al.* or *Gyalolechia lenae* (Søchting & G. Figueras) Søchting *et al.* that occasionally occur on soil on rocks. These species are easily distinguishable by the purple reaction in K of their thalli, soralia and apothecia (anthraquinones), as well as the pale yellow to orange colour of their thalli and soralia. Note also that sorediate *Flavoplaca flavocitrina* (Nyl.) Arup *et al.* and blastidiate *Gyalolechia epiphyta* (Lynge) Vondrák, which can be epigeaic, have areolate yellow to orange thalli containing anthraquinones (K+ purple). Furthermore, epigeaic species of the former genus *Fulgensia* A. Massal. & De Not. (currently included in *Gyalolechia* A. Massal.) can resemble the new species, especially when sterile, but they do not have blastidia and have a purple reaction in K. Other crustose *Teloschistaceae* occurring on soil

(e.g. *Blastenia ammiospila* (Wahlenb. ex Ach.) Arup *et al.*, *Bryoplaca* spp., *Calogaya schistidii* (Anzi) Arup *et al.* and '*Caloplaca*' *raesaenii* Bredkina) are richly fertile, have areolate or poorly developed thalli lacking vegetative propagules, and contain anthraquinones, at least in their apothecia. Other epigeaic crustose taxa known from Siberia (e.g. species of *Endocarpon*, *Psora* and *Toninia*) lack vegetative propagules and naphthopyrans.

**Ecology and distribution.** Currently the new species is known from the two nearby localities in Yakutia (right bank of the Vostochnaya Khandyga River), where it is quite common and grows on base-rich soil on exposed south-facing siliceous outcrops along small brooks at an altitude of 550–850 m a.s.l. together with species of *Collema*, *Endocarpon*, *Leptogium* and *Toninia*. The valley slopes of these brooks are covered with *Larix gmelinii* forest and the bottom of the valleys are treeless, which is probably due to the destructive impact of floods (Fig. 6A & B).


**Additional material studied.** **Russia:** *Yakutia:* same as the type, 2016, L. Konoreva J-304 (LE L-21889, hb. Frolov; GenBank Accession numbers: OQ721919 (nrITS), OQ721949 (nrLSU), Q724517 (mtSSU)); *ibid.*, along the brook crossing the Kolyma



**Figure 6.** A & B, type locality of the newly described *Neoplaca mirabilis*. A, general view. B, soil on rocks with whitish thalli of the new species. In colour online.

Highway at the c. 620th km, 63°08'42.8"N, 138°43'11.0"E, 850 m alt., 2023, *I. Frolov* 3847 & *I. Prokopiev* (hb. Frolov).

**Acknowledgements.** Prof. Mark Seaward kindly revised the manuscript. Ivan V. Frolov and Ilya A. Prokopiev were supported by the Russian Science Foundation (RSF grant no. 23-24-00207: <https://rscf.ru/en/project/23-24-00207/>).

**Author ORCIDs.**  Ivan V. Frolov, 0000-0003-4454-3229; Ilya A. Prokopiev, 0000-0001-8755-7140; Liudmila A. Konoreva, 0000-0002-4487-5154.

**Competing Interests.** The authors declare none.

**Supplementary Material.** The Supplementary Material for this article can be found at <https://doi.org/10.1017/S0024282923000567>.

## References

- Aras S and Cansaran D (2006) Isolation of DNA for sequence analysis from herbarium material of some lichen specimens. *Turkish Journal of Botany* **30**, 449–453.
- Arup U, Søchting U and Frödén P (2013) A new taxonomy of the family *Teloschistaceae*. *Nordic Journal of Botany* **31**, 16–83.
- Berger F and Aptroot A (2008) *Bactrospora flavopruinosa*, a new lichen species from Bermuda. *Lichenologist* **40**, 543–547.
- Bungartz F, Søchting U and Arup U (2020) *Teloschistaceae* (lichenized *Ascomycota*) from the Galapagos Islands: a phylogenetic revision based on morphological, anatomical, chemical, and molecular data. *Plant and Fungal Systematics* **65**, 515–576.
- Capella-Gutierrez S, Silla-Martinez JM and Gabaldon T (2009) TrimAl: a tool for automated alignment trimming in large-scale phylogenetic analyses. *Bioinformatics* **25**, 1972–1973.
- Cooke RG and Edwards JM (1981) Naturally occurring phenalenones and related compounds. In Herz W, Grisebach H and Kirby GW (eds), *Fortschritte der Chemie organischer Naturstoffe / Progress in the Chemistry of Organic Natural Products* **40**. Vienna: Springer Vienna, pp. 153–190.
- Döring H, Clerc P, Grube M and Wedin M (2000) Mycobiont-specific PCR primers for the amplification of nuclear ITS and LSU rDNA from lichenized ascomycetes. *Lichenologist* **32**, 200–204.
- Ekman S (2001) Molecular phylogeny of the *Bacidiaceae* (*Lecanorales*, lichenized *Ascomycota*). *Mycological Research* **105**, 783–797.
- Elix JA (2008) Lichen phytochemistry: additions and amendments I. *Australasian Lichenology* **63**, 20–25.
- Elix JA (2014) *A Catalogue of Standardized Chromatographic Data and Biosynthetic Relationships for Lichen Substances*, 3rd Edn. Canberra: Published by the author.
- Elix JA, Feige GB, Lumbsch HT, Mies B and Wardlaw JH (1995) The structure of simonyellin – a new lichen naphthopyran. *Australian Journal of Chemistry* **48**, 2035–2039.
- Elix JA, Wardlaw JH and Sochting U (2000) Two new anthraquinones from the lichen *Caloplaca spitsbergensis*. *Herzogia* **14**, 27–30.
- Elix JA, Øvstedal DO and Gremmen NJM (2011) A new species of *Cresponea* (*Roccellaceae*, *Ascomycota*) from Gough Island. *Nova Hedwigia* **92**, 133–136.
- Frolov I, Vondrák J, Košnar J and Arup U (2021) Phylogenetic relationships within *Pyrenodesmia* sensu lato and the role of pigments in its taxonomic interpretation. *Journal of Systematics and Evolution* **59**, 454–474.
- Gardes M and Bruns TD (1993) ITS primers with enhanced specificity for basidiomycetes – application to the identification of mycorrhizae and rusts. *Molecular Ecology* **2**, 113–118.
- Hall TA (1999) BioEdit: a user-friendly biological sequence alignment editor and analysis program for Windows 95/98/NT. *Nucleic Acids Symposium Series* **41**, 95–98.
- Huneck S (2001) New results on the chemistry of lichen substances. *Fortschritte der Chemie Organischer Naturstoffe* **81**, 1–276.
- Huneck S and Yoshimura I (1996) *Identification of Lichen Substances*. Berlin, Heidelberg: Springer-Verlag.
- Joshi Y, Wang XY, Yamamoto Y, Koh YJ and Hur JS (2010) A first modern contribution to *Caloplaca* biodiversity in South Korea: two new species and some new country records. *Lichenologist* **42**, 715–722.
- Kalb K (2004) New or otherwise interesting lichens II. *Bibliotheca Lichenologica* **88**, 301–329.
- Kantvilas G (2020) Contributions to the lichen genus *Cresponea* (*Roccellaceae*). *Lichenologist* **52**, 279–285.
- Katoh K and Standley DM (2013) MAFFT multiple sequence alignment software version 7: improvements in performance and usability. *Molecular Biology and Evolution* **30**, 772–780.
- Katoh K, Kuma K, Toh H and Miyata T (2005) MAFFT version 5: improvement in accuracy of multiple sequence alignment. *Nucleic Acids Research* **33**, 511–518.
- Khabibrakhmanova VR, Rassabina AY, Khayrullina AF and Minibayeva FV (2022) Physico-chemical characteristics and antioxidant properties of

- melanins extracted from *Leptogium furfuraceum* (Harm.). *Khimiya Rastitel'nogo Syr'ya* **4**, 115–125. (In Russian).
- Krempelhuber A** (1861) Die Lichenflora Bayerns oder Aufzählung der bisher in Bayern (diesseits des Rheins) aufgefundenen Lichenen mit besonderer Berücksichtigung der verticalen Verbreitung dieser Gewächse in den Alpen. *Denkschriften der Bayer. Botanischen Gesellschaft in Regensburg* **4**, 1–317.
- Meyer B and Printzen C** (2000) Proposal for a standardized nomenclature and characterization of insoluble lichen pigments. *Lichenologist* **32**, 571–583.
- Posada D** (2008) jModelTest: phylogenetic model averaging. *Molecular Biology and Evolution* **25**, 1253–1256.
- Ronquist F and Huelsenbeck JP** (2003) MrBayes 3: Bayesian phylogenetic inference under mixed models. *Bioinformatics* **19**, 1572–1574.
- Santesson J** (1970) Anthraquinones in *Caloplaca*. *Phytochemistry* **9**, 2149–2166.
- Sochting U** (1997) Two major anthraquinone chemosyndromes in *Teloschistaceae*. *Bibliotheca Lichenologica* **68**, 135–144.
- Sochting U** (2001) Chemosyndromes with chlorinated anthraquinones in the lichen genus *Caloplaca*. *Bibliotheca Lichenologica* **78**, 395–404.
- Sochting U and Figueras G** (2007) *Caloplaca lenae* sp. nov., and other *Caloplaca* species with caloploicin and vicanicin. *Lichenologist* **39**, 7–14.
- Sochting U and Frödén P** (2002) Chemosyndromes in the lichen genus *Teloschistes* (*Teloschistaceae*, *Lecanorales*). *Mycological Progress* **1**, 257–266.
- Sochting U, Sogaard MZ, Elix JA, Arup U, Elvebakk A and Sancho LG** (2014) *Catenarina* (*Teloschistaceae*, *Ascomycota*), a new Southern Hemisphere genus with 7-chlorocatenarin. *Lichenologist* **46**, 175–187.
- Takenaka Y, Hamada N and Tanahashi T** (2010) Structure and biosynthesis of lecanopyrone, a naphtho[1,8-cd]pyran-3-one derivative from cultured lichen mycobionts of *Lecanora leprosa*. *Zeitschrift für Naturforschung C* **65**, 637–641.
- Vilgalys R and Hester M** (1990) Rapid genetic identification and mapping of enzymatically amplified ribosomal DNA from several *Cryptococcus* species. *Journal of Bacteriology* **172**, 4239–4246.
- Vondrák J, Šoun J, Vondráková O, Fryday A, Khodosovtsev A and Davydov E** (2012) Absence of anthraquinone pigments is paraphyletic and a phylogenetically unreliable character in the *Teloschistaceae*. *Lichenologist* **44**, 401–418.
- Vondrák J, Frolov I, Arup U and Khodosovtsev A** (2013) Methods for phenotypic evaluation of crustose lichens with emphasis on *Teloschistaceae*. *Chornomorskiy Botanichniy Zhurnal* **9**, 382–405.
- Vondrák J, Frolov I, Košnar J, Arup U, Veselská T, Halıcı G, Malíček J and Sochting U** (2020) Substrate switches, phenotypic innovations and allopatric speciation formed taxonomic diversity within the lichen genus *Blastenia*. *Journal of Systematics and Evolution* **58**, 295–330.
- Wetmore CM** (1996) The *Caloplaca sideritis* group in North and Central America. *Bryologist* **99**, 292–314.
- White TJ, Bruns T, Lee S and Taylor JW** (1990) Amplification and direct sequencing of fungal ribosomal RNA genes for phylogenetics. In Innis MA, Gelfand DH, Sninsky JJ and White TJ (eds), *PCR Protocols: a Guide to Methods and Applications*. New York: Academic Press, pp. 315–322.
- Zhang YY, Wang XY, Li LJ, Sochting U, Yin AC, Wang SQ and Wang LS** (2019) *Upretia squamulosa*, a new lichen species from the arid valley of Jinsha-Jiang River, China. *Phytotaxa* **402**, 288–294.
- Zhou S and Stanosz GR** (2001) Primers for amplification of mtSSU rDNA, and a phylogenetic study of *Botryosphaeria* and associated anamorphic fungi. *Mycological Research* **105**, 1033–1044.
- Zoller S, Scheidegger S and Sperisen C** (1999) PCR primers for the amplification of mitochondrial small subunit ribosomal DNA of lichen-forming ascomycetes. *Lichenologist* **31**, 511–516.