

**Methods:** This research was a multi-stage project involving surveys and interviews with community pharmacists and Black Africans and Caribbeans as participants.

**Results:** Knowledge, attitude and beliefs around dementia and its causes appeared to be major barriers to help seeking among the Black African and Caribbean population. For example, beliefs that dementia is caused by 'the spirits' and dementia is a repercussion for past wrongdoing and therefore not amenable to medical intervention. The community pharmacists believe they are well positioned to spot initial signs of dementia among their clients and are therefore willing to offer help seeking support to this population.

**Conclusions:** To offer intervention for timely help-seeking for dementia, a culturally tailored dementia education for the Black African and Caribbean population should be considered. In addition, training on the impact of cultural beliefs on help seeking for dementia should be considered for the community pharmacists.

### **FC60: Restless legs syndrome and ferritin levels in older people with dementia: a cross-sectional study**

**Authors:** Madson Alan Maximiano-Barreto<sup>1,2</sup>; Érica Dayanne Meireles Leite<sup>1,2</sup>, Letícia Lambert<sup>3</sup>, Victor Oliveira Wercelens<sup>4</sup>; Alan Luiz Eckeli<sup>1</sup>; Marcos Hortes Nisihara Chagas<sup>1,2</sup>

1. Neurociences and Behavioral Sciences Department, University of São Paulo, Ribeirão Preto, SP, Brazil
2. Study and Research Group in Mental Health, Cognition and Aging, University of São Paulo, Ribeirão Preto, SP, Brazil
3. Bairral Psychiatry Institute, Itapira, SP, Brazil
4. Hospital São Silvestre, Aparecida de Goiânia - GO, Brazil

**Objectives:** To evaluate the relationship between Willis-Ekbom Disease/Restless Legs Syndrome and iron deficiency anemia in older people with dementia.

**Methods:** A cross-sectional study was conducted with 70 older people diagnosed with dementia and restless leg syndrome in a Psychogeriatric outpatient clinic in a city in the interior of São Paulo, Brazil. The older people filled in instruments of sociodemographic characterization, measures to evaluate the Restless Legs Syndrome, neuropsychiatric symptoms, sleep quality, sleepiness and cognition. Blood data were also collected levels of creatinine, ferritin, red blood cells, hemoglobin and hematocrit, the latter collected in the patients' medical records.

**Results:** The sample consists mostly of older people with mixed dementia (i.e., Alzheimer's disease + Vascular Dementia), with 39% of female patients and mean age of 77.80 years (9.36). This study identified a frequency of 15.7% of Restless Legs Syndrome. Patients with the syndrome present more frequency of neuropsychiatric symptoms, worse sleep quality, higher index of body mass and lower levels of ferritin ( $p < .05$ ).

**Conclusions:** A frequency of 15.7% was identified for restless leg syndrome among patients with dementia. In addition, patients with the syndrome have ferritin deficiency.

**Keywords:** dementia, aging, ferritin, restless legs syndrome; Willis-Ekbom disease