

# Searching the web: a survey on the quality of advice on postnatal sequelae of intrauterine growth restriction and the implication of developmental origins of health and disease

S. Perzel<sup>1</sup>, H. Huebner<sup>2</sup>, W. Rascher<sup>1</sup>, C. Menendez-Castro<sup>1</sup>, A. Hartner<sup>1</sup> and F. B. Fahlbusch<sup>1</sup>

<sup>1</sup>*Department of Pediatrics and Adolescent Medicine, Friedrich-Alexander-University of Erlangen-Nuremberg, Erlangen, Germany*

<sup>2</sup>*Department of Gynaecology and Obstetrics, Friedrich-Alexander-University of Erlangen-Nuremberg, Erlangen, Germany*

Intrauterine growth restriction (IUGR) and fetal growth restriction (FGR) are pregnancy complications associated with morbidity in later life. Despite a growing body of evidence from current research on developmental origins of health and disease (DOHaD), little information is currently provided to parents on long-term metabolic, cardiovascular and neurologic consequences. As parents strongly rely on internet-based health-related information, we examined the quality of information on IUGR/FGR sequelae and DOHaD in webpages used by laypersons. Simulating non-clinicians experience, we entered the terms ‘IUGR consequences’ and ‘FGR consequences’ into Google and Yahoo search engines. The quality of the top search-hits was analyzed with regard to the certification through the Health On the Net Foundation (HON), currentness of cited references, while reliability of information and DOHaD-related consequences were assessed via the DISCERN Plus score (DPS). Overall the citation status was not up-to-date and only a few websites were HON-certified. The results of our analysis showed a dichotomy between the growing body of evidence regarding IUGR/FGR-related sequelae and lack of current guidelines, leaving parents without clear directions. Furthermore, detailed information on the concept of DOHaD is not provided. These findings emphasize the responsibility of the individual physician for providing advice on IUGR/FGR-related sequelae, monitoring and follow-up.

*Accepted 19 December 2016; Revised 18 April 2017; Accepted 19 April 2017; First published online 22 May 2017*

**Key words:** consequences, DISCERN Plus score, DOHaD, FGR, HONcode, IUGR, sequelae

## Introduction

Inspired by the Barker hypothesis,<sup>1</sup> the Developmental Origins of Health and Disease (DOHaD) theorem has vastly evolved during the last decade<sup>2</sup> and along with it the knowledge of short- and long-term sequelae of intrauterine or fetal growth restriction (IUGR/FGR).<sup>3</sup> It became evident that IUGR/FGR is involved in various clinical and public health aspects.<sup>3–5</sup> There is growing evidence that the early life environment plays a pivotal role in influencing the risk to develop a wide range of non-communicable diseases (such as type-2 diabetes and cardiovascular disease) in later life.<sup>6,7</sup> Research in the field of IUGR/FGR has increased the awareness for the relevance of these pregnancy outcomes and led to the development of diagnostic screening tools to better recognize fetuses at-risk even among low-risk pregnancies.<sup>3</sup> While the current obstetric management is focused on the right time to deliver in order to prevent adverse perinatal outcomes, neonatologic care is dedicated to treatment of problems related to prematurity<sup>8</sup> – partly due to the lack of awareness regarding the relevance of IUGR/FGR long-term sequelae.<sup>5</sup> Although it is recognized that the postnatal feeding regimen and monitoring of infant growth patterns may be essential for long-term outcome,<sup>3</sup> the

establishment of aftercare and treatment guidelines to manage the medical sequelae associated with IUGR/FGR remains a current challenge.<sup>9</sup> This is especially of importance as patients diagnosed with IUGR/FGR gain increasingly more knowledge about the condition and its consequences online.

In order to supplement the information provided by their physician, women nowadays frequently obtain medical information on pregnancy and childbirth from the internet<sup>10–13</sup> and from other media.<sup>14</sup> It is conceived that this exposure might influence their decision making.<sup>10,15</sup> Unfortunately, pregnant women rarely discuss the information they accessed online with their health providers, who may not be aware of misconceptions based on potentially inaccurate information found on the web.<sup>13</sup> Postnatally, parents intensively utilize the internet as a popular resource for pediatric health-related guidance, which has been shown by numerous studies.<sup>16–18</sup> Based on Ofcom’s Adults’ Media Use and Attitudes Report 2015<sup>19</sup> the average adult in the United Kingdom currently spends more than 20 h/week online. The internet usage is currently at an all-time high with an estimated 87.9% of adults in the United Kingdom having used the internet in the last 3 months.<sup>20</sup> Health information-related internet use in the United Kingdom has remained stable at 16% weekly since 2005<sup>19</sup> among adults aged 16+. In the United States, a study from 2010 found that ~74% of English-speaking adults use the internet on a regular basis.<sup>21</sup> Thus, online health information is available to the majority of the adult population. To analyze changes in the U.S. health information environment,

\*Address for correspondence: F. B. Fahlbusch, Department of Pediatrics and Adolescent Medicine, Friedrich-Alexander-University of Erlangen-Nürnberg, Loschgstr. 15, 91054 Erlangen, Germany.  
 (Email: fabian.fahlbusch@uk-erlangen.de)

the NIH National Cancer Institute fielded the Health Information National Trends Survey (HINTS) in 2003:<sup>22</sup> It showed that 63.7% of the average adult U.S. online population had looked for health-related information (for themselves or others) at least once in the previous 12 months.<sup>22</sup> Physicians were the most highly trusted information source to patients, independent of their health literacy level.<sup>23</sup> Nevertheless, 48.6% reported going online first for specific health information, with only 10.9% going to their physicians first.<sup>22</sup> Interestingly, parents also use the internet before their contact with a healthcare specialist and in some cases the internet may be the sole source of healthcare advice.<sup>17,18</sup> Unfortunately, the quality of information available on health-related websites often varies due to the lack of editorial control.<sup>10</sup> This can lead to the presentation of inaccurate or incomplete medical information.<sup>10,24–27</sup> When Scullard *et al.*<sup>28</sup> assessed the reliability of medical advice on the web for common pediatric questions, only 39% of sites searched gave the correct information, 11% were incorrect and 49% failed to answer the question. Where an answer was available, only 78% of sites gave the correct information.<sup>28</sup> Hence we set out to assess the quality of information on postnatal consequences of IUGR/FGR and DOHaD-related consequences on websites directed at laypersons. We based our analysis on Google.com and Yahoo.com search engines to simulate the patient's perspective.<sup>10,28</sup>

## Methods

### *Selection criteria for the webpages*

Websites were considered eligible if they provided any information about IUGR/FGR, such as the etiopathology, outcome and/or maternal or perinatal short- or long-term risks associated with this pregnancy complication. Exclusion criteria were as following:<sup>10</sup> (1) IUGR-related content was limited (i.e. <100 words); (2) website not written in English; (3) complete access restricted by password; (4) repeated server unavailability for 1 week; (5) scientific libraries, book chapters and journals; (6) information mainly on animal studies; (7) promotional websites advertising healthcare institutions without providing detailed information on IUGR; (8) mere reports of personal experiences (e.g. blogs) of single individuals without any additional scientific information; (9) non-written content (e.g. videos).

### *Process of identification and selection of webpages*

The process of identification, selection and evaluation of the webpages took place from January to April 2016. Two different keywords were chosen for our internet research: 'IUGR consequences' and 'FGR consequences.' These were entered into internet search engines Google.com and Yahoo.com, with the country of origin set to Germany. In both search engines the following parameters were selected: English was set as preferred language, the browser history was erased before every search and non-institutional internet access was chosen to simulate a non-professional environment. We based our

analysis on Google.com and Yahoo.com search engines, as they ranked first and third among the most popular engines available in 2016.<sup>29</sup> Google was the most frequently used search engine among parents of attending pediatric outpatients.<sup>30</sup> Google is also among the most commonly used search engines by medical professionals when accessing academic sites.<sup>31</sup> We chose to analyze Yahoo.com over the second ranked engine Bing.com, as Yahoo.com uses Bing.com as search engine, yet additionally functions as a webportal offering links to health-related sites. Sacchetti *et al.*<sup>27</sup> showed that individuals rarely browse through more than 30 sites when looking for online information. Thus, the first three pages of results, representing 30 links, of each internet search engine and each keyword were subjected to further analysis, resulting in a total of 120 links. This approach has also been used by others, previously.<sup>10</sup> Subsequently, duplicate pages were deleted, and the remaining pages were assessed for possible inclusion by one investigator (1st author, S.P.). Websites passing the above exclusion criteria were electronically saved for later analyses by an independent investigator (last author, F.F.). Relevant information from each unique webpage was extracted using the validated tools<sup>32</sup> HONcode (Table 1)<sup>33</sup> and DISCERN Plus score (DPS, Table 2)<sup>34,35</sup> and a checklist for content specifically created for this study (see below). Only the written contents available through screen scrolling of the opening page of each website were assessed, with exception for cases where information on aims and provider of the webpages had to be accessed via a separate link. Written information provided in additional links to websites of other suppliers was not subjected to analysis. From this, categories (sponsored medical news site, medical center/university, interest group, governmental, educational) were chosen to encompass all the types of sites that had arisen. The cited references of each webpage were saved and analyzed for year of publication and overlap with references cited by all other websites examined. The category short-term sequelae of IUGR/FGR relates to the newborn period (first 28 days of life), whereas long-term refers to IUGR/FGR-related sequelae in childhood, adolescence and adulthood.

### *HONcode*

The HONcode agency was founded in 1995 by the Health of the Net Foundation.<sup>33</sup> It is approved by the United Nations and aims to enable patients, web publishers and medical professional to get access to high-value medical information by accrediting websites that publish transparent health-related information.<sup>33</sup> Webpages can apply for accreditation, which involves their examination by a professional committee based on ethical standards. The tool incorporates the assessment of eight principles, which are described in Table 1. HONcode provides a toolbar to internet users that can be added to the browsers toolbar. Whenever a website is accredited by HONcode, the user gets a visual feedback to indicate that quality health information is presented.<sup>33</sup> We utilized the toolbar and recorded whether each website was HONcode

**Table 1.** Health on the Net Foundation (HON) code of conduct for medical and health websites: list of the eight principles and the respective description

The HON code of conduct for medical and health websites (HONcode) <sup>a</sup>	
Principles	Description
1. Authoritative	Indicate the qualifications of the authors
2. Complementarity	Information should support, not replace, the doctor–patient relationship
3. Privacy	Respect the privacy and confidentiality of personal data submitted to the site by the visitor
4. Attribution	Cite the source(s) of published information, date medical and health pages
5. Justifiability	Site must back up claims relating to benefits and performance
6. Transparency	Accessible presentation, accurate email contact
7. Financial disclosure	Identify funding sources
8. Advertising policy	Clearly distinguish advertising from editorial content

<sup>a</sup>Reproduced from the HON foundation. The HON code of conduct for medical and health websites (HONcode)<sup>33</sup>

**Table 2.** List of the 16 modified questions framing the DISCERN Plus tool

Modified DISCERN Plus score	Score
1. Are the aims clear?	1–5
2. Does it achieve its aims?	1–5
3. Is it relevant?	1–5
4. Is it clear what source of information was used to compile the publication (other than the author or producer)?	1–5
5. Is it clear when the information used or reported in the publication was produced?	1–5
6. Is it balanced and unbiased?	1–5
7. Does it provide details of additional sources of support and information?	1–5
8. Does it refer to areas of uncertainty?	1–5
9. Does it describe IUGR/FGR sequelae?	1–5
10. Does it advice to contact a pediatrician for further information on IUGR/FGR-related sequelae?	1–5
11. Does the source provide information on the concept of DOHaD?	1–5
12. Does it explain the current lack of consensus guidelines regarding pediatric monitoring strategies for long-term sequelae?	1–5
13. Does it explain the current lack of consensus guidelines regarding pediatric follow-up intervals for long-term sequelae?	1–5
14. Does it explain the current lack of consensus guidelines regarding pediatric prevention strategies for long-term sequelae?	1–5
15. Does it explain the current lack of consensus guidelines regarding pediatric treatment strategies for long-term sequelae?	1–5
16. Based on the answers to all the above questions, rate the overall quality of the publication as a source of information about IUGR/FGR-related sequelae and DOHaD	1–5
	Total <sup>a</sup>

<sup>a</sup>Maximum of 80.

accredited, as described by others.<sup>34</sup> The frequency of accreditation for each website was recorded.

### DISCERN Plus

DISCERN was funded by The British Library and National Health Service of the United Kingdom and Development Programme.<sup>35</sup> It was generated to offer patients and information providers the possibility to judge the quality of written content on health-related websites. The DISCERN Plus (DP) instrument is a standardized, validated tool that determines the quality of the medical information on a website based on 16 questions.<sup>35</sup> While it was initially designed to aid in the assessment of the reliability and quality of medical treatment options<sup>35</sup> including surgical procedures,<sup>34,36</sup> it has also been

successfully used to analyze health information available on the internet.<sup>34,37</sup> Hence, we modified the DP questions for our analysis (Table 2). The first eight DP questions address the reliability of the website, while the questions 9 and 10 cover the description of IUGR/FGR-related sequelae and the advice to consult a pediatrician for further information. Questions 11–15 specifically address topics associated with DOHaD, that is concept-related information and patient information about the lack of consensus guidelines for monitoring strategies, follow-up intervals, prevention and treatment strategies (Table 2). The last question is an overall rating of the website. As provided by the DISCERN guidelines,<sup>35</sup> the first 15 questions are given a score from 1 to 5 (1 = ‘no’, 2–4 = ‘partially’ and 5 = ‘yes’). The overall quality of the website is then rated on a scale from 1 (poor quality) to 5 (excellent quality),

resulting in a maximum DP-score (DPS) of 80. The modified DP instrument was used by two independent investigators (S.P. and F.F.) to assess the top websites for each search term. DPS discrepancies were discussed and a consensus reached.

### Statistical analyses

Spearman correlation and linear regression analysis were performed using GraphPad Prism 7 (GraphPad Software Inc., La Jolla, CA, USA). Group comparisons were performed using an unpaired Student's *t*-test. A *P*-value of <0.05 was considered significant. Results are mainly presented descriptively using means, standard deviations and percentages. Graphical layout was created with Excel 2010 (Microsoft Corporation, Redmond, WA, USA) and Photoshop CS6 (Adobe Systems Inc., San Jose, CA USA).

### Results

Our search strategy retrieved a total of 120 webpages (two search engines, two key-phrases, 30 websites each), which were further assessed for eligibility. A total of 80 webpages were excluded because they did not fulfill the inclusion criteria. After exclusion of duplicates, 23 websites were selected as eligible for the final analysis, all of which were freely accessible and written in English. The complete list of the 23 URLs is available in Supplementary Table S1. The categories of the webpages were mostly sponsored medical news sites (*n* = 9), medical centers/universities (*n* = 7) and interest groups (*n* = 4) and a limited number were governmental sites (*n* = 2) (Table 3). None of the 23 websites came from a company or individual.

The overall mean DISCERN Plus score (DPS) for the 23 unique webpages was  $41.0 \pm 8.5$  (range 28–60; Table 3) regarding quality of advice, information on postnatal sequelae of IUGR/FGR and DOHaD-related content. The main focus of the analyzed websites was the provision of pregnancy-associated information. While the reliability of information (questions #1–8; Table 2) on IUGR/FGR-related sequelae was generally good (DPS  $3.7 \pm 1.4$ ; Table 3 Fig. 1D), the majority (60.9%; Table 3, Fig. 1Ad) of websites showed poor (DPS 30–43) to very poor (DPS 16–29) overall quality, mainly due to the lack of information on the concept of DOHaD and on missing consensus guidelines (questions #10–15; Table 2, Fig. 1D). In fact, only four webpages gave DOHaD-related information by mentioning the buzzwords 'Barker hypothesis' (*n* = 2), 'fetal programming' (*n* = 1), as well as 'thrifty phenotype' and 'epigenetics' (*n* = 1). Two webpages indicated that IUGR/FGR sequelae require medical follow-up as an option. However, only a single website was specific with regard to monitoring growth and neurodevelopmental outcome and further stated that the occurrence of these IUGR/FGR-related sequelae is subject to great variability, based on which guidelines outlining long-term care are pending. Only one other website discussed the likelihood of developing IUGR/FGR sequelae.

The DISCERN analysis showed that the category 'sponsored medical news site' (Table 3) had a significantly higher DPS than both the categories 'medical center/university' and 'governmental' (*P* < 0.003 and < 0.05, respectively).

Looking at the references cited by the 23 websites, 47.8% (*n* = 11) did not mention any sources of information, with 69.6% citing a maximum of five references (*n* = 16; Fig. 1Aa). Overall 336 references were cited. The mean year of reference publication was  $1997 \pm 10.6$  (Fig. 1C, dotted line, range 1947–2016), with only 12.8% (*n* = 43) of cited references after the publication of the first issue of the *Journal of DOHaD* (Issue S1, November 2009; Fig. 1C dashed line). Interestingly, 45.5% (*n* = 153) of cited references came from a single webpage (Fig. 1 Ab). The majority (*n* = 5, 22.7%) of the remaining webpages cited 1–5 references, 9.1% of websites had a reference list with citation numbers in the range of 6–10, 11–30, 31–70, respectively (*n* = 2, each). There was very little overlap (4.2%) of references (citation of the same sources) in a total of only 13.0% of the 23 webpages (maximum of three references, each). We did not find a significant difference in the number of cited references depending on the category of the webpage (Table 3, data not shown). Interestingly, we found that the number of cited references per webpage significantly correlated with its DPS (linear regression  $R^2$  0.455, *P* < 0.05; Spearman's *r* 0.74, *P* < 0.0001).

The majority (*n* = 18, 78.3%; Fig. 1 Ac) of the 23 webpages did not have a HONcode certification. Of the five HONcode-certified sites (*n* = 5, 21.7%), four came out of the category 'sponsored medical news site' and one was of 'governmental' origin (Table 3). There was no significant difference in DPS ( $46.6 \pm 8.8$  *v.*  $39.4 \pm 7.9$ ) or the number of cited references ( $32.8 \pm 67.3$  *v.*  $9.6 \pm 18.0$ ), when comparing HONcode-certified and uncertified webpages, respectively.

Despite short-comings with regard to the concept of DOHaD and its consequences (Fig. 1 Ad), 78.3% of the analyzed websites mentioned IUGR/FGR-related short- and long-term sequelae (Fig. 1 Ae). This is also reflected by the high DPS ( $3.4 \pm 1.2$ , Fig. 1d) of DISCERN question #9 (Table 2), which addresses this topic. However, only five of the 23 websites gave detailed information on both short- and long-term sequelae. Three websites mentioned neither long- nor short-term sequelae of IUGR/FGR. The categories of short- and long-term consequences as published by the websites are listed in Figure 1Ba and Bb, respectively; 26% (*n* = 6) of the analyzed webpages mentioned the existence of short-term sequelae, including one website that did not give any further related information. In detail, hypoglycemia, hypothermia and hematologic alterations (i.e. polycythemia, hyperviscosity, hyperbilirubinemia, thrombo-/leukopenia) were the top three short-term sequelae of IUGR/FGR (Fig. 1Ba). These ranks were followed by hypoxia/hypoxemia and neurodevelopmental delay (i.e. mostly reduced motor function and cerebral palsy, as well as others, such as behavioral abnormalities, immature sleep patterns, decreased visual fixation, decreased general activity, altered early mother–infant interaction and hyperactivity) (Fig. 1Ba).

**Table 3.** Overview of the quality of information on IUGR/FGR sequelae, DOHaD-related content and respective patient information<sup>a</sup>

<b>Quality of 23 websites according to total DISCERN Plus scores</b>	
Excellent (score 72–80)	0.0% ( <i>n</i> = 0)
Good (score 58–71)	4.3% ( <i>n</i> = 1)
Moderate (score 44–57)	34.8% ( <i>n</i> = 8)
Poor (score 30–43)	52.2% ( <i>n</i> = 12)
Very poor (score 16–29)	8.7% ( <i>n</i> = 2)
Mean overall DISCERN Plus scores (maximum 80)	41.0 ± 8.5
<b>Mean DISCERN Plus scores per types of questions (maximum 5)</b>	
Reliability of information (questions 1–8) <sup>b</sup>	3.7 ± 1.4
IUGR/FGR-related sequelae and pediatric care (questions 9–10)	2.3 ± 1.5
Information on the concept of DOHaD and respective patient information (questions 11–15) <sup>c</sup>	1.1 ± 0.6
Overall quality regarding IUGR/FGR-related sequelae and DOHaD (question 16)	1.3 ± 0.6
<b>Categories of analyzed webpages</b>	
Medical news site (sponsored)	<i>n</i> = 9
HONcode certification	44.4% <sup>d</sup>
Total number of references (range)	30.1 ± 51.5 (0–153)
Total DISCERN Plus score (range)	47.2 ± 6.9 (38–60)
Medical center/university	<i>n</i> = 7
HONcode certification	0.0%
Total number of references (range)	1.0 ± 2.7 (0–7)
Total DISCERN Plus score (range)	35.9 ± 5.2 (29–44)
Interest group	<i>n</i> = 4
HONcode certification	0.0%
Total number of references (range)	8.3 ± 13.4 (0–28)
Total DISCERN Plus score (range)	38.0 ± 10.8 (29–52)
Governmental	<i>n</i> = 2
HONcode certification	50.0%
Total number of references (range)	2.0 ± 1.4 (1–3)
Total DISCERN Plus score (range)	35.0 ± 5.7 (31–39)
Educational	<i>n</i> = 1
HONcode certification	0.0%
Total number of references (range)	21.0 ± 0.0
Total DISCERN Plus score (range)	45.0 ± 0.0
Company	<i>n</i> = 0
Individual	<i>n</i> = 0

<sup>a</sup>All values expressed as percentage or mean ± s.d.

<sup>b</sup>Questions related to aims, relevance for users, sources of information used, date of information, balance, additional sources and areas of uncertainty.

<sup>c</sup>Questions related to the concept of DOHaD and provision of information regarding the lack of consensus guidelines for monitoring, follow-up intervals, prevention and treatment strategies.


<sup>d</sup>Equals 80% of all HONcode certifications.

Furthermore, intestinal (necrotizing enterocolitis, feeding problems), renal immaturity (foremost hypocalcaemia), respiratory problems (i.e. respiratory failure, bronchopulmonary dysplasia), postnatal infections and others (i.e. growth retardation, delay in

dental development, retinopathy of prematurity, reduced body fat) were mentioned less frequently.

Of the websites assessed, 22% (*n* = 5) did not mention long-term sequelae of IUGR/FGR. On the remaining 18 websites,

**Fig. 1.** Quality of advice on postnatal sequelae of intrauterine/fetal growth restriction and the implications of developmental origins of health and disease. (A) Descriptive summary of relevant findings given in percent. (B) Short- and long-term sequelae of IUGR/FGR. The numbers represent the sum of the state of declaration of each consequence from all websites analyzed. Results are presented in the order most-to-least discussed. (C) Display of the year of publication for all cited references of the analyzed webpages. Black dotted line indicates the mean ± s.d., gray dashed line indicates the year of publication of the first *Journal of DOHaD* (Issue S1, November 2009). (D) Results of the modified DISCERN Plus tool (see also Tables 2 and 3) presented as score per question (mean ± s.d.). Minimal score = 1, maximum score = 5. Questions #1–8 relate to the general quality of websites, #9–10 to IUGR/FGR sequelae, #11–15 to DOHaD-related content and respective patient information, #16 represents the total score regarding information about IUGR/FGR-related sequelae and DOHaD.



metabolic syndrome (i.e. hypercholesterolemia, obesity, impaired glucose tolerance and type-2 diabetes, hyperlipidemia), cardiovascular disease (i.e. hypertension, ischemic heart disease) and problems related to longitudinal growth (i.e. short stature) were the most frequent sequelae discussed (Fig. 1Bb). These were followed by neurologic development [i.e. white/gray matter effects, IQ deficits, accelerated maturation of the hippocampus, mental health morbidity, cognitive problems (executive, concentration), clumsiness, cerebral palsy, abnormal deafness, delay of developmental milestones, hyperactivity, schizophrenia] and others (renal, autoimmune thyroid disease) (Fig. 1Bb).

A total of 52% ( $n = 12$ ) of websites name perinatal risk factors of IUGR/FGR neonates. Interestingly, while meconium aspiration was mentioned by 50.0% ( $n = 6$ ) as the only acute perinatal risk of IUGR/FGR, only 16.7% ( $n = 2$ ) of websites gave information on the risk of prematurity, oligohydramnios, higher incidence of pre-labor cesarean delivery and/or induced labor, birth asphyxia, low Apgar score and apnea. The remaining four webpages mentioned increased perinatal mortality rate ( $n = 3$ ), sepsis ( $n = 1$ ), brain damage ( $n = 1$ , i.e. intraventricular hemorrhage, periventricular leukomalacia, hypoxic ischemic encephalopathy, seizures) and possible intrauterine demise ( $n = 1$ ).

## Discussion

In the recent years, we observe a shift in the ways in which patients consume health-related information, with more patients looking for information online before consulting their physicians.<sup>22</sup> Our analysis of current web content regarding the quality of advice on postnatal sequelae of IUGR/FGR showed a significant lack of information on the concept of DOHaD and on related patient information. In cases where respective data is presented, (future) parents are left to decipher a variety of potential IUGR/FGR sequelae due to the lack of further explanation. The majority of the analyzed websites failed to mention the concept of DOHaD and the fact that consensus guidelines addressing aftercare, mitigation of risks or long-term management of children at risk for DOHaD are currently unavailable. This is reflected by the poor ranking of the majority of analyzed websites in our DISCERN Plus rating.

We were surprised to find that the category 'sponsored medical news site' had a significantly higher DPS than both the categories 'medical center/university' and 'governmental' in our DISCERN analysis, while there should be a strong incentive for these traditional institutions of patient education to improve their online services with regard to IUGR/FGR and DOHaD. In line with this finding, the majority of HONcode-certified websites were found in the category 'sponsored medical news site,' with only a minority of HONcode-certified websites present in general. The fact that we did not observe HONcode certification of webpages authored by medical centers and universities in combination with their significantly lower DPS compared to HONcode-certified 'sponsored medical news sites' might indicate that these institutional

webpages did not have the primary goal of patient education, but were rather aimed at informing the public and peers about their services and research interests in general.

Our finding of a significant positive correlation of the number of references cited per webpage and the respective DPS is limited by the fact that the majority of references was published before the beginning of the millennium and 11 of 23 websites did not cite any reference.

In our analysis we excluded scientific articles to mimic a layperson's access to medical information found online. However, there might be further restrictions present that were not included into our analysis. For example, it has to be noted that the amount of information gained might be limited by the finding that 55% of current internet users spends fewer than 15 seconds on a page, indicating the importance of webpage design to facilitate the access to data.<sup>39</sup>

Furthermore, we did not determine the quality of webpages with regard to the average reader level. It has been shown that only 1–2% were aimed at the recommended reading level,<sup>40</sup> taking into account that 16% of adults in the United Kingdom have a reading level equivalent to an 11-year old (equals 6th Grade in the U.S. school system).<sup>28</sup> Therefore, even if correct information is presented, it might not be understood or might be misinterpreted. In this respect, similar results were obtained from the national assessment of adult literacy in 2003 by the Institute of Education Sciences at the U.S. Department of Education.<sup>23</sup> It was shown that only 12% of the U.S. adults had proficient health literacy. Over 1/3 of the U.S. adults had difficulties completing common health tasks, such as following directions on a prescription drug label or adhering to a childhood immunization schedule. This limitation was prevalent among all racial and ethnic groups, as well as education levels, however, was more pronounced among non-Caucasian minorities and adults without high school education.<sup>23</sup> Noteworthy, the assessment showed that all adults, regardless of their health literacy skills, were more likely to get health information from radio/television, friends/family and health professionals, than from print media.<sup>23</sup> Adults with the most limited health literacy rarely use digital resources and written material to obtain information on health topics in this study.<sup>23</sup> Thus, while our analysis was tailored to encompass the layperson's view, it can be conceived that the general uptake of information related to IUGR/FGR and DOHaD-related consequences might be limited, especially in adults with reduced health literacy and low socioeconomic status. Unfortunately, it is known that this sub-population has the highest IUGR/FGR-rate in particular,<sup>41,42</sup> which argues for physician-based medical education of this group.

Based on our findings, we currently cannot encourage healthcare workers to recommend their patients the use of websites to research the consequences of IUGR/FGR due to the lack of information on DOHaD and the related consequences. This negative assessment is also the consequence of (i) generalization of IUGR/FGR as risk factor of a limited number of subjectively selected sequelae independent of a clear definition of the pregnancy complication, (ii) lack of consensus guidelines for IUGR/FGR aftercare and (iii) the aged data

presented online, which also includes websites run by the government or national health services, usually offering valuable medical information in other fields.<sup>28</sup>

It has to be noted, that overall IUGR/FGR-related sequelae were reported by a proportion of the analyzed websites, thereby contributing to the awareness of the relevance of the condition. Hence, it is hoped that the increasing use of the internet in healthcare will offer new opportunities for patient information in the future. Certain limitations might, however, apply regarding online medical education for laypersons with low socioeconomic status and reduced health literacy. As respective consensus guidelines are currently lacking, DOHaD-related patient education and follow-up remains within the physician's individual responsibility for now.

### Acknowledgments

Data acquisition and analysis was performed by Sandra Perzel in fulfillment of the requirements for obtaining the degree 'Dr. med.' at the Friedrich-Alexander-University of Erlangen-Nuremberg, Dept. of Pediatrics and Adolescent Medicine, Germany.

### Supplementary material

To view supplementary material for this article, please visit <https://doi.org/10.1017/S2040174417000332>

### References

- Barker DJ, Osmond C. Infant mortality, childhood nutrition, and ischaemic heart disease in England and Wales. *Lancet*. 1986; 1, 1077–1081.
- Wadhwa PD, Buss C, Entringer S, Swanson JM. Developmental origins of health and disease: brief history of the approach and current focus on epigenetic mechanisms. *Semin Reprod Med*. 2009; 27, 358–368.
- Cosmi E, Fanelli T, Visentin S, Trevisanuto D, Zanardo V. Consequences in infants that were intrauterine growth restricted. *J Pregnancy*. 2011; 2011, 364381.
- Williams TC, Drake AJ. What a general paediatrician needs to know about early life programming. *Arch Dis Child*. 2015; 100, 1058–1063.
- Barker DJ. The origins of the developmental origins theory. *J Intern Med*. 2007; 261, 412–417.
- Godfrey KM, Costello PM, Lillycrop KA. The developmental environment, epigenetic biomarkers and long-term health. *J Dev Orig Health Dis*. 2015; 6, 399–406.
- Godfrey KM, Inskip HM, Hanson MA. The long-term effects of prenatal development on growth and metabolism. *Semin Reprod Med*. 2011; 29, 257–265.
- Mandy G, Weisman L, Kim M. Infants with fetal (intrauterine) growth restriction. Retrieved 6 December 2016 from <https://www.uptodate.com/contents/infants-with-fetal-intrauterine-growth-restriction#H440406701>
- Doetsch J, Zepp F. Fetale Programmierung und frühkindliche Prägung. *Monatsschr Kinderheilkd*. 2016; 164, 89–90.
- Fioretti BT, Reiter M, Betran AP, Torloni MR. Googling caesarean section: a survey on the quality of the information available on the Internet. *BJOG*. 2015; 122, 731–739.
- Handfield B, Turnbull S, Bell RJ. What do obstetricians think about media influences on their patients? *Aust N Z J Obstet Gynaecol*. 2006; 46, 379–383.
- Lagan B, Sinclair M, Kernohan WG. Pregnancy problems: answers on the internet? *RCM Midwives*. 2007; 10, 276–278.
- Sayakhot P, Carolan-Olah M. Internet use by pregnant women seeking pregnancy-related information: a systematic review. *BMC Pregnancy Childbirth*. 2016; 16, 65.
- Moyer CA, Vishnu LO, Sonnad SS. Providing health information to women. The role of magazines. *Int J Technol Assess Health Care*. 2001; 17, 137–145.
- Lagan BM, Sinclair M, Kernohan WG. What is the impact of the Internet on decision-making in pregnancy? A global study. *Birth*. 2011; 38, 336–345.
- Wainstein BK, Sterling-Levis K, Baker SA, Taitz J, Brydon M. Use of the internet by parents of paediatric patients. *J Paediatr Child Health*. 2006; 42, 528–532.
- Khoo K, Bolt P, Babl FE, Jury S, Goldman RD. Health information seeking by parents in the Internet age. *J Paediatr Child Health*. 2008; 44, 419–423.
- Tuffrey C, Finlay F. Use of the internet by parents of paediatric outpatients. *Arch Dis Child*. 2002; 87, 534–536.
- Ofcom Adults' Media Use and Attitudes Report 2015. Retrieved 30 November 2016 from [https://www.ofcom.org.uk/\\_\\_data/assets/pdf\\_file/0014/82112/2015\\_adults\\_media\\_use\\_and\\_attitudes\\_report.pdf](https://www.ofcom.org.uk/__data/assets/pdf_file/0014/82112/2015_adults_media_use_and_attitudes_report.pdf).
- Office for National Statistics. Statistical bulletin: internet users in the UK, 2016. Retrieved 30 November 2016 from <https://www.ons.gov.uk/businessindustryandtrade/itandinternetindustry/bulletins/internetusers/2016>.
- Rainey L. Internet, broadband and cell phone statistics: the Pew Internet and American Life Center, 2010. Retrieved 14 February 2017 from <http://www.pewinternet.org/2010/01/05/internet-broadband-and-cell-phone-statistics/>.
- Hesse BW, Nelson DE, Kreps GL, et al. Trust and sources of health information: the impact of the internet and its implications for health care providers: findings from the first Health Information National Trends Survey. *Arch Intern Med*. 2005; 165, 2618–2624.
- U.S. Department of Education, Institute of Education Sciences 2003; National Assessment of Adult Literacy. America's Health Literacy: why we need accessible health information. Retrieved 14 February 2017 from <https://health.gov/communication/literacy/issuebrief/>.
- Kaicker J, Debono VB, Dang W, Buckley N, Thabane L. Assessment of the quality and variability of health information on chronic pain websites using the DISCERN instrument. *BMC Med*. 2010; 8, 59.
- Cline RJ, Haynes KM. Consumer health information seeking on the Internet: the state of the art. *Health Educ Res*. 2001; 16, 671–692.
- Hargrave D, Bartels U, Lau L, Esquembre C, Bouffet E. Quality of childhood brain tumour information on the internet in French language. *Bull Cancer*. 2003; 90, 650–655.
- Sacchetti P, Zvara P, Plante MK. The internet and patient education – resources and their reliability: focus on a select urologic topic. *Urology*. 1999; 53, 1117–1120.



28. Scullard P, Peacock C, Davies P. Googling children's health: reliability of medical advice on the internet. *Arch Dis Child*. 2010; 95, 580–582.
29. eBizMBA Inc. Top 15 Most Popular Search Engines, 2016. Retrieved 30 November 2016 from <http://www.ebizmba.com/articles/search-engines>.
30. Sim NZ, Kitteringham L, Spitz L, et al. Information on the world wide web – how useful is it for parents? *J Pediatr Surg*. 2007; 42, 305–312.
31. Giustini D. How Google is changing medicine. *BMJ*. 2005; 331, 1487–1488.
32. Gagliardi A, Jadad AR. Examination of instruments used to rate quality of health information on the internet: chronicle of a voyage with an unclear destination. *BMJ*. 2002; 324, 569–573.
33. Health on the Net Foundation. The HON code of conduct for medical and health websites (HONcode). Retrieved 30 November 2016 from <http://www.hon.ch/HONcode/Conduct.html>
34. Fast AM, Deibert CM, Hruba GW, Glassberg KI. Evaluating the quality of Internet health resources in pediatric urology. *J Pediatr Urol*. 2013; 9, 151–156.
35. Charnock D, Shepperd S, Needham G, Gann R. DISCERN: an instrument for judging the quality of written consumer health information on treatment choices. *J Epidemiol Commun Health*. 1999; 53, 105–111.
36. Wasserman M, Baxter NN, Rosen B, Burnstein M, Halverson AL. Systematic review of internet patient information on colorectal cancer surgery. *Dis Colon Rectum*. 2014; 57, 64–69.
37. Charnock D, Shepperd S. Learning to DISCERN online: applying an appraisal tool to health websites in a workshop setting. *Health Educ Res*. 2004; 19, 440–446.
38. Haile T. What you think you know about the web is wrong, 2014. Retrieved 30 November 2016 from <http://time.com/12933/what-you-think-you-know-about-the-web-is-wrong/>
39. Gurel Koybasi NA, Cagiltay K. Finding online health-related information: usability issues of health portals. *Stud Health Technol Inform*. 2012; 180, 848–852.
40. Hargrave DR, Hargrave UA, Bouffet E. Quality of health information on the Internet in pediatric neuro-oncology. *Neuro Oncol*. 2006; 8, 175–182.
41. Kramer MS. Socioeconomic determinants of intrauterine growth retardation. *Eur J Clin Nutr*. 1998; 52(Suppl. 1), S29–S32, discussion S32–S33.
42. Hendrix N, Berghella V. Non-placental causes of intrauterine growth restriction. *Semin Perinatol*. 2008; 32, 161–165.