

Glucose deprivation activates a cAMP-independent protein kinase from *Trypanosoma equiperdum*

Research Article

Cite this article: Guevara A, Lugo C, Montilla AJ, Araujo NA, Calabokis M, Bubis J (2019). Glucose deprivation activates a cAMP-independent protein kinase from *Trypanosoma equiperdum*. *Parasitology* **146**, 643–652. <https://doi.org/10.1017/S0031182018001920>

Received: 13 August 2017
Revised: 12 October 2018
Accepted: 16 October 2018
First published online: 13 November 2018

Key words:

Dourine; glucose deprivation; kemptide kinase activity; protein kinases; protein phosphorylation; signal transduction; *Trypanosoma equiperdum*

Author for correspondence:

José Bubis, E-mail: jbubis@usb.ve

Alberto Guevara^{1,2}, Cristina Lugo^{1,2}, Alejandro J. Montilla^{1,2}, Nelson A. Araujo¹, Maritza Calabokis¹ and José Bubis¹

¹Departamento de Biología Celular, Universidad Simón Bolívar, Caracas, Venezuela and ²Postgrado en Ciencias Biológicas, Universidad Simón Bolívar, Caracas, Venezuela

Abstract

Kemptide (sequence: LRRASLG) is a synthetic peptide holding the consensus recognition site for the catalytic subunit of the cAMP-dependent protein kinase (PKA). cAMP-independent protein kinases that phosphorylate kemptide were stimulated in *Trypanosoma equiperdum* following glucose deprivation. An enriched kemptide kinase-containing fraction was isolated from glucose-starved parasites using sedimentation throughout a sucrose gradient, followed by sequential chromatography on diethylaminoethyl-Sepharose and Sephacryl S-300. The trypanosome protein possesses a molecular mass of 39.07–51.73 kDa, a Stokes radius of 27.4 Å, a sedimentation coefficient of 4.06 S and a globular shape with a frictional ratio $ff_0 = 1.22$ –1.25. Optimal enzymatic activity was achieved at 37 °C and pH 8.0, and kinetic studies showed K_m values for ATP and kemptide of 11.8 ± 4.1 and $24.7 \pm 3.8 \mu\text{M}$, respectively. The parasite enzyme uses ATP and Mg^{2+} and was inhibited by other nucleotides and/or analogues of ATP, such as cAMP, AMP, ADP, GMP, GDP, GTP, CTP, β, γ -imidoadenosine 5'-triphosphate and 5'-[*p*-(fluorosulfonyl)benzoyl] adenosine, and by other divalent cations, such as Zn^{2+} , Mn^{2+} , Co^{2+} , Cu^{2+} , Ca^{2+} and Fe^{2+} . Additionally, the trypanosome kinase was inhibited by the PKA-specific heat-stable peptide inhibitor PKI- α . This study is the first biochemical and enzymatic characterization of a protein kinase from *T. equiperdum*.

Introduction

Dourine or covering sickness is a sexually transmitted infection caused by *Trypanosoma equiperdum*, a haemoparasite that belongs to the Trypanozoon subgenus of Salivarian trypanosomes (Brun *et al.*, 1998; Stevens and Brisse, 2004). This disease is characterized by genital inflammation, skin plaques and neurological signs, and mainly affects horses and other equids (Gizaw *et al.*, 2017).

The reversible phosphorylation of proteins is a ubiquitous mechanism of regulation that is vital for all living cells. In trypanosomatids, approximately 2% of their genome is composed of genes that encode for protein kinases, suggesting that protein phosphorylation plays a key role in the biology of these parasites (Naula *et al.*, 2005; Parsons *et al.*, 2005). In the genomes of *Trypanosoma brucei*, *Trypanosoma cruzi*, *Leishmania major* and *Trypanosoma evansi*, there are genes that encode for three AGC protein kinases related to the catalytic subunits of the mammalian cAMP-dependent protein kinase (*aka* protein kinase A or PKA), and a gene homologous to the gene for the mammalian PKA regulatory subunits (<http://tritrypdb.org>). It has been suggested that these PKA catalytic subunit-like protein kinases might be involved in crucial cellular processes in trypanosomatids, such as osmoregulation, DNA repair, resistance to oxidative stress, activation of metabolic enzymes, replication, growth and cellular motility (Genestra *et al.*, 2004; Parsons *et al.*, 2005; Bao *et al.*, 2008; Malki-Feldman and Jaffe, 2009). In this work, we partially purified and biochemically characterized a *T. equiperdum* protein kinase that recognizes fluorescently modified kemptide, which is a synthetic phosphate acceptor heptapeptide (sequence: LRRASLG) that resembles the local phosphorylation site sequence in pig liver pyruvate kinase and contains the consensus phosphorylation site for the PKA catalytic subunit (Kemp *et al.*, 1975; Maller *et al.*, 1978; Kemp, 1980). Therefore, kemptide serves as a specific peptide substrate for PKA and PKA-like enzymes. Moreover, the parasite enzyme was capable of phosphorylating histone type II-AS, the synthetic peptide SP20 (sequence: TTYADFIASGRTGRRNSIHD), and the α isoform of the PKA type II regulatory subunit ($\text{R}^{\text{II}}\alpha$ -subunit), which are also known to function as substrates for the PKA catalytic subunit. The ATP:phosphotransferase activity of the *T. equiperdum* kemptide kinase was inhibited by both the PKA-specific heat-stable peptide inhibitor PKI- α and the synthetic peptide IP20 (sequence: TTYADFIASGRTGRRNAIHD), which is derived from PKI- α . However, the trypanosome kinase activity was not dependent on cAMP. Interestingly, the activity of this parasite enzyme was triggered upon nutritional stress driven by glucose starvation.

Materials and methods

Parasites

Cryopreserved *T. equiperdum* parasites from the Venezuelan TeAp-N/D1 strain were inoculated into Sprague–Dawley adult albino rats. The TeAp-N/D1 strain has been shown to be closely related to the *T. equiperdum* STIB841/OVI strain (Sánchez *et al.*, 2015). When more than 100 trypanosomes per microscopic field were detected in blood smears, the infected anaesthetized rats were bled *via* heart puncture using 2% EDTA (pH 8.0) as anticoagulant. Trypanosomes were purified by ion-exchange chromatography using a fibrous diethylaminoethyl (DEAE)-cellulose column equilibrated with phosphate-buffered saline (PBS) containing glucose (PBS-G) [57 mM Na₂HPO₄, 3 mM NaH₂PO₄, 43.8 mM NaCl and 1% glucose (pH 8.0)] (Lanham and Godfrey, 1970). Parasites eluting from the column were collected by centrifugation at 1475 g, for 10 min and washed three times with PBS-G buffer. The number of parasites was measured using a haemocytometer or Neubauer chamber.

Glucose deprivation of *T. equiperdum* parasites

Freshly purified parasites were incubated in the absence of glucose with PBS [57 mM Na₂HPO₄, 3 mM NaH₂PO₄ and 43.8 mM NaCl (pH 8.0)], for various times (0, 15, 30 and 60 min), at room temperature, under gentle and constant agitation. A sample of parasites that were deprived of glucose for 60 min was reversed by further incubation with glucose using the PBS-G buffer for 30 min. In all cases, trypanosomes were centrifuged at 1475 g, for 5 min, at 4 °C and the sedimented cells were stored at –80 °C until further use.

Kemptide kinase activity using an electrophoretic gel-shift assay

Kemptide kinase activity was measured using the Pep-Tag® non-radioactive method (Promega, Madison, WI, USA) with the fluorescently labelled kemptide (PepTag® A1 Peptide-LRRASLG) as substrate. Assays were performed as instructed by the manufacturer, and the reaction mixtures were separated by electrophoresis on a 1.2% agarose gel. When the commercial PepTag® A1 Peptide-LRRASLG component had run out of the kit, the rest of the experiments were performed with either the fluram-kemptide or the fluram-kemptide-Lys8 peptide (Araujo *et al.*, 2016) at a final concentration of 0.25 mg mL⁻¹. Both kemptide and the kemptide-Lys8 peptide were chemically synthesized using the *t*-Boc methodology and were labelled with fluorescamine (fluram) as previously described (Araujo *et al.*, 2016). No preference for any of these fluorescently modified peptides was observed since all contained the specific substrate recognition site for the PKA catalytic subunit (sequence: RRXS/TΨ, where X is any amino acid and Ψ is a hydrophobic residue). Gels were revealed in a phototransilluminator (Fotodyne, Hartland, WI, USA) with UV light. The PKA catalytic subunit was purified from pig heart (Nelson and Taylor, 1981) and used as a positive control.

Kemptide kinase activity using a high-performance liquid chromatography assay

ATP:phosphotransferase activity was also determined by using a high-performance liquid chromatography (HPLC) assay on a reversed-phase octadecylsilane column (Kemp, 1980; Luzi *et al.*, 2017). HPLC conditions were as described by Luzi *et al.* (2017), but the reaction was modified by employing fluram-kemptide as substrate. Phosphorylation reactions were initiated by incorporating the *T. equiperdum* protein kinase-containing fraction, and

were terminated by the addition of 5% phosphoric acid. HPLC separations were carried out on a Waters 1525 binary pump system with a Symmetry® C₁₈ column (3.9 × 150 mm, 5 μm, Waters, Milford, MA, USA), and the absorbance was monitored on a Waters 2487 dual wavelength detector. Since fluram-kemptide possesses a maximum excitation peak at ~400 nm (Araujo *et al.*, 2016), absorbance was measured at both 210 and 400 nm. Progress of the reactions was quantified by integrating the corresponding area peaks of the dephosphorylated substrate at 400 nm. The assay was initially standardized using the purified porcine heart PKA catalytic subunit (data not shown).

Partial purification of the *T. equiperdum* kemptide kinase

Freshly purified parasites (~10⁹–10¹⁰ trypanosomes) were incubated in the absence of glucose for 1 h, as described above. Cells were extracted with lysis buffer [50 mM Tris-HCl (pH 8.0), 1 mM CaCl₂, 1 mM MgCl₂, 50 μM phenyl methyl sulfonyl fluoride (PMSF), 1 mM benzamidine, and 10 μM L-trans-epoxysuccinyl-leucylamido(4-guanidino)butane (E-64)] by sonication, using four cycles of 30 s each, with resting intervals of 2 min per cycle. The homogenate was centrifuged at 100 000 g for 1 h, at 4 °C, in order to obtain the corresponding parasite soluble fraction, which was loaded on a linear 5–20% sucrose gradient prepared in 50 mM Tris-HCl (pH 7.0), 0.1 mM EDTA, 0.15 M NH₄Cl and 0.2 mM dithiothreitol. The sample was centrifuged at 100 000 g, for 20 h, at 5 °C. Fractions were collected from the top of the tube, and a sample of each fraction was assayed for kemptide kinase activity. Three peaks of enzymatic activity were obtained (see 'Results' section).

Fractions corresponding to peak 1 of the kemptide kinase activity were pooled and loaded on a 60 ml DEAE-Sepharose column that was previously equilibrated with 50 mM Tris-HCl (pH 7.0), 0.1 mM EDTA and 0.1 mM EGTA. Proteins were eluted from the column using a 0.01–1.0 M NaCl gradient in the same buffer containing 50 μM PMSF, 1 mM benzamidine and 10 μM E-64.

Fractions containing the kemptide kinase activity peak from the DEAE-Sepharose column were pooled and separated through a 90 mL Sephacryl S-300 gel-filtration column using 50 mM Tris-HCl (pH 7.5), 150 mM NaCl, 3 mM Mg-acetate, 2 mM CaCl₂, 5 mM β-mercaptoethanol, 50 μM PMSF, 1 mM benzamidine and 10 μM E-64 as elution buffer.

Effect of glucose on the *T. equiperdum* kemptide kinase enzymatic activities

Fresh purified *T. equiperdum* parasites were incubated in the absence of glucose with PBS, for 60 min, at room temperature. Glucose-deprived parasites were then incubated for 10 min with various concentrations of glucose (0–10 mM). Following homogenization of the parasite cells with lysis buffer, the resulting whole-cell extracts were centrifuged, and kemptide kinase activity was measured on the corresponding soluble fractions by HPLC (Luzi *et al.*, 2017).

Kemptide kinase activity was also determined on the partially purified trypanosome enzyme after incubation with different concentrations of glucose (0–1 mM).

Evaluation of potential substrates for the partially purified parasite protein kinase

The electrophoretic gel-shift assay was used to analyse whether the *T. equiperdum* kemptide kinase was capable of phosphorylating histone type II-AS from calf thymus (Sigma, Saint Louis, MO, USA) and a custom synthesized 20-residue peptide, SP20 (Life Tein, Somerset, NJ, USA), which is derived from the

PKA-specific inhibitor PKI- α and possesses the following sequence: TTYADFIASGRTGRRNSIHD. In SP20, the Ala residue of PKI- α is substituted by a Ser at position 17. Kinase reactions were carried out as described by Lutz *et al.* (1994) using either 0.4 mg mL⁻¹ of histone II-AS or 0.25 mg mL⁻¹ of SP20.

Since the mammalian PKA R^{II} α -subunit contains a site that is autophosphorylated by the PKA catalytic subunit, the R^{II} α -subunit was purified from porcine heart (Taylor and Stafford, 1978; Nelson and Taylor, 1983) in order to explore its phosphorylation by the trypanosome protein kinase. Phosphorylation of the R^{II} α -subunit was performed as previously described (First *et al.*, 1988), with slight modifications. Briefly, purified R^{II} α -subunit from pig heart (5 μ g) was incubated for 30 min, at room temperature, with an aliquot (15 μ L) of the partially purified *T. equiperdum* kemptide kinase in the presence of 50 mM Tris-HCl (pH 7.5) containing 1 mM cAMP, 10 mM MgCl₂ and 1 mM ATP. A control experiment was carried out in parallel using 1 μ g of the purified PKA catalytic subunit from pig heart. Reactions were terminated by adding sodium dodecyl sulphate-polyacrylamide gel electrophoresis (SDS-PAGE) sample buffer (Laemmli, 1970), and boiling the samples for 5 min. Then, the reaction mixtures were separated by SDS-PAGE on an 8% polyacrylamide slab gel and electrotransferred to nitrocellulose sheets (Towbin *et al.*, 1979). Membranes were first incubated with rabbit polyclonal anti-PKA R^{II} α -subunit antibodies (Santa Cruz Biotechnology Dallas, TX, USA, dilution 1:5000), and then with alkaline phosphatase-conjugated antibodies against rabbit IgG (Jackson ImmunoResearch Laboratories, Inc., West Grove, PA, USA, dilution 1:1000). Polypeptide bands were visualized by the addition of 5-bromo-4-chloro-3-indolyl phosphate and nitro blue tetrazolium according to the supplier (Thermo Scientific Pierce, Waltham, MA, USA).

Biochemical characterization of the partially purified *T. equiperdum* kemptide kinase activity

Kemptide kinase activity was evaluated in the presence of 10 mM MgCl₂ and 1 mM ATP at different temperatures (-20 to 60 °C) and pH (2.0–11.0), to establish the optimal conditions for enzymatic activity. The optimal pH was determined by using suitable buffers (100 mM), which were selected based on their pKa: maleate (pH 2.0), glycine (pH 3.0), formate (pH 4.0), acetate (pH 5.0), 2-(N-morpholino)ethanesulfonic acid (pH 6.0), [3-(N-morpholino)propanesulfonic acid] (pH 7.0), Tris (pH 8.0) and N-cyclohexyl-3-aminopropanesulfonic acid (pH 9.0–11.0). In order to confirm that the effect on the enzymatic activity was produced by changes in the pH rather than to variations in the buffer system, we used a series of buffers (100 mM) with overlapping effective pH ranges, namely citrate (pH 2.0, 3.0, 4.0, 5.0 and 6.0), phosphate (pH 6.0, 7.0 and 8.0), 4-(2-hydroxyethyl)-1-piperazineethanesulfonic acid (HEPES, pH 7.0 and 8.0), Tris (pH 7.0, 8.0 and 9.0) and carbonate (pH 8.5, 9.0, 10.0 and 11.0).

The effect on the trypanosome kinase activity of cAMP and the recombinant isoform α of the PKA heat-stable inhibitor PKI from rabbit (PKI- α , Calbiochem, San Diego, CA, USA) was also evaluated. We choose two concentrations of cAMP, 5 μ M that lied in the expected physiological range, and 5 mM that was extremely high and non-physiological. The inhibitory effect of PKI- α was measured by using 0.9, 9 and 36 units of the PKA-specific inhibitor peptide. The inhibitory effect of PKI- α was corroborated by using a custom-synthesized 20-amino acid peptide, IP20 (sequence: TTYADFIASGRTGRRNAIHD, Life Tein, Somerset, NJ, USA), which is derived from PKI- α and contains its inhibitory motif. The partially purified protein kinase from *T. equiperdum* was pre-incubated with IP20 (0.25 mg mL⁻¹) in 50 mM Tris-HCl (pH 7.5), for 30 min at 4 °

C, and then the remainder components of the reaction mixture were added (0.25 mg mL⁻¹ fluram-kemptide, 10 mM MgCl₂ and 1 mM ATP). Kemptide kinase activity was monitored by using the electrophoretic gel-shift assay.

The specificity of nucleotide triphosphates and divalent cations was monitored by evaluating the enzymatic activity using 1 mM ATP, GTP or CTP (in the presence of 10 mM MgCl₂); and 10 mM MgCl₂, ZnCl₂, MnCl₂, CoCl₂, CuCl, CaCl₂ or FeSO₄ (in the presence of 1 mM ATP), respectively.

The inhibitory effect of AMP, ADP, GMP, GDP, GTP, CTP, β , γ -imido-adenosine 5'-triphosphate (AMP-PNP) and 5'-[p-(fluoro sulfonyl)benzoyl] adenosine (FSBA) on the parasite enzymatic activity was measured by pre-incubating the partially purified kemptide kinase with 2 mM of each compound for 30 min, under ice, in the presence of 10 mM MgCl₂. Subsequently, the kinase assay was performed as described above following incubation with 1 mM ATP for 1 h.

To determine the inhibitory effect of the different divalent cations on the parasite enzymatic activity, the partially purified trypanosome kemptide kinase was pre-incubated with 20 mM ZnCl₂, MnCl₂, CoCl₂, CuCl, CaCl₂ or FeSO₄, for 30 min, under ice, in the presence of 1 mM ATP. Then, the kinase assay was assayed as described above following incubation with 10 mM MgCl₂ for 1 h.

Determination of the K_m for ATP and kemptide of the *T. equiperdum* kemptide kinase

The K_m values for ATP and kemptide of the partially purified parasite enzyme were determined by HPLC (Luzi *et al.*, 2017). To measure the K_m for ATP, the reactions were performed in 50 mM Tris-HCl (pH 7.4), 10 mM MgCl₂, 30 μ M fluram-kemptide and various concentrations of ATP (0–70 μ M). To measure the K_m for kemptide, the mixtures contained 50 mM Tris-HCl (pH 7.4), 10 mM MgCl₂, 1 mM ATP and various concentrations of fluram-kemptide (0–50 μ M). Three independent experiments were performed for each concentration, and K_m values were estimated by regression analysis of double reciprocal plots (Lineweaver and Burk, 1934). Values were reported as the mean \pm the standard error of the mean (S.E.M.).

SDS-PAGE

Fractions obtained from all purification steps were separated by SDS-PAGE according to Laemmli (1970), and revealed using silver staining.

Results

Activation of kemptide kinase activities by glucose deprivation of *T. equiperdum* parasites

By using the electrophoretic gel-shift assay, no kemptide kinase activities were detected in whole-cell homogenates, particulate fractions or clarified soluble fractions from trypanosomes purified in the presence of 1% glucose, which corresponds to 55.5 mM (Fig. 1A). Equivalent results were obtained in the absence or presence of either 5 μ M or 5 mM cAMP. However, an increase in kemptide phosphorylation was achieved in homogenates and soluble fractions from parasites that underwent glucose limitation for 15, 30 and 60 min (Fig. 1B). No kemptide kinase activities were observed in the particulate fraction from glucose-starved *T. equiperdum* microbes revealing that when activated, these enzymatic activities were exclusively localized in the corresponding soluble fraction (Fig. 1B). Similar results were found by the HPLC assay when kemptide kinase activities were determined on the clarified




Fig. 1. Effect of glucose deprivation on the parasite kemptide kinase activities by using the electrophoretic gel-shift assay. (A) No kinase activities were detected when trypanosomes were purified in the presence of glucose. *T. equiperdum* parasites were purified through a DEAE-cellulose column using PBS-G (PBS containing 1% glucose), and the purified trypanosomes were immediately used (0 min). Enzymatic activities were qualitatively evaluated on the parasite whole-cell homogenate (H), particulate fraction (P) and clarified soluble fraction (S) by means of the Promega Pep-Tag® non-radioactive assay, using the PepTag® A1 kemptide as substrate. Determinations were made in the absence (–) or presence of either 5 μ M or 5 mM cAMP (+). (B) Glucose starvation activates kemptide kinase activities from *T. equiperdum*. Freshly purified parasites were incubated with PBS lacking glucose for 15, 30 and 60 min (GD = glucose deprivation). Also, a sample of the parasites deprived of glucose for 60 min was re-incubated with PBS-G for 30 min (60:30). Kemptide kinase activities were estimated on the homogenates (H), particulate fractions (P) and clarified soluble fractions (S) of all trypanosome samples by using the Promega Pep-Tag® non-radioactive assay. Measurements were made in the absence (–) or presence of 5 μ M cAMP (+). In (A) and (B), gels were revealed with UV light in a phototransilluminator. The arrowhead and arrow indicate the non-phosphorylated and phosphorylated fluorescently labelled peptide, respectively. Purified PKA catalytic subunit from porcine heart was employed as a positive control (C+). The reaction mixture without any added enzyme was used as a negative control (C–).

soluble fractions from glucose-deprived *T. equiperdum* parasites (Fig. 2A). Moreover, the activation of kemptide kinase activities elicited in glucose-deprived parasites was readily reversed, and these enzymes were inhibited upon re-addition of 1% glucose for 30 min to parasites that experienced glucose fasting for 60 min (Fig. 1B, 60:30). As also shown in Fig. 1B, no stimulation by exogenously added cAMP (5 μ M) was obtained in these experiments. Similar results were found when 5 mM cAMP was used (data not shown). If anything, a decrease in the phosphorylation signal of the fluorescently labelled kemptide substrate was observed when cAMP was present, suggesting that cAMP might be inhibiting the parasite kemptide kinase activities. Although the activity of the homogenate sample at the 60 min time point appears to be slightly higher upon addition of cAMP than in the absence of cAMP (Fig. 1B), we feel that this particular result is an artefact produced by the qualitative features of the electrophoretic gel-shift assay. Absence of stimulation by cAMP was consistent in many independent experiments that were carried out. On the basis of how PKA enzymes are commonly defined, these results established that the parasite kemptide kinase activities represent cAMP-independent enzymes.




Fig. 2. Effect of glucose deprivation on the *T. equiperdum* kemptide kinase activities by using the HPLC assay. (A) Time course of the effect of glucose fasting. Freshly purified parasites were incubated in the absence of glucose with PBS for various times (0, 15, 30 and 60 min). Parasites were then extracted with lysis buffer and the homogenates were centrifuged to obtain the corresponding clarified soluble fractions. Kemptide kinase activities were assayed by HPLC. (B) Incubation of *T. equiperdum* parasites with different concentrations of glucose. Freshly purified trypanosomes were deprived of glucose for 60 min, and then were incubated for 10 min with increasing concentrations of glucose (0–10 mM). Following homogenization and centrifugation, kemptide kinase activities were evaluated on the corresponding soluble fractions by HPLC. (C) Incubation of the partially purified kemptide kinase from *T. equiperdum* with different concentrations of glucose. The enriched kemptide kinase-containing fraction that was obtained as described in Fig. 3 was incubated with various concentrations of glucose (0–1 mM). Kemptide kinase activities were measured by HPLC. In all cases, values were reported as the mean \pm S.E.M.

Nutritionally stressed *T. equiperdum* parasites were incubated with increasing amounts of glucose in order to determine its half maximal inhibitory concentration (IC_{50}). Following homogenization and centrifugation, kemptide kinase activities were evaluated on the corresponding soluble fractions by HPLC. As seen in Fig. 2B, a concentration of ~ 10 μ M glucose was required for a 50% inactivation of these enzymes. These results showed that the inhibition by glucose of the parasite kinase activities was concentration dependent.

Partial purification of a *T. equiperdum* kemptide kinase activity

Freshly isolated *T. equiperdum* parasites were incubated in the absence of glucose for 60 min in order to stimulate the




Fig. 3. Partial purification of a protein kemptide kinase from *T. equiperdum*. (A) Sedimentation by ultracentrifugation on a sucrose gradient. The soluble fraction from glucose-starved *T. equiperdum* was subjected to sedimentation by ultracentrifugation on a 5–20% sucrose gradient. Fractions were collected from the top of the tube and assayed for kinase activity employing the fluram-kemptide-Lys8 peptide as a substrate. Purified PKA catalytic subunit from pig heart was used as a positive control (C+). The reaction mixture without any added enzyme was used as a negative control (C–). (B) Separation by anion-exchange chromatography. Fractions 4–10 from the sucrose gradient were pooled and separated on a DEAE-Sepharose column. The chromatogram (top) illustrates the protein profile (absorbance at 280 nm) and the ionic strength of the eluting fractions, which were assayed for kinase activity using the fluram-kemptide-Lys8 peptide as a substrate. The bottom panel shows the fractions containing the kemptide kinase peak, which is highlighted with an asterisk (*) in the chromatogram. (C) Separation by gel-filtration chromatography. Fractions 26–30 from the DEAE-Sepharose column were collected, concentrated and loaded on a Sephacryl S-300 column. Plotted is the absorbance at 280 nm against the elution volume of the fractions (top), which were assayed for kinase activity using the fluram-kemptide-Lys8 peptide as a substrate. The bottom panel illustrates the elution volume of the fractions containing the kemptide kinase peak, which is highlighted with an asterisk (*) in the chromatogram. In all cases, the arrowhead and arrow indicate the non-phosphorylated and phosphorylated fluorescently labelled peptide, respectively.

trypanosome kemptide kinase activities. Parasites were homogenized by sonication and centrifuged to separate the clarified soluble fraction, which contains the kemptide kinase activities, from the particulate fraction, and the soluble fraction was fractionated by sedimentation throughout a 5–20% sucrose gradient. Three peaks of kemptide kinase activity were detected (Fig. 3A): fractions 4–10 (peak 1), fractions 14–18 (peak 2) and fractions 20–22 (peak 3). We selected peak 1 for further purification.

Peak 1 from the sucrose gradient was separated by anion exchange chromatography through a DEAE-Sepharose column. Proteins were eluted by increasing the ionic strength of the washing buffer with NaCl and kemptide kinase activity was detected between fractions 26 and 30 (Fig. 3B). This kinase peak coincided with the major absorbing peak in the protein profile, which eluted at a concentration of 250 mM NaCl (Fig. 3B). Fractions 26–30 from the DEAE-Sepharose column were collected, concentrated and loaded onto a Sephacryl S-300 size exclusion chromatography column. Proteins were eluted from the column, and a peak of kemptide kinase activity was detected between elution volumes of 85.5 and 86.0 mL, with its maximum at 85.75 mL (Fig. 3C).

The purity of the kemptide kinase activity peaks from the sucrose gradient, the DEAE-Sepharose chromatography and the gel filtration column, was analysed by SDS-PAGE using a 12% polyacrylamide gel (Fig. 4A). As shown in the figure, five major polypeptide bands of 49.1, 42.8, 23.8, 22.0 and 19.4 kDa were observed in the polyacrylamide gel after the three purification steps, revealing that the *T. equiperdum* kemptide kinase enzyme was only partially purified. This enriched kemptide kinase-

containing fraction was named ppTeq-PKK for 'partially purified *T. equiperdum* protein kemptide kinase' and was employed in all further assays performed to characterize the enzyme.

As illustrated in Fig. 4B, and agreeing with the results shown in Fig. 1B, no stimulation by exogenously added cAMP was acquired on ppTeq-PKK, which confirmed that the trypanosome kinase corresponds to a cAMP-independent enzyme. As previously seen in Fig. 1B, a decrease in the phosphorylation signal of the fluorescently labelled kemptide was observed when cAMP was present (Fig. 4B), which again suggests that cAMP might be inhibiting the enzymatic activity perhaps by competing with ATP for the ATP-binding site on the parasite protein kinase.

Interestingly, the recombinant PKI- α from rabbit, which is a 77-amino acid protein that is a potent and highly specific inhibitor of PKA catalytic subunits, inhibited similarly both ppTeq-PKK and the purified PKA catalytic subunit from pig heart (Fig. 4C). Since PKI- α is not known to inhibit any other type of protein kinase, ppTeq-PKK appears to be a PKA catalytic subunit-like protein. This result was corroborated by using IP20, a peptide that contains the inhibitory sequence of PKI- α . As seen in Fig. S1, IP20 also inhibited both the purified PKA catalytic subunit from porcine heart (lane C+, IP20) and ppTeq-PKK (lane Teq-PKK, IP20).

With the intention of evaluating whether glucose inhibition was caused by a direct interaction between the monosaccharide and the enzyme, ppTeq-PKK was incubated with various concentrations of glucose. As shown in Fig. 2C, glucose did not inactivate the parasite protein kinase, implying that the sugar is not affecting




Fig. 4. (A) Separation by SDS-PAGE of the *T. equiperdum* kemptide kinase purification steps. Proteins were revealed by silver staining. Arrows indicate the molecular masses of the polypeptide bands contained in the kemptide kinase-enriched fraction. M = molecular weight markers. (B) ppTeq-PKK is a cAMP-independent kemptide kinase. Kinase activity was measured by the Promega Pep-Tag® non-radioactive assay, using the PepTag® A1 kemptide as substrate. Determinations were made in the absence (–) or presence of 5 μ M cAMP (+). (C) PKI- α inhibits ppTeq-PKK. ppTeq-PKK (Teq-PKK) was incubated with various units (U) of PKI- α (0.9, 9 and 36 units). Kinase activity was assayed using the fluram-kemptide-Lys8 peptide as substrate. In (B) and (C), the arrowhead and arrow indicate the non-phosphorylated and phosphorylated peptide, respectively. Purified PKA catalytic subunit from pig heart was utilized as a positive control (+). The reaction mixture without any added enzyme was used as a negative control (–).

directly the trypanosome enzyme but an upstream component of this kemptide kinase signalling pathway. These findings were consistent with our previous results that showed kemptide kinase activities following fractionation by sedimentation throughout a 5–20% sucrose gradient. Although the centrifuged fractions contained sucrose, three peaks of kemptide kinase activity were identified (Fig. 3A).

Evaluation of other substrates

Although kemptide has been shown to be a specific peptide substrate for PKA and PKA-like enzymes, it is also known that it can be phosphorylated by other protein kinases (MacAla et al., 1998). For that reason, we used the electrophoretic gel-shift assay to determine whether ppTeq-PKK was capable of phosphorylating two additional substrates, histone type II-AS and SP20, which are also known to serve as substrates for PKA and PKA-like enzymes. Analogous to kemptide, both histone type II-AS and SP20 were phosphorylated by the trypanosome enzymatic activity (Fig. 5A).

The PKA R^{II} α -subunit contains a phosphorylatable epitope within the catalytic subunit inhibitory sequence. As previously shown (First et al., 1988), SDS-PAGE distinguishes the phospho and dephospho forms of the pig heart R^{II} α -subunit readily. Immunoblotting using anti-R^{II} α -subunit antibodies illustrated that upon phosphorylation by the PKA catalytic subunit, the purified R^{II} α -subunit from pig heart changed its electrophoretic mobility on SDS-polyacrylamide gels from an apparent molecular mass of 55 kDa to an apparent molecular mass of 57 kDa (Fig. 5B,




Fig. 5. Phosphorylation of histone type II-AS, SP20 and the PKA R^{II} α -subunit by ppTeq-PKK. (A) The trypanosome enzyme was capable of phosphorylating kemptide, histone type II-AS and SP20 by means of the electrophoretic gel-shift assay. Kinase reactions were performed in the absence (–) or presence (+) of ppTeq-PKK. The arrowhead and arrow indicate the migration of the non-phosphorylated and phosphorylated substrates, respectively. (B) Purified R^{II} α -subunit alone (R^{II}) or in the presence of either the pig heart PKA catalytic subunit (R^{II}+C) or ppTeq-PKK (R^{II}+Teq-PKK) was incubated with 1 mM cAMP, 10 mM MgCl₂ and 1 mM ATP in 50 mM Tris-HCl (pH 7.5). Following incubation for 30 min, at room temperature, the reaction mixtures were separated by SDS-PAGE on an 8% polyacrylamide slab gel and electrotransferred to nitrocellulose. The blot was revealed by using anti-R^{II} α -subunit antibodies. Arrows indicate the migration and apparent molecular masses of the dephosphorylated (R^{II}) and phosphorylated (R^{II}-P) forms of the monomeric R^{II} α -subunit. An endogenous proteolytic fragment of the R^{II} α -subunit is also shown (COOH-R^{II}).

lane R^{II}+C). Interestingly, ppTeq-PKK was also capable of phosphorylating the purified R^{II} α -subunit from porcine cardiac muscle, as shown by the appearance of a doublet of 55 and 57 kDa (Fig. 5B, lane R^{II}+Teq-PKK). It is also known that PKA regulatory subunits possess a hinge region that is very susceptible to proteolysis and is located adjacent to the catalytic subunit inhibitory sequence. Limited proteolysis cleaves the regulatory subunits into carboxy-terminal fragments that retain the cAMP-binding properties of the native proteins but that have lost the primary contacts between the two subunits. These carboxy-terminal fragments represent about the last two-thirds of the proteins. Figure 5B also shows a band that migrates with an apparent molecular mass of ~37 kDa, and is recognized by the anti-R^{II} α -subunit antibodies. The 37 kDa polypeptide band must correspond to the R^{II} α -subunit proteolytic fragment (COOH-R^{II}) that is formed by endogenous proteolysis (Bubis and Taylor, 1987).

Determination of the molecular weight and hydrodynamic parameters of the kemptide kinase from *T. equiperdum*

ppTeq-PKK was applied to a size-exclusion column that was calibrated using markers with known molecular masses. By plotting the partition coefficient (K_{av}) of each standard vs the logarithm of its molecular weight (Laurent and Killander, 1964), a molecular mass of 39.07 kDa was determined for the partially purified trypanosome enzyme (Fig. S2A). Additionally, a linear relationship

was obtained by plotting $(-\log K_{av})^{1/2}$ against the Stokes radius of each standard protein (Siegel and Monty, 1966), and a Stokes radius of 2.74 nm was calculated for the partially purified parasite protein (Fig. S2B). A sedimentation coefficient of 4.06 S was measured for the parasite kemptide kinase by subjecting the enzyme to ultracentrifugation in the presence of a mixture of proteins with known sedimentation coefficients (Fig. S3A). As seen in Fig. S3B, a molecular mass of 51.73 kDa was obtained by plotting the sedimentation coefficient against the $(\text{molecular weight})^{2/3}$ of each standard (Martin and Ames, 1961). For spherical molecules, the molecular weight of a species can also be calculated from a combination of the measured Stokes radius and sedimentation coefficient (Siegel and Monty, 1966; Johansson *et al.*, 2001). Using this approximation, a molecular mass of 43.55 kDa was estimated for the parasite enzyme (Fig. S3C). Combining all these results, the native size of ppTeq-PKK appears to be between 39.07 and 51.73 kDa. As shown above in Fig. 4A, SDS-PAGE revealed polypeptide bands with apparent molecular masses of 42.8 and 49.1 kDa. Therefore, the parasite enzyme might correspond to one of these bands.

The frictional coefficient ff_0 can be determined from the molecular weight and the Stokes radius (Bloomfield *et al.*, 1967). The calculated frictional ratio value for the partially purified kemptide kinase was 1.22, 1.24 or 1.25 when values of the molecular mass obtained either by gel filtration (39.07 kDa), or by the combination of the measured Stokes radius and sedimentation coefficient (43.55 kDa), or by sedimentation (51.73 kDa) were employed, respectively. These findings indicate that the parasite enzyme is a globular and compact protein (Erickson, 2009). Since the frictional coefficient not only contains information about the molecular shape of the protein but also about the hydration effects, an upper boundary on the contributions of hydration can be assessed presuming that all deviations of ff_0 from unity are due to hydration (Darling *et al.*, 2000). The maximal value of hydration for the trypanosome kemptide kinase was 0.57–0.67 g of H₂O per gram of protein (depending on the experimental value used for ff_0), which slightly diverges from the 0.3–0.5 g of H₂O per gram value of a typical medium-sized and compact protein. This result also suggests that the *T. equiperdum* enzyme is globular and compact.

We were not able to determine the hydrodynamic parameters of the trypanosome kemptide kinase activities that fractionated with peaks 2 and 3 following sedimentation by ultracentrifugation (Fig. 3A). The migration volumes of these fractions were higher than that of alcohol dehydrogenase, which was the largest of the protein standards that were used for calibration. Therefore, the parasite kemptide kinase activities in peaks 2 and 3 possessed molecular masses higher than 150 kDa, and their sedimentation coefficients fell outside the linear range of the curve shown in Fig. S3A. Such high sizes for these protein kinases are suggestive of potential formation of complexes with other interacting proteins.

Biochemical characterization of the *T. equiperdum* kemptide kinase

Kemptide kinase activity was assayed at different temperatures and pH, showing optimal performance at 37 °C (Fig. 6A) and pH 8.0 (Fig. 6B). Since two variables, the pH and the buffer system, were changed in Fig. 6B, we carried out an additional experiment that incorporated chosen buffer systems on the basis of their overlapping effective pH ranges (Fig. S4). Although slight effects caused by the buffer system on the trypanosome enzymatic activity were noticed, these results ratified that pH 8.0 is the optimal pH for ppTeq-PKK.

The parasite kinase activity was only detected in the presence of ATP (Fig. 6C) and Mg²⁺ (Fig. 6D), and no activity was perceived neither when GTP or CTP, nor when other divalent cations, such as Zn²⁺, Mn²⁺, Co²⁺, Cu²⁺, Ca²⁺ and Fe²⁺ were employed as co-substrates (Fig. 6C and D). However, the enzymatic activity of the parasite kemptide kinase was inhibited by other nucleotides and/or analogues of ATP, such as AMP, ADP, GMP, GDP, GTP, AMP-PNP, FSBA and CTP, and by other divalent cations (Zn²⁺, Mn²⁺, Co²⁺, Cu²⁺, Ca²⁺ and Fe²⁺) (Fig. 6E and F). Since the trypanosome enzyme was inhibited by such a variety of nucleotides, the reduction in kinase activity that was previously seen in the presence of cAMP (Figs 1B and 4B) must be caused by inhibition rather than by a direct influence on the enzymatic activity. As the other nucleotides, cAMP is probably binding and competing with ATP for the protein kinase ATP-binding site.

Determination of the K_m for ATP and kemptide of the *T. equiperdum* kemptide kinase

Kinase activity was assayed by the HPLC methodology described by Luzi *et al.* (2017), which was modified by using fluram-kemptide as substrate. Typical Michaelis–Menten curves for ATP and fluram-kemptide were obtained for the trypanosome enzyme (Fig. 7). K_m values for ATP and kemptide of $11.8 \pm 4.1 \mu\text{M}$ (Fig. 7A) and $24.7 \pm 3.8 \mu\text{M}$ (Fig. 7B), respectively, were determined for the parasite kinase after plotting the data on double reciprocal Lineweaver–Burk plots.

Discussion

Here, we report a novel protein kinase that was partially purified from the Venezuelan TeAp-N/D1 isolate of *T. equiperdum*. This enzyme was capable of recognizing kemptide, histone II-AS, SP20 and the PKA R^{II} α -subunit, which are specific substrates for the PKA catalytic subunit, and was inhibited by PKI- α and IP20, which are specific peptide inhibitors for the PKA catalytic subunit, but surprisingly, it was not stimulated by cAMP. Noticeably, the *T. equiperdum* kinase was activated when parasites were subjected to nutritional restriction caused by lack of glucose in the medium.

Glucose is the leading source of energy in the majority of eukaryotes, being glycolysis and mitochondrial respiration the sequential glucose oxidation routes. Many trypanosomatids live in carbohydrate-rich habitats inside their hosts. For instance, *T. equiperdum*, as well as *T. evansi* and the trypomastigote slender and stumpy stages of *T. brucei*, live in the bloodstream of their mammalian host, which warrants a permanent source of glucose. Parasite trypanosomes living in carbohydrate-rich environments seem to depend on carbohydrates for their energy metabolism, and do not require a cytochrome-containing respiratory chain or an elaborate mitochondrial metabolism (Tielens and van Hellemond, 2009). Since glucose deprivation activates the *T. equiperdum* protein kinase described here, this enzyme may perhaps participate in a signalling cascade triggered by nutritional and/or energy stress. This metabolic signalling network might be implicated in controlling parasite survival during glucose fasting.

In most organisms, the PKA holoenzyme is an inactive tetrameric complex composed of two catalytic subunits bound to a dimer of regulatory subunits. Binding of cAMP to the regulatory subunits leads to the dissociation and activation of the two catalytically active catalytic subunit monomers (Taylor, 1989). Analysis of the complement of protein kinases in the completed genomes of various trypanosomatids have indicated that there are genes that encode for protein kinases related to the PKA catalytic subunits from higher eukaryotes (<http://tritrypdb.org>). The first draft of the *T. equiperdum* genome (Hébert *et al.*, 2017),

Fig. 6. Characterization of ppTeq-PKK. Measurement of the optimal temperature (A) and pH (B). Determination of the specificity for nucleotide triphosphates (C) and divalent cations (D). In (C) and (D), the reaction mixture without any added enzyme was used as a negative control (-). (E) Inhibitory effect of nucleotides and analogues of ATP in the presence of Mg^{2+} . Samples containing the parasite enzyme were pre-incubated with 2 mM of each compound, and then assayed using 1 mM ATP. Control experiments were performed in the absence (-) and presence (+) of ATP. (F) Inhibitory effect of divalent cations in the presence of ATP. Samples containing the *T. equiperdum* kinase were pre-incubated with 20 mM of each divalent cation, and then assayed using 10 mM Mg^{2+} . Control experiments were performed in the absence (-) and presence (+) of Mg^{2+} . In all cases, kinase activity was assayed employing the fluram-kemptide-Lys8 peptide as a substrate. The arrowhead and arrow indicate the non-phosphorylated and phosphorylated peptide, respectively.





Fig. 7. Determination of the K_m for ATP (A) and kemptide (B) of ppTeq-PKK. (A) The analysis of the phosphorylation of fluram-kemptide was assayed by HPLC. To determine the K_m for ATP, various concentrations of ATP (0–70 μM) were employed with a fixed concentration of fluram-kemptide (30 μM). To measure the K_m for kemptide, various concentrations of fluram-kemptide (0–50 μM) were used with a fixed concentration of ATP (1 mM). In (A) and (B), the left and right panels represent the Michaelis-Menten curves and the Lineweaver-Burk double-reciprocal plots, respectively. K_m values were reported as the mean \pm S.E.M.

also described the presence of two protein kinases that are homologous to the PKA catalytic subunits [NCBI GenBank IDs: SCU68910.1 and SCU64246.1 (<https://www.ncbi.nlm.nih.gov/genbank/>)]. A gene homologous to the gene for the mammalian PKA regulatory subunits has also been reported in trypanosomatid parasites (<http://tritypdb.org>); and recently, we have identified a 57 kDa polypeptide band corresponding to the PKA regulatory subunit-like protein from the *T. equiperdum* TeAp-N/D1 isolate (Calabokis *et al.*, 2016). Antibody cross-reactivity clearly established the high level of conservation exhibited between the *T. equiperdum* PKA regulatory subunit-like protein and its mammalian counterparts; however, the trypanosome protein differed from other PKA regulatory subunits since it is a monomer that is not capable of binding cAMP (Bubis *et al.*, 2018). It is remarkable to find that the parasite kinase enzymatic activity is inhibited or activated depending on the level of glucose in the incubation medium, which is reminiscent of the occurrence of an inactive holoenzyme, or another enzymatic form that is hindered by some inhibitory or regulatory molecule under high-glucose

conditions. Yet, preliminary results from our laboratory using co-immunoprecipitation and pull-down experiments have also shown that there is no formation of a holoenzyme type of complex between the *T. equiperdum* PKA regulatory subunit-like protein and the kemptide kinase partially purified here (data not shown). These early results suggest that another inhibitory agent is responsible for the regulation of the kemptide kinase activity when glucose is not restricted.

Free PKA catalytic subunits from mammals have K_m s for ATP and kemptide of ~ 10 – $20 \mu M$ (Smith *et al.*, 1999; Moore *et al.*, 2003; Mena-Ulecia *et al.*, 2014; Luzi *et al.*, 2017), which are similar to the K_m values determined here for ppTeq-PKK. Moreover, the molecular mass and hydrodynamic properties found for the parasite kemptide kinase were comparable to those reported for the mammalian PKA catalytic subunits (Sugden *et al.*, 1976; Rubin *et al.*, 1979; Zoller *et al.*, 1979), and for PKA-like enzymes from other trypanosomatids (Banerjee and Sarkar, 1992; Ochatt *et al.*, 1993). All these results indicate that the *T. equiperdum* enzyme is a globular protein as is the free PKA catalytic subunit from mammals, and suggest that the parasite protein is probably monomeric. On the basis of their calculated molecular weight, the gene product of either the reported SCU68910.1 or SCU64246.1 genes from *T. equiperdum* (<https://www.ncbi.nlm.nih.gov/genbank/>) might correspond to the kemptide kinase characterized here. Bioinformatics analyses using the Clustal Omega program (<http://www.ebi.ac.uk/Tools/msa/clustalo/>) of the sequences of the two *T. equiperdum* protein kinases, the three protein kinases identified in *T. evansi* [TevSTIB805.9.7920, TevSTIB805.9.7870 and TevSTIB805.10.13650 (<http://tritypdb.org>)] and the α isoform of the *Mus musculus* PKA catalytic subunit (GenBank ID: AAH54834.1, <https://www.ncbi.nlm.nih.gov/genbank/>), yielded a 47–51% identity between the parasite kinases and the mouse protein, and an expected identity of >97% between the *T. equiperdum* and *T. evansi* protein kinases (data not shown).

If the parasite kinase corresponds to a PKA type of enzyme, it is unexpected to find that cAMP is not required for its activation. However, there are several reports describing cAMP-independent activation of PKA in other systems (Zhong *et al.*, 1997; Dulin *et al.*, 2001; Ferraris *et al.*, 2002; Zhang *et al.*, 2004; Yang *et al.*, 2008; Kohr *et al.*, 2010). Even in *Leishmania donovani* that is another trypanosomatid parasite, Banerjee and Sarkar (1992) have reported the characterization of a cyclic nucleotide-independent PKA enzyme that recognized kemptide as a substrate.

Trypanosoma equiperdum parasites may use similar strategies to activate this PKA-like protein in a cAMP-independent manner.

Control of diseases caused by trypanosomatid parasites relies mostly on chemotherapy. However, there is a very narrow battery of drugs and they generally have limitation issues, such as high toxicity and emerging resistance. Accordingly, finding new drug targets is an important goal in order to develop novel anti-parasitic chemotherapeutic agents. Since phosphorylation seems to play a key role in trypanosomatid parasite biology (Parsons *et al.*, 2005), and given that protein kinases are involved in regulating cell cycle control, proliferation, differentiation and response to stress, these enzymes represent promising drug targets for diseases caused by parasitic protozoa (Naula *et al.*, 2005). Although the physiological role of the *T. equiperdum* kemptide kinase characterized here is yet to be found, this enzyme might be an attractive drug target. The findings presented here open new avenues to study signal transduction pathways in *T. equiperdum*, and is a promising starting point that may eventually lead to the development of new anti-trypanosomosis therapeutics for treatment.

Supplementary material. The supplementary material for this article can be found at <https://doi.org/10.1017/S0031182018001920>.

Author ORCIDs.  José Bubis <http://orcid.org/0000-0002-8839-6745>.

Acknowledgements. We thank Oscar Noya and María A. Lorenzo (Instituto de Medicina Tropical, Universidad Central de Venezuela, Caracas, Venezuela) for synthesizing and donating the kemptide and kemptide-Lys8 peptides that were used in this work. We also thank Susan S. Taylor (Department of Chemistry, Biochemistry and Pharmacology, University of California San Diego, La Jolla, USA) for providing the SP20 and IP20 synthetic peptides that were employed here.

Financial support. This research was supported by grant numbers S1-IC-CB-007-14, S1-IC-CB-001-17 and S1-IC-CB-008-17 from Decanato de Investigación y Desarrollo, Universidad Simón Bolívar, Caracas, Venezuela, to JB, and 2013001659 from FONACIT, Caracas, Venezuela, to NAA.

Conflict of interest. None.

Ethical standards. We declare that all the experiments in this paper were carried out in accordance with the legal and ethical standards of Venezuela.

References

- Araujo NA, Guevara A, Lorenzo MA, Calabokis M and Bubis J (2016) Fluram-kemptide-Lys8 non-radioactive assay for protein kinase A. *The Protein Journal* **35**, 247–255.
- Banerjee C and Sarkar D (1992) Isolation and characterization of a cyclic nucleotide-independent protein kinase from *Leishmania donovani*. *Molecular and Biochemical Parasitology* **52**, 195–206.
- Bao Y, Weiss LM, Braunstein VL and Huang H (2008) Role of protein kinase A in *Trypanosoma cruzi*. *Infection and Immunity* **76**, 4757–4763.
- Bloomfield V, Dalton WO and van Holde KE (1967) Frictional coefficients of multisubunit structures. I. *Theory. Biopolymers* **5**, 135–148.
- Brun R, Hecker H and Lun ZR (1998) *Trypanosoma evansi* and *T. equiperdum*: distribution, biology, treatment and phylogenetic relationship (a review). *Veterinary Parasitology* **79**, 95–107.
- Bubis J and Taylor SS (1987) Limited proteolysis alters the photoaffinity labeling of adenosine 3',5'-monophosphate dependent protein kinase II with 8-azidoadenosine 3',5'-monophosphate. *Biochemistry* **26**, 5997–6004.
- Bubis J, Martínez JC, Calabokis M, Ferreira J, Sanz-Rodríguez CE, Navas V, Escalona JL, Guo Y and Taylor SS (2018) The gene product of a *Trypanosoma equiperdum* ortholog of the cAMP-dependent protein kinase regulatory subunit is a monomeric protein that is not capable of binding cyclic nucleotides. *Biochimie* **146**, 166–180.
- Calabokis M, González Y, Merchán A, Escalona JL, Araujo NA, Sanz-Rodríguez CE, Cywiak C, Spencer LM, Martínez JC and Bubis J (2016) Immunological identification of a cAMP-dependent protein kinase regulatory subunit-like protein from the *Trypanosoma equiperdum* TeAp-N/D1 isolate. *Journal of Immunoassay and Immunochemistry* **37**, 485–514.
- Darling PJ, Holt JM and Ackers GK (2000) Coupled energetics of λ cro repressor self-assembly and site-specific DNA operator binding II: cooperative interactions of cro dimers. *Journal of Molecular Biology* **302**, 625–638.
- Dulin NO, Niu J, Browning DD, Ye RD and Voyno-Yasenetskaya T (2001) Cyclic AMP-independent activation of protein kinase A by vasoactive peptides. *Journal of Biological Chemistry* **276**, 20827–20830.
- Erickson HP (2009) Size and shape of protein molecules at the nanometer level determined by sedimentation, gel filtration, and electron microscopy. *Biological Procedures Online* **11**, 32–51.
- Ferraris JD, Persaud P, Williams CK, Chen Y and Burg MB (2002) cAMP-independent role of PKA in tonicity-induced transactivation of tonicity-responsive enhancer/osmotic response element-binding protein. *Proceedings of the National Academy of Sciences of the USA* **99**, 16800–16805.
- First EA, Bubis J and Taylor SS (1988) Subunit interaction sites between the regulatory and catalytic subunits of cAMP-dependent protein kinases. Identification of a specific interchain disulfide bond. *Journal of Biological Chemistry* **263**, 5177–5182.
- Genestra M, Cysne-Finkelstein L and Leon L (2004) Protein kinase A activity is associated with metacyclogenesis in *Leishmania amazonensis*. *Cell Biochemistry and Function* **22**, 315–320.
- Gizaw Y, Megersa M and Fayera T (2017) Dourine: a neglected disease of equids. *Tropical Animal Health and Production* **49**, 887–897.
- Hébert L, Moumen B, Madeline A, Steinbiss S, Lakhdar L, Van Reet N, Büscher P, Laugier C, Cauchard J and Petry S (2017) First draft genome sequence of the dourine causative agent: *Trypanosoma equiperdum* strain OVI. *Journal of Genomics* **5**, 1–3.
- Johansson E, Majka J and Burgers PM (2001) Structure of DNA polymerase δ from *Saccharomyces cerevisiae*. *Journal of Biological Chemistry* **276**, 43824–43828.
- Kemp BE (1980) Phosphorylation of acyl and dansyl derivatives of the peptide Leu-Arg-Arg-Ala-Ser-Leu-Gly by the cAMP-dependent protein kinase. *Journal of Biological Chemistry* **255**, 2914–2918.
- Kemp BE, Bylund DB, Huang TS and Krebs EG (1975) Substrate specificity of the cyclic AMP-dependent protein kinase. *Proceedings of the National Academy of Sciences of the USA* **72**, 3448–3452.
- Kohr MJ, Traynham CJ, Roof SR, Davis JP and Ziolo MT (2010) cAMP-independent activation of protein kinase A by the peroxy nitrite generator SIN-1 elicits positive inotropic effects in cardiomyocytes. *Journal of Molecular and Cellular Cardiology* **48**, 645–648.
- Laemmli UK (1970) Cleavage of structural proteins during the assembly of the head of bacteriophage T4. *Nature* **227**, 680–685.
- Lanham SM and Godfrey DG (1970) Isolation of salivarian trypanosomes from man and other mammals using DEAE-cellulose. *Experimental Parasitology* **28**, 521–534.
- Laurent TC and Killander J (1964) A theory of gel filtration and its experimental verification. *Journal of Chromatography A* **14**, 317–330.
- Lineweaver H and Burk D (1934) The determination of enzyme dissociation constants. *Journal of the American Chemical Society* **56**, 658–666.
- Lutz MP, Pinon DI and Miller LJ (1994) A nonradioactive fluorescent gel-shift assay for the analysis of protein phosphatase and kinase activities toward protein-specific peptide substrates. *Analytical Biochemistry* **220**, 268–274.
- Luzi NM, Lyons CE, Peterson DL and Ellis KC (2017) Characterization of PKA α enzyme kinetics and inhibition in an HPLC assay with a chromophoric substrate. *Analytical Biochemistry* **532**, 45–52.
- MacAla LJ, Hayslett JP and Smallwood JI (1998) Measurement of cAMP-dependent protein kinase activity using a fluorescent-labeled kemptide. *Kidney International* **54**, 1746–1750.
- Malki-Feldman L and Jaffe CL (2009) *Leishmania major*: effect of protein kinase A and phosphodiesterase activity on infectivity and proliferation of promastigotes. *Experimental Parasitology* **123**, 39–44.
- Maller JL, Kemp BE and Krebs EG (1978) In vivo phosphorylation of a synthetic peptide substrate of cyclic AMP-dependent protein kinase. *Proceedings of the National Academy of Sciences of the USA* **75**, 248–251.
- Martin RG and Ames BN (1961) A method for determining the sedimentation behavior of enzymes: application to protein mixtures. *Journal of Biological Chemistry* **236**, 1372–1379.
- Mena-Ulecia K, Vergara-Jaque A, Poblete H, Tiznado W and Caballero J (2014) Study of the affinity between the protein kinase PKA and peptide

- substrates derived from kemptide using molecular dynamics simulations and MM/GBSA. *PLoS ONE* **9**, e109639.
- Moore MJ, Adams JA and Taylor SS** (2003) Structural basis for peptide binding in protein kinase A. Role of glutamic acid 203 and tyrosine 204 in the peptide-positioning loop. *Journal of Biological Chemistry* **278**, 10613–10618.
- Naula C, Parsons M and Mottram JC** (2005) Protein kinases as drug targets in trypanosomes and *Leishmania*. *Biochimica et Biophysica Acta (BBA) – Proteins and Proteomics* **1754**, 151–159.
- Nelson NC and Taylor SS** (1981) Differential labeling and identification of the cysteine-containing tryptic peptides of catalytic subunit from porcine heart cAMP-dependent protein kinase. *Journal of Biological Chemistry* **256**, 3743–3750.
- Nelson NC and Taylor SS** (1983) Selective protection of sulfhydryl groups in cAMP-dependent protein kinase II. *Journal of Biological Chemistry* **258**, 10981–10987.
- Ochatt CM, Ulloa RM, Torres HN and Téllez-Iñón MT** (1993) Characterization of the catalytic subunit of *Trypanosoma cruzi* cyclic AMP-dependent protein kinase. *Molecular and Biochemical Parasitology* **57**, 73–81.
- Parsons M, Worthey EA, Ward PN and Mottram JC** (2005) Comparative analysis of the kinomes of three pathogenic trypanosomatids: *Leishmania major*, *Trypanosoma brucei* and *Trypanosoma cruzi*. *BMC Genomics* **6**, 127.
- Rubin CS, Rangel-Aldao R, Sarkar D, Erlichman J and Fleischer N** (1979) Characterization and comparison of membrane-associated and cytosolic cAMP-dependent protein kinases. Physicochemical and immunological studies on bovine cerebral cortex protein kinases. *Journal of Biological Chemistry* **254**, 3797–3805.
- Sánchez E, Perrone T, Recchimuzzi G, Cardozo I, Biteau N, Aso PM, Mijares A, Baltz T, Berthier D, Balzano-Nogueira L and Gonzatti MI** (2015) Molecular characterization and classification of *Trypanosoma* spp. Venezuelan isolates based on microsatellite markers and kinetoplast maxicircle genes. *Parasites & Vectors* **8**, 536.
- Siegel LM and Monty KJ** (1966) Determination of molecular weights and frictional ratios of proteins in impure systems by use of gel filtration and density gradient centrifugation. Application to crude preparations of sulfite and hydroxylamine reductases. *Biochimica et Biophysica Acta (BBA) – Biophysics Including Photosynthesis* **112**, 346–362.
- Smith CM, Radzio-Andzelm E, Madhusudan M, Akamine P and Taylor SS** (1999) The catalytic subunit of cAMP-dependent protein kinase: prototype for an extended network of communication. *Progress in Biophysics & Molecular Biology* **71**, 313–341.
- Stevens JR and Brisse S** (2004) Systematics of trypanosomes of medical and veterinary importance. In Maudlin I, Holmes PH and Miles MA (ed.) *The Trypanosomiasis*, Wallingford, UK: CABI Publishing, CAB International, pp. 1–23.
- Sugden PH, Holladay LA, Reimann EM and Corbin JD** (1976) Purification and characterization of the catalytic subunit of adenosine 3':5'-cyclic monophosphate-dependent protein kinase from bovine liver. *Biochemical Journal* **159**, 409–422.
- Taylor SS** (1989) cAMP-dependent protein kinase. Model for an enzyme family. *Journal of Biological Chemistry* **264**, 8443–8446.
- Taylor SS and Stafford PH** (1978) Characterization of adenosine 3':5'-monophosphate-dependent protein kinase and its dissociated subunits from porcine skeletal muscle. *Journal of Biological Chemistry* **253**, 2284–2287.
- Tielens AG and van Hellemond JJ** (2009) Surprising variety in energy metabolism within trypanosomatidae. *Trends in Parasitology* **25**, 482–490.
- Towbin H, Staehelin T and Gordon J** (1979) Electrophoretic transfer of proteins from polyacrylamide gels to nitrocellulose sheets: procedure and some applications. *Proceedings of the National Academy of Sciences of the USA* **76**, 4350–4354.
- Yang H, Lee CJ, Zhang L, Sans MD and Simeone DM** (2008) Regulation of transforming growth factor β -induced responses by protein kinase A in pancreatic acinar cells. *American Journal of Physiology – Gastrointestinal and Liver Physiology* **295**, G170–G178.
- Zhang L, Duan CJ, Binkley C, Li G, Uhler MD, Logsdon CD and Simeone DM** (2004) A transforming growth factor β -induced Smad3/Smad4 complex directly activates protein kinase A. *Molecular and Cellular Biology* **24**, 2169–2180.
- Zhong H, SuYang H, Erdjument-Bromage H, Tempst P and Ghosh S** (1997) The transcriptional activity of NF- κ B is regulated by the I κ B-associated PKAc subunit through a cyclic AMP-independent mechanism. *Cell* **89**, 413–424.
- Zoller MJ, Kerlavage AR and Taylor SS** (1979) Structural comparisons of cAMP-dependent protein kinases I and II from porcine skeletal muscle. *Journal of Biological Chemistry* **254**, 2408–2412.