

ON EQUIVALENCE RELATIONS INDUCED BY POLISH GROUPS ADMITTING COMPATIBLE TWO-SIDED INVARIANT METRICS

LONGYUN DING  AND YANG ZHENG

Abstract. Given a Polish group G , let $E(G)$ be the right coset equivalence relation $G^\omega/c(G)$, where $c(G)$ is the group of all convergent sequences in G . We first established two results:

- (1) Let G, H be two Polish groups. If H is TSI but G is not, then $E(G) \not\leq_B E(H)$.
- (2) Let G be a Polish group. Then the following are equivalent: (a) G is TSI non-archimedean; (b) $E(G) \leq_B E_0^\omega$; and (c) $E(G) \leq_B \mathbb{R}^\omega/c_0$. In particular, $E(G) \sim_B E_0^\omega$ iff G is TSI uncountable non-archimedean.

A critical theorem presented in this article is as follows: Let G be a TSI Polish group, and let H be a closed subgroup of the product of a sequence of TSI strongly NSS Polish groups. If $E(G) \leq_B E(H)$, then there exists a continuous homomorphism $S : G_0 \rightarrow H$ such that $\ker(S)$ is non-archimedean, where G_0 is the connected component of the identity of G . The converse holds if G is connected, $S(G)$ is closed in H , and the interval $[0, 1]$ can be embedded into H .

As its applications, we prove several Rigid theorems for TSI Lie groups, locally compact Polish groups, separable Banach spaces, and separable Fréchet spaces, respectively.

§1. Introduction. In recent years, logicians have achieved remarkable research outcomes in descriptive set theory, specifically in the investigation of the relative complexity among equivalence relations originating from various branches of mathematics, utilizing Borel reducibility. Polish groups and their actions play a crucial role in this research direction. Concurrently, researchers also aim to employ Borel reducibility among equivalence relations to characterize the properties of Polish groups.

The authors introduced in [5] the notion of *equivalence relations induced by Polish groups*: given a Polish group G , the equivalence relation $E(G)$ is defined on G^ω as:

$$xE(G)y \iff \lim_n x(n)y(n)^{-1} \text{ converges in } G,$$

for $x, y \in G^\omega$. These equivalence relations have shown great potential in characterizing properties of Polish groups. In fact, based on the results on Borel reducibility involving the equivalence relation $E(G)$, we can accurately determine the classes to which certain Polish groups G belong. For instance, (1) G is countable discrete iff $E(G) \sim_B E_0$; (2) G is non-archimedean iff $E(G) \leq_B =^+$; and (3) if H is CLI but G is not, then $E(G) \not\leq_B E(H)$ (see [5]). Additionally, the authors further established

Received April 16, 2024.

2020 *Mathematics Subject Classification*. Primary 03E15, 22A05, Secondary 22D05, 22E15, 46A16.

Key words and phrases. Borel reduction, TSI Polish group, equivalence relation.

© The Author(s), 2025. Published by Cambridge University Press on behalf of The Association for Symbolic Logic.
0022-4812/00/0000-0000
DOI:10.1017/jsl.2025.9



results in [6] on Borel reducibility among equivalence relations induced by locally compact abelian Polish groups, including a Rigid Theorem.

THEOREM 1.1 (Rigid Theorem, [6, Theorem 2.8]). *Let G be a compact connected abelian Polish group and H a locally compact abelian Polish group. Then $E(G) \leq_B E(H)$ iff there is a continuous homomorphism $S : G \rightarrow H$ such that $\ker(S)$ is non-archimedean.*

In this article, we shift our focus to *TSI Polish groups*, i.e., those that admit compatible complete two-sided invariant metrics. Firstly, the Borel reducibility among equivalence relations induced by Polish groups can accurately distinguish non-TSI and TSI Polish groups. Actually, we have concluded that:

THEOREM 1.2. *Let G, H be two Polish groups. If H is TSI but G is not, then $E(G) \not\leq_B E(H)$.*

To elucidate the research significance of equivalence relations induced by Polish groups, we compare them with benchmark equivalence relations such as E_0 , E_0^ω , \mathbb{R}^ω/c_0 , etc. (definitions of these benchmark equivalence relations can be found in the next section).

Surprisingly, we prove that there is NO Polish group G such that

$$E_0^\omega <_B E(G) \leq_B \mathbb{R}^\omega/c_0.$$

In stark contrast, Farah proved that the partially ordered set $P(\omega)/\text{Fin}$ can be embedded into Borel equivalence relations between E_0^ω and \mathbb{R}^ω/c_0 (see [7, Theorem 5.4]).

The above results stem from the following more precise theorem.

THEOREM 1.3. *Let G be a Polish group. Then the following are equivalent:*

- (1) G is TSI non-archimedean;
- (2) $E(G) \leq_B E_0^\omega$; and
- (3) $E(G) \leq_B \mathbb{R}^\omega/c_0$.

In particular, $E(G) \sim_B E_0^\omega$ iff G is TSI uncountable non-archimedean.

To generalize the Rigid Theorem for locally compact abelian Polish groups mentioned above, we shall first generalize [5, Theorem 6.13] to a highly technical theorem, namely the Pre-rigid Theorem (the statement of this theorem is too lengthy to include in the introduction). The Pre-rigid Theorem and all its applications involve a notion named strongly NSS Polish groups. A Polish group G is called *strongly NSS* if there exists an open neighborhood V of 1_G in G such that

$$\forall (g_n) \in G^\omega (g_n \rightarrow 1_G \Rightarrow \exists n_0 < \dots < n_k (g_{n_0} \dots g_{n_k} \notin V)).$$

We employ the Pre-rigid Theorem to prove the following theorem, where G_0 is the connected component of 1_G in G .

THEOREM 1.4. *Let G be a TSI Polish group, and let H be a closed subgroup of the product of a sequence of TSI strongly NSS Polish groups. If $E(G) \leq_B E(H)$, then there exists a continuous homomorphism $S : W \rightarrow H$ such that $\ker(S)$ is non-archimedean, where $W \supseteq G_0$ is a countable intersection of clopen subgroups in G .*

Moreover, the converse holds if $G = W$, $S(G)$ is closed in H , and the interval $[0, 1]$ can be embedded into H .

Several applications of the Pre-rigid Theorem and the above theorem are listed below.

THEOREM 1.5. *Let G be a TSI Polish group, and let H be a closed subgroup of the product of a sequence of TSI strongly NSS Polish groups. If H is totally disconnected but G is not, then $E(G) \not\leq_B E(H)$.*

Recall that a *Lie group* is a group which is also a smooth manifold such that the group operations are smooth functions. Let G be a Lie group, then G_0 is an open normal subgroup of G . A completely metrizable topological group G is called a *pro-Lie group* if every open neighborhood of 1_G contains a normal subgroup N such that G/N is a Lie group (see [9, Definition 1]).

THEOREM 1.6. *Let G, H be two TSI Polish groups such that H is a pro-Lie group. If $E(G) \leq_B E(H)$, then there exists a continuous homomorphism $S : G_0 \rightarrow H$ such that $\ker(S)$ is non-archimedean.*

Moreover, the converse holds if G is connected and $S(G)$ is closed in H .

The following theorem is a generalization of Theorem 1.1. It should be emphasized that all locally compact TSI groups are pro-Lie groups.

THEOREM 1.7 (Rigid Theorem for locally compact TSI groups). *Let G be a locally compact connected TSI Polish group and H a TSI pro-Lie Polish group. Then $E(G) \leq_B E(H)$ iff there exists a continuous homomorphism $S : G \rightarrow H$ such that $\ker(S)$ is non-archimedean.*

Note that a topological group G is a Lie group iff it is locally compact and there exists an open neighborhood V of 1_G in G such that no non-trivial subgroup of G is contained in V . This leads to a positive answer to [5, Question 7.4] as follows:

THEOREM 1.8 (Rigid Theorem for TSI Lie groups). *Let G, H be two separable TSI Lie groups such that G is connected. Then $E(G) \leq_B E(H)$ iff there exists a continuous locally injective homomorphism $S : G \rightarrow H$.*

All separable *Fréchet spaces*, i.e., separable completely metrizable topological vector spaces, can be viewed as abelian Polish groups under the addition operation.

THEOREM 1.9 (Rigid Theorem for Fréchet spaces). *Let X, Y be two separable Fréchet spaces such that Y is a closed subgroup of the product of a sequence of TSI strongly NSS Polish groups. Then $E(X) \leq_B E(Y)$ iff X is topologically isomorphic to a closed linear subspace of Y .*

All Banach spaces are Fréchet spaces. We point out that a separable Banach space X is not strongly NSS iff it has a closed linear subspace topologically isomorphic to c_0 . This implies that:

THEOREM 1.10 (Rigid Theorem for Banach spaces). *Let X, Y be two separable Banach spaces such that Y contains no closed linear subspaces topologically isomorphic to c_0 . Then $E(X) \leq_B E(Y)$ iff X is topologically isomorphic to a closed linear subspace of Y .*

In addition, we attempt to study some examples induced by totally disconnected TSI Polish groups. For $p \in [1, +\infty)$ and $a \in c_0$, let

$$I_a = \{n \in \omega : a(n) \neq 0\}, \quad A_{p,a} = \{v \in l_p : \forall n (v(n) \in a(n)\mathbb{Z})\}.$$

If I_a is infinite, then $A_{p,a}$ is a totally disconnected, strongly NSS abelian Polish group, but is not non-archimedean. In particular, we let $d(n) = 2^{-n}$, and define

$$A_p = A_{p,d} = \{v \in I_p : \forall n (v(n) \in 2^{-n}\mathbb{Z})\}.$$

THEOREM 1.11. For $p, q \in [1, +\infty)$ and $a \in c_0$, the following hold:

- (1) if I_a is a nonempty finite set, then $E(A_{p,a}) \sim_B E_0$;
- (2) if I_a is infinite, then $E(A_{p,a}) \sim_B E(A_p)$;
- (3) $E(A_p) <_B E(I_p)$;
- (4) $E(A_p) \leq_B E(I_q) \iff p = q$; and
- (5) $E(A_p) \leq_B E(A_q) \iff p = q$.

This article is organized as follows. In Section 2, we recall some notions in descriptive set theory and also recall some notions and results originated from [5] that will be repeatedly used in this article. In Section 3, we prove Theorem 1.2. It is worth noting that some notation defined in this section will continue to be used in Section 5. In Section 4, we prove Theorem 1.3. In Section 5, we prove the Pre-rigid Theorem, which will serve as the foundation for the subsequent sections. In Section 6, we prove Theorems 1.4–1.10. Finally, in Section 7, we prove Theorem 1.11.

§2. Preliminaries. In this article, all groups are assumed to contain at least two elements. Any linear space can be viewed as an abelian group. The addition operation in it, as well as in all its subgroups, is denoted by $+$, and its identity element is denoted by 0 . Unless otherwise specified, for any abstract topological group G , we use multiplicative notation to express the group operation, and 1_G to express the identity element of G .

Given a topological group G , the connected component of 1_G is denoted by G_0 , which is clearly a closed normal subgroup of G . Note that G is *totally disconnected* iff $G_0 = \{1_G\}$. It is worth noting that any open subgroup H of G is also closed, since $H = G \setminus \{gH : g \notin H\}$.

A topological space is *Polish* if it is separable and completely metrizable. For further details in descriptive set theory, we refer to [13]. We say a topological group is *Polish* if its topology is Polish. Consider a Polish group G and a Polish space X . A continuous action of G on X , denoted by $G \curvearrowright X$, is a continuous map $a : G \times X \rightarrow X$ which satisfies that $a(1_G, x) = x$ and $a(gh, x) = a(g, a(h, x))$ for $g, h \in G$ and $x \in X$. For brevity, we write gx in place of $a(g, x)$. The *orbit equivalence relation* E_G^X is defined as

$$xE_G^X y \iff \exists g \in G (gx = y).$$

A Polish group is *non-archimedean* if it has a neighborhood basis of the identity element consisting of open subgroups. A metric d on a group G is *left-invariant* if $d(gh, gk) = d(h, k)$ for all $g, h, k \in G$; we also define *right-invariant* metric similarly. We say that d is *two-sided invariant* if it is both left-invariant and right-invariant. The Birkhoff–Kakutani theorem asserts that every metrizable topological group admits a compatible left-invariant metric (see [8, Theorem 2.1.1]). We say a Polish group G is *CLI* if it admits a compatible complete left-invariant metric; and say G is *TSI* if it admits a compatible two-sided invariant metric. A compatible two-sided invariant metric on a Polish group is necessarily complete (see [8, Corollary 2.2.2]). Clearly,

every TSI Polish group is also CLI. All compact or abelian Polish groups are TSI, and all locally compact Polish groups are CLI (see [8, Exercises 2.1.8 and 2.2.5]).

Given two equivalence relations E and F on two sets X and Y respectively, we say a map $f : X \rightarrow Y$ is an (E, F) -homomorphism if

$$xEy \Rightarrow f(x)Ff(y)$$

for any $x, y \in X$. Moreover, f is called a reduction of E to F if

$$xEy \iff f(x)Ff(y)$$

for all $x, y \in X$. In particular, if both X, Y are Polish spaces, we say E is Borel reducible to F , denoted by $E \leq_B F$, if there exists a Borel reduction of E to F . We write $E \sim_B F$ if both $E \leq_B F$ and $F \leq_B E$ hold; and write $E <_B F$ if $E \leq_B F$ and $F \not\leq_B E$. We refer to [8] for background on Borel reducibility.

We recall some benchmark equivalence relations in the research of Borel reducibility. The equivalence relation E_0 on 2^ω is defined as

$$xE_0y \iff \exists m \forall n > m (x(n) = y(n)).$$

If E is an equivalence relation on a Polish space X , we define an equivalence relation E^ω on X^ω as

$$xE^\omega y \iff \forall n (x(n)Ey(n)).$$

The equivalence relation \mathbb{R}^ω / c_0 on \mathbb{R}^ω is defined as

$$x\mathbb{R}^\omega / c_0 y \iff \lim_n x(n) - y(n) = 0.$$

Now we recall the definition of equivalence relations induced by Polish groups as below, and list some relevant notions and results that will be repeatedly used throughout the rest of this article.

DEFINITION 2.1 [5, Definition 3.1]. Let G be a Polish group. We define an equivalence relation $E(G)$ on G^ω as: for $x, y \in G^\omega$,

$$xE(G)y \iff \lim_n x(n)y(n)^{-1} \text{ converges.}$$

We say $E(G)$ is the equivalence relation induced by G . Moreover, we define $c(G) = \{x \in G^\omega : \lim_n x(n) \text{ converges}\}$. Then we have

$$xE(G)y \iff xy^{-1} \in c(G) \iff c(G)x = c(G)y.$$

For TSI Polish groups G , it is more convenient to take the following $E_*(G)$ as research object than $E(G)$.

DEFINITION 2.2 [5, Definition 6.1]. Let G be a Polish group. We define an equivalence relation $E_*(G)$ on G^ω as: for $x, y \in G^\omega$,

$$xE_*(G)y \iff \lim_n x(0)x(1) \cdots x(n)y(n)^{-1} \cdots y(1)^{-1}y(0)^{-1} \text{ converges.}$$

It is trivial that $E(G) \sim_B E_*(G)$ (see [6, Proposition 2.2]). In this article, we will use these two equivalence relations interchangeably without further explanation.

For brevity, we define

$$(x, y)|_m^n = x(m) \cdots x(n)y(n)^{-1} \cdots y(m)^{-1}$$

for $x, y \in G^\omega$ and $m \leq n$. It is clear that

$$xE_*(G)y \iff \lim_n (x, y)|_0^n \text{ converges.}$$

Let G be a TSI Polish group and d a compatible complete two-sided invariant metric on G . Define for the sake of brevity

$$d(x, y)|_{[m, n+1]} = d(x, y)|_{[m, n]} = d(x(m) \cdots x(n), y(m) \cdots y(n))$$

for $x, y \in G^\omega$ and $m \leq n$. Clearly, we have $d(x, y)|_{[m, n]} = d(1_G, (x, y)|_m^n)$.

PROPOSITION 2.3 [5, Proposition 3.4]. *Let G, H be two Polish groups. If G is topologically isomorphic to a closed subgroup of H , then $E(G) \leq_B E(H)$.*

LEMMA 2.4 [5, Lemma 6.2]. *Let G be a TSI Polish group, d a compatible complete two-sided invariant metric on G .*

(1) For $g_0, \dots, g_n, h_0, \dots, h_n \in G$, we have

$$d(g_0 \cdots g_n, h_0 \cdots h_n) = d(g_0 \cdots g_n h_n^{-1} \cdots h_0^{-1}, 1_G) \leq \sum_{k=0}^n d(g_k, h_k).$$

(2) For $x, y \in G^\omega$, we have

$$xE_*(G)y \iff \limsup_m \sup_{n>m} d(x, y)|_{[m, n]} = 0.$$

(3) For $x, y \in G^\omega$, if $xE_*(G)y$, then $\lim_n d(x(n), y(n)) = 0$.

Now we recall the notion of additive reduction and a powerful lemma which converts a Borel reduction to an additive reduction.

DEFINITION 2.5 (Farah [7]).

(1) A map $\psi : \prod_n X_n \rightarrow \prod_n X'_n$ is *additive* if there exist $0 = l_0 < l_1 < \dots < l_j < \dots$ and maps $T_j : X_j \rightarrow \prod_{n \in [l_j, l_{j+1})} X'_n$ such that, for $x \in \prod_n X_n$,

$$\psi(x) = T_0(x(0)) \wedge T_1(x(1)) \wedge T_2(x(2)) \wedge \dots$$

(2) Let E and F be two equivalence relations on $\prod_n X_n$ and $\prod_n X'_n$ respectively, we say that E is *additive reducible* to F , denote by $E \leq_A F$, if there exists an additive reduction of E to F .

Let E be an equivalence relation on $\prod_n X_n$, and let $I \subseteq \omega$ be infinite. Fix an element $w \in \prod_{n \notin I} X_n$. For $x \in \prod_{n \in I} X_n$, define $x \oplus w \in \prod_n X_n$ as: $(x \oplus w)(n) = x(n)$ for $n \in I$ and $(x \oplus w)(n) = w(n)$ for $n \notin I$. We define $E|_I^w$ on $\prod_{n \in I} X_n$ as: for $x, y \in \prod_{n \in I} X_n$,

$$xE|_I^w y \iff (x \oplus w)E(y \oplus w).$$

Let (F_n) be a sequence of finite sets. A special equivalence relation $E_0(\prod_n F_n)$ defined as: for $x, y \in \prod_n F_n$,

$$xE_0\left(\prod_n F_n\right)y \iff \exists m \forall n > m (x(n) = y(n)).$$

LEMMA 2.6 [5, Lemma 6.9]. *Let E be a Borel equivalence relation on $\prod_n F_n$ with $E_0(\prod_n F_n) \subseteq E$, where all F_n are finite sets. Let H be a TSI Polish group. If $E \leq_B E_*(H)$, then there exist an infinite set $I \subseteq \omega$ and a $w \in \prod_{n \notin I} F_n$ such that $E|_I^w \leq_A E_*(H)$. In other words, there are natural numbers $0 = n_0 < n_1 < n_2 < \dots$ with $I = \{n_j : j \in \omega\}$, $0 = l_0 < l_1 < l_2 < \dots$, maps $T_{n_j} : F_{n_j} \rightarrow H^{l_{j+1}-l_j}$, and $\psi : \prod_{n \in I} F_n \rightarrow H^\omega$ with*

$$\psi(x) = T_{n_0}(x(n_0)) \frown T_{n_1}(x(n_1)) \frown T_{n_2}(x(n_2)) \frown \dots,$$

such that ψ is an additive reduction of $E|_I^w$ to $E_*(H)$.

The following notions and results are not directly presented in [5], but ideas of them have already appeared in it.

DEFINITION 2.7. Given two sets X, Y and a map $S : X \rightarrow Y$, we define two maps $S^\omega : X^\omega \rightarrow Y^\omega$ and $S^\# : X^\omega \rightarrow Y^\omega \times X^\omega$ as: for $x \in X^\omega$ and $n \in \omega$,

$$S^\omega(x)(n) = S(x(n)),$$

$$S^\#(x) = (S^\omega(x), x).$$

For any metric space (M, d) , recall that $E(M; 0)$ is an equivalence relation on M^ω (see [3, Definition 3.2]) defined as: for $x, y \in M^\omega$,

$$xE(M; 0)y \iff \lim_n d(x(n), y(n)) = 0.$$

Note that, for any Polish group G , the equivalence relation $E(G; 0)$ is independent of any choice of left-invariant compatible metric d on G , since $d(x(n), y(n)) \rightarrow 0$ iff $x(n)^{-1}y(n) \rightarrow 1_G$.

PROPOSITION 2.8. *Let G, H be two TSI Polish groups, and let $S : G \rightarrow H$. Then the following are equivalent:*

- (1) *For all $x, y \in G^\omega$, if $\lim_n x(n)^{-1}y(n) = 1_G$, then*

$$xE_*(G)y \iff S^\omega(x)E_*(H)S^\omega(y).$$

- (2) *$S^\#$ is a reduction of $E_*(G)$ to $E_*(H) \times E(G; 0)$.*

PROOF. This is an easy corollary of Lemma 2.4(3). ⊢

LEMMA 2.9. *Let G, H be two TSI Polish groups such that $E_*(G) \leq_B E_*(H) \times E(G; 0)$. If the interval $[0, 1]$ can be embedded into H , or $E(G; 0) \leq_B E(H; 0)$ holds, then we have $E_*(G) \leq_B E_*(H)$.*

PROOF. By [3, Theorem 3.4(ii)], we have $E(G; 0) \leq_B E([0, 1]; 0)$.

Let $f : [0, 1] \rightarrow H$ be an embedding. By the uniform continuity of f and $f^{-1} : f([0, 1]) \rightarrow [0, 1]$, we see that $E([0, 1]; 0) \sim_B E(f([0, 1]); 0) \leq_B E(H; 0)$. So $E_*(G) \leq_B E_*(H) \times E(H; 0)$.

Finally, by [5, Lemma 6.3], we obtain $E_*(H) \times E(H; 0) \leq_B E_*(H)$. ⊢

§3. Non-TSI vs TSI Polish groups. In this section, we show that we can use Borel reducibility among equivalence relations induced by Polish groups to characterize the class of TSI Polish groups.

Let G be a CLI Polish group, H a TSI Polish group. Suppose that d'_G is a compatible complete left-invariant metric on G and d_H is a compatible complete two-sided invariant metric on H . Put $d_G(g, g') = d'_G(g^{-1}, (g')^{-1})$ for $g, g' \in G$. It is trivial to check that d_G is a compatible complete right-invariant metric on G .

Assume that $E_*(G) \leq_B E_*(H)$. Let (F_n) be a sequence of finite subsets of G such that

- (i) $1_G \in F_n = F_n^{-1}$;
- (ii) $F_{n-1}^n \subseteq F_n$; and
- (iii) $\bigcup_n F_n$ is dense in G .

Denote by E the restriction of $E_*(G)$ on $\prod_n F_n$. By Lemma 2.6, there exist a $w \in \prod_{n \notin I} F_n$ and natural numbers $0 = n_0 < n_1 < n_2 < \dots$ with $I = \{n_j : j \in \omega\}$, $0 = l_0 < l_1 < l_2 < \dots$, maps $T_{n_j} : F_{n_j} \rightarrow H^{l_{j+1}-l_j}$, and $\psi : \prod_{n \in I} F_n \rightarrow H^\omega$ with

$$\psi(x) = T_{n_0}(x(n_0)) \wedge T_{n_1}(x(n_1)) \wedge T_{n_2}(x(n_2)) \wedge \dots,$$

such that ψ is an additive reduction of $E|_I^w$ to $E_*(H)$.

Put $u_{n_0} = 1_G$, and for $j > 0$, let

$$u_{n_j} = w(n_{j-1} + 1) \cdots w(n_j - 1).$$

By (i) and (ii), $u_{n_j}^{-1} \in F_{n_j}$ for each $j \in \omega$.

Define $x_0 \in \prod_{n \in I} F_n$ as

$$x_0(n_j) = u_{n_j}^{-1} \quad (\forall j \in \omega).$$

We may assume that $\psi(x_0) = 1_{H^\omega}$, otherwise we can replace ψ with the following ψ' : for $x \in \prod_{n \in I} F_n$ and $k \in \omega$,

$$\psi'(x)(k) = \psi(x_0)(0) \cdots \psi(x_0)(k-1) \psi(x)(k) \psi(x_0)(k)^{-1} \cdots \psi(x_0)(0)^{-1}.$$

Clearly, $(\psi(x), \psi(y))|_0^k = (\psi'(x), \psi'(y))|_0^k$, so we have

$$\psi(x)E_*(H)\psi(y) \iff \psi'(x)E_*(H)\psi'(y)$$

holds for $x, y \in \prod_{n \in I} F_n$. Hence ψ' is an additive reduction of $E|_I^w$ to $E_*(H)$ with $\psi'(x_0) = 1_{H^\omega}$, as desired.

For $s = (h_0, \dots, h_{l-1})$ and $t = (h'_0, \dots, h'_{l-1})$ in H^l , we define

$$d_H^\infty(s, t) = \max_{0 \leq i \leq m < l} d_H(h_i \cdots h_m, h'_i \cdots h'_m).$$

For any integer $j > 0$ and $g \in F_{n_{j-1}}$, it is worth noting that

$$u_{n_j}^{-1}g = w(n_j - 1)^{-1} \cdots w(n_{j-1} + 1)^{-1}g \in F_{n_{j-1}}^{n_j} \subseteq F_{n_j}.$$

LEMMA 3.1. *For any $q \in \omega$, there exists a $\delta_q > 0$ such that*

$$\forall^\infty n \in I \forall g, g' \in F_{n-1} (d_G(g, g') < \delta_q \Rightarrow d_H^\infty(T_n(u_n^{-1}g), T_n(u_n^{-1}g')) < 2^{-q}).$$

PROOF. If not, then there exist an $\varepsilon_0 > 0$, a strictly increasing sequence (j_k) , and $g_k, g'_k \in F_{n_{j_k-1}}$ for each k , such that

$$d_G(g_k, g'_k) < 2^{-(k+1)}, \quad d_H^\infty(T_{n_{j_k}}(u_{n_{j_k}}^{-1} g_k), T_{n_{j_k}}(u_{n_{j_k}}^{-1} g'_k)) \geq \varepsilon_0.$$

Since d_G is right-invariant, we have $\lim_k g'_k g_k^{-1} = 1_G$. We shall inductively find a strictly increasing sequence of natural numbers $k_0 < k_1 < \dots$ so that

$$d_G(g'_{k_0} \cdots g'_{k_{p-1}} g_{k_{p-1}}^{-1} \cdots g_{k_0}^{-1}, g'_{k_0} \cdots g'_{k_p} g_{k_p}^{-1} \cdots g_{k_0}^{-1}) < 2^{-p}$$

for each integer $p > 0$. Put $k_0 = 0$ first. Assume that k_0, \dots, k_{p-1} have been found. By the continuity of group operations, there is some $\delta > 0$ such that, for $g, g' \in G$ with $d_G(1_G, g'g^{-1}) < \delta$, we have

$$d_G(g'_{k_0} \cdots g'_{k_{p-1}} 1_G g_{k_{p-1}}^{-1} \cdots g_{k_0}^{-1}, g'_{k_0} \cdots g'_{k_{p-1}} g' g^{-1} g_{k_{p-1}}^{-1} \cdots g_{k_0}^{-1}) < 2^{-p}.$$

Then we can find a $k_p > k_{p-1}$ large enough so that $d_G(1_G, g'_{k_p} g_{k_p}^{-1}) < \delta$. This completes the induction. It follows that, for $m > p$, we have

$$d_G(g'_{k_0} \cdots g'_{k_p} g_{k_p}^{-1} \cdots g_{k_0}^{-1}, g'_{k_0} \cdots g'_{k_m} g_{k_m}^{-1} \cdots g_{k_0}^{-1}) < \sum_{i=p+1}^m 2^{-i} < 2^{-p}.$$

Since d_G is complete, we get that

$$\lim_p g'_{k_0} \cdots g'_{k_p} g_{k_p}^{-1} \cdots g_{k_0}^{-1} \text{ converges.}$$

For each $n \in I$, put

$$x(n) = \begin{cases} u_n^{-1} g'_{k_p}, & n = n_{j_{k_p}}, \\ u_n^{-1}, & \text{otherwise,} \end{cases} \quad y(n) = \begin{cases} u_n^{-1} g_{k_p}, & n = n_{j_{k_p}}, \\ u_n^{-1}, & \text{otherwise.} \end{cases}$$

For any $n \geq n_{j_{k_0}}$, let p_n be the largest p such that $n_{j_{k_p}} \leq n$. Note that

$$(x \oplus w, y \oplus w)|_0^n = g'_{k_0} \cdots g'_{k_{p_n}} g_{k_{p_n}}^{-1} \cdots g_{k_0}^{-1},$$

so $(x \oplus w)E_*(G)(y \oplus w)$. But for each $p \in \omega$, if $k = k_p$, then

$$\max_{l_{j_k} \leq i \leq m < l_{j_{k+1}}} d_H(\psi(x), \psi(y))|_{[i,m]} = d_H^\infty(T_{n_{j_k}}(u_{n_{j_k}}^{-1} g'_k), T_{n_{j_k}}(u_{n_{j_k}}^{-1} g_k)) \geq \varepsilon_0.$$

So $\psi(x)E_*(H)\psi(y)$ fails, contradicting that ψ is a reduction. -1

DEFINITION 3.2. For any $g \in \bigcup_n F_n$, if $g \in F_n$ for some $n < n_j$, then $u_{n_j}^{-1}g \in F_{n_j}$, so $T_{n_j}(u_{n_j}^{-1}g) \in H^{l_{j+1}-l_j}$. This allows us to define

$$S_{n_j}(g) = T_{n_j}(u_{n_j}^{-1}g)(0) \cdots T_{n_j}(u_{n_j}^{-1}g)(l_{j+1} - l_j - 1) \in H.$$

Recall that $\psi(x_0) = 1_{H^\omega}$, this implies that

$$S_{n_j}(1_G) = T_{n_j}(u_{n_j}^{-1})(0) \cdots T_{n_j}(u_{n_j}^{-1})(l_{j+1} - l_j - 1) = 1_H$$

for all $j \in \omega$. It is clear that, for $g, g' \in F_n$ with $n < n_j$,

$$d_H(S_{n_j}(g), S_{n_j}(g')) \leq d_H^\infty(T_{n_j}(u_{n_j}^{-1}g), T_{n_j}(u_{n_j}^{-1}g')).$$

LEMMA 3.3. Let $x, y \in \prod_{n \in I} F_n$ such that $\lim_j d_G(x(n_j), y(n_j)) = 0$ and $x(n_j), y(n_j) \in F_{n_j-1}$ for all $j > 0$. Then

$$\psi(x')E_*(H)\psi(y') \iff \lim_j (\psi(x'), \psi(y'))|_0^{l_j-1} \text{ converges,}$$

where $x'(n) = u_n^{-1}x(n), y'(n) = u_n^{-1}y(n)$ for all $n \in I$.

PROOF. If $\psi(x')E_*(H)\psi(y')$, then $\lim_p (\psi(x'), \psi(y'))|_0^p$ converges. In particular, $\lim_j (\psi(x'), \psi(y'))|_0^{l_j-1}$ converges.

Conversely, suppose $\lim_j (\psi(x'), \psi(y'))|_0^{l_j-1}$ converges. For any $q \in \omega$, by Lemma 3.1, there is a $\delta_q \in \omega$ such that

$$\forall^\infty n \in I \forall g, g' \in F_{n-1} (d_G(g, g') < \delta_q \Rightarrow d_H^\infty(T_n(u_n^{-1}g), T_n(u_n^{-1}g')) < 2^{-q}).$$

Since $\lim_j d_G(x(n_j), y(n_j)) = 0$ and $x(n_j), y(n_j) \in F_{n_j-1}$ for any integer $j > 0$, there exists a $j_0 \in \omega$ such that

$$\forall j > j_0 (d_H^\infty(T_{n_j}(u_{n_j}^{-1}x(n_j)), T_{n_j}(u_{n_j}^{-1}y(n_j))) < 2^{-q}).$$

Then for any $j, k \in \omega$ with $j > j_0$ and $l_j \leq k < l_{j+1}$, since d_H is two-sided invariant, we have

$$\begin{aligned} d_H((\psi(x'), \psi(y'))|_0^{l_j-1}, (\psi(x'), \psi(y'))|_0^k) &= d_H(\psi(x'), \psi(y'))|_{[l_j, k]} \\ &\leq d_H^\infty(T_{n_j}(x'(n_j)), T_{n_j}(y'(n_j))) \\ &< 2^{-q}. \end{aligned}$$

So by the convergency of $\lim_j (\psi(x'), \psi(y'))|_0^{l_j-1}$, we obtain that

$$\lim_k (\psi(x'), \psi(y'))|_0^k \text{ converges,}$$

and hence $\psi(x')E_*(H)\psi(y')$. -1

The following proposition may be well known. We provide a proof here for the convenience of readers.

PROPOSITION 3.4 (folklore). Let G be a Polish group. Then G is TSI iff for all sequences (g_p) and (g'_p) in G , we have

$$\lim_p g_p g'_p = 1_G \iff \lim_p g'_p g_p = 1_G.$$

PROOF. Suppose G is TSI, and let d_G be a compatible complete two-sided invariant metric on G . Then we have $d_G(1_G, g_p g'_p) = d_G(g_p^{-1}, g'_p) = d_G(1_G, g'_p g_p)$, so $\lim_p g_p g'_p = 1_G \iff \lim_p g'_p g_p = 1_G$.

On the other hand, fix an open neighborhood basis (V_n) of 1_G . We claim that

$$\forall n \exists m_n \forall g \in G (gV_{m_n}g^{-1} \subseteq V_n).$$

If not, there exist an n_0 and two sequences $(g_p), (h_p)$ in G such that $\lim_p h_p = 1_G$, but $g_p h_p g_p^{-1} \notin V_{n_0}$ for each $p \in \omega$. Put $g'_p = h_p g_p^{-1}$. It is clear that $\lim_p g'_p g_p = 1_G$, but $g_p g'_p \not\rightarrow 1_G$, which is a contradiction.

Now we put $U_n = \bigcup_{g \in G} gV_{m_n}g^{-1}$. Note that $U_n \subseteq V_n$ for each $n \in \omega$. Therefore, (U_n) is also an open neighborhood basis of 1_G with $gU_n g^{-1} = U_n$ for all $g \in G$ and $n \in \omega$. By [8, Exercise 2.1.4], G admits a compatible two-sided invariant metric. Hence G is TSI (see [2, Corollary 1.2.2]). \dashv

THEOREM 3.5. *Let G, H be two Polish groups. If H is TSI but G is not, then $E(G) \not\leq_B E(H)$.*

PROOF. Assume toward a contradiction that $E(G) \leq_B E(H)$. By [5, Theorem 4.3], it suffices to consider the case that G is CLI. Let d_G be a compatible complete right-invariant metric on G and d_H a compatible complete two-sided invariant metric on H .

We use the notation defined earlier in this section.

Since G is not TSI, by Proposition 3.4, there are two sequences $(g_p), (g'_p)$ of elements of G such that $\lim_p g'_p g_p = 1_G$, but $g_p g'_p \not\rightarrow 1_G$. Since d_G is right-invariant, $\lim_p d_G(g'_p, g_p^{-1}) = 0$. By transferring to a subsequence, we may assume that, there is a $\delta > 0$ such that $\inf_p d_G(1_G, g_p g'_p) > \delta$.

Since $\bigcup_n F_n$ is dense, by perturbation, we may assume that $\{g_p, g'_p : p \in \omega\} \subseteq \bigcup_n F_n$. By Lemma 3.1 and Definition 3.2, we can find two sequence $(p(i))$ and $(q(i))$ of natural numbers so that

- (i) $g_{p(i)}, g'_{p(i)} \in F_{n_{q(i)-1}}$;
- (ii) $d_H(S_{n_{q(i)+1}}(g'_{p(i)}), S_{n_{q(i)+1}}(g_{p(i)}^{-1})) < 2^{-i}$; and
- (iii) for each $i \in \omega$, $0 < p(i) < q(i) < q(i) + 1 < p(i + 1)$.

For each $n \in I$, define

$$x(n) = \begin{cases} u_n^{-1} g_{p(i)}, & n = n_{q(i)}, \\ u_n^{-1} g'_{p(i)}, & n = n_{q(i)+1}, \\ u_n^{-1}, & \text{otherwise,} \end{cases} \quad y(n) = \begin{cases} u_n^{-1} g_{p(i)}, & n = n_{q(i)}, \\ u_n^{-1} g_{p(i)}^{-1}, & n = n_{q(i)+1}, \\ u_n^{-1}, & \text{otherwise.} \end{cases}$$

For $j \in \omega$, by letting $h'_j = S_{n_j}(u_{n_j} x(n_j))$ and $h_j = S_{n_j}(u_{n_j} y(n_j))$, we have

$$(\psi(x), \psi(y))|_0^{j+1-1} = h'_0 \cdots h'_j h_j^{-1} \cdots h_0^{-1}.$$

For $i \in \omega$ and $m > j > q(i) + 1$, it follows from (ii) and Lemma 2.4(1) that

$$d_H(h'_j \cdots h'_m, h_j \cdots h_m) < \sum_{k>i} 2^{-k} < 2^{-i}.$$

Now by Lemma 2.4(2),

$$\lim_j (\psi(x), \psi(y))|_0^{j+1-1} = \lim_j h'_0 \cdots h'_j h_j^{-1} \cdots h_0^{-1} \text{ converges.}$$

Note that $\lim_i d_G(g'_{p(i)}, g_{p(i)}^{-1}) = 0$. So by Lemma 3.3, we have

$$\psi(x)E_*(H)\psi(y),$$

and hence $(x \oplus w)E_*(G)(y \oplus w)$. In particular,

$$\lim_i (x \oplus w, y \oplus w)_0^{n_{q(i)+1}} = \lim_i g_{p(0)}g'_{p(0)} \cdots g_{p(i)}g'_{p(i)} \text{ converges.}$$

Therefore, it follows that

$$\lim_i g_{p(i)}g'_{p(i)} = \lim_i (g_{p(0)}g'_{p(0)} \cdots g_{p(i-1)}g'_{p(i-1)})^{-1} g_{p(0)}g'_{p(0)} \cdots g_{p(i)}g'_{p(i)} = 1_G.$$

We obtain a contradiction. ⊥

§4. A gap between E_0^ω and \mathbb{R}^ω/c_0 . In this section, we prove that there is NO Polish group G satisfying that

$$E_0^\omega <_B E(G) \leq_B \mathbb{R}^\omega/c_0.$$

To do so, we need the following notions. The author [4] defined equivalence relations $E(X, (x_n))$ and \mathbb{R}^ω/cs . Let (x_n) be a sequence in a Banach space X . We define an equivalence relation $E(X, (x_n))$ on \mathbb{R}^ω as: for any $a, b \in \mathbb{R}^\omega$,

$$(a, b) \in E(X, (x_n)) \iff \sum_n (a(n) - b(n))x_n \text{ converges in } X.$$

The equivalence relation \mathbb{R}^ω/cs is defined as: for any $a, b \in \mathbb{R}^\omega$,

$$(a, b) \in \mathbb{R}^\omega/cs \iff \sum_n (a(n) - b(n)) \text{ converges in } \mathbb{R}.$$

The following lemma is an easy corollary of [4, Lemma 4.2].

LEMMA 4.1. *Let E be a Borel equivalence relation on $\prod_n F_n$ with $E_0(\prod_n F_n) \subseteq E$, where all (F_n) are finite sets. If $E \leq_B \mathbb{R}^\omega/c_0$, then there exist an infinite set $I \subseteq \omega$ and a $w \in \prod_{n \notin I} F_n$ such that $E|_I^w \leq_A \mathbb{R}^\omega/c_0$.*

PROOF. Let $e_n = (0, \dots, 0, \overset{n}{1}, 0, \dots)$ for each integer $n > 0$. Then (e_n) is the canonical Schauder basis of c_0 . Note that $E(c_0, (e_n)) = \mathbb{R}^\omega/c_0$. By applying [4, Lemma 4.2] to $E(c_0, (e_n))$, we conclude the proof. ⊥

THEOREM 4.2. *Let G be a Polish group. If $E(G) \leq_B \mathbb{R}^\omega/c_0$, then G is TSI non-archimedean.*

PROOF. Suppose $E(G) \leq_B \mathbb{R}^\omega/c_0$. Note that $\mathbb{R}^\omega/c_0 <_B \mathbb{R}^\omega/cs = E_*(\mathbb{R})$ (see [4, Theorem 5.9(i)]). By Theorem 3.5, G is also TSI. Let d_G be a compatible complete two-sided invariant metric on G . Put $V_k = \{g \in G : d_G(1_G, g) < 2^{-k}\}$.

Assume for contradiction that G is not non-archimedean. Then there is a $K \in \omega$ such that V_K contains no open subgroups of G . It is clear that $V_k = V_k^{-1}$ and $\bigcup_m V_k^m$ is an open subgroup which is not contained in V_K . So there exists $m_k > 0$ with $V_k^{m_k} \not\subseteq V_K$ for each k . We can find $g_{k,0}, \dots, g_{k,m_k-1} \in V_k$ such that $g_{k,0} \cdots g_{k,m_k-1} \notin V_K$. Put $h_j = g_{k,i}$ for $j = \sum_{l < k} m_l + i$ with $i < m_k$. Then we have $\lim_j h_j = 1_G$. By Cauchy Criterion,

$$\lim_j h_0 \cdots h_j \text{ diverges.}$$

Assume that $E_*(G) \leq_B \mathbb{R}^\omega / c_0$. Let (F_n) be a sequence of finite subsets of G such that

- (i) $1_G, h_0 \in F_n = F_n^{-1}$;
- (ii) $F_{n-1} \subseteq F_n$; and
- (iii) for $n > 0, h_n \in F_{n-1}$.

Denote by E the restriction of $E_*(G)$ on $\prod_n F_n$. By Lemma 4.1, there exist an infinite set $I \subseteq \omega$ and a $w \in \prod_{n \notin I} F_n$ such that $E|_I^w \leq_A \mathbb{R}^\omega / c_0$. So there are natural numbers $0 = n_0 < n_1 < n_2 < \dots$ with $I = \{n_j : j \in \omega\}, 0 = l_0 < l_1 < l_2 < \dots$, maps $T_{n_j} : F_{n_j} \rightarrow \mathbb{R}^{l_{j+1}-l_j}$, and $\psi : \prod_{n \in I} F_n \rightarrow \mathbb{R}^\omega$ with

$$\psi(x) = T_{n_0}(x(n_0)) \wedge T_{n_1}(x(n_1)) \wedge T_{n_2}(x(n_2)) \wedge \dots,$$

such that ψ is an additive reduction of $E|_I^w$ to \mathbb{R}^ω / c_0 .

Put $u_{n_0} = 1_G$, and for $j > 0$, let

$$u_{n_j} = w(n_{j-1} + 1) \cdots w(n_j - 1).$$

Assume again for contradiction that there are $\varepsilon_0 > 0$ and natural numbers $0 < j(0) < j(1) < \dots < j(p) < \dots$ such that

$$\max_{0 \leq k < l_{j(p)+1} - l_{j(p)}} |T_{n_{j(p)}}(u_{n_{j(p)}}^{-1} h_{j(p)})(k) - T_{n_{j(p)}}(u_{n_{j(p)}}^{-1})(k)| \geq \varepsilon_0.$$

By $\lim_p h_{j(p)} = 1_G$, we can find natural numbers $p_0 < p_1 < \dots < p_i < \dots$ so that $d_G(1_G, h_{j(p_i)}) < 2^{-i}$ for each $i \in \omega$. Thus $(h_{j(p_0)} \cdots h_{j(p_i)})$ is d_G -Cauchy, so $\lim_i h_{j(p_0)} \cdots h_{j(p_i)}$ converges. For each $n \in I$, put $x_0(n) = u_n^{-1}$ and

$$y'_0(n) = \begin{cases} u_n^{-1} h_{j(p_i)}, & n = n_{j(p_i)}, \\ u_n^{-1}, & \text{otherwise.} \end{cases}$$

For any $n > n_{j(p_0)}$, let i_n be the largest i with $n_{j(p_i)} \leq n$. Then

$$(y'_0 \oplus w, x_0 \oplus w)|_0^n = h_{j(p_0)} \cdots h_{j(p_{i_n})},$$

and thus $(y'_0 \oplus w)E_*(G)(x_0 \oplus w)$. Note that for all $i \in \omega$,

$$\max_{l_{j(p_i)} \leq k < l_{j(p_i)+1}} |\psi(y'_0)(k) - \psi(x_0)(k)| \geq \varepsilon_0.$$

So $\psi(y'_0)\mathbb{R}^\omega / c_0 \psi(x_0)$ fails, which is a contradiction.

Therefore, for all $\varepsilon > 0$, we have

$$\forall^\infty j \forall k \in [0, l_{j+1} - l_j) (|T_{n_j}(u_{n_j}^{-1} h_j)(k) - T_{n_j}(u_{n_j}^{-1})(k)| < \varepsilon).$$

For each $n \in I$, put $y_0(n) = u_n^{-1} h_j$, where $n = n_j$. Then we have

$$\psi(y_0)\mathbb{R}^\omega / c_0 \psi(x_0),$$

and hence $(y_0 \oplus w)E_*(G)(x_0 \oplus w)$. In particular,

$$\lim_j (y_0 \oplus w, x_0 \oplus w)|_0^{n_j} = \lim_j h_0 \cdots h_j \text{ converges.}$$

This leads to a contradiction. ⊖

Building upon previous results, we establish the following theorem.

THEOREM 4.3. *Let G be a Polish group. Then the following are equivalent:*

- (1) G is TSI non-archimedean;
- (2) $E(G) \leq_B E_0^\omega$; and
- (3) $E(G) \leq_B \mathbb{R}^\omega/c_0$.

In particular, $E(G) \sim_B E_0^\omega$ iff G is TSI uncountable non-archimedean.

PROOF. (1) \Rightarrow (2) follows from [5, Theorem 3.5]. (2) \Rightarrow (3) follows from [8, Lemma 8.5.3]. (3) \Rightarrow (1) follows from Theorem 4.2.

Again by [5, Theorem 3.5], we see that $E(G) \sim_B E_0^\omega$ iff G is TSI uncountable non-archimedean. ◻

The equivalence relation \mathbb{R}^ω/c on \mathbb{R}^ω is defined as

$$(a, b) \in \mathbb{R}^\omega/c \iff \lim_n a(n) - b(n) \text{ exists.}$$

Note that $\mathbb{R}^\omega/c_0 <_B \mathbb{R}^\omega/c_s = E_*(\mathbb{R})$ (see [4, Theorem 5.9(i)]) and $E(\mathbb{R}) = \mathbb{R}^\omega/c$. The results of this section make the following question very interesting.

QUESTION 4.4. *For any Polish group G , does it hold that*

$$E(G) <_B E(\mathbb{R}) \iff E(G) \leq_B E_0^\omega?$$

§5. The Pre-rigid Theorem on TSI Polish groups. In this section, we prove a highly technical theorem, namely the Pre-rigid Theorem, which will serve as the foundation for the subsequent sections.

We say that a topological group G has no small subgroups, or is NSS, if there exists an open subset $V \ni 1_G$ in G such that no non-trivial subgroup of G is contained in V .

To generalize [5, Theorem 6.13], we introduce the following definition.

DEFINITION 5.1. A Polish group G is called *strongly NSS* if there exists an open set $V \ni 1_G$ in G such that

$$\forall (g_n) \in G^\omega (g_n \rightarrow 1_G \Rightarrow \exists n_0 < \dots < n_k (g_{n_0} \dots g_{n_k} \notin V)),$$

where the set V is called an *unenclosed set* of G .

PROPOSITION 5.2. *Let G be a Polish group. The following hold:*

- (1) if G is strongly NSS, then G is NSS; and
- (2) if G is locally compact, then G is strongly NSS iff G is NSS.

PROOF. (1) Let V be an unenclosed set of G . We have

$$\forall g \in G (g \neq 1_G \Rightarrow \exists m (g^m \notin V)).$$

Thus V contains no non-trivial subgroups of G , so G is NSS.

(2) By (1), we only need to prove another direction. Suppose that G is NSS and locally compact. Let $V \ni 1_G$ be an open subset of G such that \bar{V} is compact and contains no non-trivial subgroups of G . Then

$$\forall g \in \bar{V} (g \neq 1_G \Rightarrow \exists m (g^m \notin \bar{V})).$$

We claim that V is an unenclosed set of G . Fix a $(g_n) \in G^\omega$ with $g_n \not\rightarrow 1_G$. By the definition of unenclosed set, it suffices to consider the case that $(g_n) \in V^\omega$. Since \bar{V} is compact, there exist a subsequence (g_{n_i}) of (g_n) and an element $1_G \neq h \in \bar{V}$ such that $\lim_i g_{n_i} = h$. Then we can find an $m \in \omega$ with $h^m \notin \bar{V}$. By the continuity of group operations, there are $i_0 < i_1 < \dots < i_{m-1}$ such that $g_{n_{i_0}} g_{n_{i_1}} \dots g_{n_{i_{m-1}}} \notin \bar{V}$. Therefore G is strongly NSS. \dashv

The addition group \mathbb{R}^ω with the product topology is not strongly NSS because it is not NSS. Let (e_n) be the canonical basis in c_0 . For any $r > 0$, the sequence $(\frac{r}{2}e_n)$ witnesses that the open set $\{x \in c_0 : \|x\| < r\}$ is not an unenclosed set of c_0 . On the other hand, for any $x \in c_0$ with $\|x\| \neq 0$, we have $\lim_n \|nx\| \rightarrow +\infty$. Thus the Banach space c_0 is NSS, but not strongly NSS, under the addition operation.

Now, we are ready to prove the Pre-rigid Theorem. Let us recall that the definitions of the maps $(\pi_m^S)^\omega$ and $S^\#$ appearing in the following theorem can be found in Definition 2.7.

THEOREM 5.3 (Pre-rigid Theorem). *Let G be a TSI Polish group, and let H be a closed subgroup of the product of a sequence of TSI strongly NSS Polish groups (H_m) . If $E(G) \leq_B E(H)$, then for each $m \in \omega$, there exist an open subgroup W_m of G and a continuous map $\pi_m^S : W_m \rightarrow H_m$ with $\pi_m^S(1_G) = 1_{H_m}$ satisfying the following:*

- (i) *The map $(\pi_m^S)^\omega : W_m^\omega \rightarrow H_m^\omega$ is an $(E_*(W_m), E_*(H_m))$ -homomorphism.*
- (ii) *Define $S : W = \bigcap_m W_m \rightarrow H$ as: for $g \in W$,*

$$S(g) = (\pi_0^S(g), \pi_1^S(g), \dots, \pi_m^S(g), \dots),$$

then the map $S^\# : W^\omega \rightarrow H^\omega \times W^\omega$ is a reduction of $E_(W)$ to $E_*(H) \times E(W; 0)$.*

Moreover, the converse holds if $G = W$ and the interval $[0, 1]$ can be embedded into H .

PROOF. Let $d_G, d_{H_m} \leq 1$ be compatible complete two-sided invariant metrics on G and H_m respectively. A compatible complete two-sided invariant metric d_H on H is defined as

$$d_H(h, h') = \sum_{m=0}^\infty 2^{-m} d_{H_m}(h(m), h'(m))$$

for $h, h' \in H$. Assume that $E_*(G) \leq_B E_*(H)$.

We use the notation defined in the arguments before Lemma 3.1 and before Lemma 3.3.

For $m \in \omega$, let $\pi_m : \prod_n H_n \rightarrow H_m$ be the canonical projection map, so $\pi_m(h) = h(m)$ for each h . It is clear that every π_m is a continuous homomorphism.

It is worth noting that

$$\forall h, h' \in H (d_{H_m}(\pi_m(h), \pi_m(h')) \leq 2^m d_H(h, h')).$$

For $m \in \omega$, let X_m be the set of all $g \in \bigcup_n F_n$ such that:

$$\forall \delta > 0 \exists j_\delta (g \in F_{n_{j_\delta-1}} \wedge \forall j' > j \geq j_\delta (d_{H_m}(1_{H_m}, \pi_m(S_{n_{j'}}(g)S_{n_{j'}}(g^{-1}))) < \delta)).$$

Note that $S_{n_j}(1_G) = 1_H$ for all $j \in \omega$, so $1_G \in X_m$.

CLAIM 1. For any $m \in \omega$, there is an $r_m > 0$ such that

$$\forall g, g' \in \bigcup_n F_n ((d_G(g, g') < r_m \wedge g \in X_m) \Rightarrow g' \in X_m).$$

PROOF OF CLAIM 1. If not, there are an $m_0 \in \omega$ and two sequences $(g_i), (g'_i) \in \bigcup_n F_n$ such that

- (i) $d_G(g_i, g'_i) < 2^{-i}$; and
- (ii) $g_i \in X_{m_0}$ and $g'_i \notin X_{m_0}$.

Note that H_{m_0} is strongly NSS. Let open set $V \ni 1_{H_{m_0}}$ be an unenclosed set of H_{m_0} . Put $p(-1, 2M_{-1} - 1) = 0$. For any $i \in \omega$, we will find an $M_i \in \omega$ and natural numbers $0 < p(i, 0) < p(i, 1) < \dots < p(i, 2M_i - 1)$ so that

- (1) $p(i - 1, 2M_{i-1} - 1) < p(i, 0)$;
- (2) $g_i, g'_i \in F_{n_{p(i,0)-1}}$;
- (3) for $i \in \omega$,

$$\pi_{m_0}(S_{n_{p(i,0)}}(g'_i)S_{n_{p(i,1)}}((g'_i)^{-1}) \dots S_{n_{p(i,2M_i-2)}}(g'_i)S_{n_{p(i,2M_i-1)}}((g'_i)^{-1})) \notin V;$$

- (4) for $i \in \omega$,

$$\sum_{l=0}^{M_i-1} d_{H_{m_0}}(1_{H_{m_0}}, \pi_{m_0}(S_{n_{p(i,2l)}}(g_i)S_{n_{p(i,2l+1)}}(g_i^{-1}))) < 2^{-i}.$$

Let us begin with $i = 0$. Since $g'_0 \notin X_{m_0}$, there are a $\delta_0 > 0$ and natural numbers $p(0) < p(1) < \dots < p(j) < \dots$ such that $g'_0 \in F_{n_{p(0)-1}}$ and for each $j \in \omega$,

$$d_{H_{m_0}}(1_{H_{m_0}}, \pi_{m_0}(S_{n_{p(2j)}}(g'_0)S_{n_{p(2j+1)}}((g'_0)^{-1}))) \geq \delta_0.$$

By the definition of X_{m_0} and $g_0 \in X_{m_0}$, there exists a strictly increasing sequence (j_k) of natural numbers such that $g_0 \in F_{n_{p(j_0)-1}}$ and for each $k \in \omega$,

$$d_{H_{m_0}}(1_{H_{m_0}}, \pi_{m_0}(S_{n_{p(2j_k)}}(g_0)S_{n_{p(2j_k+1)}}(g_0^{-1}))) < 2^{-(k+1)}.$$

Let $h \in H_{m_0}^\omega$ be such that

$$h(k) = \pi_{m_0}(S_{n_{p(2j_k)}}(g'_0)S_{n_{p(2j_k+1)}}((g'_0)^{-1}))$$

for each $k \in \omega$. Then $h(k) \rightarrow 1_{H_{m_0}}$. Since V is an unenclosed set of H_{m_0} , there are finitely many natural numbers $k_0 < k_1 < \dots < k_q$ such that $h(k_0) \dots h(k_q) \notin V$, i.e.,

$$\pi_{m_0}(S_{n_{p(2j_{k_0})}}(g'_0)S_{n_{p(2j_{k_0+1})}}((g'_0)^{-1}) \dots S_{n_{p(2j_{k_q})}}(g'_0)S_{n_{p(2j_{k_q+1})}}((g'_0)^{-1})) \notin V.$$

Now we put $M_0 = q + 1$, $p(0, 2l) = p(2j_{k_l})$, and $p(0, 2l + 1) = p(2j_{k_l} + 1)$ for all $l < q + 1$. Then we have $g_0, g'_0 \in F_{n_{p(j_0)-1}} \subseteq F_{n_{p(0,0)-1}}$, and

$$\sum_{l=0}^q d_{H_{m_0}}(1_{H_{m_0}}, \pi_{m_0}(S_{n_{p(0,2l)}}(g_0)S_{n_{p(0,2l+1)}}(g_0^{-1}))) < \sum_{l=0}^q 2^{-(k_l+1)} < 1.$$

We can see that clauses (1)–(4) hold for $i = 0$.

Now assume that M_0, \dots, M_i and $p(0, 0) < \dots < p(0, 2M_0 - 1) < \dots < p(i, 0) < \dots < p(i, 2M_i - 1)$ have been defined. Using similar arguments as those presented in the preceding paragraph, pick a $p' > p(i, 2M_i - 1)$ with $g_{i+1}, g'_{i+1} \in F_{n_{p'-1}}$, then we can find an $M_{i+1} \in \omega$ and natural numbers $p' \leq p(i + 1, 0) < \dots < p(i + 1, 2M_{i+1} - 1)$ so that clauses (1)–(4) hold.

For each $n \in I$, define

$$x(n) = \begin{cases} u_n^{-1} g'_i, & n = n_{p(i,2k)}, 0 \leq k < M_i, \\ u_n^{-1} (g'_i)^{-1}, & n = n_{p(i,2k+1)}, 0 \leq k < M_i, \\ u_n^{-1}, & \text{otherwise,} \end{cases}$$

$$y(n) = \begin{cases} u_n^{-1} g_i, & n = n_{p(i,2k)}, 0 \leq k < M_i, \\ u_n^{-1} g_i^{-1}, & n = n_{p(i,2k+1)}, 0 \leq k < M_i, \\ u_n^{-1}, & \text{otherwise.} \end{cases}$$

Then for $q \in \omega$, the following equality holds:

$$(x \oplus w, y \oplus w)|_0^q = \begin{cases} g'_i g_i^{-1}, & n_{p(i,2k)} \leq q < n_{p(i,2k+1)}, 0 \leq k < M_i, \\ 1_G, & \text{otherwise.} \end{cases}$$

By (i), we have

$$\lim_q (x \oplus w, y \oplus w)|_0^q = 1_G.$$

So $(x \oplus w)E_*(G)(y \oplus w)$, and thus $\psi(x)E_*(H)\psi(y)$ holds. It follows from Lemma 2.4(2) that

$$\lim_i d_H(\psi(x), \psi(y))|_{[I_{p(i,0)}, I_{p(i,2M_i-1)+1})} = 0.$$

For each $j \in \omega$, let $h'_j = S_{n_j}(u_{n_j}x(n_j))$ and $h_j = S_{n_j}(u_{n_j}y(n_j))$. Note that $S_{n_j}(1_G) = 1_H$ for all $j > 0$, so

$$(\psi(x), \psi(y))|_{I_{p(i,0)}}^{I_{p(i,2M_i-1)+1}} = h'_{p(i,0)} h'_{p(i,1)} \cdots h'_{p(i,2M_i-1)} (h_{p(i,0)} h_{p(i,1)} \cdots h_{p(i,2M_i-1)})^{-1}.$$

Now we have

$$\lim_i d_H(h'_{p(i,0)} h'_{p(i,1)} \cdots h'_{p(i,2M_i-1)}, h_{p(i,0)} h_{p(i,1)} \cdots h_{p(i,2M_i-1)}) = 0,$$

and hence

$$\lim_i h'_{p(i,0)} h'_{p(i,1)} \cdots h'_{p(i,2M_i-1)} (h_{p(i,0)} h_{p(i,1)} \cdots h_{p(i,2M_i-1)})^{-1} = 1_H.$$

Let $V_1 \ni 1_{H_{m_0}}$ be an open subset of H_{m_0} with $V_1^2 \subseteq V$. Since π_{m_0} is a continuous homomorphism, there exists an $i_0 \in \omega$ such that

$$\pi_{m_0}(h'_{p(i,0)} h'_{p(i,1)} \cdots h'_{p(i,2M_i-1)}) \pi_{m_0}(h_{p(i,0)} h_{p(i,1)} \cdots h_{p(i,2M_i-1)})^{-1} \in V_1$$

holds for any $i > i_0$. By (4), we have

$$\lim_i d_{H_{m_0}}(1_{H_{m_0}}, \pi_{m_0}(h_{p(i,0)} h_{p(i,1)} \cdots h_{p(i,2M_i-1)})) \leq \lim_i 2^{-i} = 0.$$

Therefore, for i large enough,

$$\pi_{m_0}(h_{p(i,0)}h_{p(i,1)} \cdots h_{p(i,2M_i-1)}) \in V_1.$$

It follows that

$$\pi_{m_0}(h'_{p(i,0)}h'_{p(i,1)} \cdots h'_{p(i,2M_i-1)}) \in V,$$

i.e.,

$$\pi_{m_0}(S_{n_{p(i,0)}}(g'_i)S_{n_{p(i,1)}}((g'_i)^{-1}) \cdots S_{n_{p(i,2M_i-2)}}(g'_i)S_{n_{p(i,2M_i-1)}}((g'_i)^{-1})) \in V,$$

contradicting (3). ⊥

For any $m \in \omega$, let $V_m = \{g \in G : d_G(1_G, g) < r_m\}$ and $W_m = \bigcup_i V_m^i$. Note that $V_m^{-1} = V_m$. It is clear that W_m is an open, and thus a clopen subgroup of G . We claim that $W_m \cap \bigcup_n F_n \subseteq X_m$. For $i = 0$, note that $1_G \in X_m$, so Claim 1 gives that $V_m \cap \bigcup_n F_n \subseteq X_m$. Assume that $V_m^i \cap \bigcup_n F_n \subseteq X_m$. Let $h \in V_m^{i+1} \cap \bigcup_n F_n$ with $h = gg'$, where $g \in V_m^i$ and $g' \in V_m$. Note that $\bigcup_n F_n$ is a dense subgroup of G . We can find $\hat{g}, \hat{g}' \in \bigcup_n F_n$ such that $d_G(h, \hat{g}\hat{g}') < r_m$ with $\hat{g} \in V_m^i$ and $\hat{g}' \in V_m$. Since $d_G(\hat{g}, \hat{g}\hat{g}') = d_G(1_G, \hat{g}') < r_m$ and $\hat{g} \in X_m$, we have $\hat{g}\hat{g}' \in X_m$, and thus $h \in X_m$. So $W_m \cap \bigcup_n F_n \subseteq X_m$.

Let $g \in W_m \cap \bigcup_n F_n \subseteq X_m$. Pick a $j_g \in \omega$ with $g \in F_{n_{j_g-1}}$. For any $\delta > 0$, by the definition of X_m , there exists a $j_\delta \geq j_g$ such that

$$\forall j' > j \geq j_\delta (d_{H_m}(1_{H_m}, \pi_m(S_{n_j}(g)S_{n_{j'}}(g^{-1}))) < \delta).$$

Let $j, k \in \omega$ with $j_\delta < j < k$. Fix a $k' > k$. Since π_m is a homomorphism, we have

$$d_{H_m}(1_{H_m}, \pi_m(S_{n_j}(g))\pi_m(S_{n_{k'}}(g^{-1}))) < \delta,$$

$$d_{H_m}(1_{H_m}, \pi_m(S_{n_k}(g))\pi_m(S_{n_{k'}}(g^{-1}))) < \delta.$$

So $d_{H_m}(\pi_m(S_{n_j}(g)), \pi_m(S_{n_k}(g))) < 2\delta$. Thus $(\pi_m(S_{n_j}(g)))$ is a Cauchy sequence in H_m . By the completeness of d_{H_m} , we can define

$$\pi_m^S(g) = \lim_j \pi_m(S_{n_j}(g)) \in H_m.$$

Note that for $g, g' \in F_n$ with $n < n_j$,

$$d_H(S_{n_j}(g), S_{n_j}(g')) \leq d_H^\infty(T_{n_j}(u_{n_j}^{-1}g), T_{n_j}(u_{n_j}^{-1}g')),$$

and

$$\forall h, h' \in H (d_{H_m}(\pi_m(h), \pi_m(h')) \leq 2^m d_H(h, h')).$$

By Lemma 3.1, π_m^S is uniformly continuous on $W_m \cap \bigcup_n F_n$, which can be uniquely extended to a uniformly continuous map from W_m to H_m , still denoted by π_m^S . Put $W = \bigcap_m W_m$, for any $g \in W$, we let

$$S(g) = (\pi_0^S(g), \pi_1^S(g), \dots, \pi_m^S(g), \dots).$$

We claim that $S(g) \in H$. Indeed, fix a $g \in W$ and an arbitrary neighborhood O of $S(g)$, then there are $\varepsilon > 0$ and natural number L such that

$$\left\{ h \in \prod_m H_m : \forall m \leq L (d_{H_m}(\pi_m(h), \pi_m^S(g)) < \varepsilon) \right\} \subseteq O.$$

Clearly $g \in \bigcap_{m \leq L} W_m$. Since $\bigcup_n F_n$ is dense in G , by the continuity of π_m^S , we can find some $g' \in \bigcap_{m \leq L} W_m \cap \bigcup_n F_n$ so that $d_{H_m}(\pi_m^S(g'), \pi_m^S(g)) < \varepsilon$ for $m \leq L$. Note that $\pi_m^S(g') = \lim_j \pi_m(S_{n_j}(g')) \in H_m$ holds for any $m \leq L$. So there is a $j' \in \omega$ such that $S_{n_{j'}}(g') \in H \cap O$. Thus $S(g) \in H$ by the fact that H is closed in $\prod_m H_m$. The map S is a continuous map from W to H .

It is worth noting that $S(1_G) = 1_H$ and W is a closed subgroup of G .

Now we prove clause (ii) first:

CLAIM 2. *The map $S^\#$ is a reduction of $E_*(W)$ to $E_*(H) \times E(W; 0)$.*

PROOF OF CLAIM 2. Let $x, y \in W^\omega$. Note that $x E_*(G) y \iff x E_*(W) y$ as W is a closed subgroup of G . By Proposition 2.8, we only need to show that

$$x E_*(G) y \iff S^\omega(x) E_*(H) S^\omega(y)$$

holds whenever $\lim_p d_G(x(p), y(p)) = 0$. Since $\bigcup_n F_n$ is dense, by the definitions of d_H and S , for each $p \in \omega$, we can find a sufficiently large $j(p) \in \omega$ and $g_p, g'_p \in \bigcup_n F_n$ so that

- (1) $0 < j(0) < j(1) < \dots < j(p) < \dots$;
- (2) $g_p, g'_p \in F_{n_{j(p)}-1} \cap \bigcap_{m \leq p} W_m$;
- (3) $d_G(x(p), g'_p) < 2^{-p}$ and $d_G(y(p), g_p) < 2^{-p}$; and
- (4) for all $m \leq p$,

$$\begin{aligned} d_{H_m}(\pi_m(S(x(p))), \pi_m(S_{n_{j(p)}}(g'_p))) &< 2^{-(p+1)}, \\ d_{H_m}(\pi_m(S(y(p))), \pi_m(S_{n_{j(p)}}(g_p))) &< 2^{-(p+1)}. \end{aligned}$$

Now we define, for each $n \in I$,

$$\hat{x}(n) = \begin{cases} u_n^{-1} g'_p, & n = n_{j(p)}, \\ u_n^{-1}, & \text{otherwise,} \end{cases} \quad \hat{y}(n) = \begin{cases} u_n^{-1} g_p, & n = n_{j(p)}, \\ u_n^{-1}, & \text{otherwise.} \end{cases}$$

For any $i < k$, by (3) and Lemma 2.4(1),

$$d_G((x, y)|_i^k, g'_i \dots g'_k g_k^{-1} \dots g_i^{-1}) < \sum_{q=i}^k 2^{-q+1} < 2^{-i+2}.$$

Then by Lemma 2.4(2),

$$x E_*(G) y \iff \lim_p g'_0 \dots g'_p g_p^{-1} \dots g_0^{-1} \text{ converges.}$$

For any $k > n_{j(0)}$, let p_k be the largest p with $n_{j(p)} \leq k$. Note that

$$(\hat{x} \oplus w, \hat{y} \oplus w)|_0^k = g'_0 \dots g'_{p_k} g_{p_k}^{-1} \dots g_0^{-1}.$$

It follows that

$$xE_*(G)y \iff (\hat{x} \oplus w)E_*(G)(\hat{y} \oplus w) \iff \psi(\hat{x})E_*(H)\psi(\hat{y}).$$

By (4), we have

$$\begin{aligned} d_H(S(x(p)), S_{n_{j(p)}}(g'_p)) &= \sum_{m=0}^\infty 2^{-m} d_{H_m}(\pi_m(S(x(p))), \pi_m(S_{n_{j(p)}}(g'_p))) \\ &< \sum_{m=0}^p 2^{-(m+p+1)} + \sum_{m=p+1}^\infty 2^{-m} \\ &< 2^{-(p-1)}, \\ d_H(S(y(p)), S_{n_{j(p)}}(g_p)) &= \sum_{m=0}^\infty 2^{-m} d_{H_m}(\pi_m(S(y(p))), \pi_m(S_{n_{j(p)}}(g_p))) \\ &< \sum_{m=0}^p 2^{-(m+p+1)} + \sum_{m=p+1}^\infty 2^{-m} \\ &< 2^{-(p-1)}. \end{aligned}$$

Note that $S_{n_j}(1_G) = 1_H$ for all $j \in \omega$, by similar arguments as above, we see that

$$\begin{aligned} &S^\omega(x)E_*(H)S^\omega(y) \\ \iff &\lim_p S_{n_{j(0)}}(g'_0) \cdots S_{n_{j(p)}}(g'_p) S_{n_{j(p)}}(g_p)^{-1} \cdots S_{n_{j(0)}}(g_0)^{-1} \text{ converges.} \end{aligned}$$

For any $i > j(0)$, let p_i be the largest p with $j(p) \leq i$. Then we have

$$(\psi(\hat{x}), \psi(\hat{y}))|_0^{i+1-1} = S_{n_{j(0)}}(g'_0) \cdots S_{n_{j(p_i)}}(g'_{p_i}) S_{n_{j(p_i)}}(g_{p_i})^{-1} \cdots S_{n_{j(0)}}(g_0)^{-1}.$$

This implies that

$$\lim_i (\psi(\hat{x}), \psi(\hat{y}))|_0^{i+1-1} \text{ converges} \iff S^\omega(x)E_*(H)S^\omega(y).$$

Note that $\lim_p d_G(g'_p, g_p) = 0$, it follows from Lemma 3.3 that

$$\lim_i (\psi(\hat{x}), \psi(\hat{y}))|_0^{i+1-1} \text{ converges} \iff \psi(\hat{x})E_*(H)\psi(\hat{y}).$$

So $S^\omega(x)E_*(H)S^\omega(y) \iff \psi(\hat{x})E_*(H)\psi(\hat{y}) \iff xE_*(G)y$. □

Subsequently, we prove clause (i) as follows:

CLAIM 3. For $m \in \omega$, if $x, y \in W_m^\omega$, then

$$xE_*(G)y \implies (\pi_m^S)^\omega(x)E_*(H_m)(\pi_m^S)^\omega(y).$$

PROOF OF CLAIM 3. Let $x, y \in W_m^\omega$ with $x E_*(G) y$. Then for each $p \in \omega$, we can find a $j(p) \in \omega$ and $g_p, g'_p \in W_m \cap \bigcup_n F_n$ so that

- (1) $0 < j(0) < j(1) < \cdots < j(p) < \cdots$;
- (2) $g_p, g'_p \in F_{n_{j(p)}-1} \cap W_m$;
- (3) $d_G(x(p), g'_p) < 2^{-p}$ and $d_G(y(p), g_p) < 2^{-p}$; and
- (4) we have

$$\begin{aligned} d_{H_m}(\pi_m^S(x(p)), \pi_m(S_{n_{j(p)}}(g'_p))) &< 2^{-p}, \\ d_{H_m}(\pi_m^S(y(p)), \pi_m(S_{n_{j(p)}}(g_p))) &< 2^{-p}. \end{aligned}$$

Now define, for each $n \in I$,

$$\hat{x}(n) = \begin{cases} u_n^{-1} g'_p, & n = n_{j(p)}, \\ u_n^{-1}, & \text{otherwise,} \end{cases} \quad \hat{y}(n) = \begin{cases} u_n^{-1} g_p, & n = n_{j(p)}, \\ u_n^{-1}, & \text{otherwise.} \end{cases}$$

Following a similar approach to the proof of Claim 2, we obtain

$$xE_*(G)y \iff (\hat{x} \oplus w)E_*(G)(\hat{y} \oplus w) \iff \psi(\hat{x})E_*(H)\psi(\hat{y}).$$

So we have

$$\lim_p (\psi(\hat{x}), \psi(\hat{y}))|_0^{l_{j(p)+1}^{-1}} \text{ converges,}$$

and thus

$$\lim_p S_{n_{j(0)}}(g'_0) \cdots S_{n_{j(p)}}(g'_p) S_{n_{j(p)}}(g_p)^{-1} \cdots S_{n_{j(0)}}(g_0)^{-1} \text{ converges.}$$

Since π_m is a continuous homomorphism, we obtain

$$\lim_p \pi_m(S_{n_{j(0)}}(g'_0) \cdots S_{n_{j(p)}}(g'_p) S_{n_{j(p)}}(g_p)^{-1} \cdots S_{n_{j(0)}}(g_0)^{-1}) \text{ converges.}$$

Following again the similar arguments as in the proof of Claim 2, we get $(\pi_m^S)^\omega(x)E_*(H_m)(\pi_m^S)^\omega(y)$. ⊣

It follows from Claims 2 and 3 that W_m, π_m^S, W , and S are as desired. Finally, by Lemma 2.9, we can conclude that the converse is also true if $G = W$ and if the interval $[0, 1]$ can be embedded into H . ⊣

REMARK 5.4. Recall that G_0 is the connected component of 1_G in G . Since each W_m in the preceding theorem is an open subgroup of G , it is also clopen in G . So $G_0 \subseteq W_m$ as it is connected, and thus $G_0 \subseteq W$.

From the proof of the preceding theorem, it follows that W can be chosen to be clopen when H is TSI strongly NSS. Similarly, by [5, Theorem 6.13], W can be also clopen when H is locally compact, and W can be chosen to be $W = G$ when H is compact.

§6. Rigid theorems. In this section, we use the Pre-rigid Theorem to prove several Rigid theorems for various classes of TSI Polish groups.

LEMMA 6.1. *Let G, H be two Polish groups and $S : G \rightarrow H$ a continuous map with $S(1_G) = 1_H$. Suppose H is NSS. Then the following are equivalent:*

- (1) *There exists an open subgroup W of G such that the map $S \upharpoonright W : W \rightarrow H$ is a continuous homomorphism.*
- (2) *There exists an open subgroup W' of G such that the map S^ω is an $(E_*(W'), E_*(H))$ -homomorphism.*

PROOF. (1) \Rightarrow (2). Let W be an open subgroup of G such that $S \upharpoonright W : W \rightarrow H$ is a continuous homomorphism. Then

$$\forall x, y \in W^\omega \forall g \in W (\lim_n (x, y)|_0^n = g \Rightarrow \lim_n (S^\omega(x), S^\omega(y))|_0^n = S(g)).$$

Since W is also closed, we see that S^ω is an $(E_*(W), E_*(H))$ -homomorphism. Hence $W' = W$ is as required.

(2) \Rightarrow (1). Let W' be an open subgroup of G such that the map S^ω is an $(E_*(W'), E_*(H))$ -homomorphism, and let d_G, d_H be compatible left-invariant metrics on G and H respectively.

Define $X = \{g \in W' : S(g^{-1}) = S(g)^{-1}\}$. Then $X \neq \emptyset$, since $1_G \in X$.

CLAIM 1. *There exists an $r_0 > 0$ such that*

$$\forall g, g' \in W' ((d_G(g, g') < r_0 \wedge g \in X) \Rightarrow g' \in X).$$

PROOF OF CLAIM 1. If not, then we can find two sequences (g_q) and (g'_q) in W' so that

- (i) $g_q \in X$ and $g'_q \notin X$;
- (ii) $\lim_q (g'_q)^{-1}g_q = 1_G$.

Let $h_q = S((g'_q)^{-1})S(g'_q)$, then $h_q \neq 1_H$ for $q \in \omega$. Since H has no small subgroups, there exists some $D > 0$ such that, for each $q \in \omega$, we can find an $m_q > 0$ with $d_H(h_q^{m_q}, 1_H) \geq D$.

Put $M_{-1} = 0$ and $M_q = m_0 + \dots + m_q$ for $q \in \omega$. For each $p \in \omega$, define

$$x(p) = \begin{cases} (g'_q)^{-1}, & p = 2(M_{q-1} + i), 0 \leq i < m_q, \\ g'_q, & p = 2(M_{q-1} + i) + 1, 0 \leq i < m_q, \end{cases}$$

$$y(p) = \begin{cases} g_q^{-1}, & p = 2(M_{q-1} + i), 0 \leq i < m_q, \\ g_q, & p = 2(M_{q-1} + i) + 1, 0 \leq i < m_q. \end{cases}$$

For any $k \in \omega$, let q_k be the largest q such that $M_{q-1} \leq k$. Clearly,

$$(x, y)|_0^{2k+1} = 1_G, \quad (x, y)|_0^{2k} = (g'_{q_k})^{-1}g_{q_k}.$$

By (ii), we have $x E_*(G) y$. Since S^ω is an $(E_*(G), E_*(H))$ -homomorphism, $S^\omega(x) E_*(H) S^\omega(y)$ holds, i.e.,

$$\lim_k (S^\omega(x), S^\omega(y))|_0^k \text{ converges.}$$

Then by Lemma 2.4(2) and $g_q \in X$, we have

$$\lim_q h_q^{m_q} = \lim_q (S^\omega(x), S^\omega(y))|_{2M_{q-1}}^{2M_q-1} = 1_H.$$

A contradiction! ⊥

Put $V = \{g \in W' : d_G(1_G, g) < r_0\}$. Note that $V = V^{-1}$. Define $W_1 = \bigcup_i V^i$, which is an open subgroup of W' . We claim that $W_1 \subseteq X$. Note that $1_G \in X$, so by Claim 1, we have $V \subseteq X$. Assume that $V^i \subseteq X$. For any $g = g_1 g_2 \in V^{i+1}$ with $g_1 \in V^i$ and $g_2 \in V$, we note that $g_1 \in X$ and $d_G(g_1, g) = d_G(g_1, g_1 g_2) = d_G(1_G, g_2) < r_0$. Again by Claim 1, we have $g \in X$. This shows that $W_1 \subseteq X$.

CLAIM 2. *There exists $0 < r_1 < r_0$ such that*

$$\forall g, g' \in W_1 (d_G(1_G, g) < r_1 \Rightarrow S(gg') = S(g)S(g')).$$

PROOF OF CLAIM 2. If not, there are two sequences $(g_q), (g'_q)$ in W_1 such that $\lim_q g_q = 1_G$ and $S(g_q g'_q) \neq S(g_q)S(g'_q)$ for $q \in \omega$.

For each $q \in \omega$, let $h_q = S(g_q g'_q)S((g'_q)^{-1})S(g_q^{-1})$. It follows from $W_1 \subseteq X$ that

$$h_q = S(g_q g'_q)S(g'_q)^{-1}S(g_q)^{-1} \neq 1_H.$$

Since H has no small subgroups, there exists some $D > 0$ such that, for each $q \in \omega$, we can find an $m_q \in \omega$ with $d_H(h_q^{m_q}, 1_H) \geq D$.

Put $M_{-1} = 0$ and $M_q = m_0 + \dots + m_q$ for $q \in \omega$. For each $p \in \omega$, define

$$x(p) = \begin{cases} g_q g'_q, & p = 3(M_{q-1} + i), 0 \leq i < m_q, \\ (g'_q)^{-1}, & p = 3(M_{q-1} + i) + 1, 0 \leq i < m_q, \\ g_q^{-1}, & p = 3(M_{q-1} + i) + 2, 0 \leq i < m_q, \end{cases}$$

$$y(p) = \begin{cases} g'_q, & p = 3(M_{q-1} + i), 0 \leq i < m_q, \\ (g'_q)^{-1}, & p = 3(M_{q-1} + i) + 1, 0 \leq i < m_q, \\ 1_G, & p = 3(M_{q-1} + i) + 2, 0 \leq i < m_q. \end{cases}$$

For any $k \in \omega$, let q_k be the largest q such that $M_{q-1} \leq k$. Clearly,

$$(x, y)|_0^{3k} = (x, y)|_0^{3k+1} = g_{q_k}, \quad (x, y)|_0^{3k+2} = 1_G.$$

Note that $\lim_q g_q = 1_G$, so $x E_*(G) y$ holds. Since S^ω is an $(E_*(G), E_*(H))$ -homomorphism, $S^\omega(x) E_*(H) S^\omega(y)$ holds, i.e.,

$$\lim_k (S^\omega(x), S^\omega(y))|_0^k \text{ converges.}$$

By Lemma 2.4(2) and $g'_q \in W_1 \subseteq X$, we have $S(g'_q)S((g'_q)^{-1}) = 1_H$. So

$$\lim_q h_q^{m_q} = \lim_q (S^\omega(x), S^\omega(y))|_{3M_{q-1}}^{3M_{q-1}} = 1_H.$$

A contradiction! ⊥

Finally, let $V_1 = \{g \in W' : d_G(1_G, g) < r_1\}$ and $W = \bigcup_i V_1^i$. Note that $V_1 = V_1^{-1}$ and $V_1 \subseteq V$, so W is an open subgroup of W_1 . For any $g, g' \in W$, we shall check that $S(gg') = S(g)S(g')$. There are $g_0, g_1, \dots, g_m \in V_1$ such that $g = g_0 g_1 \dots g_m$. Note that $W \subseteq W_1$, so by Claim 2, we have

$$S(gg') = S(g_0)S(g_1 \dots g_m g') = S(g_0) \dots S(g_m)S(g') = S(g)S(g').$$

Then W is as required. ⊥

Consider a Polish group G and a sequence (g_n) in G . Recall that (g_n) is ι -Cauchy if it is d -cauchy for some compatible left-invariant metric d on G . This definition is independent of the choice of d (see [1, Proposition 3.B.1]).

Let G, H be two Polish groups and $\varphi : G \rightarrow H$ a continuous homomorphism. We define $\text{IPC}(\varphi)$ as the set of all $x \in G^\omega$ that satisfies:

$$(x(0) \dots x(p)) \text{ is } \iota\text{-Cauchy} \iff (\varphi(x(0)) \dots \varphi(x(p))) \text{ is } \iota\text{-Cauchy.}$$

LEMMA 6.2. *Let G, H be two Polish groups and $\varphi : G \rightarrow H$ a continuous homomorphism. If $x \in \text{IPC}(\varphi)$ for any $x \in G^\omega$ with $\lim_p x(p) = 1_G$, then $\ker(\varphi)$ is non-archimedean.*

Moreover, if $\varphi(G)$ is closed in H , then the converse is also true.

PROOF. Let d_G, d_H be compatible left-invariant metrics on G and H respectively. Define $V_k = \{g \in \ker(\varphi) : d_G(1_G, g) < 2^{-k}\}$.

First, suppose $x \in \text{IPC}(\varphi)$ for any $x \in G^\omega$ with $\lim_p x(p) = 1_G$.

Assume toward a contradiction that $\ker(\varphi)$ is not non-archimedean. Then there exists a $K \in \omega$ such that V_K contains no open subgroups of $\ker(\varphi)$. Since $\bigcup_m V_k^m$ is an open subgroup of $\ker(\varphi)$, we have $\bigcup_m V_k^m \not\subseteq V_K$. Thus for each $k \in \omega$, we can find an $m_k \in \omega$ and $g_{k,0}, \dots, g_{k,m_k-1} \in V_k$ such that $g_{k,0}g_{k,1} \dots g_{k,m_k-1} \notin V_K$.

Put $M_{-1} = 0$ and $M_k = m_0 + \dots + m_k$ for $k \in \omega$. Let $x \in G^\omega$ be defined as $x(p) = g_{k,i}$ for $p = M_{k-1} + i$ with $0 \leq i < m_k$. Note that $\lim_p x(p) = 1_G$, so $x \in \text{IPC}(\varphi)$. Since d_G is left-invariant, we have

$$\forall k (d_G(x(0) \cdots x(M_{k-1} - 1), x(0) \cdots x(M_k - 1)) \geq 2^{-K}).$$

Thus $(x(0) \cdots x(p))$ is not ι -Cauchy. But $(\varphi(x(0)) \cdots \varphi(x(p)))$ is ι -Cauchy since $\varphi(x(p)) = 1_H$ for all $p \in \omega$, contradicting that $x \in \text{IPC}(\varphi)$.

Now assume that $\varphi(G)$ is closed in H and $\ker(\varphi)$ is non-archimedean. Let $x \in G^\omega$ with $\lim_p x(p) = 1_G$. We prove $x \in \text{IPC}(\varphi)$ as follows:

On the one hand, suppose that $(x(0) \cdots x(p))$ is ι -Cauchy. For any $\varepsilon > 0$, since φ is continuous, there is a $\delta > 0$ such that

$$\forall g \in G (d_G(1_G, g) < \delta \Rightarrow d_H(1_H, \varphi(g)) < \varepsilon).$$

Since $(x(0) \cdots x(p))$ is ι -Cauchy, we can find an $N \in \omega$ such that

$$\forall m \geq n > N (d_G(1_G, x(n) \cdots x(m)) < \delta).$$

Thus we have that, for any $m \geq n > N$,

$$d_H(1_H, \varphi(x(n)) \cdots \varphi(x(m))) = d_H(1_H, \varphi(x(n) \cdots x(m))) < \varepsilon.$$

This shows that $(\varphi(x(0)) \cdots \varphi(x(p)))$ is ι -Cauchy.

On the other hand, suppose that $(\varphi(x(0)) \cdots \varphi(x(p)))$ is ι -Cauchy. Let (W_p) be a decreasing neighborhood basis of $1_{\ker(\varphi)}$ such that each W_p is an open subgroup of $\ker(\varphi)$. Let $\tilde{\varphi} : G/\ker(\varphi) \rightarrow H$ be defined as

$$\tilde{\varphi}(\ker(\varphi)g) = \varphi(g).$$

Note that $\varphi(G)$ is closed, so it is also a Polish group under the topology inherited from H . Thus $\tilde{\varphi}$ is a topological isomorphism of $G/\ker(\varphi)$ onto $\varphi(G)$ (see [8, Corollary 2.3.4]). Note that $\mu = \tilde{\varphi}^{-1} \circ \varphi$ is the canonical projection map, where $\mu(g) = \ker(\varphi)g$ for $g \in G$. Since $\tilde{\varphi}$ is a topological isomorphism, we have $(\mu(x(0)) \cdots \mu(x(p)))$ is ι -Cauchy.

Let $d_\mu(\ker(\varphi)g, \ker(\varphi)g') = \inf\{d_G(hg, h'g') : h, h' \in \ker(\varphi)\}$, then d_μ is a compatible left-invariant metric on $G/\ker(\varphi)$ (see [8, Lemma 2.2.8]).

For any $\varepsilon > 0$, we can find $0 < \varepsilon' < \varepsilon$ and $p_0 \in \omega$ satisfying

$$\{g \in \ker(\varphi) : d_G(1_G, g) < 3\varepsilon'\} \subseteq W_{p_0} \subseteq \{g \in \ker(\varphi) : d_G(1_G, g) < \varepsilon\}.$$

Note that $\lim_p x(p) = 1_G$ and $(\mu(x(0)) \cdots \mu(x(p)))$ is ι -Cauchy. Thus there is an $N \in \omega$ such that

$$\begin{aligned} d_G(1_G, x(n)) &< \varepsilon', \\ d_\mu(\mu(x(N)) \cdots x(n), \ker(\varphi)1_H) &< \varepsilon' \end{aligned}$$

for each $n > N$. By the definition of d_μ , there exists some $g_n \in \ker(\varphi)$ for each $n > N$ such that

$$d_G(x(N) \cdots x(n), g_n) < \varepsilon'.$$

For any $n > N$, we have

$$d_G(x(N) \cdots x(n), x(N) \cdots x(n+1)) = d_G(1_G, x(n+1)) < \varepsilon'.$$

All these together imply that $d_G(1_G, g_n^{-1}g_{n+1}) = d_G(g_n, g_{n+1}) < 3\varepsilon'$. So $g_n^{-1}g_{n+1} \in W_{p_0}$. Since W_{p_0} is a subgroup, it follows that $g_n^{-1}g_m \in W_{p_0}$ for any $m \geq n \geq N$. This gives that

$$\begin{aligned} d_G(x(0) \cdots x(n), x(0) \cdots x(m)) &= d_G(x(N) \cdots x(n), x(N) \cdots x(m)) \\ &< 2\varepsilon' + d_G(g_n, g_m) \\ &< 2\varepsilon' + \varepsilon < 3\varepsilon. \end{aligned}$$

Therefore, $(x(0) \cdots x(p))$ is ι -Cauchy. ⊖

PROPOSITION 6.3. *Let G, H be two TSI Polish groups and $\varphi : G \rightarrow H$ a continuous homomorphism. Then the following are equivalent:*

- (1) for $x \in G^\omega$, if $\lim_p x(p) = 1_G$, then $x \in \text{IPC}(\varphi)$; and
- (2) the map $\varphi^\#$ is a reduction of $E_*(G)$ to $E_*(H) \times E(G; 0)$.

PROOF. Since G and H are both TSI, for $x \in G^\omega$, we have

$$\begin{aligned} (x(0) \cdots x(p)) \text{ is } \iota\text{-Cauchy} &\iff xE_*(G)1_{G^\omega}, \\ (\varphi(x(0)) \cdots \varphi(x(p))) \text{ is } \iota\text{-Cauchy} &\iff \lim_p \varphi^\omega(x)E_*(H)1_{H^\omega}. \end{aligned}$$

So (2) \Rightarrow (1) follows from Proposition 2.8. We prove (1) \Rightarrow (2) as follows:

Let d_G be a compatible complete two-sided invariant metric on G .

Given $x, y \in G^\omega$. Suppose $x \in \text{IPC}(\varphi)$. Since φ is a continuous homomorphism, we have $\varphi^\omega(x)E_*(H)\varphi^\omega(y)$ holds and $\lim_p d_G(x(p), y(p)) = 0$.

On the other hand, suppose $\varphi^\omega(x)E_*(H)\varphi^\omega(y)$ holds and $\lim_p d_G(x(p), y(p)) = 0$. Define $z \in G^\omega$ as:

$$z(p) = y(0) \cdots y(p-1)x(p)y(p)^{-1}y(p-1)^{-1} \cdots y(0)^{-1}$$

for $p \in \omega$. Since d_G is two-sided invariant,

$$\lim_p d_G(z(p), 1_G) = \lim_p d_G(x(p), y(p)) = 0.$$

So $z \in \text{IPC}(\varphi)$. Since φ is homomorphism, it holds that

$$\varphi(z(p)) = \varphi(y(0)) \cdots \varphi(y(p-1))\varphi(x(p))\varphi(y(p))^{-1}\varphi(y(p-1))^{-1} \cdots \varphi(y(0))^{-1}.$$

Therefore,

$$\begin{aligned} xE_*(G)y &\iff (z(0) \cdots z(p)) \text{ is } \iota\text{-Cauchy}, \\ \varphi^\omega(x)E_*(H)\varphi^\omega(y) &\iff (\varphi(z(0)) \cdots \varphi(z(p))) \text{ is } \iota\text{-Cauchy}. \end{aligned}$$

So (1) \Rightarrow (2) follows from Proposition 2.8 again. ⊖

Now we are ready to prove the following theorem, which is crucial for the rest of this article.

THEOREM 6.4. *Let G be a TSI Polish group, and let H be a closed subgroup of the product of a sequence of TSI strongly NSS Polish groups (H_m) . If $E(G) \leq_B E(H)$, then there exists a continuous homomorphism $S : W \rightarrow H$ such that $\ker(S)$ is non-archimedean, where $W \supseteq G_0$ is a countable intersection of clopen subgroups in G .*

Moreover, the converse holds if $G = W$, $S(G)$ is closed in H , and the interval $[0, 1]$ can be embedded into H .

PROOF. Suppose $E(G) \leq_B E(H)$. It follows from Theorem 5.3 that, for each $m \in \omega$, there exists an open subgroup W'_m of G and a continuous map $\pi_m^S : W'_m \rightarrow H_m$ with $\pi_m^S(1_G) = 1_{H_m}$ such that (1) $(\pi_m^S)^\omega : (W'_m)^\omega \rightarrow H_m^\omega$ is an $(E_*(W'_m), E_*(H_m))$ -homomorphism; and (2) let $W' = \bigcap_m W'_m$ and define $S : W' \rightarrow H$ as $S(g) = (\pi_0^S(g), \pi_1^S(g), \dots)$, then $S^\#$ is a reduction of $E_*(W')$ to $E_*(H) \times E(W'; 0)$.

Note that H_m is NSS. By Lemma 6.1, there is an open subgroup W_m of W'_m such that $\pi_m^S \upharpoonright W_m$ is a homomorphism. We put $W = \bigcap_m W_m$, and still denote $S \upharpoonright W$ by S for brevity. Then S is a continuous homomorphism. It is clear that $G_0 \subseteq W \subseteq W'$.

As both G and H are TSI, according to Lemma 6.2 and Proposition 6.3, $\ker(S)$ is non-archimedean. Thus, W and S fulfill the requirements.

On the other hand, let S be a continuous homomorphism from G to H such that $\ker(S)$ is non-archimedean. Suppose that $S(G)$ is closed in H and the interval $[0, 1]$ can be embedded into H . Again by Lemma 6.2 and Proposition 6.3, the map $S^\#$ is a reduction of $E_*(G)$ to $E_*(H) \times E(G; 0)$. Finally, it follows from Lemma 2.9 that $E(G) \leq_B E(H)$. ←

REMARK 6.5. Let G be a TSI Polish group and H a TSI strongly NSS Polish group. Suppose that $E(G) \leq_B E(H)$. Then it follows from the proof of Theorem 6.4 that there exist an open subgroup $W \supseteq G_0$ of G and a continuous homomorphism $S : W \rightarrow H$ such that $\ker(S)$ is non-archimedean. Moreover, the map $S^\#$ is a reduction of $E_*(W)$ to $E_*(H) \times E(W; 0)$.

This observation will be crucial in Section 7.

The following is an immediate corollary.

COROLLARY 6.6. *Let G be a TSI Polish group and H a closed subgroup of the product of a sequence of TSI strongly NSS Polish groups. If H is totally disconnected but G is not, then $E(G) \not\leq_B E(H)$.*

PROOF. Assume for contradiction that $E(G) \leq_B E(H)$. By Theorem 6.4, there is a continuous homomorphism $S : G_0 \rightarrow H$ such that $\ker(S)$ is non-archimedean. Note that $S(G_0)$ is also connected, so $S(G_0) \subseteq H_0$. Since H is totally disconnected, we have $S(G_0) = \{1_H\}$. So $G_0 = \ker(S)$, which is connected and non-archimedean. This implies that $G_0 = \{1_G\}$, contradicting that G is not totally disconnected. ←

Note that the converses of Lemma 6.2 and Theorem 6.4 require $S(G)$ to be closed in H . Next we present a lemma in which this requirement can be avoided.

LEMMA 6.7. *Let G, H be two TSI Polish groups. Suppose $S : G \rightarrow H$ is a continuous homomorphism, and $U \ni 1_G$ is an open subset of G such that $S \upharpoonright U : U \rightarrow S(U)$ is a homeomorphism. Then the map $S^\#$ is a reduction of $E_*(G)$ to $E_*(H) \times E(G; 0)$.*

PROOF. Let d_G, d_H be compatible complete two-sided invariant metrics on G and H respectively. By Proposition 2.8, we only need to show that

$$xE_*(G)y \iff S^\omega(x)E_*(H)S^\omega(y)$$

holds whenever $\lim_p d_G(x(p), y(p)) = 0$.

Suppose $xE_*(G)y$ holds. Since $S : G \rightarrow H$ is a continuous homomorphism, it follows trivially that $S^\omega(x)E_*(H)S^\omega(y)$.

On the other hand, suppose $\lim_p d_G(x(p), y(p)) = 0$ and $S^\omega(x)E_*(H)S^\omega(y)$. We prove $x E_*(G)y$ as follows:

If not, by Lemma 2.4(2), $\sup_{q>p} d_G(x, y)|_{[p,q]} \not\rightarrow 0$. So there exist $r > 0$ and two strictly increasing sequences of natural numbers $(p_k), (q_k)$ such that $p_k < q_k$ and $d_G(1_G, (x, y)|_{[p_k, q_k]}^{q_k}) = d_G(x, y)|_{[p_k, q_k]} > r$ for each $k \in \omega$. Let $0 < r_0 < r$ satisfying that $\{g \in G : d_G(1_G, g) < r_0\} \subseteq U$. There exists a $K \in \omega$ such that $d_G(x(p), y(p)) < r_0/2$ for $p > p_K$. For $p > p_k$ and $k > K$, since d_G is two-sided invariant, we have

$$d_G((x, y)|_{p_k}^p, (x, y)|_{p_k}^{p-1}) = d_G(x(p)y(p)^{-1}, 1_G) < r_0/2,$$

and hence

$$d_G(1_G, (x, y)|_{p_k}^p) < d_G(1_G, (x, y)|_{p_k}^{p-1}) + r_0/2.$$

For each $k > K$, we can find $p_k < p'_k < q_k$ such that

$$r_0/2 \leq d_G(1_G, (x, y)|_{p'_k}^{p'_k}) < r_0,$$

so $(x, y)|_{p'_k}^{p'_k} \in U$. By $S^\omega(x)E_*(H)S^\omega(y)$, it is clear that

$$S((x, y)|_{p'_k}^{p'_k}) = (S^\omega(x), S^\omega(y))|_{p'_k}^{p'_k} \rightarrow 1_H.$$

Since $S \upharpoonright U$ is a homeomorphism from U to $S(U)$, we have $(x, y)|_{p'_k}^{p'_k} \rightarrow 1_G$. This contradicts that $d_G(1_G, (x, y)|_{p'_k}^{p'_k}) \geq r_0/2$. ⊥

6.1. Applications on Lie groups, locally compact groups, and pro-Lie groups.

Recall that a *Lie group* is a group which is also a smooth manifold such that the group operations are smooth functions. A topological group is a Lie group iff it is locally compact NSS (see [11, p. 159]). Clearly, a Lie group is Polish iff it is separable iff it has only countably many connected components. Let G be a Lie group, then G_0 is an open normal subgroup of G . For more details on Lie groups, we refer to [19].

A completely metrizable topological group G is called a *pro-Lie group* if every open neighborhood of 1_G contains a normal subgroup N such that G/N is a Lie group (see [9, Definition 1]). For more details on pro-Lie groups, we refer to [9].

Applying Theorem 6.4, we obtain the following result.

THEOREM 6.8. *Let G, H be two TSI Polish groups such that H is a pro-Lie group. If $E(G) \leq_B E(H)$, then there exists a continuous homomorphism $S : G_0 \rightarrow H$ such that $\ker(S)$ is non-archimedean.*

Moreover, the converse holds if G is connected and $S(G)$ is closed in H .

PROOF. Since H is a pro-Lie group, there exist a countable open neighborhood basis (U_m) of 1_H and a sequence of normal subgroups (N_m) of H such that $N_m \subseteq U_m$ and H/N_m is a Lie group for each $m \in \omega$. Note that N_m is closed in H , since H/N_m is Hausdorff. Without loss of generality, assume that $N_m \supseteq N_{m+1}$. We define a map $f : H \rightarrow \prod_m H/N_m$ as $f(h)(m) = hN_m$ for $h \in H, m \in \omega$. By [12, Proposition 2.3], f is a topologically isomorphic embedding and $f(H)$ is closed in $\prod_m H/N_m$.

By [8, Exercise 2.2.8], all H/N_m are separable TSI Lie groups. As mentioned above, all Lie groups are locally compact NSS. By Proposition 5.2(2), all Lie groups are strongly NSS. So the first part of the theorem follows from Theorem 6.4.

For proving the second part of the theorem, by Theorem 6.4, we only need to show that the interval $[0, 1]$ can be embedded into H . Since $S(G)$ is a closed subgroup of H , it is also a pro-Lie Polish group. Since $\ker(S)$ is non-archimedean and G is connected, $\ker(S) \neq G$. So $S(G)$ is non-singleton and connected. Note that, for any non-singleton connected pro-Lie Polish group K , there exists a nontrivial continuous homomorphism $\gamma : \mathbb{R} \rightarrow K$ (see [9, Proposition 19 and Definition 2.6]). All of these together allow us to embed the interval $[0, 1]$ into $S(G)$, and also into H . \dashv

A topological group G is called a *SIN-group* if G admits arbitrarily small invariant identity neighborhoods, or equivalently, G has a neighborhood basis $(U_i)_{i \in I}$ of 1_G such that $gU_i g^{-1} = U_i$ for all $g \in G$ (see [11, Definition 2.1]). It follows from [8, Exercise 2.1.4] that a Polish group is SIN iff it is TSI. By [11, Theorem 3.6], every locally compact TSI Polish group is a pro-Lie group.

We point out that, in [6, Theorem 1.2], groups G and H are required to be abelian with G being compact. Following is a generalized theorem that removes these requirements.

THEOREM 6.9 (Rigid Theorem for locally compact TSI groups). *Let G be a locally compact connected TSI Polish group, H a TSI pro-Lie Polish group. Then $E(G) \leq_B E(H)$ iff there exists a continuous homomorphism $S : G \rightarrow H$ such that $\ker(S)$ is non-archimedean.*

PROOF. (\Rightarrow) . It follows from Theorem 6.8 and $G = G_0$.

(\Leftarrow) . Let $\varphi : G \rightarrow G/\ker(S)$ be the canonical projection. Note that φ is a continuous surjective homomorphism with $\varphi(g) = \ker(S)g$ for $g \in G$. Clearly, $\ker(\varphi) = \ker(S)$ is non-archimedean. By Lemma 6.2 and Proposition 6.3, the map $\varphi^\#$ is a continuous reduction of $E_*(G)$ to $E_*(G/\ker(S)) \times E(G; 0)$.

Let $S^* : G/\ker(S) \rightarrow H$ be the map defined as $S^*(\ker(S)g) = S(g)$ for $g \in G$. It is clear that S^* is a continuous injective homomorphism. Let d_G be a compatible complete two-sided invariant metric on G , and define

$$d^*(\ker(S)g, \ker(S)g') = \inf\{d_G(kg, k'g') : k, k' \in \ker(S)\}.$$

Then d^* is a compatible metric on $G/\ker(S)$ (see [8, Lemma 2.2.8]). Since G is locally compact, we can find some $r > 0$ such that the closure \overline{V} of the open set $V = \{g \in G : d_G(1_G, g) < r\}$ is compact. Let

$$U = \{\ker(S)g : d^*(1_{G/\ker(S)}, \ker(S)g) < r\}.$$

Since d_G is two-sided invariant, from the definition of d^* , we see that $U \subseteq S(V)$. Note that $\overline{U} \subseteq S(\overline{V})$, so \overline{U} is also compact. This implies that $S^* \upharpoonright \overline{U} : \overline{U} \rightarrow S^*(\overline{U})$ is a homeomorphism. By Lemma 6.7, the map $(S^*)^\#$ is a continuous reduction of $E_*(G/\ker(S))$ to $E_*(H) \times E(G/\ker(S); 0)$.

Given $x, y \in G^\omega$ with $\lim_p d_G(x(p), y(p)) = 0$. It follows from Proposition 2.8 that

$$xE_*(G)y \iff \varphi^\omega(x)E_*(G/\ker(S))\varphi^\omega(y).$$

The continuity of φ implies $\lim_p d^*(\varphi(x(p)), \varphi(y(p))) = 0$, so we also have

$$\varphi^\omega(x)E_*(G/\ker(S))\varphi^\omega(y) \iff (S^*)^\omega(\varphi^\omega(x))E_*(H)(S^*)^\omega(\varphi^\omega(y)).$$

Note that $S = S^* \circ \varphi$, so $S^\omega = (S^*)^\omega \circ \varphi^\omega$. It is clear that

$$xE_*(G)y \iff S^\omega(x)E_*(H)S^\omega(y).$$

Thus the map $S^\#$ is a continuous reduction of $E_*(G)$ to $E_*(H) \times E(G; 0)$.

Since G is connected, $\ker(S) \neq G$, and so $\{1_H\} \neq S(G) \subseteq H_0$. So H_0 is non-singleton. As a closed subgroup of H , H_0 is also a TSI pro-Lie Polish group. Similar to the last paragraph of the proof of Theorem 6.8, we can embed the interval $[0, 1]$ into H_0 , and also in H . Therefore, the (\Leftarrow) part follows from Lemma 2.9. \dashv

By restricting our analysis to Lie groups, we provide an affirmative response to [5, Question 7.4] as follows:

THEOREM 6.10 (Rigid Theorem for TSI Lie groups). *Let G, H be two separable TSI Lie groups such that G is connected. Then $E(G) \leq_B E(H)$ iff there exists a continuous locally injective homomorphism $S : G \rightarrow H$.*

PROOF. By Theorem 6.9, we only need to show that $\ker(S)$ is discrete. Note that any non-archimedean subgroup of a Lie group is discrete, as all Lie groups are NSS. \dashv

6.2. Applications on Banach spaces and Fréchet spaces. Now we focus on infinite dimensional vector spaces. Let us recall some elementary notions. All separable Fréchet spaces, i.e., separable, completely metrizable topological vector spaces (see [20, 5-1]), can be viewed as abelian Polish groups under the addition operation. In this article, all vector spaces are assumed to be real, i.e., over the field \mathbb{R} . This is because, in accordance with view of Borel reducibility, the equivalence relation induced by a separable Fréchet space is independent of the choice of the field of scalars.

Let X be a vector space. A map $\|\cdot\| : X \rightarrow \mathbb{R}$ is called a *total paranorm* (see [20, 2-1]) if

- (1) $\|v\| \geq 0$, $\| -v \| = \|v\|$, and $\|v\| = 0 \iff v = 0$;
- (2) $\|v + v'\| \leq \|v\| + \|v'\|$;
- (3) for $(t_n) \in \mathbb{R}^\omega$, $(v_n) \in X^\omega$, if $t_n \rightarrow t$ and $\|v_n - v\| \rightarrow 0$, then $\|t_n v_n - tv\| \rightarrow 0$.

A total paranorm $\|\cdot\|$ is called a *norm* if $\|tv\| = |t|\|v\|$ for $t \in \mathbb{R}$ and $v \in X$. Any Fréchet space admits a compatible complete two-sided invariant metric d , which is given by a total paranorm $\|\cdot\|$ as $d(v, v') = \|v - v'\|$.

In particular, all separable Banach spaces are separable Fréchet spaces. The following are some classical separable Banach spaces:

$$c_0, C[0, 1], l_p, L_p[0, 1] \quad (\forall p \in [1, +\infty)).$$

The classical Banach–Mazur theorem [21, II.B] asserts that every separable Banach space is isometric to a subspace of $C[0, 1]$. Thus for any separable Banach space X , we have $E(X) \leq_B E(C[0, 1])$. For more details on Fréchet spaces and Banach spaces, we refer to [20, 21].

THEOREM 6.11 (Rigid Theorem for Fréchet spaces). *Let X, Y be two separable Fréchet spaces such that Y is a closed subgroup of the product of a sequence of TSI strongly NSS Polish groups. Then $E(X) \leq_B E(Y)$ iff X is topologically isomorphic to a closed linear subspace of Y .*

PROOF. (\Leftarrow). It follows from Proposition 2.3.

(\Rightarrow). Suppose $E(X) \leq_B E(Y)$. Note that X is connected. By Theorem 6.4, Theorem 5.3(ii), and (2) \Rightarrow (1) of Proposition 6.3, there exists a continuous homomorphism $S : X \rightarrow Y$ such that $\ker(S)$ is non-archimedean, and for $x \in X^\omega$, if $\lim_p x(p) = 0$, then

$$\sum_p x(p) \text{ converges} \iff \sum_p S(x(p)) \text{ converges.}$$

For any integers m, n with $n > 0$ and $v \in X$, we have that $nS(\frac{1}{n}v) = S(v)$, and thus $S(\frac{mv}{n}) = \frac{m}{n}S(v)$. By its continuity, S is a \mathbb{R} -linear map. Note that $\ker(S)$ is a non-archimedean closed linear subspace of X , so $\ker(S) = \{0\}$. Therefore S is injective. We will show that S is a topological isomorphism from X onto $S(X)$.

Let $\|\cdot\|_X, \|\cdot\|_Y$ be total paranorms on X and Y respectively. Assume for contradiction that there exist a sequence (v_q) in X and a $\delta > 0$ such that

- (1) $\lim_q \|S(v_q)\|_Y = 0$, and
- (2) $\inf_q \|v_q\|_X > \delta$.

By transferring to a subsequence, we may assume that $\|S(v_q)\|_Y < 2^{-q}$ for all $q \in \omega$. For each $q \in \omega$, we can find an integer $m_q > 0$ such that $\|\frac{v_q}{m_q}\|_X < 2^{-q}$.

Put $M_{-1} = 0$ and $M_q = m_0 + \dots + m_q$ for $q \in \omega$. For each $p \in \omega$, define $x \in X^\omega$ with $x(p) = \frac{v_q}{m_q}$ for $p = M_{q-1} + i$ and $0 \leq i < m_q$. It follows that $\lim_p x(p) = 0$, so

$$\sum_p x(p) \text{ converges} \iff \sum_p S(x(p)) \text{ converges.}$$

Fix a $k \in \omega$. By [10, Exercise E7.12(ii)], there is an open set $V \ni 0$ in Y such that

$$\forall r \in [0, 1] \forall h \in V (\|rh\|_Y < 2^{-(k+2)}).$$

By (1), there is a $Q > k$ such that

$$\forall q (q > Q \Rightarrow S(v_q) \in V).$$

For any $p' > p > M_Q$, there are $q' \geq q > Q$ such that $M_{q-1} \leq p < M_q$ and $M_{q'-1} \leq p' < M_{q'}$. It follows that

$$S(x(p)) + \dots + S(x(M_q - 1)) = \frac{M_q - p}{m_q} S(v_q), \quad S(v_q) \in V,$$

$$S(x(M_{q'-1})) + \dots + S(x(p')) = \frac{p' - M_{q'-1} + 1}{m_{q'}} S(v_{q'}), \quad S(v_{q'}) \in V.$$

These imply that

$$\begin{aligned} \left\| \sum_{i=p}^{p'} S(x(i)) \right\|_Y &\leq \|S(x(p)) + \cdots + S(x(M_q - 1))\|_Y + \sum_{i=q+1}^{q'-1} \|S(v_i)\|_Y \\ &\quad + \|S(x(M_{q'-1})) + \cdots + S(x(p'))\|_Y \\ &\leq 2^{-(k+2)} + \sum_{i=q+1}^{q'-1} 2^{-i} + 2^{-(k+2)} < 2^{-k}. \end{aligned}$$

This shows that $(S(x(0)) + \cdots + S(x(p)))$ is a Cauchy sequence. So the series $\sum_p S(x(p))$ converges.

For any $q \in \omega$, we have

$$\|x(M_{q-1}) + x(M_{q-1} + 1) + \cdots + x(M_q - 1)\|_X = \|v_q\|_X > \delta,$$

and thus $\sum_p x(p)$ diverges. A contradiction!

Now we see that

$$\forall (v_q) \in X^\omega (\lim_q S(v_q) = 0 \Rightarrow \lim_q v_q = 0).$$

This implies that $S^{-1} : S(X) \rightarrow X$ is continuous. So S is a topological linear isomorphism from X onto $S(X)$, and hence $S(X)$ is closed. \dashv

The following theorem characterizes strongly NSS separable Banach spaces.

THEOREM 6.12. *A separable Banach space X is not strongly NSS iff it has a closed linear subspace topologically isomorphic to c_0 .*

PROOF. (\Leftarrow). It follows from the fact that any closed subgroup of a strongly NSS Polish group is strongly NSS, whereas c_0 is not strongly NSS.

(\Rightarrow). Suppose X is not strongly NSS. Denote the norm of X by $\|\cdot\|$. Let $V = \{v \in X : \|v\| < 1\}$. Then V is not an unenclosed set of X , and thus there exists a sequence (v_i) in X such that $v_i \not\rightarrow 0$ and

$$\forall j < k \forall \theta \in \{-1, 1\}^{\{j, \dots, k\}} \|\theta(j)v_j + \cdots + \theta(k)v_k\| < 2.$$

For any integer $n > 0$ and $z^0, z^1, \dots, z^m \in \mathbb{R}^n$, let

$$\text{Con}(\{z^0, z^1, \dots, z^m\}) = \left\{ \sum_{i=0}^m \lambda_i z^i : \sum_{i=0}^m \lambda_i = 1, \forall i \leq m (\lambda_i \geq 0) \right\}.$$

Define $\mathbb{D}(n) = \{z \in \mathbb{R}^n : \forall i < n (|z(i)| \leq 1)\}$. It is clear that

$$\mathbb{D}(n) = \text{Con}(\{\theta \in \mathbb{R}^n : \forall i < n (\theta(i) = \pm 1)\}).$$

Now let $(t_i) \in c_0$. We claim that $\sum_i t_i v_i$ converges. For any $\varepsilon > 0$, we can find an $i_0 \in \omega$ so that $|t_i| < \varepsilon$ for $i > i_0$. Let $r = \sup\{|t_i| : i > i_0\}$. We may assume $r > 0$. Then for any $i_0 < j < k$, we have $(t_j, \dots, t_k)/r \in \mathbb{D}(k - j + 1)$. So there is a $\lambda : \{-1, 1\}^{\{j, \dots, k\}} \rightarrow \mathbb{R}$ with $\sum_{\theta \in \{-1, 1\}^{\{j, \dots, k\}}} \lambda(\theta) = 1$ and $\lambda(\theta) \geq 0$ for each θ such

that

$$(t_j, \dots, t_k)/r = \sum_{\theta \in \{-1,1\}^{\{j,\dots,k\}}} \lambda(\theta)(\theta(j), \dots, \theta(k)).$$

So

$$\sum_{i=j}^k \frac{t_i}{r} v_i = \sum_{\theta \in \{-1,1\}^{\{j,\dots,k\}}} \lambda(\theta) \sum_{i=j}^k \theta(i) v_i.$$

Then we have

$$\begin{aligned} \left\| \sum_{i=j}^k t_i v_i \right\| &\leq \sum_{\theta \in \{-1,1\}^{\{j,\dots,k\}}} r \lambda(\theta) \left\| \sum_{i=j}^k \theta(i) v_i \right\| \\ &< \sum_{\theta \in \{-1,1\}^{\{j,\dots,k\}}} 2r \lambda(\theta) = 2r \leq 2\varepsilon. \end{aligned}$$

So by Cauchy Criterion, $\sum_i t_i v_i$ converges.

Therefore, $\sum_i t_i v_i$ converges whenever $(t_i) \in c_0$. Note that $\sum_i v_i$ diverges, since $v_i \not\rightarrow 0$. By [14, Proposition 2.e.4] and its remark, X has a closed linear subspace topologically isomorphic to c_0 . ⊣

Applying previous results to separable Banach spaces, we get:

THEOREM 6.13 (Rigid Theorem for Banach spaces). *Let X, Y be two separable Banach spaces such that Y contains no closed linear subspaces topologically isomorphic to c_0 . Then $E(X) \leq_B E(Y)$ iff X is topologically isomorphic to a closed linear subspace of Y .*

PROOF. It follows from Theorems 6.11 and 6.12. ⊣

REMARK 6.14. It is well known that l_p and $L_p[0, 1]$ contain no closed linear subspaces topologically isomorphic to c_0 , where $p \in [1, +\infty)$. Note that l_p is topologically isomorphic to a closed linear subspace of l_q iff $p = q$ (see [17, Theorem 5.1]). Let X be a separable Fréchet space and let Y be the space l_p or $L_p[0, 1]$. Then $E(X) \leq_B E(Y)$ iff X is topologically isomorphic to a closed linear subspace of Y . In particular,

$$E(l_p) \leq E(l_q) \iff p = q.$$

Without the assuming of strongly NSS for TSI Polish groups G and H , we do not know how to compare $E(G)$ and $E(H)$ with respect to Borel reducibility so far. For instance, it is not known whether $E(C[0, 1]) \leq_B E(c_0)$, or whether $E(l_p) \leq_B E(c_0)$ for $p \in (0, +\infty)$.

A worthwhile question is:

QUESTION 6.15. *Let X, Y be two separable Fréchet spaces. Does it hold that $E(X) \leq_B E(Y)$ iff X is topologically isomorphic to a closed linear subspace of Y ?*

§7. Uniformly NSS Polish groups. Recall that a topological group G is *uniformly NSS* (see [16, Definition 11]) if there is an open subset $V \ni 1_G$ of G such that, for any open subset $U \ni 1_G$ of G ,

$$\exists n \forall g \in G (g \notin U \Rightarrow \exists m \leq n (g^m \notin V)).$$

Clearly, a locally compact Polish group is uniformly NSS iff it is NSS.

A topological group is a *Banach-Lie group* if it is a Banach manifold such that the group operations are smooth functions (see [18, Section 6]). All Banach spaces and Lie groups are Banach-Lie groups, while all Banach-Lie groups are uniformly NSS (see [15, Theorem 2.7]).

Under the assumption of uniformly NSS, we show that every Borel reduction of $E(G)$ to $E(H)$ results in a continuous homomorphism, which is also a local homeomorphism.

THEOREM 7.1. *Let G, H be two TSI Polish groups. Assume that G is uniformly NSS and H is strongly NSS. If $E(G) \leq_B E(H)$, then there exist an open subgroup W of G , an open neighborhood $U \subseteq W$ of 1_G , and a continuous homomorphism $S : W \rightarrow H$ such that $S \upharpoonright \bar{U} : \bar{U} \rightarrow S(\bar{U})$ is a homeomorphism and $S(\bar{U})$ is closed in H .*

Moreover, the converse is also true if $G = W$ and the interval $[0, 1]$ can be embedded into H .

PROOF. Let d_G, d_H be compatible complete two-sided invariant metrics on G and H respectively. Assume that $E(G) \leq_B E(H)$. It follows from Remark 6.5 that there exist an open subgroup W' of G and a continuous homomorphism $S : W' \rightarrow H$ such that, for $x, y \in (W')^\omega$, if $\lim_p x(p)y(p)^{-1} = 1_G$, then we have

$$xE_*(G)y \iff S^\omega(x)E_*(H)S^\omega(y).$$

Let $V \ni 1_G$ be an open subset of G witnessing that G is uniformly NSS. For each $k \in \omega$, define $V_k = \{g \in W' : d_G(1_G, g) < 2^{-k}\}$.

We claim that there exists a $k_0 \in \omega$ such that

$$\forall (g_q) \in V_{k_0}^\omega (\lim_q S(g_q) = 1_H \Rightarrow \lim_q g_q = 1_G). \quad (*)$$

If not, then for each $k \in \omega$, there are a $j(k) \in \omega$ and a sequence $(g_{k,q})$ of elements of V_k such that $\lim_q S(g_{k,q}) = 1_H$ and $g_{k,q} \notin V_{j(k)}$ for all $q \in \omega$. Then by the definition of V , there exists an $n_k > 0$ such that

$$\forall g \in G (g \notin V_{j(k)} \Rightarrow \exists m \leq n_k (g^m \notin V)).$$

Pick a large enough $q_k \in \omega$ so that $d_H(1_H, S(g_{k,q_k})) < 2^{-k}/n_k$. Then we can find an $m_k \leq n_k$ with $g_{k,q_k}^{m_k} \notin V$.

Put $M_{-1} = 0$ and $M_k = m_0 + \dots + m_k$ for $k \in \omega$. Let $x \in G^\omega$ be defined as $x(p) = g_{k,q_k}$ for $p = M_{k-1} + i$ with $0 \leq i < m_k$. It is clear that $\lim_p x(p) = 1_G$. This implies that

$$xE_*(G)1_{G^\omega} \iff S^\omega(x)E_*(H)S^\omega(1_{G^\omega}).$$

For any $k \in \omega$, we have

$$x(M_{k-1})x(M_{k-1} + 1) \dots x(M_k - 1) = g_{k,q_k}^{m_k} \notin V.$$

By Lemma 2.4(2), we see that $x E_*(G) 1_{G^\omega}$ fails.

For any $p < p'$, if $M_{k-1} \leq p < M_k$ and $M_{k'-1} \leq p' < M_{k'}$, then we have

$$\begin{aligned} d_H(1_{H^\omega}, S^\omega(x))|_{[p,p']} &\leq \sum_{i=p}^{p'} d_H(1_H, S(x(i))) \\ &= \sum_{i=p}^{M_k-1} d_H(1_H, S(g_{k,q_k})) + \sum_{i=k+1}^{k'-1} m_i d_H(1_H, S(g_{i,q_i})) + \sum_{i=M_{k'-1}}^{p'} d_H(1_H, S(g_{k',q_{k'}})) \\ &< 2^{-k} m_k/n_k + \sum_{i=k+1}^{k'-1} 2^{-i} m_i/n_i + 2^{-k'} m_{k'}/n_{k'} \\ &< 3 \cdot 2^{-k}. \end{aligned}$$

This implies that $\lim_p \sup_{p \leq p'} d_H(1_H, S^\omega(x))|_{[p,p']} = 0$. Then it follows from Lemma 2.4(2) that $S^\omega(x) E_*(H) S^\omega(1_{G^\omega})$. A contradiction! So (*) holds.

Now put $U = V_{k_0+2}$. Then $\bar{U} \subseteq V_{k_0+1} = V_{k_0+1}^{-1}$, so $\bar{U}^{-1} \bar{U} \subseteq V_{k_0}$. For $g \in \bar{U}$ and $(g_p) \in \bar{U}^\omega$, since $g_p^{-1} g \in V_{k_0}$ for each p , we have

$$\lim_p S(g_p) = S(g) \Rightarrow \lim_p S(g_p^{-1} g) = 1_H \Rightarrow \lim_p g_p^{-1} g = 1_G \Rightarrow \lim_p g_p = g.$$

This implies that $S \upharpoonright \bar{U}$ is a topological embedding.

Then we only need to show that $S(\bar{U})$ is closed. Let $(g_p) \in \bar{U}^\omega$ and $h \in H$ with $\lim_p S(g_p) = h$. Assume for contradiction that (g_p) does not converge in \bar{U} . Then it is not d_G -Cauchy, we can find two strictly increasing sequences of natural numbers $(p_k), (q_k)$ with $p_k < q_k$ such that $d_G(g_{p_k}, g_{q_k}) \not\rightarrow 0$. By the fact that $\lim_k S(g_{p_k}^{-1} g_{q_k}) = \lim_k S(g_{p_k})^{-1} S(g_{q_k}) = 1_H$, we have $\lim_k g_{p_k}^{-1} g_{q_k} = 1_G$, and we get a contradiction as desired. Therefore, $\lim_p g_p = g$ for some $g \in \bar{U}$, and hence $h = \lim_p S(g_p) = S(g) \in S(\bar{U})$. So $S(\bar{U})$ is closed.

Finally, if $G = W$ and the interval $[0, 1]$ can be embedded into H , by lemmas 6.7 and 2.9, we see that the converse is also true. ⊖

COROLLARY 7.2. *Let G, H be two TSI Polish groups. Assume that G is uniformly NSS and H is strongly NSS. If $E(G) \leq_B E(H)$, then G is strongly NSS too.*

PROOF. By Theorem 7.1, we can get an open subset $U \ni 1_G$ of G and a homeomorphism $S : \bar{U} \rightarrow H$ such that $S(gg') = S(g)S(g')$ for $g, g', gg' \in \bar{U}$. Let V be an unenclosed set of H . It is clear that $S^{-1}(V) \cap U$ is an unenclosed set of G , so G is strongly NSS. ⊖

REMARK 7.3. The assumption of uniformly NSS in Theorem 7.1 and Corollary 7.2 can not be avoided. For example, for p -adic solenoid \mathbb{T}_p , we have $E(\mathbb{T}_p) \leq_B E(\mathbb{T})$, where \mathbb{T} is strongly NSS but \mathbb{T}_p is not (see [5, Theorem 6.24]).

In the rest of this article, we attempt to study some examples that are induced by totally disconnected TSI Polish groups.

EXAMPLE 7.4. Let $G = \{v \in c_0 : \forall n (v(n) \in 2^{-n}\mathbb{Z})\}$. Then G is uniformly NSS but not strongly NSS with the subspace topology inherited from c_0 . By Theorem 6.12, l_p is strongly NSS. So $E(G) \not\leq_B E(l_p)$ for $p \in [1, +\infty)$.

Recall that a sequence (v_j) in a Banach space X is called a *Schauder basis* of X if

$$\forall v \in X \exists! (t_j) \in \mathbb{R}^\omega (v = \sum_j t_j v_j).$$

Two sequences (v_j) and (v'_j) are called *equivalent* in X if

$$\sum_j t_j v_j \text{ converges} \iff \sum_j t_j v'_j \text{ converges}.$$

For $p \in [1, +\infty)$ and $a \in c_0$, we define

$$I_a = \{n \in \omega : a(n) \neq 0\}, \quad A_{p,a} = \{v \in l_p : \forall n (v(n) \in a(n)\mathbb{Z})\}.$$

If I_a is infinite, then $A_{p,a}$ equipped the relative topology inherited from l_p is a totally disconnected, strongly NSS abelian Polish group, but is not non-archimedean. In particular, we put $d(n) = 2^{-n}$, and let

$$A_p = A_{p,d} = \{v \in l_p : \forall n (v(n) \in 2^{-n}\mathbb{Z})\}.$$

Let $e_j = (0, \dots, 0, \overset{j}{1}, 0, \dots)$, then (e_j) is a Schauder basis of l_p .

THEOREM 7.5. For $p, q \in [1, +\infty)$ and $a \in c_0$, the following hold:

- (1) if I_a is a nonempty finite set, then $E(A_{p,a}) \sim_B E_0$;
- (2) if I_a is infinite, then $E(A_{p,a}) \sim_B E(A_p)$;
- (3) $E(A_p) <_B E(l_p)$;
- (4) $E(A_p) \leq_B E(l_q) \iff p = q$; and
- (5) $E(A_p) \leq_B E(A_q) \iff p = q$.

PROOF. (1) If I_a is a nonempty finite set, then $A_{p,a}$ is a nontrivial countable discrete group. So $E(A_{p,a}) \sim_B E_0$ (see [5, Theorem 3.5(1)]).

(2) Suppose that I_a is infinite. We can find a strictly increasing sequence of $j(n) \in \omega$ for each $n \in I_a$ such that $2^{-j(n)} \leq |a(n)|$. Clearly, there is an $m_n \in \omega$ with

$$|a(n)| \leq m_n 2^{-j(n)} \leq 2|a(n)|.$$

Let $\varphi : A_{p,a} \rightarrow A_p$ be defined as

$$\varphi(v)(k) = \begin{cases} \frac{m_n v(n)}{2^{j(n)} a(n)}, & k = j(n), n \in I_a, \\ 0, & \text{otherwise.} \end{cases}$$

It is obvious that φ is a continuous homomorphism. For any $v \in A_{p,a}$, we have

$$\|v\|_p = \left(\sum_{n \in I_a} |v(n)|^p \right)^{\frac{1}{p}} \leq \left(\sum_{n \in I_a} \left| \frac{m_n v(n)}{2^{j(n)} a(n)} \right|^p \right)^{\frac{1}{p}} = \|\varphi(v)\|_p \leq 2\|v\|_p.$$

Thus φ is a topological group isomorphism from $A_{p,a}$ onto $\varphi(A_{p,a})$. It follows from Proposition 2.3 that $E(A_{p,a}) \leq_B E(A_p)$. Similarly, $E(A_p) \leq_B E(A_{p,a})$.

(3) Since A_p is a closed subgroup of l_p , we get that $E(A_p) \leq_B E(l_p)$. By Corollary 6.6, we have that $E(A_p) <_B E(l_p)$.

(4) The (\Leftarrow) part is from (3). We prove the (\Rightarrow) part as follows:

Suppose that $E(A_p) \leq_B E(l_q)$. Then by Theorem 7.1, there exist an open subgroup W of A_p and a continuous homomorphism $S : W \rightarrow l_q$ such that $S \upharpoonright \overline{U} : \overline{U} \rightarrow S(\overline{U})$ is a homeomorphism and $S(\overline{U})$ is closed in l_q , where $U \subseteq W$ is an open neighborhood of 0.

For $k \in \omega$, let $V_k = \{v \in A_p : \|v\|_p \leq 2^{-k}\}$. Since S is continuous, we can find an integer $k_0 > 0$ so that

$$V_{k_0} \subseteq U, \quad \forall v \in V_{k_0} (\|S(v)\|_q \leq 1).$$

For any $k \in \omega$, we define $v_k \in V_{k_0}$ as

$$v_k(n) = \begin{cases} 2^{-k_0-1}, & n = k + k_0, \\ 0, & \text{otherwise.} \end{cases}$$

Note that for any $k \neq k'$, we have

$$\|v_k\|_p = 2^{-k_0-1} < \|v_k - v_{k'}\|_p = 2^{1/p} 2^{-k_0-1} \leq 2^{-k_0},$$

and thus both v_k and $v_k - v_{k'}$ are in V_{k_0} . So

$$\sup_k \|S(v_k)\|_q \leq 1, \quad \sup_{k \neq k'} \|S(v_k) - S(v_{k'})\|_q \leq 1.$$

Since $S \upharpoonright \overline{U}$ is a homeomorphism, there are $D_1, D_2 > 0$ such that

$$\inf_k \|S(v_k)\|_q > D_1, \quad \inf_{k \neq k'} \|S(v_k) - S(v_{k'})\|_q > D_2.$$

Since $\sup_k \|S(v_k)\|_q \leq 1$, by the compactness of $[-1, 1]^\omega$, there are a subsequence $(S(v_{k_i}))$ of $(S(v_k))$ and a $w \in [-1, 1]^\omega$ such that

$$\forall n (\lim_i S(v_{k_i})(n) = w(n)).$$

It is clear that $\|w\|_q \leq 1$, and thus $w \in l_q$. Put $w_i = S(v_{k_{2i}}) - S(v_{k_{2i+1}}) \in l_q$ for each $i \in \omega$. We have

$$D_2 < \inf_i \|w_i\|_q \leq \sup_i \|w_i\|_q \leq 1, \quad \forall n (\lim_i w_i(n) = 0).$$

It follows from [14, Proposition 1.a.12] that there is a subsequence (w_{i_j}) of (w_i) which is equivalent to a block basis (see [14, Definition 1.a.10]) (w'_j) of (e_j) . In other words, we get that, for any $(t_j) \in \mathbb{R}^\omega$,

$$\sum_j t_j w_{i_j} \text{ converges} \iff \sum_j t_j w'_j \text{ converges.}$$

Then it is routine to check that $0 < \inf_j \|w'_j\|_q \leq \sup_j \|w'_j\|_q < \infty$. By similar arguments in proof of [14, Proposition 2.a.1(i)], we see that (w'_j) is equivalent to (e_j) in l_q . Therefore, (w_{i_j}) is equivalent to (e_j) in l_q .

Let $x(j) = v_{k_{2i_j}} - v_{k_{2i_j+1}}$ for each $j \in \omega$. Then $x \in V_{k_0}^\omega \subseteq U^\omega \subseteq W^\omega$. It is trivial to check that $(x(j))$ is equivalent to (e_j) in l_p . Note that $2^{-k_0-1} < \|x(j)\|_p \leq 2^{-k_0}$

and $D_2 \leq \|w_{i_j}\|_q \leq 1$ for $j \in \omega$. Thus for any $(t_j) \in \mathbb{R}^\omega$, we have

$$\lim_j t_j x(j) = 0 \iff \lim_j t_j = 0 \iff \lim_j t_j w_{i_j} = 0.$$

Put $X = \{(t_j) \in \mathbb{R}^\omega : \forall j (t_j \in 2^{-j}\mathbb{Z} \cap [0, 1])\}$. Then for any $(t_j) \in X$, $j \in \omega$, we have $\|t_j x(j)\|_p \leq \|x(j)\|_p \leq 2^{-k_0}$. So $t_j x(j) \in V_{k_0} \subseteq U \subseteq W$. Note that $x(j) \in U$. Then we have $S(t_j x(j)) = t_j w_{i_j}$ for each $(t_j) \in X$ (This is because that $w_{i_j} = S(v_{k_{2i_j}}) - S(v_{k_{2i_j+1}}) = S(v_{k_{2i_j}} - v_{k_{2i_j+1}}) = S(x(j))$).

For any $t = (t_j) \in X$, let $\bar{x}^t = (t_j x(j)) \in W^\omega$. By Lemma 6.7, $S^\#$ is a reduction of $E_*(W)$ to $E_*(I_q) \times E(W; 0)$. Let $y = 0 \in W^\omega$. Again by Proposition 2.8, we can get the following conclusion that, if $\lim_j \bar{x}^t(j) = 0$, then

$$\sum_j \bar{x}^t(j) - y(j) \text{ converges} \iff \sum_j S(\bar{x}^t(j)) - S(y(j)) \text{ converges.}$$

Because both sides of the above formula imply $\lim_j \bar{x}^t(j) = 0$. So for any $(t_j) \in X$,

$$\sum_j t_j x(j) \text{ converges} \iff \sum_j t_j w_{i_j} \text{ converges.}$$

From the fact that $(x(j))$ is equivalent to (e_j) in l_p and (w_{i_j}) is equivalent to (e_j) in l_q , we see that

$$\forall (t_j) \in X ((t_j) \in l_p \iff (t_j) \in l_q),$$

this gives that $p = q$.

(5) It follows immediately from (3) and (4). ◻

Acknowledgements. The authors would like to thank the anonymous referee for helpful comments and suggestions.

Funding. This research is partially supported by the National Natural Science Foundation of China (Grant No. 12271264).

REFERENCES

- [1] H. BECKER, *Polish group actions: Dichotomies and generalized elementary embeddings*. *Journal of the American Mathematical Society*, vol. 11 (1998), pp. 397–449.
- [2] H. BECKER and A. S. KECHRIS, *The Descriptive Set Theory of Polish Group Actions*, London Mathematical Society Lecture Note Series, 232, Cambridge University Press, Cambridge, 1996.
- [3] L. DING, *Borel reducibility and Hölder(α) embeddability between Banach spaces*. *The Journal of Symbolic Logic*, vol. 77 (2012), pp. 224–244.
- [4] ———, *On equivalence relations generated by Schauder bases*. *The Journal of Symbolic Logic*, vol. 82 (2017), pp. 1459–1481.
- [5] L. DING and Y. ZHENG, *On equivalence relations induced by Polish groups*, preprint, 2022, [arXiv:2204.04594](https://arxiv.org/abs/2204.04594), Unpublished.
- [6] ———, *On equivalence relations induced by locally compact abelian Polish groups*. *The Journal of Symbolic Logic*, (2023), pp. 1–16. <https://doi.org/10.1017/jsl.2023.35>.
- [7] I. FARAH, *Basis problem for turbulent action II: c_0 -equalities*. *Proceedings of the London Mathematical Society*, vol. 82 (2001), pp. 1–30.
- [8] S. GAO, *Invariant Descriptive Set Theory*, Monographs and Textbooks in Pure and Applied Mathematics, 293, CRC Press, Boca Raton, 2009.

- [9] K. H. HOFMANN and S. A. MORRIS, *The Lie Theory of Connected Pro-Lie Groups*, EMS Tracts in Mathematics, 2, European Mathematical Society, Zürich, 2007.
- [10] ———, *The Structure of Compact Groups*, De Gruyter Studies in Mathematics, 25, De Gruyter, Berlin, 2013.
- [11] K. H. HOFMANN, S. A. MORRIS, and M. STROPPEL, *Varieties of topological groups, Lie groups, and SIN-groups*. *Colloquium Mathematicum*, vol. 70 (1996), pp. 151–163.
- [12] ———, *Locally compact groups, residual Lie groups, and varieties generated by Lie group*. *Topology and its Applications*, vol. 71 (1996), pp. 63–91.
- [13] A. S. KECHRIS, *Classical Descriptive Set Theory*, Graduate Texts in Mathematics, 156, Springer-Verlag, New York, 1995.
- [14] J. LINDENSTRAUSS and L. TZAFRIRI, *Classical Banach Spaces I: Sequence Spaces*, Ergebnisse der Mathematik und ihrer Grenzgebiete, 92, Springer-Verlag, Berlin, 1977.
- [15] S. A. MORRIS and V. PESTOV, *On Lie groups in varieties of topological groups*. *Colloquium Mathematicae*, vol. 78 (1998), pp. 39–47.
- [16] C. ROSENDAL, *Lipschitz structure and minimal metrics on topological groups*. *Arkiv for Matematik*, vol. 56 (2018), pp. 185–206.
- [17] W. J. STILES, *On properties of subspaces of l_p , $0 < p < 1$* . *Transactions of the American Mathematical Society*, vol. 149 (1970), pp. 405–415.
- [18] H. UPMEIER, *Symmetric Banach Manifolds and Jordan C^* -Algebras*, North-Holland Mathematics Studies, 104, North-Holland, Amsterdam, 1985.
- [19] V. S. VARADARAJAN, *Lie Groups, Lie Algebras, and their Representations*, Graduate Texts in Mathematics, 102, Springer-Verlag, New York, 1974.
- [20] A. WILANSKY, *Modern Methods in Topological Vector Spaces*, McGraw-Hill International Book Company, New York, 1978.
- [21] P. WOJTAŚCZYK, *Banach Spaces for Analysts*, Cambridge Studies in Advanced Mathematics, 25, Cambridge University Press, Cambridge, 1991.

SCHOOL OF MATHEMATICAL SCIENCES AND LPMC
 NANKAI UNIVERSITY
 TIANJIN, 300071
 P.R. CHINA

E-mail: dingly@nankai.edu.cn

E-mail: 1120200015@mail.nankai.edu.cn