

Original Article

Cite this article: Yu B, Steptoe A, Chen Y, Jia X (2021). Social isolation, rather than loneliness, is associated with cognitive decline in older adults: the China Health and Retirement Longitudinal Study. *Psychological Medicine* 51, 2414–2421. <https://doi.org/10.1017/S0033291720001014>

Received: 20 September 2019
Revised: 30 January 2020
Accepted: 26 March 2020
First published online: 27 April 2020


Key words:

Cognitive function; loneliness; older adults; social isolation

Author for correspondence:

Bin Yu, E-mail: yubin80@tmu.edu.cn

Social isolation, rather than loneliness, is associated with cognitive decline in older adults: the China Health and Retirement Longitudinal Study

Bin Yu¹ , Andrew Steptoe², Yongjie Chen³ and Xiaohua Jia^{4,5}

¹Department of Psychiatry and Psychology, School of Basic Medical Sciences, Tianjin Medical University, Tianjin, China; ²Department of Behavioural Science and Health, University College London, London, UK; ³Department of Epidemiology and Statistics, School of Public Health, Tianjin Medical University, Tianjin, China; ⁴Department of Ultrasound, General Hospital of People's Liberation Army, Beijing, China and ⁵Key Laboratory of Molecular Imaging of Chinese Academy of Sciences, Institute of Automation, Chinese Academy of Sciences, Beijing, China

Abstract

Background. Social isolation and loneliness have each been associated with cognitive decline, but most previous research is limited to Western populations. This study examined the relationships of social isolation and loneliness on cognitive function among Chinese older adults. **Methods.** This study used two waves of data (2011 and 2015) from the China Health and Retirement Longitudinal Study (CHARLS) and analyses were restricted to those respondents aged 50 and older. Social isolation, loneliness, and cognitive function were measured at baseline. Follow-up measures on cognitive function were obtained for 7761 participants (mean age = 60.97, s.d. = 7.31; male, 50.8%). Lagged dependent variable models adjusted for confounding factors were used to evaluate the association between baseline isolation, loneliness, and cognitive function at follow-up.

Results. Loneliness was significantly associated with the cognitive decline at follow-up (episodic memory: $\beta = -0.03$, $p < 0.01$; mental status: $\beta = -0.03$, $p < 0.01$) in the partially adjusted models. These associations became insignificant after additional confounding variables (chronic diseases, health behaviors, disabilities, and depressive symptoms) were taken into account (all $p > 0.05$). By contrast, social isolation was significantly associated with decreases in all cognitive function measures at follow-up (episodic memory: $\beta = -0.05$, $p < 0.001$; mental status: $\beta = -0.03$, $p < 0.01$) even after controlling for loneliness and all confounding variables.

Conclusions. Social isolation is associated with cognitive decline in Chinese older adults, and the relationships are independent of loneliness. These findings expand our knowledge about the links between social relationships and the cognitive function in non-Western populations.

Introduction

Population aging is one of the major challenges worldwide. China has the world's largest aging population of older adults. By 2050, the number of Chinese people aged 60 and older is expected to reach 479 million (DESA, 2017). In parallel with this aging profile, the number of Chinese older adults with dementia is projected to reach 18 million in 2030 (Chan et al., 2013). Dementia makes a major contribution to disability and health care needs among older people. The enormous disease burden has made the prevention and treatment of dementia public health priorities for China.

Increasing efforts have been made to identify modifiable factors that may prevent or slow the progression of cognitive decline in older age, and impoverished social relationships – defined as social isolation or loneliness – have received considerable attention. Accumulated evidence has shown that both social isolation (Evans, Martyr, Collins, Brayne, & Clare, 2018; Kuiper et al., 2016) and loneliness (Boss, Kang, & Branson, 2015; Cacioppo & Hawkley, 2009) affect cognitive function. Although loneliness and social isolation are sometimes perceived to be synonymous, they are conceptually different. In essence, social isolation refers to the objective aspects of isolation, such as living alone, having few contacts, or little involvement in social activities (Holt-Lunstad, Smith, Baker, Harris, & Stephenson, 2015). In contrast, loneliness refers to a subjective feeling resulting from a discrepancy between actual and desired social relationships (Peplau & Perlman, 1982). Even socially connected individuals can feel lonely, just as socially isolated individuals can be satisfied with their social relationships. Studies have found that correlations between loneliness and social isolation are generally moderate (Cornwell & Waite, 2009; McHugh, Kenny, Lawlor, Steptoe, & Kee, 2017).

It has been suggested that analyzing both objective and subjective aspect of social relationship in the same study can allow us to better understand how these two social constructs affect

health (Cornwell & Waite, 2009; Holt-Lunstad et al., 2015; Newell & Menec, 2019). However, researchers have mostly examined the relationship with cognitive impairment of only one construct, either loneliness or social isolation (Boss et al., 2015; Evans et al., 2018; Kuiper et al., 2016). Among the few studies in which loneliness and social isolation were examined concurrently, there were mixed results. Some findings support the greater association of loneliness (Holwerda et al., 2014) and some findings support the greater association of social isolation in predicting with cognitive decline (Beller & Wagner, 2018; Griffin, Mezuk, Williams, Perrin, & Rybarczyk, 2018). Synergistic effects of loneliness and social isolation on cognitive function were also found by some researchers. For example, analysis from the Rush Memory and Aging Project found that participants who were lonely and those with limited participation in social activities were more likely to develop Alzheimer's disease (Wilson et al., 2007). A longitudinal study with English older adults also indicated that both social isolation and loneliness were significantly associated with poorer cognitive function (Shankar, Hamer, McMunn, & Steptoe, 2013). More information is needed regarding both the relative and synergistic influences of social isolation and loneliness on cognitive function.

Meanwhile, another gap in knowledge is the lack of studies on the cognitive consequences of social isolation and loneliness in non-Western countries. Most studies on this topic have been conducted in North America and European countries (Courtin & Knapp, 2017; Evans et al., 2018; Kuiper et al., 2016), and it is not established whether similar patterns occur in other cultures. The importance of such research is underscored by the fact that isolation and loneliness are equally prevalent in non-Western as in Western countries. The proportion of empty-nest families (refers to those older people with no children or whose children have already left home) in China is estimated to reach 90% by 2030 (Rafnsson, Orrell, d'Orsi, Hogervorst, & Steptoe, 2017). It has been suggested that the association between social relationships and health could be more salient in Chinese populations, for whom cultural tradition emphasizes the family system and collectivism (Yang & Victor, 2008). However, only a few studies have investigated the impact of social relationships on cognitive function among older adults and most were limited by small sample size or cross-sectional design (Fung, Lee, Cheng, & Lam, 2019; Wang et al., 2012). Two recent publications based on the Chinese Longitudinal Healthy Longevity Survey using relatively large sample size have reported the negative impact of poor social relationships on cognitive function. However, both of these studies mainly focused on loneliness rather than social isolation (Zhong, Chen, & Conwell, 2016; Zhong, Chen, Tu, & Conwell, 2017).

Given that very few studies examined loneliness and social isolation simultaneously in relation to cognition in non-Western populations, and the inconclusive findings in this area, we used data from the China Health and Retirement Longitudinal Study (CHARLS) to examine the impact of social isolation and loneliness, individually and simultaneously, on cognitive function among Chinese older adults. We hypothesize that both social isolation and loneliness would be significantly associated with cognitive decline over 4 years. However, we made no specific assumption about which of these two factors might play a stronger role in the light of existing mixed results. Regarding previous evidence indicating possible gender differences in the association between social relationship factors and health (Shumaker & Hill, 1991; Shye, Mullooly, Freeborn, & Pope, 1995), we also tested the

interaction of gender with social isolation and loneliness on cognitive function for our sample.

Methods

Participants

Data are from the CHARLS, a nationally representative longitudinal survey sampled residents from 150 counties across 28 provinces in China, with a response rate of 80.5% (Zhao, Hu, Smith, Strauss, & Yang, 2012). The CHARLS is one of the most up-to-date longitudinal data sets collected in China to study the health and well-being of older adults. The survey assigned 23 422 dwelling units to interviewers at baseline in 2011. After excluding empty or non-resident dwellings, 12 740 were age-eligible for CHARLS (Zhao et al., 2012). In the present study, we used data from two waves of the CHARLS collected in 2011 and 2015. The baseline sample included 17 708 respondents. Our analytic sample was restricted to those respondents aged 50 and older ($n = 13\ 649$). We excluded those respondents who had missing values on any predictor or cognitive test at baseline ($n = 3506$) or cognitive test at follow-up ($n = 2382$), which resulted in a final sample size of 7761 respondents (mean age = 60.97, *s.d.* = 7.31; male, 50.8%).

Measures

Loneliness

In our study, loneliness was measured with one single item included in the Center for Epidemiological Studies Depression Scale (CESD): 'In the last week, how often did you feel lonely?'. The respondent chose among four ordinal responses scored 1–4: rarely or none of the time (<1 day), some or a little of the time (1–2 days), occasionally or a moderate amount of the time (3–4 days), most or all of the time (5–7 days). This one-item measure correlates highly with multi-item loneliness scales and has been used in a number of previous studies (Gow, Corley, Starr, & Deary, 2013; Holwerda et al., 2014; Luo & Waite, 2014; Nummela, Seppanen, & Uutela, 2011; Tilvis, Pitkala, Jolkkonen, & Strandberg, 2000). Loneliness was dichotomized into two categories [0 (not lonely) = those who reported feeling lonely rarely or none of the time, and 1 (lonely) = those who felt lonely sometimes, occasionally or most of the time] (Teguo et al., 2016).

Social isolation

Three items were combined to create an index of social isolation, which was adapted from previous research (Glei, Goldman, Ryff, Lin, & Weinstein, 2012; Steptoe, Shankar, Demakakos, & Wardle, 2013). One point was assigned if participants were not married; had less than weekly contact (by phone, in person, or by e-mail) with children, not participating in any social activities over the last month (e.g. interacted with friends; played chess or cards; going to the community club; went to a sport, social, or other clubs; did voluntary or charity work). Scores of social isolation ranged from 0 to 3, with higher scores indicating greater isolation.

Cognitive assessment

CHARLS included similar items for cognitive function as those used in the American Health and Retirement Study (HRS), which were the components of the Telephone Interview for Cognitive Status (TICS) (Crimmins, Kim, Langa, & Weir, 2011).

McArdle, Fisher, and Kadlec (2007)'s study of the HRS data suggested two factors to adequately capture cognitive function (McArdle et al., 2007). Following McArdle et al. and based upon previous studies using the CHARLS data (Lei, Hu, McArdle, Smith, & Zhao, 2012; Pan, Luo, & Roberts, 2018), we constructed two measures of cognitive function. The first is an episodic memory measurement. In CHARLS, memory was assessed through an immediate word recall based on respondents' capacity to immediately repeat in any order a list of 10 Chinese nouns just read to them, followed by a delayed recall that tests respondents' ability to repeat the same list of words 4–10 min later (Crimmins et al., 2011). The episodic memory measure was created by averaging the immediate and delayed recall scores, and scores ranged from 0 to 10. The second cognitive measure is based on the components of the mental status questions of the TICS established to capture the intactness or mental status of individuals. Orientation was assessed by asking respondents to name today's date (month, day, year), and identify the correct day of the week. Visuospatial ability was assessed by asking respondents to accurately re-draw a previously shown picture. Numeric ability was assessed through the serial subtraction of 7 from 100 (up to five times). Scores on these items were aggregated into a single score that ranged from 0 to 10 and was labeled as mental status, as recommended by McArdle et al. (2007). For both measures, higher scores indicate better cognitive function.

Control variables in the baseline survey

The analyses were adjusted for several demographic variables and behavioral, psychological, and clinical risk factors. Age, gender, education, and area of residence (urban/rural) were measured by self-report. Education was dichotomized as lower than secondary school and secondary school or above. Health habits, including drinking and smoking, were collected using a standardized questionnaire (Zhao et al., 2012). Respondents were asked whether they were current smokers and whether they consumed alcohol in the past 12 months. Two measures of functional limitation were considered. CHARLS asked respondents if they required assistance with any of six activities of daily living (ADLs: walking, dressing, bathing, eating, getting into and out of bed, and toileting) or five instrumental ADLs (IADLs: preparing a hot meal, shopping for groceries, making telephone calls, taking medicines, and managing money) (Lei et al., 2014). Both ADLs and IADLs were treated as binary predictors (1 = functional limitations present; 0 = no functional limitations present) due to most respondents (ADL: 83.8%; IADL: 80%) denying any limitations. Chronic diseases including hypertension, diabetes, and heart diseases were obtained by asking respondents if a physician had ever told them that they had the condition. Depressive symptoms were measured with the 10-item Center for Epidemiological Studies Depression Scale (CESD-10). Modified CESD scoring was used to exclude the loneliness question in order to derive a separate depression score that was calculated as the sum of the remaining nine questions (CESD-9, range 0–27).

Statistical analysis

Characteristics of the overall sample at baseline were described using means and standard deviations for continuous data and percentages for categorical data. Rank order correlations were used to assess correlations between key predictors including loneliness (continuous scores ranged from 1 to 4 were used) and social isolation and other variables included in the analyses. To examine

the association between isolation, loneliness, and cognitive function, we used lagged dependent variable regression models with Ordinary Least Squares estimation. The lagged dependent variable model is superior for analyzing the effects of predictor variables on an outcome with two-wave panel data while controlling for the influence of time-invariant variables (Johnson, 2005). Seven models were fitted for both cognitive measurements. Model 1A was constructed to examine the associations between loneliness at baseline and cognitive function at follow-up by partially adjusting for control variables including age, gender, education, area of residence, and baseline cognitive function. Other control variables including chronic diseases, smoking and drinking status, ADL and IADL disabilities were then added in Model 1B. Model 1C added CESD-9 scores in the fully adjusted model. Three similar models were fitted to test the independent associations of isolation with cognitive function (Model 2A, B, and C). Model 3 added both isolation and loneliness into the fully adjusted model. We tested whether there was an interaction effect between isolation and loneliness on cognitive function by including appropriate interaction terms into the fully adjusted model. The interactions between isolation, loneliness, and gender were also assessed. For all regression analyses, standardized regression coefficients (β) were reported as variables under study were measured on different scales. Analyses were carried out using SPSS version 20.0.

Results

Table 1 shows the characteristics of the participants under study at baseline and their correlation with isolation and loneliness. The mean age of participants was just over 60 years, and more than half were men. When compared with those who were excluded from the original sample ($n = 5888$), participants included in the study sample ($n = 7661$) were more likely to be male (50.8% *v.* 47.9%; $p = 0.001$), younger (mean age in years: 60.97 *v.* 64.05, $p < 0.001$), and living in a rural area (62.3% *v.* 56.8%; $p < 0.001$). They were more likely to be a current smoker (32.9% *v.* 25.4%; $p < 0.001$) and drinker (34.1% *v.* 30.2%; $p < 0.001$), were less likely to have hypertension (28.4% *v.* 31.5%; $p = 0.001$), ADL disabilities (16.2% *v.* 24.1%; $p < 0.001$), and IADL disabilities (20.0% *v.* 30.3%; $p < 0.001$). In addition, individuals who were included into the study sample were more likely to feel lonely (35.7% *v.* 28.1%, $p < 0.001$), had a lower level of depressive symptoms (mean CESD-9 score: 6.80 *v.* 7.71, $p < 0.001$), lower level of isolation (mean score: 0.76 *v.* 0.90; $p < 0.001$), and both their episodic memory (mean score: 3.50 *v.* 3.09; $p < 0.001$) and mental status (mean score: 6.57 *v.* 5.50; $p < 0.001$) were better than those who were excluded from the analysis.

Of all the participants, the mean scores (s.d.) for loneliness and social isolation was 1.52 (0.94) and 0.76 (0.67) respectively, with 28.1% reported that they felt lonely sometimes, occasionally, or most of the time. In total, 15.3% of the participants were unmarried. Unadjusted correlations between baseline control variables included in the analyses and isolation and loneliness were evaluated (Table 1). Being more socially isolated or lonelier was associated with being females, older, less educated, living in rural area, being a drinker, having more ADL and IADL disabilities, and a higher level of depressive symptoms. Diagnosed hypertension was positively associated with loneliness but negatively associated with isolation, as was diagnosed heart diseases. Diagnosed diabetes was only negatively associated with isolation. Loneliness was positively associated with every sub-dimension of isolation and moderately correlated with the total score of isolation ($\rho =$

Table 1. Baseline characteristics of the participants ($n = 7761$)

Characteristic	Mean (s.d.), or, %	Correlation with loneliness	Correlation with social isolation
Age, M (s.d.), y	60.97 (7.31)	0.060***	0.106***
Gender (male, %)	50.8	-0.100***	-0.060***
Education (less than lower secondary education, %)	88.7	-0.080***	-0.131***
Residence (live in the urban area, %)	37.7	0.102***	0.095***
Hypertension, %	28.4	0.048***	-0.002
Diabetes, %	6.8	0.005	-0.042***
Heart diseases, %	14.0	0.058***	-0.029*
Smoking, %	32.9	-0.022	-0.024*
Drinking, %	34.1	-0.047***	-0.061***
ADL disability, %	16.2	0.190***	0.102***
IADL disability, %	20.0	0.175***	0.102***
CESD-9, M (s.d.)	6.80 (5.79)	0.454***	0.152***
Baseline cognitive function			
Episodic memory, M (s.d.)	3.50 (1.63)	-0.118***	-0.143***
Mental status, M (s.d.)	6.57 (2.84)	-0.184***	-0.172***
Loneliness, M (s.d.)	1.52 (0.94)	-	0.165***
Social isolation, M (s.d.)	0.76 (0.67)	0.187***	-
Not married, %	15.3	0.225***	0.487***
Less than weekly contact with children, %	8.2	0.048***	0.379***
Not participate in social activities, %	52.5	0.044***	0.768***

M, mean; s.d., standard deviation; ADL, activities of daily living; IADL, instrumental activities of daily living; CESD, Center for Epidemiologic Studies Depression scale. * $p < 0.05$, ** $p < 0.01$, *** $p < 0.001$.

0.17, $p < 0.001$). After 4 years, the mean scores on both episodic memory (from 3.50 to 3.12) and mental status (from 6.57 to 6.11) were significantly lower ($t = 18.07$, $p < 0.001$ for memory; $t = 14.93$, $p < 0.001$ for mental status).

Tables 2 and 3 show the results of lagged regression on episodic memory and mental status, respectively. Loneliness was significantly associated with cognitive decline (episodic memory: $\beta = -0.03$, $p = 0.002$; mental status: $\beta = -0.03$, $p = 0.002$) after 4 years when five control variables including age, gender, education, area of residence, and baseline episodic memory or mental status score were adjusted (Model 1A). These associations remain significant when other covariates such as chronic diseases, health behaviors, and disabilities were included in the models (Model 1B). However, these associations were no longer significant (episodic memory: $\beta = -0.02$, $p = 0.179$; mental status: $\beta = -0.02$, $p = 0.088$) when CESD-9 were adjusted (Model 1C). In contrast, greater isolation was significantly associated with lower scores on episodic memory ($\beta = -0.04$, $p < 0.001$) and mental status ($\beta = -0.03$, $p = 0.003$) 4 years later even after all the covariates including CESD-9 scores were taken into account (Model 2C). Adding loneliness to the model did not reduce the association with isolation (Model 3). The interaction terms between isolation and loneliness were not statistically significant for episodic memory (p for interaction = 0.690) or mental status (p for interaction = 0.050), indicating that there was no synergistic effect between isolation and loneliness on cognitive decline. Isolation \times gender interaction was tested by adding the terms into the fully adjusted model including isolation and loneliness. This interaction term

was neither significant for episodic memory (p for interaction = 0.934) nor for mental status (p for interaction = 0.224). No significant interaction between gender and loneliness was found for either cognitive measure (all p for interaction > 0.5).

To reduce the risk of reverse causation in analyses, we reran Model 3 after excluding the respondents with very low cognition scores at the baseline survey (bottom 10%), on the grounds that people with very impaired cognition may find it difficult to engage socially with others. The results were unchanged [for episodic memory: isolation ($\beta = -0.04$, $p = 0.001$); loneliness ($\beta = -0.02$, $p = 0.064$); for mental status: isolation ($\beta = -0.03$, $p = 0.013$); loneliness ($\beta = -0.02$, $p = 0.111$)].

Discussion

In this study, a large representative sample of Chinese older adults was selected to examine simultaneously the associations of social isolation and loneliness with cognitive decline over a 4-year follow-up period. In accordance with our hypotheses, both loneliness and social isolation showed association with decreases in episodic memory and mental status. However, this association becomes less significant for loneliness when all the other confounding variables, especially depressive symptoms, were taken into account. While the association for social isolation seems independent of loneliness and other confounding variables. Our results seem to indicate a stronger negative relationship between social isolation than loneliness and cognitive function, which is consistent with two recent studies. For instance, Griffin et al.

Table 2. Predicting episodic memory at follow-up

	Model 1			Model 2			Model 3
	A (β)	B (β)	C (β)	A (β)	B (β)	C (β)	(β)
Age	-0.22***	-0.22***	-0.22***	-0.22***	-0.22***	-0.22***	-0.22***
Gender	0.01	0.02	0.02	0.01	0.02	0.02	0.02
Education	0.13***	0.13***	0.13***	0.13***	0.13***	0.12***	0.12***
Residence	-0.11***	-0.10***	-0.10***	-0.11***	-0.10***	-0.10***	-0.10***
Baseline EM	0.33***	0.32***	0.32***	0.33***	0.32***	0.32***	0.32***
Loneliness	-0.03**	-0.03*	-0.02	-	-	-	-0.01
Isolation	-	-	-	-0.05***	-0.04***	-0.04***	-0.04***
Hypertension	-	-0.00	-0.00	-	-0.00	-0.00	-0.00
Diabetes	-	-0.00	-0.00	-	-0.01	-0.00	-0.00
Heart diseases	-	0.05***	0.05***	-	0.05***	0.05***	0.05***
Smoking	-	-0.03*	-0.03*	-	-0.03*	-0.03*	-0.03*
Drinking	-	0.01	0.01	-	0.01	0.01	0.01
ADL disability	-	0.00	0.01	-	0.00	0.01	0.01
IADL disability	-	-0.06***	-0.06***	-	-0.06***	-0.06***	-0.06***
CESD-9	-	-	-0.03*	-	-	-0.03*	-0.03*

EM, episodic memory; ADL, activities of daily living; IADL, instrumental activities of daily living; CESD, Center for Epidemiologic Studies Depression scale.

β = standardized regression coefficient.

* $p < 0.05$, ** $p < 0.01$, *** $p < 0.001$.

Table 3. Predicting mental status at follow-up

	Model 1			Model 2			Model 3
	A (β)	B (β)	C (β)	A (β)	B (β)	C (β)	(β)
Age	-0.11***	-0.11***	-0.11***	-0.11***	-0.11***	-0.11***	-0.11***
Gender	0.12***	0.14***	0.13***	0.12***	0.14***	0.14***	0.13***
Education	0.08***	0.08***	0.08***	0.08***	0.08***	0.08***	0.08***
Residence	-0.08***	-0.08***	-0.07***	-0.08***	-0.08***	-0.08***	-0.07***
Baseline MS	0.49***	0.49***	0.48***	0.49***	0.48***	0.48***	0.48***
Loneliness	-0.03**	-0.03**	-0.02	-	-	-	-0.02
Isolation	-	-	-	-0.03**	-0.03**	-0.03**	-0.02**
Hypertension	-	0.02	0.02	-	0.01	0.01	0.01
Diabetes	-	-0.00	-0.00	-	-0.00	-0.00	-0.00
Heart diseases	-	0.02*	0.03**	-	0.02*	0.02*	0.02*
Smoking	-	-0.03*	-0.03*	-	-0.03**	-0.03**	-0.03*
Drinking	-	0.01	0.01	-	0.00	0.00	0.00
ADL disability	-	0.01	0.01	-	0.01	0.01	0.01
IADL disability	-	-0.03**	-0.03**	-	-0.04***	-0.03**	-0.03***
CESD-9	-	-	-0.03*	-	-	-0.03**	-0.02*

MS, mental status; ADL, activities of daily living; IADL, instrumental activities of daily living; CESD, Center for Epidemiologic Studies Depression scale.

β = standardized regression coefficient.

* $p < 0.05$, ** $p < 0.01$, *** $p < 0.001$.

found that social isolation was longitudinally associated with worse cognitive performance in older Americans whereas loneliness only cross-sectionally correlated with lower cognitive

function (Griffin et al., 2018). Another longitudinal study with German older adults also found that social isolation rather than loneliness can predict cognitive health (Beller & Wagner, 2018).

Substantial evidence indicated that loneliness is associated with a higher risk of cognitive decline at older ages (Boss et al., 2015), and may contribute to the development of dementia (Rafnsson et al., 2017; Wilson et al., 2007). However, contradictory evidence also exists that indicates no independent association between loneliness and cognitive decline (Okely & Deary, 2018; Rawtaer et al., 2017). Two studies with Chinese older adults found a significant association between loneliness and cognitive decline. Although loneliness was measured with one single item in these two studies as ours, depressive symptoms were not controlled in their models (Zhong et al., 2016, 2017). Our study does indicate negative associations between loneliness and cognitive function. However, these relationships become non-significant when all the confounding variables including depressive symptoms were controlled. It is notable that in our study, the associations between loneliness and baseline poor health was stronger than the associations between social isolation and the same health index, especially for cardiovascular health and depressive symptoms. Therefore, our finding that loneliness no longer was associated with cognitive decline after these confounding variables had been controlled may reflect its association with baseline health. Accounting for depressive symptoms did more to reduce the association between cognitive function than accounting for the clinical and behavioral covariates. Loneliness was measured by one item from CESD in our study, and the two are significantly correlated ($\rho = 0.45$, $p < 0.001$). Thus, it is not a surprise that controlling for depressive symptoms would reduce the size of the association. Such a result is consistent with previous studies. In a longitudinal study of Scottish aging adults, the associations between loneliness and cognitive ability were no longer significant when the model included depressive symptoms as covariates (Gow et al., 2013). Similar results were also found for studies of mortality (Steptoe et al., 2013; Teguio et al., 2016). In a cohort study adopted a same measurement of loneliness as ours, the association between loneliness and mortality also became insignificant when depressive symptoms (CESD scores excluding loneliness item) were adjusted (Teguio et al., 2016). These results may not imply that loneliness is not important but rather indicate that the experience of loneliness may be characteristic of people who already have mental health problems. Future studies with more independent measurements of loneliness and depressive symptoms are warranted to clarify this issue.

Nevertheless, convincing evidence, including the present study, highlights the associations between social isolation and cognitive decline (Kuiper et al., 2016). Several theories have been proposed to explain the association of social isolation and cognitive function. One is the 'use it or lose it' theory (Hultsch, Hertzog, Small, & Dixon, 1999), which argues that engagement in intellectual, physical, and social activities stimulates the brain. Decrease in engagement in social activities may result in the lack of use of mental faculties that may in turn lead to a decline of cognitive ability. Another theory is stress-buffering, proposing that social relationships are beneficial in stressful situations (Fratiglioni, Paillard-Borg, & Winblad, 2004). Stress has been associated with cognitive decline due to structural changes in the hippocampus (Wilson et al., 2003). Social relationships may prevent or modulate responses to stressful events that are damaging to health.

Although previous studies provide evidence for the negative associations between social relationships and cognitive function, it should be noted that most findings are based on a Western sample and hence a limited cultural context (Courtin & Knapp, 2017;

Evans et al., 2018; Kuiper et al., 2016). Westerners tend to have higher objective social isolation compared with non-Western populations. Taking the USA as an example, the percentage of single-person households within the same historical period is almost three times that of China (Hu & Peng, 2015). Furthermore, Western countries tend to be more individualistic from a cultural viewpoint (Hofstede, Hofstede, & Minkov, 2005). Thus, our study expands knowledge about the role of social relationships in a more collectivistic culture.

Strengths of our study include the large sample size and the fact that CHARLS is designed to be representative of the community-dwelling Chinese population aged ≥ 50 . Cognitive function was assessed with a series of objective tests, and the study benefited from the detailed measures of potential confounders. The study also has limitations. First, loneliness was assessed with only one direct question regarding the perception of loneliness in the last week. Despite wide use in the literature and strong correlations with several established multiple-item scales, this measure may be less reliable than a composite measure that taps multiple aspects of loneliness (Holwerda et al., 2014; Petersen et al., 2016; Victor, Grenade, & Boldy, 2005). The effects of loneliness became insignificant when depressive symptoms were adjusted, which also may be because the loneliness item is derived from the CESD-10. However, another study using a more complex and independent measure of loneliness has reported similar findings as ours (Griffin et al., 2018). Second, the present study investigated the consequences of loneliness and social isolation longitudinally with a relative short-term of 4 years. This prevents us from examining the long-term association of social relationships with cognitive function, and whether certain aspects of social relationships become more or less important over time. Future work could test the relationship using longer-term panel data of Chinese population. Third, CHARLS does not give full neuropsychological testing for the large sample at baseline, and thus cannot provide formal diagnoses of mild cognitive impairment or dementia. However, we have applied a widely used way to exclude the respondents with very low cognition scores (bottom 10%) at the baseline survey (Ganguli et al., 1993; Weuve et al., 2004). A sensitivity analysis after the exclusion yielded similar results as the primary analysis. Finally, this is an observational study, and causal conclusions cannot be drawn. Although we took multiple covariates into account, there may be other unmeasured factors responsible for the associations recorded here.

Conclusion

In this prospective study, high levels of social isolation were associated with an increased risk of cognitive decline. By contrast, loneliness was not linked with the cognitive decline when other confounding variables were controlled. These findings expand our knowledge about the association of social relationships with cognitive function in non-Western populations. Cognitive decline is a strong risk factor for the development of dementia (Bennett et al., 2002). Efforts to reduce isolation may therefore have substantial benefits in terms of preventing dementia among older adults in China.

Acknowledgments. This work was supported by the National Social Science Foundation, China (grant numbers 18BSH118).

Conflict of interest. None.

References

- Beller, J., & Wagner, A. (2018). Disentangling loneliness: Differential effects of subjective loneliness, network quality, network size, and living alone on physical, mental, and cognitive health. *Journal of Aging and Health, 30*(4), 521–539.
- Bennett, D. A., Wilson, R. S., Schneider, J. A., Evans, D. A., Beckett, L. A., Aggarwal, N. T., ... Bach, J. (2002). Natural history of mild cognitive impairment in older persons. *Neurology, 59*(2), 198–205. doi: 10.1212/wnl.59.2.198
- Boss, L., Kang, D.-H., & Branson, S. (2015). Loneliness and cognitive function in the older adult: A systematic review. *International Psychogeriatrics, 27*(4), 541–553.
- Cacioppo, J. T., & Hawkey, L. C. (2009). Perceived social isolation and cognition. *Trends in Cognitive Sciences, 13*(10), 447–454.
- Chan, K. Y., Wang, W., Wu, J. J., Liu, L., Theodoratou, E., Car, J., ... Campbell, H. (2013). Epidemiology of Alzheimer's disease and other forms of dementia in China, 1990–2010: A systematic review and analysis. *The Lancet, 381*(9882), 2016–2023.
- Cornwell, E. Y., & Waite, L. J. (2009). Social disconnectedness, perceived isolation, and health among older adults. *Journal of Health and Social Behavior, 50*(1), 31–48.
- Courtin, E., & Knapp, M. (2017). Social isolation, loneliness and health in old age: A scoping review. *Health & Social Care in the Community, 25*(3), 799–812.
- Crimmins, E. M., Kim, J. K., Langa, K. M., & Weir, D. R. (2011). Assessment of cognition using surveys and neuropsychological assessment: The Health and Retirement Study and the Aging, Demographics, and Memory Study. *Journals of Gerontology Series B: Psychological Sciences and Social Sciences, 66*(suppl_1), i162–i171.
- Evans, I. E., Martyr, A., Collins, R., Brayne, C., & Clare, L. (2018). Social isolation and cognitive function in later life: A systematic review and meta-analysis. *Journal of Alzheimer's disease, 70*, S119–S144.
- Fratiglioni, L., Paillard-Borg, S., & Winblad, B. (2004). An active and socially integrated lifestyle in late life might protect against dementia. *The Lancet Neurology, 3*(6), 343–353.
- Fung, A. W., Lee, A. T., Cheng, S.-T., & Lam, L. C. (2019). Loneliness interacts with family relationship in relation to cognitive function in Chinese older adults. *International Psychogeriatrics, 31*(4), 467–475.
- Ganguli, M., Belle, S., Ratcliff, G., Seaberg, E., Huff, F. J., von der Porten, K., & Kuller, L. H. (1993). Sensitivity and specificity for dementia of population-based criteria for cognitive impairment: The MoVIES project. *Journal of Gerontology, 48*(4), M152–M161.
- Glei, D. A., Goldman, N., Ryff, C. D., Lin, Y.-H., & Weinstein, M. (2012). Social relationships and inflammatory markers: An analysis of Taiwan and the US. *Social Science & Medicine, 74*(12), 1891–1899.
- Gow, A. J., Corley, J., Starr, J. M., & Deary, I. J. (2013). Which social network or support factors are associated with cognitive abilities in old age? *Gerontology, 59*(5), 454–463.
- Griffin, S. C., Mezuk, B., Williams, A. B., Perrin, P. B., & Rybarczyk, B. D. (2018). Isolation, not loneliness or cynical hostility, predicts cognitive decline in older Americans. *Journal of Aging and Health, 32*(1), 52–60.
- Hofstede, G., Hofstede, G. J., & Minkov, M. (2005). *Cultures and organizations: Software of the mind*. New York: McGraw-Hill.
- Holt-Lunstad, J., Smith, T. B., Baker, M., Harris, T., & Stephenson, D. (2015). Loneliness and social isolation as risk factors for mortality: A meta-analytic review. *Perspectives on Psychological Science, 10*(2), 227–237.
- Holwerda, T. J., Deeg, D. J., Beekman, A. T., van Tilburg, T. G., Stek, M. L., Jonker, C., & Schoevers, R. A. (2014). Feelings of loneliness, but not social isolation, predict dementia onset: Results from the Amsterdam Study of the Elderly (AMSTEL). *Journal of Neurology Neurosurgery & Psychiatry, 85*(2), 135–142.
- Hu, Z., & Peng, X. (2015). Household changes in contemporary China: An analysis based on the four recent censuses. *The Journal of Chinese Sociology, 2*(1), 9.
- Hultsch, D. F., Hertzog, C., Small, B. J., & Dixon, R. A. (1999). Use it or lose it: Engaged lifestyle as a buffer of cognitive decline in aging? *Psychology and Aging, 14*(2), 245–263.
- Johnson, D. (2005). Two-wave panel analysis: Comparing statistical methods for studying the effects of transitions. *Journal of Marriage and Family, 67*(4), 1061–1075.
- Kuiper, J. S., Zuidersma, M., Zuidema, S. U., Burgerhof, J. G., Stolk, R. P., Oude Voshaar, R. C., & Smidt, N. (2016). Social relationships and cognitive decline: A systematic review and meta-analysis of longitudinal cohort studies. *International Journal of Epidemiology, 45*(4), 1169–1206.
- Lei, X., Hu, Y., McArdle, J. J., Smith, J. P., & Zhao, Y. (2012). Gender differences in cognition among older adults in China. *Journal of Human Resources, 47*(4), 951–971.
- Lei, X., Sun, X., Strauss, J., Zhao, Y., Yang, G., Hu, P., ... Yin, X. (2014). Health outcomes and socio-economic status among the mid-aged and elderly in China: Evidence from the CHARLS national baseline data. *The Journal of the Economics of Ageing, 3*, 29–43. doi: <https://doi.org/10.1016/j.jeoa.2014.05.001>
- Luo, Y., & Waite, L. J. (2014). Loneliness and mortality among older adults in China. *Journals of Gerontology Series B: Psychological Sciences and Social Sciences, 69*(4), 633–645.
- McArdle, J. J., Fisher, G. G., & Kadlec, K. M. (2007). Latent variable analyses of age trends of cognition in the Health and Retirement Study, 1992–2004. *Psychology and Aging, 22*(3), 525–545.
- McHugh, J., Kenny, R., Lawlor, B., Steptoe, A., & Kee, F. (2017). The discrepancy between social isolation and loneliness as a clinically meaningful metric: Findings from the Irish and English longitudinal studies of ageing (TILDA and ELISA). *International Journal of Geriatric Psychiatry, 32*(6), 664–674.
- Newall, N. E., & Menec, V. H. (2019). Loneliness and social isolation of older adults: Why it is important to examine these social aspects together. *Journal of Social and Personal Relationships, 36*(3), 925–939.
- Nummela, O., Seppanen, M., & Uutela, A. (2011). The effect of loneliness and change in loneliness on self-rated health (SRH): A longitudinal study among aging people. *Archives of Gerontology and Geriatrics, 53*(2), 163–167. doi: 10.1016/j.archger.2010.10.023
- Okely, J. A., & Deary, I. J. (2018). Longitudinal associations between loneliness and cognitive ability in the Lothian Birth Cohort 1936. *The Journals of Gerontology: Series B, 74*(8), 1376–1386.
- Pan, X., Luo, Y., & Roberts, A. R. (2018). Secondhand smoke and women's cognitive function in China. *American Journal of Epidemiology, 187*(5), 911–918.
- Peplau, L. A., & Perlman, D. (1982). *Loneliness: A Sourcebook of Current Theory, Research and Therapy*. New York: Wiley-Interscience.
- Petersen, J., Kaye, J., Jacobs, P. G., Quinones, A., Dodge, H., Arnold, A., & Thielke, S. (2016). Longitudinal relationship between loneliness and social isolation in older adults: Results from the cardiovascular health study. *Journal of Aging and Health, 28*(5), 775–795. doi: 10.1177/0898264315611664
- Rafnsson, S. B., Orrell, M., d'Orsi, E., Hogervorst, E., & Steptoe, A. (2017). Loneliness, social integration, and incident dementia over 6 years: Prospective findings from the English Longitudinal Study of Ageing. *The Journals of Gerontology: Series B, 75*(1), 114–124.
- Rawtaer, I., Gao, Q., Nyunt, M. S. Z., Feng, L., Chong, M. S., Lim, W. S., ... Ng, T. P. (2017). Psychosocial risk and protective factors and incident mild cognitive impairment and dementia in community dwelling elderly: Findings from the Singapore Longitudinal Ageing Study. *Journal of Alzheimer's Disease, 57*(2), 603–611.
- Shankar, A., Hamer, M., McMunn, A., & Steptoe, A. (2013). Social isolation and loneliness: Relationships with cognitive function during 4 years of follow-up in the English Longitudinal Study of Ageing. *Psychosomatic Medicine, 75*(2), 161–170.
- Shumaker, S. A., & Hill, D. R. (1991). Gender differences in social support and physical health. *Health Psychology, 10*(2), 102–111.
- Shye, D., Mullooly, J. P., Freeborn, D. K., & Pope, C. R. (1995). Gender differences in the relationship between social network support and mortality: A longitudinal study of an elderly cohort. *Social Science & Medicine, 41*(7), 935–947.
- Steptoe, A., Shankar, A., Demakakos, P., & Wardle, J. (2013). Social isolation, loneliness, and all-cause mortality in older men and women. *Proceedings of the National Academy of Sciences, 110*(15), 5797–5801.

- Teguo, M. T., Simo-Tabue, N., Stoykova, R., Meillon, C., Cogne, M., Amiéva, H., & Dartigues, J.-F. (2016). Feelings of loneliness and living alone as predictors of mortality in the elderly: The PAQUID study. *Psychosomatic Medicine*, 78(8), 904–909.
- Tilvis, R. S., Pitkala, K. H., Jolkkonen, J., & Strandberg, T. E. (2000). Social networks and dementia. *Lancet (London, England)*, 356(9223), 77–78.
- UN DESA (2017). World population prospects, United Nations Department of Economic and Social Affairs. Population Division.
- Victor, C., Grenade, L., & Boldy, D. (2005). Measuring loneliness in later life: A comparison of differing measures. *Reviews in Clinical Gerontology*, 15(01), 63–70.
- Wang, H.-X., Jin, Y., Hendrie, H. C., Liang, C., Yang, L., Cheng, Y., ... Murrell, J. R. (2012). Late life leisure activities and risk of cognitive decline. *Journals of Gerontology Series A: Biomedical Sciences and Medical Sciences*, 68(2), 205–213.
- Weuve, J., Kang, J. H., Manson, J. E., Breteler, M. M., Ware, J. H., & Grodstein, F. (2004). Physical activity, including walking, and cognitive function in older women. *JAMA*, 292(12), 1454–1461.
- Wilson, R., Evans, D., Bienias, J., De Leon, C. M., Schneider, J., & Bennett, D. (2003). Proneness to psychological distress is associated with risk of Alzheimer's disease. *Neurology*, 61(11), 1479–1485.
- Wilson, R. S., Krueger, K. R., Arnold, S. E., Schneider, J. A., Kelly, J. F., Barnes, L. L., ... Bennett, D. A. (2007). Loneliness and risk of Alzheimer disease. *Archives of general psychiatry*, 64(2), 234–240.
- Yang, K., & Victor, C. R. (2008). The prevalence of and risk factors for loneliness among older people in China. *Ageing & Society*, 28(3), 305–327.
- Zhao, Y., Hu, Y., Smith, J. P., Strauss, J., & Yang, G. (2012). Cohort profile: The China health and retirement longitudinal study (CHARLS). *International Journal of Epidemiology*, 43(1), 61–68.
- Zhong, B.-L., Chen, S.-L., & Conwell, Y. (2016). Effects of transient versus chronic loneliness on cognitive function in older adults: Findings from the Chinese Longitudinal Healthy Longevity Survey. *The American Journal of Geriatric Psychiatry*, 24(5), 389–398.
- Zhong, B.-L., Chen, S.-L., Tu, X., & Conwell, Y. (2017). Loneliness and cognitive function in older adults: Findings from the Chinese longitudinal healthy longevity survey. *The Journals of Gerontology: Series B*, 72(1), 120–128.