

Original Article

Interventions to improve healthcare workers' hand hygiene compliance: A systematic review of systematic reviews

Lesley Price PhD¹, Jennifer MacDonald PhD¹, Lucyna Gozdziewska BA¹, Tracey Howe PhD², Paul Flowers PhD¹, Lesley Shepherd MSc³, Yvonne Watt MSc² and Jacqui Reilly PhD^{1,3}

¹Safeguarding Health through Infection Prevention Research Group, School of Health and Life Sciences, Glasgow Caledonian University, Glasgow, Scotland, United Kingdom, ²School of Health and Life Sciences, Glasgow Caledonian University, Glasgow, Scotland, United Kingdom and ³Health Protection Scotland, Glasgow, Scotland, United Kingdom

Abstract

Objective: To synthesize the existing evidence base of systematic reviews of interventions to improve healthcare worker (HCW) hand hygiene compliance (HHC).

Methods: PRISMA guidelines were followed, and 10 information sources were searched in September 2017, with no limits to language or date of publication, and papers were screened against inclusion criteria for relevance. Data were extracted and risk of bias was assessed.

Results: Overall, 19 systematic reviews (n=20 articles) were included. Only 1 article had a low risk of bias. Moreover, 15 systematic reviews showed positive effects of interventions on HCW HHC, whereas 3 reviews evaluating monitoring technology did not. Findings regarding whether multimodal rather than single interventions are preferable were inconclusive. Targeting social influence, attitude, self-efficacy, and intention were associated with greater effectiveness. No clear link emerged between how educational interventions were delivered and effectiveness.

Conclusions: This is the first systematic review of systematic reviews of interventions to improve HCW HHC. The evidence is sufficient to recommend the implementation of interventions to improve HCW HHC (except for monitoring technology), but it is insufficient to make specific recommendations regarding the content or how the content should be delivered. Future research should rigorously apply behavior change theory, and recommendations should be clearly described with respect to intervention content and how it is delivered. Such recommendations should be tested for longer terms using stronger study designs with clearly defined outcomes.

(Received 15 May 2018; accepted 10 September 2018)

Healthcare-associated infections (HAIs) have serious consequences for patients and healthcare systems, leading to longer hospital stays, increased mortality and morbidity, and financial burden.^{1,2} In Europe, ~80,000 hospital patients suffer at least 1 HAI on any given day, yielding an overall prevalence of 5.7%.³ HAIs also affect millions of patients worldwide annually.^{3–7}

Organisms that cause HAIs can be transmitted to patients through healthcare worker (HCW) hands contaminated by patient contact or touching the patient environment.² Effective hand hygiene (HH) is thus critical to preventing HAIs,⁸ which is reflected by the emphasis on HCW compliance with HH guidelines. Such HH guidance relates to both opportunity and technique. Opportunity concerns *when* to do HH, and the World

Health Organization (WHO) specify Five Moments.² Technique relates to *how* to enact HH, with 2 main procedures internationally: the Six Step² and Three Step⁹ techniques. However, HCW compliance with this guidance is suboptimal.^{2,10–17}

A major challenge within healthcare systems is how to improve hand hygiene compliance (HHC) among HCWs. An early systematic review identified 21 primary studies evaluating HH interventions for HCWs.¹⁸ A plethora of primary studies and a growing number of systematic reviews have since assessed the effectiveness of interventions to improve HCW HHC. To assist practitioners in optimizing HHC among HCWs, this review aims to synthesize the evidence base of systematic reviews of interventions to improve HCW HHC.

Author for correspondence: Lesley Price, Safeguarding Health through Infection Prevention Research Group, School of Health and Life Sciences, Glasgow Caledonian University, Cowcaddens Road, Glasgow, G4 0BA, UK. E-mail: L.Price@gcu.ac.uk

PREVIOUS PRESENTATION: Results based on a search of the literature up to June 2016 were presented on September 19, 2017, at the Annual Conference of the Infection Prevention Society, held in the Manchester Central Convention Complex, England, United Kingdom. This presentation resulted in the following abstract: Price L, MacDonald J, Gozdziewska L, et al. An overview of systematic reviews of interventions designed to improve healthcare workers' hand hygiene compliance. *J Infect Prev* 2017;18: S56–S57.

Cite this article: Price L, et al. (2018). Interventions to improve healthcare workers' hand hygiene compliance: A systematic review of systematic reviews. *Infection Control & Hospital Epidemiology* 2018, 39, 1449–1456. doi: 10.1017/ice.2018.262

© 2018 by The Society for Healthcare Epidemiology of America. All rights reserved.

Methods

Protocol and registration

This systematic review followed a published protocol¹⁹ and is reported according to the PRISMA guidelines.²⁰

Inclusion criteria

Systematic reviews were included if they evaluated any intervention to improve HHC among HCWs. Interventions could

have no comparator or be compared to usual care, another intervention, or historical control. Systematic reviews were required to report HHC as the primary outcome. Other outcomes of interest included bacterial load on HCW hands, HAI rates, organizational culture, and psychological variables. No restrictions were placed on the design of primary studies within systematic reviews.

Information sources and search

In September 2017, we searched 4 databases (CINAHL, EMBASE, MEDLINE, and PsycINFO) and 6 specialist registers (Cochrane Database of Systematic Reviews, Database of Abstracts of Reviews of Effectiveness, Epistemonikos, Joanna Briggs Institute Database of Systematic Reviews and Implementation Reports, Health Technology Assessment Database, and PROSPERO). No language or date of publication restrictions were applied. The search included index terms and text words relating to HH²¹ and systematic review methods.²² Database searches were broadly similar; modifications were made to account for minor differences in functionality (see Supplemental Table 1 for MEDLINE search). Due to limited functionality of specialist registers (except the Cochrane Database of Systematic Reviews), these searches were restricted to HH text words. We also manually searched the reference lists of included systematic reviews.

Systematic review selection

Systematic review selection was conducted in 2 stages, with all papers assessed by 2 independent reviewers. First, titles and abstracts of included papers were screened against the inclusion criteria. Second, papers that appeared to meet the inclusion criteria or lacked sufficient information to allow an informed judgement on relevance underwent full-text review. Disagreements were resolved via discussion or referral to a third reviewer.

Data collection and risk of bias within systematic reviews

A standardized tool was devised for data extraction (Supplemental Table 2). Risk of bias within systematic reviews was assessed using the ROBIS tool (Supplemental Table 3).²³ Data were extracted and risk of bias was assessed by 2 independent reviewers for 25% (n = 5) of systematic reviews. The remaining systematic reviews were data extracted and assessed for risk of bias by 1 reviewer and checked by another. Disagreements were resolved through discussion or referral to a third reviewer.

Synthesis

Findings were synthesized following the Economic and Social Research Council's guidance for narrative synthesis.²⁴

Results

Systematic review selection

The search yielded 993 papers (Fig. 1). Following the removal of duplicates, 566 unique papers remained; all were screened against the inclusion criteria. Most papers (n = 481) were discarded at the title or abstract stage, and 65 were excluded by full-text review (Fig. 1). Overall, 19 systematic reviews (n = 20 articles) were included.^{18,25–43} Reference list checks did not identify any further papers.

Systematic review characteristics

The characteristics of the 19 systematic reviews are summarized in Supplemental Table 4. Overall, 15 narrative syntheses,^{18,25–29,31,32,35,36/37,39–43} 3 meta-analyses,^{33,34,38} and 1 network meta-analysis³⁰ were published between 2001 and 2017, with 15 published after 2010.^{25–33,38–43} Primary studies in included systematic reviews were published from 1986 to 2016 and ranged in number from 3⁴¹ to 73.³² Collectively,¹ 236 unique primary studies were cited. However, some primary studies were included in >1 systematic review. The degree of overlap has been quantified and presented in a transparent manner^{44,45}: 139 (58.9%) primary studies were cited once; 46 (19.5%) were cited twice; 25 (10.6%) were cited 3 times; 17 (7.2%) were cited 4 times; 7 (3.0%) were cited 5 times; and 2 (<1%) were cited 6 times.

Countries and healthcare settings

In all but 2 systematic reviews where inclusion was limited to primary studies conducted in developed³² or low- or middle-income countries,⁴² systematic reviews were open to primary studies from all countries. Regarding healthcare settings, 13 systematic reviews included primary studies conducted in hospitals.^{25,26,28,30,31,33–35,39–43} In addition, 6 systematic reviews included primary studies conducted in hospitals in addition to elder care homes;²⁹ nursing homes;³² long-term care facilities;^{27,32,36/37,38} care homes for people with disabilities;¹⁸ and/or primary care facilities.^{27,29}

Population

All systematic reviews were open to primary studies of any type of HCW, with the exception of Doronina *et al*,²⁶ which specified a particular professional group (nurses). Most systematic reviews included data from a range of HCWs (eg, nurses, doctors, healthcare assistants, and students), and 6 systematic reviews included at least 1 primary study (n = 1,^{28,30,33,43} n = 3,⁴⁰ and n = 6³²) with data from patients or visitors or relatives, but the proportions of the overall samples that were not HCWs were not reported.

Interventions

With regard to types of interventions, 11 systematic reviews took an inclusive approach. Others focused on the introduction of alcohol-based hand rub (ABHR),³⁴ ABHR accessibility,⁴¹ educational interventions,²⁵ interventions using psychological theory,³⁹ monitoring technology,^{31,40,43} or quality improvement strategies.³⁵ Supplemental Table 5 illustrates how the content of interventions evaluated in primary studies of each systematic review mapped onto the WHO multimodal strategy for HH.² The most frequent component was 'observation and feedback,' which was mapped in all but 1 systematic review,⁴¹ followed by 'training and education' (n = 16)^{18,25–39,42} and 'reminders' (n = 15).^{18,25–33,35,38,40,43} The least common component was 'safety climate,' which was mapped in 10 systematic reviews.^{25–30,32,35,38,43}

¹Excluding primary studies in Ward *et al*⁴³ because it is unclear exactly how many reported HHC and/or HAI outcomes and only including 8 primary studies in Kingston *et al*²⁹ with baseline and post-intervention HHC data, upon which conclusions about effectiveness were based.




Fig. 1. PRISMA flow diagram outlining the systematic review selection process.

Outcomes

Hand hygiene compliance was measured by direct observation (n = 13),^{18,26–30,32,33,36/37,38,39,41,42} unobtrusive observation (n = 2),^{26,28} video camera (n = 4),^{27,30,38,43} mobile handheld devices (n = 1),⁴³ electronic monitoring (n = 9),^{32,36/37,42} or self-report (n = 3).^{32,36/37,42} Proxy measures, such as rate or number of HH events (n = 4),^{27,28,30,40} ABHR consumption or soap use (n = 11),^{18,26,31,36/37,38,40,41} and procurement of ABHR or soap (n = 2),^{27,39} were also employed. A lack of longer-term evaluation of HHC was observed in 11 systematic reviews.^{18,25–30,32,33,36/37,42} HAI rates were reported in 11 systematic reviews.^{25,27,30,31,33,35,36/37,39,40,42,43} Bacterial load on HCW hands, organizational culture, and psychological variables were not reported in any systematic reviews.

Study designs

In 3 systematic reviews,^{26,27,30} primary studies were required to meet the Cochrane Effective Practice and Organization of Care methodological criteria for randomized controlled trials (RCTs), non-randomized trials, controlled before-and-after studies, or interrupted time series (ITS).⁴⁶ Across the other systematic reviews, before-and-after studies were common,^{18,28,31–33,35,36/37,38–40}

although 5 reviewers did not specify the design of the included primary studies.^{25,34,41–43} Although Kingston et al²⁹ described all primary studies as ‘clinical trials,’ this term was not defined, and ITS and before-and-after studies were included.⁴⁷

Risk of bias within systematic reviews

Overall, 13 systematic reviews had a high risk of bias,^{18,25,26,29,31–34,36/37,38,41–43} 5 systematic reviews had an unclear risk of bias,^{28,30,35,39,40} and 1 systematic review had a low risk of bias (Table 1).²⁷ The most common methodological weaknesses within systematic reviews related to synthesis and findings, for example, not reporting individual study results, not including all primary studies in the synthesis, or not addressing biases in the synthesis. Other common methodological weaknesses within systematic reviews were not reporting a risk of bias assessment or the process for data extraction and risk of bias assessment.

Effectiveness of HH interventions

Hand hygiene compliance. In total, 18 systematic reviews reported the overall effectiveness of interventions in improving HCW

Table 1. Risk of Bias Within Systematic Reviews

| First Author (Year) | Level of Concern | | | | | | | |
|---|--------------------------------|---|---|----------------------------------|--|---|--|----------------------------------|
| | Domain 1: Eligibility Criteria | Domain 2: Identification and Selection of Studies | Domain 3: Data Collection and Study Appraisal | Domain 4: Synthesis and Findings | All concerns identified in domains 1 to 4 addressed in interpretation of findings? | Relevance of identified studies to the review's research question appropriately considered? | Emphasizing results based on statistical significance avoided? | Overall Judgment of Risk of Bias |
| Cherry et al (2012) ²⁵ | Unclear | Unclear | Low | High | No | Probably yes | Probably no | High |
| Doronina et al (2017) ²⁶ | High | Unclear | Unclear | High | No | Yes | Probably yes | High |
| Gould et al (2017) ²⁷ | Low | Low | Low | Low | Yes | Yes | Yes | Low |
| Huis et al (2012) ²⁸ | Unclear | Unclear | Low | Low | Probably no | Yes | Yes | Unclear |
| Kingston et al (2016) ²⁹ | High | High | High | High | No | Probably yes | Yes | High |
| Luangasanatip et al (2015) ³⁰ | High | Low | Unclear | Low | Probably no | Probably yes | Yes | Unclear |
| Mitchell et al (2014) ³¹ | Unclear | High | High | High | No | Probably no | Yes | High |
| Naikoba & Hayward (2001) ¹⁸ | Unclear | High | High | High | No | Probably yes | Yes | High |
| Neo et al (2016) ³² | Unclear | High | High | High | No | Probably yes | Probably yes | High |
| Ofek Shlomai et al (2015) ³³ | Unclear | High | Low | Unclear | No | Probably yes | No | High |
| Picheansathian et al (2004) ³⁴ | High | High | High | Unclear | No | No | Probably no | High |
| Ranji et al (2007) ³⁵ | Unclear | Low | Unclear | Low | Probably no | Probably yes | Yes | Unclear |
| Ritchie et al (2005) ³⁶ and Stout et al (2007) ³⁷ | Unclear | Unclear | High | High | No | Yes | Yes | High |
| Schweizer et al (2014) ³⁸ | Unclear | Unclear | High | High | Probably no | Probably yes | Yes | High |
| Srigley et al (2015) ³⁹ | High | Low | Low | Low | Probably no | Yes | Yes | Unclear |
| Srigley et al (2015) ⁴⁰ | High | Unclear | Low | Low | Probably no | Yes | Yes | Unclear |
| Stiller et al (2016) ⁴¹ | High | Low | Unclear | High | No | Probably yes | Probably yes | High |
| Vindigni et al (2011) ⁴² | Unclear | High | High | High | No | Probably yes | No | High |
| Ward et al (2014) ⁴³ | High | High | High | High | No | No | Yes | High |

HHC (Supplemental Table 6). Also, 2 meta-analyses showed similar increases (odds ratio [OR], 2.04; 95% confidence interval [CI], 1.40–2.97³³; and Peto OR, 1.96; CI, 1.56–2.46³⁴). Luangasanatip et al³⁰ found that 18 of 22 pairwise comparisons (82%) showed both

stepwise increases in HHC during intervention implementation and a trend for increasing HHC postintervention. In 8 narrative syntheses, most or all primary studies reported significant improvements in HHC.^{25,26,32,35,36/37–39,41} Overall effectiveness was further

supported by 4 narrative syntheses, which described positive findings, largely without reference to statistical significance.^{27–29,42} Furthermore, 3 narrative syntheses on monitoring technology found scarce evidence for effectiveness in improving HHC in general.^{31,40,43}

Healthcare-associate infection rates. In 4 systematic reviews, most or all primary studies that measured HAI rates showed a reduction in HAI rates, although significance levels were not always stated.^{25,30,33,36/37} Results were more mixed in 4 other systematic reviews reporting HAI data,^{27,31,35,43} and there were no significant decreases in HAI rates in relevant primary studies included within 2 systematic reviews.^{39,40} The final systematic review did not report HAI results.⁴²

Intervention content and effectiveness

Supplemental Table 7 summarizes findings of 10 systematic reviews that considered the relationship between intervention content and effectiveness. One meta-analysis³⁸ indicated that interventions with more components, as conceptualized by WHO in their multimodal strategy for HH,² did not see larger increases in HHC.² Conversely, within the same systematic review, 2 further meta-analyses of primary studies assessing the same combination of components showed that using all components of the WHO multimodal strategy for HH (OR, 1.82; 95% CI, 1.69–1.97) seems more effective in improving HHC than including only feedback, education, and reminders (OR, 1.47; 95% CI, 1.12–1.94).³⁸ Additionally, in a network meta-analysis,³⁰ interventions that supplemented the WHO multimodal strategy for HH with incentives, goal setting, or accountability produced further improvements in HHC than ‘training and education’ or ‘system change’ (OR not reported) and the WHO multimodal strategy for HH alone (OR, 1.82; 95% CI, 0.2–12.2). Doronina et al²⁶ reached a similar conclusion in their narrative synthesis.

Naikoba and Hayward¹⁸ emphasized that combining education with written material, reminders, and continued performance feedback can have a marked effect on HHC compared to single interventions comprising reminders or regular performance feedback, which in turn are more effective than one-off education and ABHR provision. Neo et al³² also proposed that effectiveness may be enhanced by multimodal interventions, as well as facilities design and planning and financial rewards. In another narrative synthesis, multimodal interventions supplying ABHR were as conducive to improving HHC as those without.^{36/37} Meta-analytic findings suggest that providing (OR, 2.81; 95% CI, 1.32–5.96) in contrast to not providing (OR, 1.55; 95% CI, 1.13–2.11) performance feedback in a multimodal intervention is more likely to improve HHC.³³ However, the Cochrane authors concluded that it is unclear whether multimodal over single interventions are preferable, or which components add the most value.²⁷

With regard to monitoring technology, Mitchell et al³¹ reported that devices delivering a real-time reminder that HH was indicated but not actioned were consistently linked to increased HHC, while systems with periodic feedback by managers produced variable findings. However, in assessing devices that give reminders without feedback, aggregate feedback without reminders, or individual feedback and reminders, Srigley et al⁴⁰ found limited evidence to recommend any specific technology.

²Increase in HHC for interventions with 1–2 components (OR, 3.44; 95% CI, 1.11–10.68), 3–4 components (OR, 2.16; 95% CI, 1.82–2.55), and ≥ 5 components (OR, 2.49; 95% CI, 1.74–3.56).

Delivery of educational interventions and effectiveness

Cherry et al²⁵ considered the relationship between how the educational interventions were delivered and effectiveness. Delivery of education was separated into 6 groups of education with (1) demonstration; (2) no demonstration; (3) self-study; (4) video; (5) demonstration and video, and (6) an online element. However, they were unable to identify a method of delivery that was more effective than another.²⁵

Use of theoretical frameworks

Huis et al²⁸ found a significant positive correlation between the effectiveness of interventions tested in controlled studies and the number of theoretical determinants of behavior ($n=1-5$) addressed ($r=.961$; $P=.009$).³ They also noted that less commonly addressed determinants (ie, social influence, attitude, self-efficacy, and intention) were mainly targeted in interventions addressing ≥ 4 determinants.²⁸ Gould et al²⁷ reported interventions lacked convincing theoretical underpinning. Likewise, Srigley et al³⁹ observed that it was often unclear how theory informed interventions, that typically not all theoretical constructs were represented, and that measures of theoretical constructs were not always consistent with guidelines. No theoretical approach appeared more effective at improving HHC than another.³⁹

Discussion

This systematic review has, for the first time, identified, described, and synthesized the existing evidence base of systematic reviews of interventions to improve HHC among HCWs. In addition, the systematic review was conducted in a transparent and rigorous manner and benefited from a comprehensive literature search, spanning a wide period with no language restrictions. The results of 15 of 18 systematic reviews that reported overall effectiveness showed positive effects of interventions on HCW HHC, across various healthcare settings for different professional groups. However, 6 of 11 systematic reviews that extracted HAI data described mixed or nonsignificant findings. Several reviewers advocated multimodal interventions, incorporating performance feedback and extending the WHO multimodal strategy for HH over single interventions to elicit improvements in HCW HHC. Still, this conclusion was not unanimous. Regarding theory, targeting higher numbers of theoretical determinants of behavior (up to 5) appears to increase effectiveness, with interventions that address social influence, attitude, self-efficacy, and intention especially effective. There was no clear link between how educational interventions were delivered and effectiveness.

Limits of the evidence and recommendations for practice

Although a substantial number of systematic reviews showed positive effects of interventions to improve HHC among HCW, only 1 systematic review had a low risk of bias.²⁷ This systematic review concluded that there was sufficient evidence to

³One theoretical determinant ($n=3$): median relative difference (improvement), 17.6 (range, –8.8 to 61). Two theoretical determinants ($n=1$): relative difference (improvement), 25.7. Three theoretical determinants ($n=3$): median relative difference (improvement), 42.3 (range, 19.5–82.7). Four theoretical determinants ($n=2$): median relative difference (improvement), 43.9 (range, 14.8–73). Five theoretical determinants ($n=3$): median relative difference (improvement), 49.5 (range, –8.6 to 429). And 7 theoretical determinants ($n=1$): relative difference (improvement), 9.7.

recommend interventions to improve hand hygiene. However, the evidence on the optimum content and how it is should be delivered remain unclear. In addition, evidence for the use of monitoring technology is insufficient to recommend its use.^{31,40,43} Theory-informed interventions are recommended by the Cochrane systematic review.²⁷ The significant findings of Huis *et al*²⁸ suggest that targeting social influence, attitude, self-efficacy, and intention may enhance effectiveness, but the evidential basis for this is somewhat lacking.

Limits of the evidence and recommendations for future research

Caution is required when interpreting these findings because only 1 systematic review was at low risk of bias. Reviewers should use quality assessment tools and follow best-practice review guidelines.^{48–58} Transparency in reporting of methods should be ensured to minimize bias in future studies. Systematic reviews were restricted in that they largely comprised before-and-after studies; more robust study designs are required moving forward. Also, reviewers rarely conveyed the total sample size, and the type of HCW was often not further defined. These reporting omissions likely reflect the absence of this detail in primary studies. However, this information is necessary to interpret generalizability; what works in one population may differ in another.⁵⁹ Primary studies tended to run over short time periods; the longer-term effect⁶⁰ of HH interventions has not yet been established. Researchers should consider this when designing studies.

In conclusion, in this systematic review of systematic reviews, we found predominantly low-quality evidence that interventions to improve HCW HHC are effective. The evidence is sufficient to recommend the implementation of interventions to improve HCW HHC (with the exception of monitoring technology), but it is insufficient to make specific recommendations about the content and how the content should be delivered. To fill existing research gaps and to develop a more viable evidence base to enable the generation of recommendations for practice, systematic reviews should follow reporting guidelines, and primary studies need to utilize more robust research designs.

Acknowledgments.

Financial support. The Scottish Healthcare Associated Infection Prevention Institute (SHAIPi) is a research consortium set up through a grant from the Chief Scientist Office of the Scottish Infection Research Network. SHAIPi supports a program of work carried out by members of the Safeguarding Health through Infection Prevention Research Group at Glasgow Caledonian University. L.P., J.M., P.F., and J.R. conducted the work for this systematic review of systematic reviews in their SHAIPi-funded time.

Conflicts of interest. L.P. is the Director of Studies for a PhD student (LG) whose study is being supported by SureWash. L.P. is leading, and L.G. is working on, another study supported by SureWash. L.P. and L.G. have 2 SureWash Elite machines on loan for data collection purposes for these 2 studies. All other authors report no conflicts of interest relevant to this article.

Supplementary material. To view supplementary material for this article, please visit <https://doi.org/10.1017/ice.2018.262>

References

1. Marchetti A, Rossiter R. Economic burden of healthcare-associated infection in US acute care hospitals: societal perspective. *J Med Econ* 2013;16:1399–1404.
2. World Health Organization. WHO guidelines on hand hygiene in health care. World Health Organization website. <http://www.who.int/gpsc/5may/tools/9789241597906/en/>. Published 2009. Accessed January 11, 2018.
3. European Centre for Disease Prevention and Control. Summary: Point prevalence survey of healthcare associated infections and antimicrobial use in European acute care hospitals. European Centre for Disease Prevention and Control website. https://ecdc.europa.eu/sites/portal/files/media/en/healthtopics/Healthcare-associated_infections/point-prevalence-survey/Documents/healthcare-associated-infections-antimicrobial-use-PPS-summary.pdf. Published 2009. Accessed January 11, 2018.
4. Allegranzi B, Nejad SB, Combescure C, *et al*. Burden of endemic healthcare-associated infection in developing countries: SYSTEMATIC review and meta-analysis. *Lancet* 2011;377:228–241.
5. Cassini A, Plachouras D, Eckmanns T, *et al*. Burden of six healthcare-associated infections on European population health: estimating incidence-based disability-adjusted life years through a population prevalence-based modelling study. *PLoS Med* 2016;13:e1002150.
6. Ling ML, Apisarnthanarak A, Madriaga G. The burden of healthcare-associated infections in Southeast Asia: a systematic literature review and meta-analysis. *Clin Infect Dis* 2015;60:1690–1699.
7. Nejad SB, Allegranzi B, Syed SB, Ellis B, Pittet D. Health-care-associated infection in Africa: a systematic review. *Bull World Health Organ* 2011;89:757–765.
8. Allegranzi B, Pittet D. Role of hand hygiene in healthcare-associated infection prevention. *J Hosp Infect* 2009;73:305–315.
9. Boyce JM, Pittet D, Healthcare Infection Control Practices Advisory Committee, HICPAC/SHEA/APIC/IDSA Hand Hygiene Task Force. Guideline for hand hygiene in health-care settings: recommendations of the Healthcare Infection Control Practices Advisory Committee and the HICPAC/SHEA/APIC/IDSA Hand Hygiene Task Force. *MMWR Recomm Rep* 2002;51:1–45.
10. Arias AV, Garcell HG, Ochoa YR, Arias KF, Miranda FR. Assessment of hand hygiene techniques using the World Health Organization's six steps. *J Infect Public Health* 2016;9:366–369.
11. Erasmus V, Daha TJ, Brug H, *et al*. Systematic review of studies on compliance with hand hygiene guidelines in hospital care. *Infect Control Hosp Epidemiol* 2010;31:283–294.
12. Laustsen S, Lund E, Bibby BM, Kristensen B, Thulstrup AM, Møller J. Effect of correctly using alcohol-based hand rub in a clinical setting. *Infect Control Hosp Epidemiol* 2008;29:954–956.
13. Park H, Kim S, Lim Y, *et al*. Assessment of the appropriateness of hand surface coverage for healthcare workers according to World Health Organization hand hygiene guidelines. *Am J Infect Control* 2014;42:559–561.
14. Reilly JS, Price L, Lang S, *et al*. A pragmatic randomized controlled trial of 6-step vs 3-step hand hygiene technique in acute hospital care in the United Kingdom. *Infect Control Hosp Epidemiol* 2016;37:661–666.
15. Thivichon-Prince B, Barsotti O, Girard R, Morrier J. Hand hygiene practices in a dental teaching center: measures and improve. *Eur J Dent* 2014;8:481–486.
16. Tschudin-Sutter S, Sepulcri D, Dangel M, Schuhmacher H, Widmer AF. Compliance with the world health organization hand hygiene technique: a prospective observational study. *Infect Control Hosp Epidemiol* 2015;36:482–483.
17. Widmer AF, Conzelmann M, Tomic M, Frei R, Stranden AM. Introducing alcohol-based hand rub for hand hygiene: the critical need for training. *Infect Control Hosp Epidemiol* 2007;28:50–54.
18. Naikoba S, Hayward A. The effectiveness of interventions aimed at increasing handwashing in healthcare workers: a systematic review. *J Hosp Infect* 2001;47:173–180.
19. Currie K, Duguid C, Flowers P, *et al*. An overview of systematic reviews and meta-analyses of interventions designed to improve healthcare workers' hand hygiene compliance. York: PROSPERO 2016 CRD42016040027. http://www.crd.york.ac.uk/PROSPERO/display_record.asp?ID=CRD42016040027. Published 2016. Accessed January 11, 2018.

20. Moher D, Liberati A, Tetzlaff J, Altman DG, PRISMA Group. Preferred reporting items for systematic reviews and meta-analyses: the PRISMA statement. *Ann Intern Med* 2009;151:264–269.
21. Price L, MacDonald J, Melone L, *et al.* Effectiveness of national and sub-national infection prevention and control interventions in high and upper-middle income countries: outcomes of a systematic literature review. *Lancet Infect Dis* 2018;18:e159–e171.
22. Search filters. Edinburgh: Scottish Intercollegiate Guidelines Network website. <http://www.sign.ac.uk/methodology/filters.html>. Published 2013. Accessed January 11, 2018.
23. Whiting P, Savović J, Higgins JP, *et al.* ROBIS: a new tool to assess risk of bias in systematic reviews was developed. *J Clin Epidemiol* 2016;69:225–234.
24. Popay J, Roberts H, Sowden A, *et al.* Guidance on the conduct of narrative synthesis in systematic reviews. ESRC Research Methods Programme website. <http://citeseerx.ist.psu.edu/viewdoc/download?doi=10.1.1.178.3100&rep=rep1&type=pdf>. Published 2006. Accessed January 11, 2018.
25. Cherry MG, Brown JM, Bethell GS, Neal T, Shaw NJ. Features of educational interventions that lead to compliance with hand hygiene in healthcare professionals within a hospital care setting. A BEME systematic review: BEME Guide No. 22. *Med Teach* 2012;34:e406–e420.
26. Doronina O, Jones D, Martello M, Biron A, Lavoie-Tremblay M. A systematic review on the effectiveness of interventions to improve hand hygiene compliance of nurses in the hospital setting. *J Nurs Scholarsh* 2017;49:143–152.
27. Gould DJ, Moralejo D, Drey N, Chudleigh JH, Taljaard M. Interventions to improve hand hygiene compliance in patient care. *Cochrane Database Syst Rev* 2017;9:CD005186.
28. Huis A, van Achterberg T, de Bruin M, Grol R, Schoonhoven L, Hulscher M. A systematic review of hand hygiene improvement strategies: a behavioural approach. *Implement Sci* 2012;7:92.
29. Kingston L, O'Connell NH, Dunne CP, O'Connell NH. Hand hygiene-related clinical trials reported since 2010: a systematic review. *J Hosp Infect* 2016;92:309–320.
30. Luangsanatip N, Hongsuwan M, Limmathurotsakul D, *et al.* Comparative efficacy of interventions to promote hand hygiene in hospital: systematic review and network meta-analysis. *BMJ* 2015;351:h3728.
31. Mitchell MD, Betesh J, Ravitz N, Runyan D, Umscheid CA. *Automated Hand Hygiene Monitoring Systems*. Philadelphia, PA: Center for Evidence-based Practice, 2014.
32. Neo JRJ, Sagha-Zadeh R, Vilemeyer O, Franklin E. Evidence-based practices to increase hand hygiene compliance in health care facilities: an integrated review. *Am J Infect Control* 2016;44:691–704.
33. Ofek Shlomai N, Rao S, Patole S. Efficacy of interventions to improve hand hygiene compliance in neonatal units: a systematic review and meta-analysis. *Eur J Clin Microbiol Infect Dis* 2015;34:887–897.
34. Picheansathian W. A systematic review on the effectiveness of alcohol-based solutions for hand hygiene. *Int J Nurs Pract* 2004;10:3–9.
35. Ranji SR, Shetty K, Posley KA, *et al.* *Prevention of Healthcare-Associated Infections*. Vol 6 of Shojania KG, McDonald KM, Wachter RM, and Owens DK (editors). *Closing the Quality Gap: A Critical Analysis of Quality Improvement Strategies*. Technical Review 9 (Prepared by the Stanford University-UCSF Evidence-based Practice Center under Contract No. 290-02-0017). AHRQ Publication No. 04(07)-0051-6. Rockville, MD: Agency for Healthcare Research and Quality, 2007. <https://www.ahrq.gov/downloads/pub/evidence/pdf/qualgap6/hainfgap.pdf>. Published 2007. Accessed January 11, 2018.
36. Ritchie K, Iqbal K, Macpherson K, Riches E, Stout A. The provision of alcohol based products to improve compliance with hand hygiene. Healthcare Improvement Scotland website. http://www.healthcareimprovementscotland.org/previous_resources/hta_report/hta_7.aspx. Published 2005. Accessed January 11, 2018.
37. Stout A, Ritchie K, Macpherson K. Clinical effectiveness of alcohol-based products in increasing hand hygiene compliance and reducing infection rates: a systematic review. *J Hosp Infect* 2007;66:308–312.
38. Schweizer ML, Reisinger HS, Ohl M, *et al.* Searching for an optimal hand hygiene bundle: A meta-analysis. *Clin Infect Dis* 2014;58:248–259.
39. Srigley JA, Corace K, Hargadon DP, *et al.* Applying psychological frameworks of behaviour change to improve healthcare worker hand hygiene: a systematic review. *J Hosp Infect* 2015;91:202–210.
40. Srigley JA, Gardam M, Fernie G, Lightfoot D, Lebovic G, Muller MP. Hand hygiene monitoring technology: a systematic review of efficacy. *J Hosp Infect* 2015;89:51–60.
41. Stiller A, Salm F, Bischoff P, Gastmeier P. Relationship between hospital design and healthcare-associated infection rates: a systematic review and meta-analysis. *Antimicrob Resist Infect Control* 2016;5:51.
42. Vindigni SM, Riley PL, Jhung M. Systematic review: handwashing behaviour in low- to middle-income countries: outcome measures and behaviour maintenance. *Trop Med Int Health* 2011;16:466–477.
43. Ward MA, Schweizer ML, Polgreen PM, Gupta K, Reisinger HS, Perencevich EN. Automated and electronically assisted hand hygiene monitoring systems: a systematic review. *Am J Infect Control* 2014;42:472–478.
44. Cheung A, Weir W, Mayhew A, Kozloff N, Brown K, Grimshaw J. Overview of systematic reviews of the effectiveness of reminders in improving healthcare professional behavior. *Syst Rev* 2012;1:36.
45. Lorimer K, Kidd L, Lawrence M, McPherson K, Cayless S, Cornish F. Systematic review of reviews of behavioural HIV prevention interventions among men who have sex with men. *AIDS Care* 2012;25:133–150.
46. Cochrane Effective Practice and Organisation Care. What study designs should be included in an EPOC review and what should they be called? EPOC Resources for review authors website. http://epoc.cochrane.org/sites/epoc.cochrane.org/files/public/uploads/Resources-for-authors2017/what_study_designs_should_be_included_in_an_epoc_review.pdf. Published 2017. Accessed January 11, 2018.
47. Chudleigh J, Drey N, Moralejo D, Gould DJ. Systematic reviews of hand hygiene in patient care post 2010. *J Hosp Infect* 2016;94:110–111.
48. Cochrane Effective Practice and Organisation Care. Suggested risk of bias criteria for EPOC reviews. EPOC Resources for review authors website. http://epoc.cochrane.org/sites/epoc.cochrane.org/files/public/uploads/Resources-for-authors2017/suggested_risk_of_bias_criteria_for_epoc_reviews.pdf. Published 2107. Accessed January 11, 2018.
49. Zingg W, Castro-Sanchez E, Secci FV, *et al.* Innovative tools for quality assessment: integrated quality criteria for review of multiple study designs (ICROMS). *Public Health* 2016;133:19–37.
50. Wells GA, Shea B, O'Connell D, *et al.* The Newcastle-Ottawa Scale (NOS) for assessing the quality of nonrandomised studies in meta-analyses. University of Ottawa website. http://www.ohri.ca/programs/clinical_epidemiology/oxford.asp Published 2000. Accessed January 11, 2018.
51. Jadad AR, Moore RA, Carroll D, *et al.* Assessing the quality of reports of randomized clinical trials: is blinding necessary? *Control Clin Trials* 1996;17:1–12.
52. Rosella L, Bowman C, Pach B, Morgan S, Fitzpatrick T, Goel V. The development and validation of a meta-tool for quality appraisal of public health evidence: meta Quality Appraisal Tool (MetaQAT). *Public Health* 2016;136:57–65.
53. Anderson LA, Sharpe PA. Improving patient and provider communication: A synthesis and review of communication interventions. *Patient Educ Couns* 1991;17:99–134.
54. Downs SH, Black N. The feasibility of creating a checklist for the assessment of the methodological quality both of randomised and non-randomised studies of health care interventions. *J Epidemiol Community Health* 1998;52:377–384.
55. Higgins JPT, Altman DG, Sterne JAC. Chapter 8: Assessing risk of bias in included studies. In: Higgins JPT, Green S, editors. *Cochrane Handbook for Systematic Reviews of Interventions*, version 5.1.0 (updated March 2011). The Cochrane Collaboration website. <http://handbook.cochrane.org/>. Published 2011. Accessed January 11, 2018.
56. Systematic reviews: CRD's guidance for undertaking reviews in healthcare. Centre for Reviews and Dissemination website. <http://www.york.ac.uk/crd/guidance/>. Published 2009. Accessed January 11, 2018.
57. Higgins JPT, Green S. Cochrane handbook for systematic reviews of interventions Version 5.1.0 (updated March 2011). The Cochrane Collaboration website. <http://handbook.cochrane.org/>. Published 2011. Accessed January 11, 2018.
58. SIGN 50: A guideline developer's handbook. Edinburgh: Scottish Intercollegiate Guidelines Network website. <http://www.sign.ac.uk/sign-50.html>. Published 2015. Accessed January 11, 2018.

59. Michie S, West R. Behaviour change theory and evidence: a presentation to Government. *Health Psychol Rev* 2013;7:1–22.
60. NICE guidelines PH49 Behaviour change: Individual approaches (Recommendation 10 Ensure behaviour change is maintained for at least a year). National Institute for Health and Care Excellence website. <https://www.nice.org.uk/guidance/ph49/chapter/1-recommendations#recommendation-10-ensure-behaviour-change-is-maintained-for-at-least-a-year>. Published 2014. Accessed January 11, 2018.