

Angiostrongylus cantonensis: a review of its distribution, molecular biology and clinical significance as a human pathogen

JOEL BARRATT^{1,2*}†, DOUGLAS CHAN^{1,2,3†}, INDY SANDARADURA^{3,4}, RICHARD MALIK⁵, DEREK SPIELMAN⁶, ROGAN LEE⁷, DEBORAH MARRIOTT³, JOHN HARKNESS³, JOHN ELLIS² and DAMIEN STARK³

¹ *i3 Institute, University of Technology Sydney, Ultimo, NSW, Australia*

² *School of Life Sciences, University of Technology Sydney, Ultimo, NSW, Australia*

³ *Department of Microbiology, SydPath, St. Vincent's Hospital, Victoria St., Darlinghurst, NSW, Australia*

⁴ *School of Medicine, University of Notre Dame, Sydney, Australia*

⁵ *Centre for Veterinary Education, B22, The University of Sydney, NSW 2006, Australia*

⁶ *Faculty of Veterinary Science, The University of Sydney, Sydney, NSW 2006, Australia*

⁷ *ICPMR, Westmead Clinical School, Westmead Hospital, Westmead, NSW 2065, Australia*

(Received 6 August 2015; revised 15 March 2016; accepted 17 March 2016; first published online 26 May 2016)

SUMMARY

Angiostrongylus cantonensis is a metastrongyloid nematode found widely in the Asia-Pacific region, and the aetiological agent of angiostrongyliasis; a disease characterized by eosinophilic meningitis. *Rattus* rats are definitive hosts of *A. cantonensis*, while intermediate hosts include terrestrial and aquatic molluscs. Humans are dead-end hosts that usually become infected upon ingestion of infected molluscs. A presumptive diagnosis is often made based on clinical features, a history of mollusc consumption, eosinophilic pleocytosis in cerebral spinal fluid, and advanced imaging such as computed tomography. Serological tests are available for angiostrongyliasis, though many tests are still under development. While there is no treatment consensus, therapy often includes a combination of anthelmintics and corticosteroids. Angiostrongyliasis is relatively rare, but is often associated with morbidity and sometimes mortality. Recent reports suggest the parasites' range is increasing, leading to fatalities in regions previously considered *Angiostrongylus*-free, and sometimes, delayed diagnosis in newly invaded regions. Increased awareness of angiostrongyliasis would facilitate rapid diagnosis and improved clinical outcomes. This paper summarizes knowledge on the parasites' life cycle, clinical aspects and epidemiology. The molecular biology of *Angiostrongylus* spp. is also discussed. Attention is paid to the significance of angiostrongyliasis in Australia, given the recent severe cases reported from the Sydney region.

Key words: Angiostrongyliasis, *Angiostrongylus*, Diagnosis, Eosinophilic meningitis, Molluscs, Rat, Lungworm, Clinical significance, Epidemiology, Molecular biology.

INTRODUCTION

Angiostrongylus cantonensis (common name: the rat lungworm) is a metastrongyloid nematode originally identified in the Guangzhou region of China in the brown rat, *Rattus norvegicus* (Chen, 1935). It was first assigned to the genus *Pulmonema* which was later synonymized with *Angiostrongylus* (Dougherty, 1946) resulting in the current combination, *A. cantonensis*. It was identified as a human pathogen in 1945 (Beaver and Rosen, 1964) and is now recognized as the leading cause of eosinophilic meningitis worldwide (Wang *et al.* 2008; Graeff-Teixeira *et al.* 2009; Murphy and Johnson, 2013). It was the aetiological agent in over 2877 human cases of eosinophilic meningitis and is gaining recognition as an emerging zoonosis (Wang *et al.* 2012). Dogs and certain wildlife species are

important biosentinels for angiostrongyliasis, highlighting the potential risk for humans living in regions where these species are affected (Ma *et al.* 2013).

While *A. cantonensis* is the leading cause of eosinophilic meningitis, several other aetiological agents must be considered (Graeff-Teixeira *et al.* 2009). Serological tests for angiostrongyliasis are commercially available, though have not been widely adopted for routine use. Consequently, diagnosis is often intuitive, relying on an accurate patient history and non-specific tests such as computed tomography (CT) or magnetic resonance imaging (MRI). A patient's travel history or history of mollusc consumption greatly assist the diagnostic endeavour. Microscopic examination of cerebrospinal fluid (CSF) can be helpful, though the low sensitivity of this technique limits its use (Eamsobhana and Yong, 2009). Given the potential lethality of angiostrongyliasis, it must be given due consideration in all cases of eosinophilic meningitis.

Traditionally, *A. cantonensis* is endemic to the temperate and tropical parts of the Far East (York *et al.* 2014), though its current range includes

* Corresponding author: University of Technology Sydney, School of Life Sciences/iThree Institute, PO Box 123 Broadway, NSW 2007, Australia. E-mail: joel.barratt-1@uts.edu.au

† These authors contributed equally to this manuscript.

Southeast Asia, the Pacific Islands, parts of South and Central America and the Caribbean (Wang *et al.* 2008). Relatively new epidemiological data have emerged from Australia and the USA, suggesting that the parasite's geographical range is expanding (Teem *et al.* 2013; Chan *et al.* 2015; Iwanowicz *et al.* 2015). Australian cases of angiostrongyliasis have been reported since the 1970s, possibly as early as 1959 (Prociv and Carlisle, 2001), though interest in this parasite was renewed due to recent reports of lethal cases on Australia's eastern coast.

A review summarizing current knowledge on various aspects of *A. cantonensis* is provided. This includes its life cycle and transmission, the global epidemiology of angiostrongyliasis, clinical manifestations, current diagnostic approaches, therapy, and measures for control and prevention. The molecular biology of *A. cantonensis* is also discussed. Particular attention is paid to the significance of angiostrongyliasis in Australia given the increasing reports of angiostrongyliasis in Australian humans and wildlife (Blair *et al.* 2013; Ma *et al.* 2013; Morton *et al.* 2013; Spratt, 2015). An increased awareness of this emerging zoonosis is paramount to improving the prognosis for affected patients and reducing human, and companion animal suffering as the range of the parasite increases.

CLASSIFICATION

Angiostrongylus cantonensis is placed in the family Angiostrongylidae in the superfamily Metastrongyloidea which includes over 180 species across 45 genera. *Angiostrongylus cantonensis* was originally described as *Pulmonema cantonensis* by Chen (1935) and was subsequently placed in the genus *Angiostrongylus* by Dougherty (1946). Later, Uebelaker (1986) split *Angiostrongylus* into five genera: *Angiostrongylus* (in carnivores), *Parastrongylus* (murids), *Angiocaulus* (mustelids), *Gallegostrongylus* (gerbils and one murid) and *Sterfanskostrongylus* (insectivores), though this classification, placing *A. cantonensis* in *Parastrongylus*, is now rarely used (Cowie, 2013a).

Twenty-one species of *Angiostrongylus* are currently recognized, with the most significant being *A. costaricensis* and *A. cantonensis*, given their role as zoonotic pathogens (Spratt, 2015). Other significant species include *Angiostrongylus malaysiensis*, *Angiostrongylus vasorum* and *Angiostrongylus mackerrasae*. These species are considered animal pathogens but their zoonotic potential is yet to be demonstrated (Spratt, 2015).

LIFE CYCLE AND TRANSMISSION

Elucidation of the *A. cantonensis* life cycle was initially accredited to Mackerras and Sandars (1955) who studied what they believed to be *A. cantonensis*

in *R. norvegicus* captured on river banks in Queensland, Australia. These parasites were later identified as *A. mackerrasae*, although the life cycles of *A. cantonensis*, *A. mackerrasae* and *A. malaysiensis* are extremely similar, differing mostly in their host preferences (Bhaibulaya, 1975; Spratt, 2015). *Angiostrongylus costaricensis* also has a similar life cycle, though possesses a tropism for the mesenteric arteries of its definitive host, where egg production takes place (Mota and Lenzi, 2005). In contrast, *A. cantonensis* produces eggs in the pulmonary arteries of its definitive hosts (Thiengo *et al.* 2013; Spratt, 2015). The life cycle of *A. cantonensis* (Fig. 1) involves one of many potential intermediate hosts (molluscs) and definitive hosts (various rat species), and a large number of potential paratenic hosts.

Rattus rattus (the black rat) and *R. norvegicus* (the brown rat) are the favoured definitive hosts of *A. cantonensis*, though at least 17 rodent species may behave as definitive hosts, capable of passing first-stage (L₁) larvae (Fig. 2) in their feces (Yong and Eamsobhana, 2013). Simoes *et al.* (2014) postulated that the movement of *A. cantonensis* to new regions is mediated mostly by male rats that often have greater dispersal ability than females, while females are important for maintenance of *A. cantonensis* on a small, local scale. Rats become infected by ingestion of an intermediate or a paratenic host, containing third-stage (L₃) *A. cantonensis* larvae (Fig. 2). A few hours after ingestion, L₃ larvae penetrate the rats' intestinal wall and enter its blood stream where they are dispersed via the circulation. Larvae that reach the brain undergo an additional moult to become fourth-stage (L₄) larvae. A fifth moult ensues in the subarachnoid space, where larvae enter the young adult (L₅) stage. Young adult worms leave the central nervous system (CNS) and migrate through the circulation, and at 25 dpi, are found in the pulmonary arteries. At 35 dpi, adults have reached sexual maturity and females begin producing eggs that hatch in the terminal branches of the pulmonary arteries, liberating L₁ larvae. These larvae penetrate the alveolae, migrate to the pharynx and are swallowed by the rat. The L₁ larvae then travel through the digestive tract and appear in the rats' feces approximately 42 days after the initial exposure (Thiengo *et al.* 2013).

Definitive host competency may vary between rat species. For example, the Australian indigenous rat, *Rattus fuscipes*, seems a poor definitive host for *A. cantonensis* but is thought to be the preferred host of *A. mackerrasae* (Prociv and Carlisle, 2001). Similarly, the prevalence of *A. cantonensis* is usually much higher in *R. norvegicus* compared with *Rattus flavipectus* (Zhang *et al.* 2008a; Deng *et al.* 2012). This could be attributable to the biological competencies of different rat species, or simply, due to differences in the dietary habits of




Fig. 1. Life cycle of *A. cantonensis*. (A) Male (♂) and female (♀) adult *A. cantonensis* live in the pulmonary arteries of *Rattus* rats, their preferred definitive host. Females lay eggs that hatch in the terminal branches of the pulmonary arteries, liberating first-stage (L₁) larvae. The L₁ larvae penetrate the alveolae, migrate to the pharynx and are swallowed. (B) The L₁ larvae travel through the digestive tract and are passed in the rat feces. (C) A terrestrial or freshwater mollusc ingests the rat feces containing L₁ larvae, or L₁ larvae actively penetrate the mollusc tegument. The L₁ larvae undergo two moults in the mollusc to become third-stage (L₃) larvae. Infected molluscs are then ingested by a rat. The L₃ larvae penetrate the rats' intestine and migrate via the circulation to the brain where they undergo two additional moults to become young adult (L₅) worms. The L₅ worms leave the CNS and travel through the circulation to the pulmonary arteries where they mature to adulthood and reproduce. (D) Paratenic hosts eat molluscs infected with L₃ larvae, and the larvae become quiescent in these hosts. Infected paratenic hosts remain infectious to accidental hosts such as humans. (E) Fresh produce contaminated with mollusc slime may also represent a source of human infection, though direct ingestion of raw or undercooked molluscs (F) is the most common route of human infection. (G) Once ingested, L₃ larvae penetrate the intestinal wall and travel through the blood stream to the central nervous system. (H) The larvae enter the brain and in accidental hosts such as humans, eventually die. A granulomatous inflammatory reaction in the CNS is caused in response to dead worms, which manifests as eosinophilic meningitis.

different rat species, though this is yet to be ascertained (Zhang *et al.* 2008a). Non-*Rattus* rats such as the bandicoot rat (*Bandicota indica*) and the white-toothed rat (*Berylmys berdmorei*), are also definitive hosts of *A. cantonensis* (Pipitgool *et al.* 1997; Deng *et al.* 2012; Yong and Eamsobhana, 2013). The finding of adult *A. cantonensis* in the Ryukyu Islands tree rat (*Diplothrix legata*) (Okano *et al.* 2014) also implicates this species as a definitive host, though this requires further investigation. Experimental infections in Mongolian gerbils (*Meriones unguiculatus*) confirmed they are highly susceptible to angiostrongyliasis yet behave as poor definitive hosts; worms developed to sexual maturity in gerbils though very few L₁ larvae were shed in their feces (Wei *et al.* 2014).

Molluscs become infected by ingesting rat feces or by active penetration of L₁ larvae through their tegument (Morassutti *et al.* 2014). First-stage larvae remain viable for several days after being excreted, though viability drops sharply after this (Yousif and Lammler, 1975a). The efficiency of




Fig. 2. Microscopic observation of L₁ (A, B) and L₃ (C, D) *A. cantonensis* larvae from rat feces and mollusc tissue, respectively. Each L₁ larvae is approximately 320 μm in length, while L₃ larvae are slightly larger, at approximately 400 μm in length. The L₁ larvae were imaged under bright-field microscopy, while the L₃ larvae were imaged under phase-contrast microscopy.

mollusc infection is temperature dependent; it is greater at 26 °C compared with 24 °C (Yousif and Lammler, 1975a). Within the mollusc, L₁ larvae undergo two moults to become L₃ larvae; a roughly 20-day process marked by distinct morphological changes (Lv *et al.* 2009c; Thiengo *et al.* 2013; Zeng *et al.* 2013b). Larvae often migrate to the muscular layer of the foot given its excellent vascular supply, which provides favourable conditions for metastrongyloid larvae development (Mendonca *et al.* 1999; Giannelli *et al.* 2015). Larvae also migrate to the lung tissues (Yousif *et al.* 1980; Lv *et al.* 2009c). Consequently, mollusc lungs are often examined microscopically in field studies, for the characteristic nodules that contain L₃ larvae (Lv *et al.* 2009c; Hu *et al.* 2011; Qvarnstrom *et al.* 2013). In the aquatic snail *Pila polita*, ingested larvae migrate from the digestive gland to the intestine and then to the mantle, where the highest parasite burdens were recorded (Tesana *et al.* 2008). In *Parmarion martensi* semi-slugs (semi-slugs are molluscs with a shell that is too small to retract into), the midsection and tail had the highest larval densities; few larvae were found in the head (Jarvi *et al.* 2012). *Angiostrongylus cantonensis* is also pathogenic to molluscs, inducing biochemical disturbances, inflammation and reduction in reproductive capacity (Yousif *et al.* 1980; Tunholi-Alves *et al.* 2011, 2012, 2014, 2015).

Intermediate hosts of *A. cantonensis* include species from as many as 51 mollusc families, though some species may harbour more L₃ larvae than others (Table 1) (Yousif and Lammler, 1975b; Kim *et al.* 2014). When the aquatic snails *Pomacea canaliculata* and *P. polita* were experimentally infected with *A. cantonensis*, higher larval burdens were observed in *P. polita* (Tesana *et al.* 2008). The giant African land snail, *Achatina fulica*, may harbour thousands of L₃ larvae, and is thought to be a major contributor to the spread of *A. cantonensis* globally (Alicata, 1965a), though this is controversial (Civeyrel and Simberloff, 1996; Cowie, 2013a). The effect of mollusc size on larval burden has also been considered, though correlations were only observed in some species (Yousif and Lammler, 1975a; Pipitgool *et al.* 1997; Ibrahim, 2007; Tesana *et al.* 2009). It is suggested that parasite burdens are more closely linked to the number of L₁ larvae that the molluscs have been exposed to (Yousif and Lammler, 1975a). The prevalence of *A. cantonensis* in mollusc populations also varies between locations (Table 2), probably due to environmental factors such as temperature, humidity, the distribution of molluscs and the behaviours of local rats (Kim *et al.* 2014). Water salinity is also a contributing factor, with higher prevalences observed in aquatic molluscs living in water with lower salinity (Ibrahim, 2007).

In China and parts of Southeast Asia, *P. canaliculata*, and in Thailand, *Pila* spp., are considered

important intermediate hosts of *A. cantonensis* (Lv *et al.* 2009b; Tesana *et al.* 2009; Eamsobhana, 2013; Yang *et al.* 2013a). The number of known intermediate host species is increasing, with recent reports describing infections in new species from mollusc families such as the Achatinellidae, Assimineidae, Oxychilidae, and the species *Theba pisana*, *Plutonia lamarckii*, *Zachrysis provisorica*, *Cornu aspersum* and *Cryptozona siamensis* (Table 2) (Kim *et al.* 2014; Vitta *et al.* 2016). In Australia, two terrestrial snail species, *C. aspersum* and *Bradybaena similis* (Fig. 3) were recently confirmed as natural intermediate hosts of *A. cantonensis* (Table 2). Third-stage larvae of *A. cantonensis* also infect a multitude of paratenic hosts including freshwater prawns & shrimp, land crabs, frogs, toads, monitor lizards and planarians (Cowie, 2013b; Qvarnstrom *et al.* 2013; Eamsobhana, 2014). The L₃ larvae remain infective to definitive and accidental hosts that eat infected paratenic hosts.

Humans are one of many accidental hosts that become infected via one of the three possible transmission pathways (Fig. 1D–F). In Australia, naturally acquired infections in other accidental hosts have involved dogs, horses, brushtail possums (Fig. 4), Bennett's wallabies, rufous bettongs, tamarins (in captivity), black and grey-headed flying foxes, tawny frogmouths (Fig. 4, online Supplementary materials), gang-gang cockatoos and yellow-tailed black cockatoos (McKenzie *et al.* 1978; Wright *et al.* 1991; Higgins *et al.* 1997; Carlisle *et al.* 1998; Barrett *et al.* 2002; Monks *et al.* 2005; Lunn *et al.* 2012; Reece *et al.* 2013; Aghazadeh *et al.* 2015a; Walker *et al.* 2015). It has been suggested that brush-tail possums and tawny frogmouths are susceptible dead-end hosts native to Australia that serve as bio-sentinels for the presence of *A. cantonensis* (Ma *et al.* 2013; Spratt, 2015). In the USA, infections in several dead-end hosts have also been confirmed, including a privately owned orangutan with a history of snail consumption (Emerson *et al.* 2013), and a captive gibbon from a zoo in Miami (Duffy *et al.* 2004). Burns *et al.* (2014) described a fatal case in a captive African pygmy falcon from San Diego Zoo, Southern California, USA. Kottwitz *et al.* (2014) reported cases of lethal angiostrongyliasis in captive Geoffroy's tamarins from a zoo in Alabama, USA.

Deliberate ingestion of infected raw or undercooked molluscs is the most common route of infection for humans in Asian countries, such as Thailand, where known intermediate hosts, such as *Pila* spp., are regularly eaten raw as part of the local diet (Eamsobhana, 2013). Human *A. cantonensis* infections have also been linked to several paratenic host species. Reports describing the ingestion of monitor lizards in Sri Lanka, Thailand & India, raw frogs in China and, freshwater shrimp, fish & crabs in the Pacific Islands, implicate ingestion of these disparate hosts as a route for acquiring

Table 1. Larval parasite loads (infection intensities) in naturally infected intermediate and paratenic hosts of *A. cantonensis*

Mollusc species	Location	Number of larvae identified per infected mollusc	Reference
Terrestrial snails			
<i>Achatina fulica</i>	Kamphaeng Phet Province, Thailand	Min = 5, Max = 15	Vitita <i>et al.</i> (2016)
	Brazil	Mean = 220.5–833.6, Max = 3108	Oliveira <i>et al.</i> (2015) ^a
	Hawaii Island, Hawaii, USA	Mean = 5, Median = 1, Max = 18	Qvarnstrom <i>et al.</i> (2013) ^b
	Oahu, Hawaii, USA	Mean = 7, Median = 3, Max = 25	Qvarnstrom <i>et al.</i> (2013) ^b
	Tahiti, French Polynesia	Min = 11, Mean = 18.2, Median = 18, Max = 28	Fontanilla and Wade (2012)
	Panyu region, Guangzhou City, China	Mean = 57	Chen <i>et al.</i> (2012a)
	Northeast Thailand	Min = 5, Mean = 849.8, Median = 140.5, Max = 3113	Tesana <i>et al.</i> (2009)
	Northeast Thailand	Mean \geq 6	Pipitgool <i>et al.</i> (1997) ^c
	Sydney, Australia	Max = 0.01–1	Chan <i>et al.</i> (2015) ^d
<i>Bradybaena similaris</i>	Brisbane, Australia	Max = 2	Aghazadeh <i>et al.</i> (2015b)
<i>Cornu aspersum</i>	Sydney, Australia	Max = 0.01 to 1	Chan <i>et al.</i> (2015) ^d
<i>Cryptosona siamensis</i>	Kamphaeng Phet Province, Thailand	Min = 1, Max = 4858	Vitita <i>et al.</i> (2016)
Aquatic snails			
<i>Biomphalaria alexandrina</i>	North Sinai, Egypt	Mean = 1.5–3	Ibrahim (2007) ^e
<i>Cleopatra cyclostomoides</i>	North Sinai, Egypt	Mean = 1.75–7	Ibrahim (2007) ^e
<i>Cleopatra bulimoides</i>	North Sinai, Egypt	Mean = 1–3.36	Ibrahim (2007) ^e
<i>Lanistes carinatus</i>	North Sinai, Egypt	Mean = 2–37.62	Ibrahim (2007) ^e
<i>Lymnaea natalensis</i>	North Sinai, Egypt	Mean = 2	Ibrahim (2007)
<i>Melanooides tuberculata</i>	North Sinai, Egypt	Mean = 1.33–3	Ibrahim (2007) ^e
<i>Pila pesmei</i>	Northeast Thailand	Min = 1, Mean = 20.2, Median = 3, Max = 159	Tesana <i>et al.</i> (2009)
<i>Pomacea canaliculata</i>	Panyu region, Guangzhou City, China	Mean = 1.64	
	Southern Taiwan	Mean values: A = 60, B = 49, C = 65, D = 36, E = 53	Wang <i>et al.</i> (2011) ^f
	Guangdong Province, China	6000 in one snail, >1000 in two others	Li <i>et al.</i> (2008)
	Louisiana, USA	Mean = 19, Median = 2, Max = 71	Qvarnstrom <i>et al.</i> (2013) ^b
<i>Pomacea maculata</i>	Hawaii Island, Hawaii, USA	Mean = 445, Median = 110, Max \geq 1000	Qvarnstrom <i>et al.</i> (2013) ^b
Semi-slugs	Hawaii, USA	Mean = 0.10–60	Jarvi <i>et al.</i> (2012) ^g
<i>Parmarion martensi</i>	Hawaii Island, Hawaii, USA	Mean = 205, Median = 2, Max = 819	Qvarnstrom <i>et al.</i> (2013) ^b
Terrestrial slugs	Hawaii Island, Hawaii, USA	Mean \leq 1, Median \leq 1, Max \leq 1	Qvarnstrom <i>et al.</i> (2013) ^b
<i>Laevicaulis alte</i>	Hawaii Island, Hawaii, USA	Mean = 35, Median \leq 1, Max = 468	Qvarnstrom <i>et al.</i> (2013) ^b
<i>Pallifera</i> spp.	Hawaii Island, Hawaii, USA		
<i>Veronicella cubensis</i>	Hawaii Island, Hawaii, USA		
Other			
Other mollusc species	Hawaii Island, Hawaii, USA	Mean \leq 1, Median \leq 1, Max \leq 1	Qvarnstrom <i>et al.</i> (2013) ^b
	Northeast Thailand	Min = 1, Mean = 6.7, Median = 6, Max = 13	Tesana <i>et al.</i> (2009)
Planarians (paratenic hosts)	Hawaii Island, Hawaii, USA	Mean = 4, Median \leq 1, Max = 30	Qvarnstrom <i>et al.</i> (2013) ^b

^a Mean infection intensity varied between months.

^b Mean infection intensity is per 25 mg of tissue.

^c The mean infection intensity for L₃ larvae from all worm species was 13.6. Infection of Wistar rats with these larvae resulted in a recovery rate of 48.3% adult worms and 91.7% of these (~ 6) were *A. cantonensis*.

^d Mean infection intensity is per 50 mg of tissue.

^e Mean infection intensity varied between seasons and/or locations.

^f Snails from irrigation canals in five different locations (designated A–E).

^g Mean infection intensity is per 1 mg of tissue. Also, the mean infection intensity varied depending on the mollusc tissue sample screened (mollusc head, mid-section, tail or slime), and the location from which molluscs were collected.

Table 2. Prevalence of *A. cantonensis* in definitive, intermediate and paratenic hosts

Host species	Location	Prevalence	Reference	
Definitive hosts				
<i>Bandicota indica</i>	Guangdong Province, China	1/6 (16.7%)	Deng <i>et al.</i> (2012)	
	Northeast Thailand	1/69 (1.5%)	Pipitgool <i>et al.</i> (1997)	
<i>Rattus exulans</i>	Hawaii Island, Hawaii, USA	10/10 (100%)	Qvarnstrom <i>et al.</i> (2013) ^a	
<i>Rattus flavipectus</i>	Shenzhen, Guangdong Province, China	6/121 (4.9%)	Zhang <i>et al.</i> (2008a)	
	Guangdong Province, China	3/118 (2.5%)	Deng <i>et al.</i> (2012)	
<i>Rattus fuscipes</i>	Brisbane, Queensland, Australia	14/53 (26%)	Aghazadeh <i>et al.</i> (2015b)	
<i>Rattus lutreolus</i>	Brisbane, Queensland, Australia	2/5 (40%)	Aghazadeh <i>et al.</i> (2015b)	
<i>Rattus norvegicus</i>	Brisbane, Queensland, Australia	4/15 (27%)	Aghazadeh <i>et al.</i> (2015b)	
	New Orleans, Louisiana, USA	2/36 (5.6%)	York <i>et al.</i> (2015)	
	Trindade, São Gonçalo, Brazil	78/110 (70.9%)	Simoes <i>et al.</i> (2014)	
	Hawaii Island, Hawaii, USA	1/1 (100%)	Qvarnstrom <i>et al.</i> (2013)	
	Islands in the Ogasawara Archipelago, Japan	46 /57 (80.7%)	Tokiwa <i>et al.</i> (2013)	
	Guangdong Province, China	52/263 (19.8%)	Deng <i>et al.</i> (2012)	
	Guangzhou, Guangdong Province, China	20/220 (9.1%)	Yang <i>et al.</i> (2012)	
	Hainan, China	13/118 (11.0%)	Hu <i>et al.</i> (2011)	
	Guangdong Province, China	310/1835 (16.9%)	Pan <i>et al.</i> (2011)	
	São Gonçalo, Rio de Janeiro, Brazil	19/27 (70.4%)	Simoes <i>et al.</i> (2011)	
	Grenada, West Indies	45/192 (23.4%)	Chikweto <i>et al.</i> (2009)	
	Shenzhen, Guangdong Province, China	31/187 (16.6%)	Zhang <i>et al.</i> (2008a)	
	Jamaica	20/78 (25.6%)	Lindo <i>et al.</i> (2002)	
	Northeast Thailand	2/52 (3.8%)	Pipitgool <i>et al.</i> (1997)	
	Northeast Puerto Rico	19/40 (47.5%)	Andersen <i>et al.</i> (1986)	
	<i>Rattus rattus</i>	Brisbane, Queensland, Australia	52/325 (16%)	Aghazadeh <i>et al.</i> (2015b)
		Hawaii Island, Hawaii, USA	26/26 (100%)	Qvarnstrom <i>et al.</i> (2013) ^a
Tenerife, Canary Islands		30/54 (55.6%)	Martin-Alonso <i>et al.</i> (2011) ^b	
Phitsanulok Province, Thailand		1/62 (1.6%)	Vitta <i>et al.</i> (2011)	
Tenerife, Canary Islands		10/67 (15%)	Foronda <i>et al.</i> (2010)	
Jamaica		4/31 (12.9%)	Lindo <i>et al.</i> (2002)	
Northeast Puerto Rico		10/63 (15.9%)	Andersen <i>et al.</i> (1986)	
<i>Sigmodon hispidus</i> ^c	McCurain County, Oklahoma, USA	1/30 (3.3%)	York <i>et al.</i> (2015)	
Intermediate hosts				
<i>Achatina fulica</i>	Kamphaeng Phet Province, Thailand	3/275 (1.1%)	Vitta <i>et al.</i> (2016)	
	Miami, Florida, USA	18/50 (36%)	Iwanowicz <i>et al.</i> (2015)	
	São Gonçalo, Rio de Janeiro, Brazil	78.7% of 153	Oliveira <i>et al.</i> (2015)	
	China	21.5% of 21 138	Song <i>et al.</i> (2016) ^d	
	Hawaii Island, Hawaii, USA	7/8 (87.5%)	Qvarnstrom <i>et al.</i> (2013)	
	Oahu, Hawaii, USA	5/9 (55.6%)	Qvarnstrom <i>et al.</i> (2013)	
	Miami, Florida	4/140 (2.9%)	Teem <i>et al.</i> (2013)	
	Panyu region, Guangzhou, China	83/367 (22.6%)	Chen <i>et al.</i> (2012a)	
	Guangdong Province, China	223/1354 (16.5%)	Deng <i>et al.</i> (2012)	
	Qushui Village, Suixi County, Zhanjiang City	13%	Shen <i>et al.</i> (2012)	
	Guangzhou, China	111/795 (14%)	Yang <i>et al.</i> (2012)	
	Hainan, China	121/534 (22.7%)	Hu <i>et al.</i> (2011)	
	Hawaii, USA	5/6 (83.3%)	Qvarnstrom <i>et al.</i> (2010) ^e	
	Pernambuco, Northeast Brazil	14/33 (42.4%)	Thiengo <i>et al.</i> (2010)	
	China ^f	13.4% of 3549	Lv <i>et al.</i> (2009b)	
	Northeast Thailand	4/53 (7.6%)	Tesana <i>et al.</i> (2009)	
	Shenzhen, Guangdong Province, China	31/302 (10.3%)	Zhang <i>et al.</i> (2008a)	
	Jiangmen, Guangdong Province, China	45% of 695	Zhang <i>et al.</i> (2008b)	
	Santa Catarina Island, Florianopolis, Brazil	1/244 (0.4%)	Neuhauss <i>et al.</i> (2007)	
	<i>Achatina</i> spp.	Phitsanulok Province, Thailand	38/307 (12.4%)	Vitta <i>et al.</i> (2011)
		Southern Florida, USA	1/19 (5.3%)	Stockdale-Walden <i>et al.</i> (2015) ^g
	<i>Alcadia striata</i>	Suixi County, Zhanjiang, China	10%	Shen <i>et al.</i> (2012)
	<i>Ampularum crossean</i>	Fuzhou and Xiamen, Fujian Province, China	31/19 843 (0.2%)	Li <i>et al.</i> (2013a)
	<i>Bellamyia aeruginosa</i>	Guangdong Province, China	1/70 (1.4%)	Deng <i>et al.</i> (2012)
	<i>Bellamyia</i> spp.	Guangdong Province, China	3/91 (3.3%)	Deng <i>et al.</i> (2012)
	<i>Biomphalaria alexandrina</i>	North Sinai, Egypt	3/294 (1%)	Ibrahim (2007)
		Sydney, Australia	1/10 (10%)	Chan <i>et al.</i> (2015)
<i>Bradybaena similaris</i>	São Gonçalo, Rio de Janeiro, Brazil	24.6% of 245	Oliveira <i>et al.</i> (2015)	
	Southern Florida, USA	3/7 (42.9%)	Stockdale-Walden <i>et al.</i> (2015) ^g	
<i>Cepaea</i> spp.	Hainan, China	9/43 (20.9%)	Hu <i>et al.</i> (2011)	
<i>Cleopatra bulimoides</i>	North Sinai, Egypt	41/3588 (1.1%)	Ibrahim (2007)	
<i>Cleopatra cyclostomoides</i>	North Sinai, Egypt	45/3401 (1.3%)	Ibrahim (2007)	
<i>Cornu aspersum</i>	Brisbane, Queensland, Australia	1/87 (1.2%)	Aghazadeh <i>et al.</i> (2015b) ^h	
	Sydney, Australia	14/312 (4.5%)	Chan <i>et al.</i> (2015)	
<i>Cryptozonia siamensis</i>	Tenerife, Canary Islands, Spain	9/18 (50%)	Martin-Alonso <i>et al.</i> (2015) ⁱ	
	Kamphaeng Phet Province, Thailand	45/425 (10.6%)	Vitta <i>et al.</i> (2016)	
<i>Hemiplecta distincta</i>	Northeast Thailand	2/10 (20%)	Tesana <i>et al.</i> (2009)	

Table 2. (Cont.)

Host species	Location	Prevalence	Reference
<i>Laevicaulis alte</i>	Hawaii Island, Hawaii, USA	4/5 (80%)	Qvarnstrom <i>et al.</i> (2013)
	Hawaii, USA	4/5 (80%)	Qvarnstrom <i>et al.</i> (2010) ^e
<i>Lanistes carinatus</i>	North Sinai, Egypt	56/2490 (2.2%)	Ibrahim (2007)
<i>Lymnaea natalensis</i>	North Sinai, Egypt	2/218 (0.9%)	Ibrahim (2007)
<i>Melanoides tuberculata</i>	North Sinai, Egypt	26/4112 (0.6%)	Ibrahim (2007)
<i>Pallifera</i> spp.	Hawaii Island, Hawaii, USA	2/5 (40%)	Qvarnstrom <i>et al.</i> (2013)
<i>Parmarion martensi</i>	Hawaii Island, Hawaii, USA	77/97 (79.4%)	Qvarnstrom <i>et al.</i> (2013)
	Hawaii, USA	83/112 (74.1%)	Qvarnstrom <i>et al.</i> (2010) ^e
	Hawaii, USA	25/34 (73.5%)	Qvarnstrom <i>et al.</i> (2007)
<i>Phlegm bilineatus</i>	Hainan, China	146/517 (28.2%)	Hu <i>et al.</i> (2011)
<i>Pila pesmei</i>	Northeast Thailand	11/454 (2.4%)	Tesana <i>et al.</i> (2009)
<i>Pila polita</i>	Northeast Thailand	9/181 (5%)	Tesana <i>et al.</i> (2009)
	Northeast Thailand	4/423 (0.9%)	Pipitgool <i>et al.</i> (1997)
<i>Plutonia lamarckii</i>	Tenerife, Canary Islands, Spain	43/60 (71.7%)	Martin-Alonso <i>et al.</i> (2015) ⁱ
<i>Pomacea canaliculata</i>	China	7.6% of 41 299	Song <i>et al.</i> (2016) ^d
	Fuzhou and Xiamen, Fujian Province, China	753/5744 (13.8%)	Li <i>et al.</i> (2013a)
	Panyu region, Guangzhou City, China	3.08% of 357	Chen <i>et al.</i> (2012a)
	Guangdong Province, China	172/2929 (5.87%)	Deng <i>et al.</i> (2012)
	Guangzhou, China	11/734 (1.5%)	Yang <i>et al.</i> (2012)
	Hainan, China	64/518 (12.4%)	Hu <i>et al.</i> (2011)
	Southern Taiwan	109/535 (20.4%)	Wang <i>et al.</i> (2011)
	China ^f	6.8% of 11 709	Lv <i>et al.</i> (2009b)
	Shenzhen, Guangdong Province, China	65/314 (20.7%)	Zhang <i>et al.</i> (2008a)
	Jiangmen, Guangdong Province, China	1.8% of 720	Zhang <i>et al.</i> (2008b)
	Louisiana, USA	5/31 (16.1%)	Qvarnstrom <i>et al.</i> (2010) ^e
<i>Pomacea lineata</i>	Pernambuco, Northeastern Brazil	5/95 (5.3%)	Thiengo <i>et al.</i> (2010)
<i>Pomacea maculata</i>	New Orleans, Louisiana, USA	5/31 (16.1%)	Qvarnstrom <i>et al.</i> (2013)
	Gretna, Louisiana, USA	8/60 (13.3%)	Teem <i>et al.</i> (2013)
	Mandeville, Louisiana, USA	1/40 (2.5%)	Teem <i>et al.</i> (2013)
<i>Theba pisana</i>	Tenerife, Canary Islands, Spain	7/20 (35%)	Martin-Alonso <i>et al.</i> (2015) ⁱ
<i>Thelidomus asper</i>	Mandeville, Jamaica	4/10 (40%)	Lindo <i>et al.</i> (2002)
<i>Veronicella cubensis</i>	Hawaii Island, Hawaii, USA	27/71 (38%)	Qvarnstrom <i>et al.</i> (2013)
	Hawaii, USA	23/50 (46%)	Qvarnstrom <i>et al.</i> (2010) ^e
<i>Zachrysis provisorica</i>	Southern Florida, USA	7/34 (20.6%)	Stockdale-Walden <i>et al.</i> (2015) ^g
Other aquatic snails	China ^f	0.05% of 7287	Lv <i>et al.</i> (2009b)
	Northeast Thailand	4/841 (0.5%)	Tesana <i>et al.</i> (2009)
Other terrestrial snails	China ^f	0.3% of 1421	Lv <i>et al.</i> (2009b)
Other terrestrial slugs	China ^f	6.5% of 5370	Lv <i>et al.</i> (2009b)
Other molluscs	Hawaii Island, Hawaii, USA	6/14 (42.9%)	Qvarnstrom <i>et al.</i> (2013)
	Hawaii, USA	5/16 (31.3%)	Qvarnstrom <i>et al.</i> (2010) ^e
Paratenic hosts			
Planarians	Hawaii Island, Hawaii, USA	8/12 (66.7%)	Qvarnstrom <i>et al.</i> (2013)
	Hawaii, USA	2/2 (100%)	Qvarnstrom <i>et al.</i> (2010) ^e

^a This study compared the necropsy of rats with real-time PCR, for confirmation of an *A. cantonensis* infection. This table shows the higher prevalence value.

^b This is a study on the seroprevalence of *A. cantonensis* in rats and indicates exposure to worms rather than an active, current infection.

^c *Sigmodon hispidus* is an important definitive host of *A. costaricensis* (Graeff-Teixeira *et al.* 1990), though it has not yet been described as a definitive host of *A. cantonensis* (Yong and Eamsobhana, 2013).

^d A meta-analysis involving 38 studies from different regions in China, over a 10-year period.

^e This study compared conventional and real-time PCR for detection of *A. cantonensis* in mollusc tissues. This table shows the higher prevalence value.

^f This study involved 164 counties from 19 provinces, predominantly along southeastern coastline of China.

^g This study compared microscopic identification of worms from pepsin digested mollusc tissue to PCR, for detection of *A. cantonensis* in mollusc tissues. This table shows the higher prevalence value.

^h The prevalence shown for this study was generated from a group of molluscs containing mixed species, in which one *C. aspersum* was infected only.

ⁱ This study compared microscopic identification of worms from pepsin digested mollusc tissue to a loop-mediated isothermal amplification assay, for detection of *A. cantonensis*. This table shows the higher prevalence value.

Angiostrongylus infections (Hidelaratchi *et al.* 2005; Malvy *et al.* 2008; Tsai *et al.* 2011; Cowie, 2013b; Pai *et al.* 2013; Eamsobhana, 2014).

The shedding of *A. cantonensis* larvae in infected mollusc mucus onto fresh produce may represent

another route of transmission (Fig. 1E). This pathway is relevant to those that regularly eat uncooked, inadequately washed, plant material, i.e. in salads (Barrow *et al.* 1996; Lindo *et al.* 2002; Slom *et al.* 2002; Waugh *et al.* 2005; Yeung *et al.* 2013).



Fig. 3. Two terrestrial mollusc species that were identified as natural intermediate hosts of *A. cantonensis* in Australia. *Cornu aspersum* and *Bradybaena similaris* are introduced species in Australia, and are common inhabitants of gardens and parks in metropolitan areas along the eastern coast of the continent.



Fig. 4. Clinical presentation of angiostrongyliasis in some Australian wildlife species, including a brushtail possum (*Trichosurus vulpecula*) (A) and multiple tawny frogmouths (*Podargus strigoides*) (B–H). These animals were described as biosentinels for *A. cantonensis* in the Sydney region (Ma *et al.* 2013). Panel (A) shows a juvenile brushtail possum that had been in care for 6 weeks. Food supplied to its large cage was gradually being left uneaten, but food offered to the possum in its nest box was eaten ravenously. Closer examination showed an inability to ambulate. Clinical examination showed hind limb and tail paralysis. Pinching the hind paws elicited no pain response but a strong and even exaggerated withdrawal reflex, typical for spinal cord damage. Panels (B–H) show tawny frogmouths displaying signs typical for spinal cord damage, prior to a diagnosis of angiostrongyliasis. The birds present with varied clinical signs, from moribund in advanced cases (B), to reduced or normal mentation (C–H). Some birds may be alert and aware but unable to fly or stand (H).

An outbreak has also been linked to drinking raw vegetable juices (Tsai *et al.* 2004, 2013). Independent studies that included several mollusc species, reported *A. cantonensis* in mollusc mucus,

though the small number of larvae shed may be negligible or insufficient to represent a major source of human infection (Qvarnstrom *et al.* 2007; Jarvi *et al.* 2012; Chan *et al.* 2015). However, mollusc slime was




Fig. 5. Countries where *A. cantonensis* has been detected in naturally infected hosts. Shaded countries are those in which *A. cantonensis* was identified in studies screening naturally infected animals, or where humans have acquired infections. Unshaded countries include those that are yet to find evidence of *A. cantonensis*, or countries where studies examining potential hosts for *A. cantonensis* infection have not been carried out. This map does not include countries that have reported sporadic clinical cases of angiostrongyliasis that were probably acquired abroad. Mainland Australia is shaded on the map, though the island state of Tasmania is not. Native rats in Tasmania are known hosts of *A. mackerrasae* (Prociv *et al.* 2000), though *A. cantonensis* has not been reported in Tasmania. Mainland Spain is also not shaded on the map, although *A. cantonensis* was detected in the Canary Islands, off the western coast of Africa, which are part of Spain. Currently *A. cantonensis* is not known to be present in mainland Spain.

the predicted mode of transmission in two cases of severe *Angiostrongylus* eosinophilic meningitis (AEM) recently reported in young children (Morton *et al.* 2013). In these cases, a history of mollusc consumption was denied, though this is difficult to confirm in young children. Freshwater may represent another source of infection as *A. cantonensis* L₃ larvae reportedly survive in freshwater for up to 72 h (Cheng and Alicata, 1964). Studies of the feline metastrongyloid lungworms *Aelurostrongylus abstrusus* and *Troglostrongylus brevior* demonstrated that L₃ larvae were also shed in mollusc mucus, and could be found in sediments from tap water containing *C. aspersum* that had been experimentally infected and drowned (Giannelli *et al.* 2015).

EPIDEMIOLOGY AND GLOBAL DISTRIBUTION

Angiostrongylus cantonensis is the most common cause of eosinophilic meningitis. It has been reported in over 30 countries since its identification as a human pathogen in 1945, predominantly in the tropics and sub-tropics (Beaver and Rosen, 1964; Wang *et al.* 2012; Eamsobhana *et al.* 2013a; Eamsobhana, 2014) (Tables 1 and 2, Fig. 5). While more than 2877 human cases of angiostrongyliasis have been described, many have probably gone unreported because of a lack of awareness and difficulties associated with its diagnosis (Qvarnstrom *et al.* 2007). Reports of angiostrongyliasis in Europe and other regions where it was not known previously are becoming more frequent, with travel to parts of Asia, the Pacific Islands and Latin America noted as the likely route of exposure for affected individuals (Bartschi *et al.* 2004; Chancellor, 2007; Leone *et al.* 2007; Ali *et al.* 2008; Malvy *et al.*

2008; Luessi *et al.* 2009; Maretic *et al.* 2009; Lammers *et al.* 2015).

While it is traditionally considered a disease of the Far East, reports of locally acquired angiostrongyliasis are becoming increasingly common in sub-tropical and temperate regions (Gutteridge *et al.* 1972; Prociv *et al.* 2000; Senanayake *et al.* 2003; Blair *et al.* 2013; Morton *et al.* 2013). It may be that the range of *A. cantonensis* has expanded recently, possibly as a consequence of global warming or other environmental factors (York *et al.* 2014). Hochberg *et al.* (2007) attribute the spread of *A. cantonensis* to the sheer diversity of its intermediate hosts, and the efficient dispersion of ship-borne rats. A similar expansion in geographic range has not been reported for *A. costaricensis*; the causative agent of human abdominal angiostrongyliasis (Spratt, 2015). The first documented human *A. costaricensis* infection was from Costa Rica (Morera and Cespedes, 1971; Morera, 1973), and later reports confirmed its presence in Venezuela, Ecuador, Honduras, Mexico, Nicaragua, Brazil, Guatemala, Columbia, the Caribbean Islands and Southern USA (Morera *et al.* 1983; Incani *et al.* 2007; Palominos *et al.* 2008; Spratt, 2015). Apparently, only one case of human *A. costaricensis* infection has been reported outside the Americas; an isolated case in an African man (Baird *et al.* 1987). The relatively restricted range of *A. costaricensis* is probably related to the limited range of its preferred definitive host; the hispid cotton rat (*Sigmodon hispidus*) (Graeff-Teixeira *et al.* 1990), which is generally found only in southern North America and parts of Central and South America.

Most human cases of angiostrongyliasis (~1300 cases; 47% of all cases worldwide) have been reported in Thailand (Wang *et al.* 2008), where between 0·3

and 2 people per 100 000 become infected annually (Suankratay *et al.* 2001; Eamsobhana, 2013). This high prevalence is attributable to the dietary habits of the local populace, involving the regular consumption of dishes like 'koi-hoi' that contain raw or undercooked molluscs such as *Pomacea maculata*, *P. canaliculata* and *Pila* spp.; common intermediate hosts of *A. cantonensis* (Schmutzhard *et al.* 1988; Lv *et al.* 2009a; Eamsobhana *et al.* 2010; Odermatt *et al.* 2010; Cowie, 2013b; Eamsobhana, 2014; Kim *et al.* 2014). The consumption of undercooked monitor lizard livers was linked to a fatal case of AEM in Thailand (Eamsobhana, 2014), where very high rates of monitor lizard infection have been reported; as high as 96% in one population (Eamsobhana, 2013). Angiostrongyliasis has also been reported in the neighbouring Southeast Asian Nations of Laos (Harinasuta, 1983), Cambodia (Brumpt *et al.* 1968) and Vietnam (Chau *et al.* 2003).

While the majority of AEM cases are reported from Thailand, most epidemiological data on angiostrongyliasis, including intermediate and definitive host surveys, comes from China. Several outbreaks of angiostrongyliasis have occurred in China, predominantly in provinces along the eastern coast (Lin *et al.* 2005; Wang *et al.* 2007, 2010; Lv *et al.* 2008, 2009a; Zhang *et al.* 2008a; Zhou *et al.* 2009; Hu *et al.* 2011). Only three cases of AEM were reported in China between 1984 and 1996, and the recent increase in cases is attributable to widespread changes in human dietary patterns, where snails have become a popular food item (Chen *et al.* 2005; Zhou *et al.* 2009; Wang *et al.* 2010). *Achatina fulica* and *P. canaliculata* are considered the most important species for *A. cantonensis* transmission in China, where both are introduced species, farmed as a food source (Wang *et al.* 2007; Lv *et al.* 2008, 2009b). Multiple Chinese outbreaks were directly linked to the consumption of raw or undercooked *P. canaliculata*, which is more popular as a Chinese food item than *A. fulica* (Wang *et al.* 2007; Lv *et al.* 2009a, b; Zhou *et al.* 2009). Consequently, *P. canaliculata* is aquacultured intensively for human consumption; it is often sold in local markets and served regularly in restaurants (Zhang *et al.* 2008a; Lv *et al.* 2009a, b; Zhou *et al.* 2009; Wang *et al.* 2010; Li *et al.* 2013a). *Pomacea canaliculata* is also more widespread in China than the terrestrial *A. fulica*, possibly because of its efficient spread via waterways during flooding (Lv *et al.* 2009b).

Serological surveys suggest human exposure to *A. cantonensis* is common in China. On Hainan Island, 92 of 459 subjects (20.04%) had antibodies to *A. cantonensis* (Hu *et al.* 2011). In another study from Hainan Island, anti-*A. cantonensis* immunoglobulin G (IgG) was detected in 20.6% of 393 participants, 12.5% of whom habitually ate raw snails (Li *et al.* 2011). In Guangdong Province, 42 of 300 people

(14%) had IgG antibodies to *A. cantonensis*, five of whom had been recently exposed based on the detection of circulating IgM (Zhang *et al.* 2008b). The prevalence of *A. cantonensis* on Hainan Island was equal for both genders, though subjects under the age of 14 were more likely to be seropositive (Hu *et al.* 2011). Chen *et al.* (2011c) detected circulating *A. cantonensis* antigen at a prevalence of 0.8% in members of the general Chinese population though differences were observed between certain groups. Males had a higher prevalence than females and those involved in the aquaculture or processing of molluscs for human consumption were more likely to have circulating antigen (Chen *et al.* 2011c). The prevalence of *A. cantonensis* in molluscs and rats has also been investigated in several regions in China (Table 2).

Hundreds of AEM cases have been reported in Taiwan, many of which were linked to the handling or consumption of *P. canaliculata* and *A. fulica* (Hwang and Chen, 1991; Tsai *et al.* 2004, 2013; Wan and Weng, 2004; Wang *et al.* 2007). The aquatic snail, *Bellamya chinensis*, is also considered an important intermediate host of *A. cantonensis* in Taiwan (Lv *et al.* 2009b). Early Taiwanese reports of AEM described cases predominantly in children and indigenous Taiwanese (Yii *et al.* 1975; Hwang and Chen, 1991), though recent reports more commonly describe infections in adults (Tseng *et al.* 2011, 2013). In those earlier cases, infections peaked during the rainy season, and those affected often reported eating thoroughly cooked *A. fulica* prior to disease (Yii *et al.* 1975). This suggests that *A. cantonensis* may have been ingested inadvertently during preparation of the snails for cooking.

Cases of AEM in Hawaii have been reported from as early as 1958 (Wallace, 2013). Between 2001 and 2005, 24 cases were reported though only one case was parasitologically confirmed; a case involving a young child. Most of these cases occurred in Honolulu, on the main island of Hawaii (Hochberg *et al.* 2007, 2011). Deliberate consumption of raw or undercooked molluscs is rare in Hawaii and most exposures were attributed to inadvertent ingestion of molluscs or their mucus. Patients sometimes recalled finding slugs in their food or drink, and the mother of another patient recalled her infant child putting grass in her mouth (Hochberg *et al.* 2011). *Parmarion martensi* is considered important for maintaining the endemicity of *A. cantonensis* in Hawaii, given its high prevalence in Hawaiian populations, and the remarkably high parasite burdens observed in some specimens (Tables 1 and 2).

In Australia, virtually all native and introduced terrestrial and freshwater molluscs can be experimentally infected with *A. cantonensis* (Prociv and Carlisle, 2001). Despite this, the frequency of infected molluscs in Australia is low compared to

what is observed in Southeast Asia (Chan *et al.* 2015). It was suggested that the mollusc species in Australia may not be the preferred hosts of *A. cantonensis* (Chan *et al.* 2015). Furthermore, the climate in Sydney is temperate to sub-tropical, whereas transmission of *A. cantonensis* is favoured by tropical climates which experience warm temperatures, high humidity and high rainfall (Hu *et al.* 2011). The minimum temperature threshold (the temperature at which larvae halt development in their mollusc hosts), is approximately 15 °C for *A. cantonensis* (Lv *et al.* 2006; Morley, 2010). In Sydney, temperatures regularly fall below 15 °C in winter, probably restricting *A. cantonensis* transmission to the warmer months of spring and summer. In Eastern Australia the prevalence of *Angiostrongylus* spp. in *Rattus* spp. is high compared with its prevalence in molluscs (Table 2).

Human cases of AEM have been diagnosed in multiple cities along the eastern coast of Australia including Brisbane, Sydney and Melbourne (Prociv *et al.* 2000). The first Australian case of human AEM was reported in 1971 in the Brisbane region (Gutteridge *et al.* 1972), and several human cases have been reported in Australia since (Prociv and Carlisle, 2001; Senanayake *et al.* 2003; Blair *et al.* 2013; Morton *et al.* 2013). Australian clinical cases mostly involved young children or infants, with or without a known history of mollusc consumption (Prociv *et al.* 2000; Morton *et al.* 2013). Two cases involved young adults who knowingly ingested slugs from the Sydney area (Senanayake *et al.* 2003; Blair *et al.* 2013). The rate of human infections is proportionally lower in Australia compared with China and Thailand, probably because snails are not widely eaten in Australia. Given the rarity of human angiostrongyliasis in Australia and its mostly self-limited nature, some infections have probably gone unreported. Regardless, the recent cases of severe angiostrongyliasis and increasing range of *A. cantonensis* underpin the need for increased awareness of angiostrongyliasis in Australia, particularly given that the aetiological agent sometimes remains unidentified until post-mortem (Morton *et al.* 2013).

Angiostrongylus cantonensis has also been detected in Japan, parts of Southern USA, the Caribbean Islands [though it was absent in rats from Barbados (Levett *et al.* 2004)], Tenerife, Brazil, Papua New Guinea, French Polynesia, Fiji, the Philippines, Indonesia, Sri Lanka, India, South Africa and Egypt (Tables 1 and 2, Fig. 5) (Alicata, 1965b; Kliks and Palumbo, 1992; Uga *et al.* 1996; Asato *et al.* 2004; Batmanian and O'Neill, 2004; Lindo *et al.* 2004; Abo-Madyan *et al.* 2005; Owen, 2005; Waugh *et al.* 2005; Caldeira *et al.* 2007; Chikweto *et al.* 2009; Dorta-Contreras *et al.* 2009; Archer *et al.* 2011; Constantino-Santos *et al.* 2014; Oehler *et al.* 2014; Okano *et al.* 2014; Lammers *et al.* 2015; Stockdale-Walden *et al.* 2015).

PATHOPHYSIOLOGY AND CLINICAL MANIFESTATIONS

Angiostrongylus cantonensis larvae are neurotropic, preferentially infecting CNS tissue, and causing inflammation at these sites. Autopsies of AEM patients indicate that the external surfaces of the brain and the spinal cord are generally normal and gross haemorrhage is uncommon, though has been reported in severe human cases (Wang *et al.* 2008; Morton *et al.* 2013). Infiltration of lymphocytes, plasma cells and eosinophils is common in the meninges (Wang *et al.* 2008) (Fig. 6). Third-stage larvae may be observed in association with the meninges and nerve roots (Chen *et al.* 2005). Eosinophilic pleocytosis in the CSF and increased CSF protein are common (Schmutzhard *et al.* 1988; Dorta-Contreras *et al.* 2011). Cellular infiltration around live worms is unusual though dead worms precipitate granuloma formation, infiltration by eosinophils and occasionally Charcot–Leyden crystals (Wang *et al.* 2008; Martins *et al.* 2015). Physical tracks and microcavities due to the burrowing movement of larvae may be observed in the brain and spinal cord (Morton *et al.* 2013; Murphy and Johnson, 2013).

Angiostrongyliasis initiates a Th2 immune response in the CNS, characterized by the infiltration of eosinophils into the subarachnoid space, and CSF eosinophilia (Intapan *et al.* 2008; Graeff-Teixeira *et al.* 2009; Murphy and Johnson, 2013; Martins *et al.* 2015). The role of eosinophils in the immune response to helminth infections is incompletely understood (Klion and Nutman, 2004). However, the classical view of eosinophils as effectors for killing worms has given way to the idea that they are probably not directly involved in killing them, but are actually important regulators of the cellular immune response (Gebreselassie *et al.* 2012; Gosnell and Kramer, 2013). As supported by rodent studies of AEM, microglia excrete eosinophil chemoattractants such as eotaxin and macrophage inflammatory protein in response to dead worms, to recruit eosinophils to the brain parenchyma (Chang *et al.* 2004; Chang and Yen, 2004; Intapan *et al.* 2007; Gosnell and Kramer, 2013; Zhao *et al.* 2013; Li *et al.* 2014a; Wei *et al.* 2015). Once in the brain, eosinophils enhance this Th2 response by producing Th2 cytokines, excreting chemoattractants, and presenting antigen to CD4⁺ T cells (Spencer and Weller, 2010). Consequently, levels of Th2 cytokines such as interleukin (IL) 4, IL5, IL10 and IL13 increase in the brain and CSF (Intapan *et al.* 2008; Yu *et al.* 2015). Th2 cytokines may also become elevated in the periphery (Diao *et al.* 2009). Angiostrongyliasis can increase CNS levels of IL33; an important mediator of eosinophil infiltration, Th2 cell differentiation and expression of IL5 and IL13 (Du *et al.* 2013; Peng *et al.* 2013;




Fig. 6. Haematoxylin and eosin-stained tissue sections from Australian wildlife showing histopathological changes caused by *A. cantonensis* infection. (A) Spinal cord section from a sub-adult female brushtail possum that had hind limb paralysis. Focal haemorrhage of the spinal cord is apparent, along with greatly thickened, congested and inflamed meninges. (B) The same spinal cord section shown in (A), though at a higher magnification. The focal haemorrhage and thickened, congested, inflamed meninges can be seen more clearly, associated with a cross-section of a larval nematode identified as *A. cantonensis* in this animal. The meninges are infiltrated by plasma cells, lymphocytes, macrophages and neutrophils (marked non-suppurative meningitis). (C) Spinal cord from the same brushtail possum showing tissue damage and granulomatous inflammation in the white matter where presumably a larval *Angiostrongylus* travelled through. (D) Brain section from an affected tawny frogmouth showing cross-sections of larval *Angiostrongylus* associated with granulomatous meningitis. (E) Brain and meninges from the same tawny frogmouth as (D), showing more obvious granulomatous meningitis. (F) Brain from another affected tawny frogmouth showing a larval *Angiostrongylus* migrating through the white matter before an inflammatory response has occurred.

Chuang *et al.* 2016; Saluja *et al.* 2015). IL5 and IL13 are two of several important regulators of IgE production (Deo *et al.* 2010). As a result, intrathecal IgE may be elevated in AEM (Dorta-Contreras *et al.* 2005, 2011; Padilla-Docal *et al.* 2008). Monocyte chemoattractant protein 1 is also expressed in the brain of mice in response to injury caused by migrating larvae (Yu *et al.* 2015).

Blood brain barrier (BBB) dysfunction is a feature of AEM, associated with the activity of host matrix metalloproteinase-9 (MMP-9); a protease that degrades extracellular matrix proteins such as

fibronectin and elastin (Hsu and Lai, 2007; Wei *et al.* 2011). In healthy brain tissue, MMP-9 expression is low, though becomes elevated in response to certain stimuli, including brain tissue damage (Vafadari *et al.* 2016). In AEM, MMP-9 expression increases, possibly due to the damage inflicted by migrating worms (Tsai *et al.* 2008). Mouse studies of AEM suggest that eosinophils release MMP-9 into the subarachnoid space (Tseng *et al.* 2004), activating a proteolytic cascade that disrupts the BBB (Chen *et al.* 2006; Tsai *et al.* 2008; Chiu and Lai, 2013, 2014). In mice with AEM, MMP-9 was also

observed within endothelial cells lining the vascular spaces of the brain and in leucocytes within the subarachnoid space (Lai *et al.* 2004). *Angiostrongylus cantonensis* also excretes MMPs and other proteases involved in the pathogenesis of AEM (discussed in a later section). Dysfunction of the BBB in AEM may also be mediated by vascular endothelial growth factor; an inducer of vascular permeability and mediator of brain oedema (Tsai *et al.* 2007a). Levels of pro-apoptotic proteins, plasminogen activators, and reactive oxygen species are also elevated in the CNS of mice with AEM (Hou *et al.* 2004; Chen *et al.* 2006, 2008). Expression of the 14-3-3 β protein; a marker of neuronal damage, is also elevated in CSF and serum from AEM patients (Tsai *et al.* 2014a).

The clinical manifestations of angiostrongyliasis occur partly as a result of increased intracranial pressure (ICP); a common clinical sign of AEM (Graeff-Teixeira *et al.* 2009; Murphy and Johnson, 2013). Increased ICP may result from vasodilation in the subarachnoid space and brain parenchyma, decreased absorption of CSF, or brain oedema (Murphy and Johnson, 2013). As a consequence of high ICP, AEM patients often present with mild to severe headaches, though prolonged high ICP can eventuate in more serious neurological sequelae (Wang *et al.* 2008, 2010; Tseng *et al.* 2011; Murphy and Johnson, 2013).

Eosinophilic meningitis is the most common manifestation of angiostrongyliasis (Wang *et al.* 2012). Rarely, severe sequelae including coma, convulsion, epilepsy, amentia, hypomnesia and even death can occur (Wang *et al.* 2010; Howe, 2013; Morton *et al.* 2013). However, patients more often present with headache, neck stiffness, paraesthesia, muscle weakness, Brudzinski's sign, fever, vomiting and nausea. Symptoms such as face/limb paralysis, memory loss, confusion, dizziness, conscious disturbance, tinnitus, hyperesthesia, dystonia, urinary retention, photophobia, pneumonitis, peritonitis, abdominal pain, bowel dysfunction, orbital/retro-orbital pain and paralysis of the extra-ocular muscles, are also less common (Chau *et al.* 2003; Podwall *et al.* 2004; Furugen *et al.* 2006; Jin *et al.* 2008; Li *et al.* 2008; Wang *et al.* 2008, 2010; Tseng *et al.* 2011; Kwon *et al.* 2013). Symptoms can be protracted, taking months to disappear (Hochberg *et al.* 2011). In humans and animals, neurological damage is sometimes irreversible (Chau *et al.* 2003; Batmanian and O'Neill, 2004) (Fig. 4, online Supplementary materials). Encephalitic angiostrongyliasis is a rarer manifestation that is generally fatal (Sawanyawisuth, 2008). Elderly patients who become infected with *A. cantonensis* and experience fever and prolonged headaches are at greater risk of developing encephalitic angiostrongyliasis (Sawanyawisuth, 2008; Sawanyawisuth *et al.* 2009).

Myelitis, sacral myeloradiculitis and inflammation of the nerve roots can occur in AEM (Hsu *et al.* 2009; Murphy and Johnson, 2013; Ueda *et al.* 2015), though this is more often associated with gnathostomiasis; the disease caused by *Gnathostoma* spp. worms (Schmutzhard *et al.* 1988). Intraparenchymal cerebral haemorrhage has also been reported (Lilic and Addison, 2013). Presumably, disease severity and incubation period vary depending on the number of larvae consumed (Murphy and Johnson, 2013). Incubation periods range from as little as 1 day to several months, with a median of 11 days following ingestion of L₃ larvae (Zhou *et al.* 2009; Tseng *et al.* 2011; Wang *et al.* 2012; Murphy and Johnson, 2013; Sawanyawisuth *et al.* 2013). As human angiostrongyliasis is often self-limiting, mortality rates are usually low (Wang *et al.* 2008; Graeff-Teixeira *et al.* 2009).

While there is little supporting evidence, children seem especially predisposed to angiostrongyliasis, as evidenced by their disproportionate representation in some outbreaks (Yii, 1976), and the particularly severe cases reported in infants and children (Prociv *et al.* 2000; Li *et al.* 2001; Lindo *et al.* 2004; Morton *et al.* 2013; Murphy and Johnson, 2013; Evans-Gilbert *et al.* 2014). The reason for this remains unclear. However, in cases such as those reported from Taiwan, infected children were thought to have been playing with live snails and possibly eating them (Yii *et al.* 1975; Yii, 1976), which is something adults are unlikely to do. This may expose children to greater numbers of larvae, resulting in severer manifestations. Wang *et al.* (2008) and Sawanyawisuth *et al.* (2013) also note that the clinical manifestations of AEM differ between adults and children; somnolence, fever, constipation, abdominal pain, vomiting, nausea, hepatomegaly, neck stiffness and cranial nerve palsies were more common in children.

In approximately 1.1% of cases *A. cantonensis* causes ocular disease, which may or may not be concurrent with AEM (Patikulasila *et al.* 2003; Sawanyawisuth *et al.* 2007; Baheti *et al.* 2008; Chi *et al.* 2014). Patients with ocular angiostrongyliasis may present with blepharospasm, diplopia, strabismus, blurred vision, loss of colour vision or complete vision loss (Liu *et al.* 2006; Wang *et al.* 2006b; Sawanyawisuth *et al.* 2007; Sinawat *et al.* 2008; Qi *et al.* 2009). The route taken by *A. cantonensis* to enter the eye is unknown, though worms may travel from the brain via the optic nerve, through the circulation via the retinal artery, or enter the eye directly from the environment (Martins *et al.* 2015). Worms may be found in the anterior chamber, the vitreous cavity or the subretinal space (Kumar *et al.* 2005; Malhotra *et al.* 2006; Sawanyawisuth *et al.* 2007; Crane *et al.* 2013; Sinawat and Yospaiboon, 2013; Galor and Eberhard, 2014). Clinical signs of ocular angiostrongyliasis are diverse and may include uveitis, macular

oedema, retinal oedema, necrotic retinitis, panophthalmitis, papilledema, optic neuritis, optic nerve compression, orbital inflammation, increased intraocular pressure, retinal oedema, macular oedema and a pale optic disc (Kumar *et al.* 2005; Liu *et al.* 2006; Wang *et al.* 2006b; Sawanyawisuth and Kitthaweesin, 2008; Sinawat *et al.* 2008; Qi *et al.* 2009; Feng *et al.* 2013; Sinawat and Yospaiboon, 2013; Chi *et al.* 2014). Altered epithelial pigment and subretinal tracks may also be observed (Sinawat *et al.* 2008).

DIAGNOSIS

Diagnosis of AEM is often overlooked, particularly in regions previously considered non-endemic. The lack of standardization in diagnostic procedures for angiostrongyliasis often results in a presumptive diagnosis; based on patient history and suggestive clinical findings (i.e. eosinophilic meningitis) (Murphy and Johnson, 2013). A history of residence or travel to endemic regions and/or a history of raw mollusc consumption are integral in establishing the diagnosis (Tsai *et al.* 2003; Cowie, 2013b). A history of eating unwashed fresh produce, such as lettuce, is also informative (Lindo *et al.* 2002; Waugh *et al.* 2005). An accurate patient history may differentiate AEM from neural gnathostomiasis, which can also manifest as eosinophilic meningitis. While AEM patients often have a history of eating raw molluscs, gnathostomiasis patients usually recall eating undercooked poultry or fish (Senthong *et al.* 2013).

Microscopic detection of L₃ larvae (Fig. 2) from a patient's CSF or eye provides a definitive diagnosis. However, the sensitivity of CSF microscopy depends on sample volume, which is generally limited, often leading to poor sensitivity and false negative results (Prociv *et al.* 2000; Chen *et al.* 2005; Graeff-Teixeira *et al.* 2009). Molecular and immunological tests offer greater sensitivity and several of these have been described for angiostrongyliasis (discussed in later sections). Peripheral blood and CSF eosinophil counts, and CT or MRI imaging can aid the diagnosis (Graeff-Teixeira *et al.* 2009). In ocular angiostrongyliasis, worms may be observed by slit lamp examination of the eye (Malhotra *et al.* 2006; Mattis *et al.* 2009).

During a helminth infection, the proportion of eosinophils may reach 7–36% of the total white blood cell count in peripheral blood (normal range is 0.5–5%) and 10% or more (100–1000 eosinophils per μL) of the white cell count in CSF (normal range is <10 eosinophils/ μL^{-1}) (Schulte *et al.* 2002; Wang *et al.* 2008; Sawanyawisuth and Chotmongkol, 2013). Eosinophils can be difficult to differentiate from neutrophils using some staining techniques such as the toluidine blue wet film. In aseptic meningitis, particularly associated with peripheral

eosinophilia, more specific Romanowsky-based stains such as the May–Grünwald–Giemsa stain or Wright stain should be performed (Senanayake *et al.* 2003; Graeff-Teixeira *et al.* 2009). The Diff-Quik stain may also be useful, having been used to confirm eosinophilia in cases of canine angiostrongyliasis (Lunn *et al.* 2012). While *A. cantonensis* is the leading cause of eosinophilic meningitis, other aetiological agents must be considered in the differential diagnosis (Table 3).

Brain MRI or CT scans may reveal some abnormalities in AEM patients, though lesions resulting from gnathostomiasis are often distinct from those seen in AEM (Senthong *et al.* 2013). In gnathostomiasis with CNS involvement, abnormal CT or MRI findings are common (Kanpittaya *et al.* 2012). Conversely, AEM patients may have normal CT or MRI findings though non-specific cerebral oedema, focal oedematous changes, nodular enhancing lesions, meningeal/leptomeningeal enhancement and mild ventricular dilatation may be apparent (Tsai *et al.* 2003, 2012; Jin *et al.* 2008; Wang *et al.* 2010; Tseng *et al.* 2011; Kanpittaya *et al.* 2012; Nalini *et al.* 2013). Intracerebral haemorrhage and myelitis are more suggestive of gnathostomiasis (Kanpittaya *et al.* 2012). Lesions may also be present on the spinal cord of AEM patients (Diao *et al.* 2010). Brain abnormalities may become apparent in MRI scans as early as 3 weeks after the onset of AEM symptoms (Jin *et al.* 2005).

Immunodiagnostic assays

Immunodiagnostic tests for angiostrongyliasis using purified antigens or monoclonal antibodies have been available for decades. The earliest enzyme-linked immunosorbant assays (ELISAs) were developed using crude or partially purified adult *A. cantonensis* antigens (Chuan-Min and Eng-Rin, 1991). Two immunodominant *A. cantonensis* antigens were identified; a 29 and a 31 kDa antigen, and most immunoassays described target these (Wilkins *et al.* 2013) (Table 4). Immunoblots targeting the 29 kDa antigen cross-reacted with sera from patients infected with other tissue-invading helminths, while tests targeting the 31 kDa antigen exhibited greater specificity (Wilkins *et al.* 2013).

As most cases of AEM originate in Thailand, routine serological testing has been implemented there in several regional hospital laboratories (Eamsobhana and Yong, 2009). These laboratories utilize an in-house dot-blot assay based on the 31 kDa antigen, purified from *A. cantonensis* worms (Eamsobhana *et al.* 2003, 2006; Eamsobhana, 2013; Wilkins *et al.* 2013). This 31 kDa antigen consists of up to four distinct glycoproteins that react to sera from angiostrongyliasis patients (Morassutti *et al.* 2012a). As purification of these glycoproteins from worms is laborious, attempts were made to

Table 3. Causative agents to be considered in the differential diagnosis of eosinophilic meningitis

Disease	Aetiology/causal species
Parasitic infections	
Gnathostomiasis	<i>Gnathostoma spinigerum</i>
Schistosomiasis	<i>Schistosoma mansoni</i> , <i>Schistosoma haematobium</i> , <i>Schistosoma japonicum</i> , <i>Schistosoma mekongi</i> , <i>Schistosoma malayensis</i>
Cysticercosis	<i>Taenia Solium</i>
Toxocarosis	<i>Toxocara canis</i> , <i>Toxocara cati</i>
Baylisascariasis	<i>Baylisascaris procyonis</i>
Paragonimiasis	<i>Paragonimus miyazakki</i> , <i>Paragonimus skrjabini</i> , <i>Paragonimus heterotremus</i> , <i>Paragonimus africanus</i> , <i>Paragonimus uterobilateralis</i> , <i>Paragonimus kellicoti</i> , <i>Paragonimus mexicanus</i> , <i>Paragonimus westermani</i>
Trichinellosis	<i>Trichinella spiralis</i>
Hydatidosis	<i>Enchinococcus granulosus</i> , <i>Enchinococcus multilocularis</i>
Coenurosis	<i>Taenia multiceps</i>
Strongyloidiasis	<i>Strongyloides stercoralis</i>
Filariasis	<i>Loa loa</i> , <i>Wucheria bancrofti</i> , <i>Onchocerca volvulus</i> , <i>Meningonema peruzzi</i>
Arthropod infections	
Myiasis	Maggots
Fungal Infections	
Coccidioidomycosis	<i>Coccidioides immitis</i> <i>Coccidioides posadasii</i>
Allergic aspergillus sinusitis	<i>Aspergillus</i> spp.
Bacterial Infections	
Streptococcal newborn infection	Group B Streptococcal Infection
Neurosyphilis	<i>Treponema pallidum</i>
Rocky Mountain spotted fever, Rickettsial illness	<i>Rickettsia</i> spp.
Viral infections	
Coxsackie viral meningitis	Coxsackie virus
Lymphocytic choriomeningitis	<i>Arenaviridae</i>
Other causes of eosinophilic meningitis	
Neoplasms	Chronic Eosinophilic Leukaemia, Systemic Mastocytosis, Chronic Myelomonocytic Leukaemia, Juvenile Myelomonocytic Leukaemia, Chronic Myeloid Leukaemia, Atypical Chronic Myeloid Leukaemia, Idiopathic Hypereosinophilic Syndrome
Drugs	Ibuprofen, Ciprofloxacin, Vancomycin, Gentamicin, Rifampicin, Minocycline
Reagents and plastic devices	Intravenous Catheters, Intracranial Hardware, Iodized oils, Illicit drugs

Note: This table was prepared using information presented by Graeff-Teixeira *et al.* (2009) and Diaz (2009).

generate recombinants of them for immunodiagnostic purposes (Morassutti *et al.* 2013a). This endeavour has been met with limited success so far, which may be attributable to the choice of expression system (Morassutti *et al.* 2013a). Eamsobhana *et al.* (2015b) recently described a rapid dot-immunogold filtration assay for detecting serum antibodies reactive to the 31 kDa antigen. This assay boasts 100% sensitivity and specificity and produces a result in 3–5 min, compared to the 3 h required for an immunoblot test result (Eamsobhana *et al.* 2015b).

Currently, Shenzhen Combined Biotech Co. Ltd, China, provides the only commercial immunoassays for detecting anti-*A. cantonensis* antibodies, though these have not been implemented for routine diagnostic use based on current literature. The company produces three ELISA tests; one for detecting human IgG, another for detecting human IgM, and a third for detecting rat IgG (Hu *et al.* 2011).

Molecular assays

Several nucleic acid amplification assays have been described for the detection of *A. cantonensis* DNA, most targeting the ribosomal RNA genes (rDNA) (Table 4). Given the limited genetic variation in the rDNA between the various Angiostrongylidae, new tests that differentiate them with improved resolution must focus on different markers (Chan *et al.* 2015). The development of molecular tests that differentiate closely related Angiostrongylidae was initially hampered by the limited availability of sequence data for them. A technique for distinguishing certain *Angiostrongylus* spp. was developed several years ago; a PCR restriction fragment length polymorphism technique targeting the mitochondrial *cytochrome c oxidase subunit I (COI)* gene, and the internal transcribed spacer 2 (ITS2) DNA (Caldeira *et al.* 2003). It is unknown however, whether this technique can differentiate between the closely related *A. cantonensis* and *A. mackerrasae*. With the current

Table 4. Assays recently evaluated for the specific detection of *A. cantonensis*

Diagnostic method	Target	Sensitivity	Specificity	Reference
Microscopy Microscopy of CSF	Third-stage larvae	Low – dependent on the volume of CSF tested	High	Graeff-Teixeira <i>et al.</i> (2009); Rai <i>et al.</i> (2014)
Immunological Rapid dot-immunogold - filtration assay	Detects serum antibodies against purified 31 kDa glycoprotein of <i>A. cantonensis</i> in 3–5 min	100%	100%	Eamsobhana <i>et al.</i> (2014, 2015b)
Protein Microarray	Serum antibodies to semi-purified antigens of <i>A. cantonensis</i>	92.1%	100%	Chen <i>et al.</i> (2012b)
ELISA	Serum antibodies to semi-purified antigens of <i>A. cantonensis</i>	92.1%	100%	Chen <i>et al.</i> (2012b)
Sandwich ELISA	Excreted/secreted antigens from adult <i>A. cantonensis</i> in CSF and serum	100% – Minimum detectability was 0.5 ng of antigen per ml of serum	100% – No cross-reactivity with sera from patients with any of nine other parasites	Chen <i>et al.</i> (2011b)
Immunoblot	Serum antibodies to a 29 kDa antigen of <i>A. cantonensis</i>	55.6%	100%	Sawanyawisuth <i>et al.</i> (2011)
Gold immunochromatography	A 55 kDa protein from adult worms of <i>A. cantonensis</i> in patient sera	100%	100% – No cross-reactivity with sera from patients with any of five other parasites	Huang <i>et al.</i> (2010)
Dot-blot ELISA	Serum antibodies to purified crude antigen of <i>A. cantonensis</i>	81.3% of eosinophilic meningitis patients tested positive	100% – No cross-reactivity with sera from patients with other parasites (<i>n</i> = 18) or patients with other causes of meningitis (<i>n</i> = 52)	Tomanakan <i>et al.</i> (2008)
ELISA	CSF antibodies of different IgG subclasses to young adult female <i>A. cantonensis</i> antigen	13–47% – IgG subclass dependent	90–100% – IgG subclass dependent	Kittimongkolma <i>et al.</i> (2007)
Dot-blot assay	Serum antibodies to antigens between 3 and 100 kDa from adult <i>A. cantonensis</i>	100%	86%	Eamsobhana <i>et al.</i> (2006)
Sandwich ELISA	Circulating antigen of adult <i>A. cantonensis</i> in patient sera	86.4%	100% – No cross-reactivity with sera from patients with any of four other parasites	Liang <i>et al.</i> (2005)
Immuno-PCR	Circulating antigen from L ₅ stage <i>A. cantonensis</i> in patient sera	98%	100%	Chye <i>et al.</i> (2004)
Dot-blot assay	Serum antibodies to a 31 kDa glycoprotein antigen from adult <i>A. cantonensis</i>	100%	100%	Eamsobhana <i>et al.</i> (2003, 2004)
Immunoblot	Serum antibody of the IgG4 class to a 29 kDa from young adult somatic extracts of <i>A. cantonensis</i>	75%	95%	Intapan <i>et al.</i> (2003)
ELISA	Antibodies to somatic antigen from young adult, female <i>A. cantonensis</i>	0–100% – Antibody class dependent; 100% for IgG1	88.9–100% – Antibody class dependent; 100% for IgG1	Intapan <i>et al.</i> (2002)

Table 4. (Cont.)

Diagnostic method	Target	Sensitivity	Specificity	Reference
Immunoblot	CSF and serum IgG to a 29 kDa antigen from young adult female <i>A. cantonensis</i>	55.6%	99.4%	Maleewong <i>et al.</i> (2001)
ELISA	Antibodies to a 204 kDa protein antigen from sub-adult (L ₅) <i>A. cantonensis</i>	91% in sera and 83% in CSF	98% in sera and 100% in CSF	Chye <i>et al.</i> (2000)
Nucleotide amplification				
Real-Time PCR (with high-resolution melt curve)	ITS1 DNA	A single <i>A. cantonensis</i> L ₃ larvae	No cross-reactivity with DNA from <i>Clonorchis sinensis</i> and <i>Gnathostoma spinigerum</i>	Ke <i>et al.</i> (2014)
Real-time PCR (TaqMan)	ITS2 DNA	At least a single L ₁ larvae in a grain of rat feces	No cross-reactivity with DNA from four other rat parasites	Fang <i>et al.</i> (2012)
LAMP ^a	18S rRNA gene	1 fg of DNA	No cross-reactivity with DNA from six other parasites	Chen <i>et al.</i> (2011d)
LAMP ^a	ITS1 DNA	0.01 ng of DNA (0.32 larva per 0.1 g of snail tissue)	No cross-reactivity with DNA from five other parasites	Liu <i>et al.</i> (2011)
Multiplex PCR	18S rRNA gene	93.8%	80.6%	Wei <i>et al.</i> (2010)
Real-time PCR (TaqMan)	ITS1	Ten plasmid copies	No cross-reactivity with several DNA samples containing DNA from other parasitic and free living nematodes	Qvarnstrom <i>et al.</i> (2010)
Conventional PCR	18S rRNA gene	At least one L ₃ larva per mg of mollusc tissue	Cross-reactivity reported with DNA from another metastrongyloid nematode	Qvarnstrom <i>et al.</i> (2007)
Conventional PCR	<i>A. cantonensis</i> gene sequence encoding a 66 kDa protein	55% – 18.5 ng μL^{-1} genomic DNA from <i>A. costaricensis</i>	100% – Failed to amplify DNA from CSF samples from patients with cerebral gnathostomiasis and neurocysticercosis	da Silva <i>et al.</i> (2003); Eamsobhana <i>et al.</i> (2013b); Rodriguez <i>et al.</i> (2014)

Note: This table only lists assays described in the literature during or after the year 2000. For information on earlier assays see the reviews by Graeff-Teixeira *et al.* (2009), Eamsobhana and Yong (2009) and Wilkins *et al.* (2013).

^a LAMP, Loop-mediated isothermal amplification.

availability of several *Angiostrongylus* spp. mitochondrial genomes (discussed in a later section), the development of improved, species-specific assays may now be achievable. Molecular detection, though currently not used routinely, holds potential for the future diagnosis of angiostrongyliasis, enabling specific detection of *A. cantonensis* DNA in patient CSF (Wilkins *et al.* 2013; Qvarnstrom *et al.* 2016).

One conventional PCR assay amplifying a 1134 bp fragment of the 18S rDNA was used to detect *A. cantonensis* DNA in molluscs from Hawaii, in response to an angiostrongyliasis outbreak (Qvarnstrom *et al.* 2007). Sequencing of PCR products obtained in that study confirmed three false positives resulting from cross-reactivity with other nematode species; a

consequence of the highly conserved nature of 18S rDNA (Qvarnstrom *et al.* 2007). Another PCR assay targeting a 66 kDa protein from *A. cantonensis/A. costaricensis* is also available. This assay detected *A. costaricensis* DNA in patient sera and in paraffin-embedded tissues (da Silva *et al.* 2003; Rodriguez *et al.* 2014). It also detected *A. cantonensis* DNA in patient CSF, though with comparatively low sensitivity (Table 4) (Eamsobhana *et al.* 2013b).

A TaqMan qPCR assay was developed for detecting *A. cantonensis* DNA in molluscs collected in the field in Hawaii (Qvarnstrom *et al.* 2010). This assay amplifies the ITS1 DNA region, which is less conserved than the 18S rDNA among helminth species (Qvarnstrom *et al.* 2010). The assay has also been

trialed on human CSF for diagnosing AEM, with promising results (Thyssen *et al.* 2013; Qvarnstrom *et al.* 2016). However, some patients returned intermittently positive and negative results, indicating that the amount of *A. cantonensis* DNA in CSF changes over the course of infection (Qvarnstrom *et al.* 2016). It has also been used to screen molluscs in field studies from various other locations (Tables 1 and 2), and to detect *A. cantonensis* DNA in the blood and peripheral tissues of experimentally infected, wild-type *R. rattus* from Hawaii (Jarvi *et al.* 2015). However, this assay was recently found to cross-react with DNA from *A. mackerrasae* (Chan *et al.* 2015). Consequently, a more specific test is required to confirm the prevalence of *A. cantonensis* in Australia, where *A. mackerrasae* is endemic (Mackie *et al.* 2013). *Angiostrongylus mackerrasae* causes disease in animals though its zoonotic potential remains unknown (Spratt, 2015). Some human cases of angiostrongyliasis may have been attributable to *A. mackerrasae* in Australia, though the absence of an assay that differentiates it from *A. cantonensis* makes this possibility difficult to explore. Mitochondrial genes such as the *COI* gene and *NADH dehydrogenase subunit 6* gene represent possible targets for differentiating closely related Angiostrongylidae as they possess greater variability between populations compared with the chromosomal rDNA sequences (Gasser *et al.* 2012; Lv *et al.* 2012, 2014; Tokiwa *et al.* 2012; Aghazadeh *et al.* 2015c; Yong *et al.* 2015c).

Two loop-mediated isothermal amplification (LAMP) assays have been developed for *A. cantonensis*, one targeting the ITS1 DNA (Liu *et al.* 2011) and the other targeting the 18S rDNA (Chen *et al.* 2011d). These assays were up to ten times more sensitive than conventional PCR (Chen *et al.* 2011d; Liu *et al.* 2011) and have the added benefit of being performed at a single temperature (~65 °C), on a simple heat block or in a warm water bath.

MOLECULAR BIOLOGY

Several wide-scale genomic, transcriptomic and proteome studies have contributed to our current understanding of *Angiostrongylus* spp. molecular biology. These studies will potentially underpin the development of improved diagnostics, therapeutics and novel molecular tools for differentiating species and strains within the Angiostrongylidae. As more data becomes available, it is becoming increasingly apparent that *A. cantonensis* genetically diverse, probably attributable to its sexual reproductive cycle. Sex determination occurs via an XX/XO system, whereby males receive a single X-chromosome and females receive two, resulting in a diploid chromosome number of 11 and 12, for males and females, respectively (Eamsobhana *et al.*

2009). Genetically distinct strains of *A. cantonensis* can possess significant differences in infectivity, fecundity and virulence (Lee *et al.* 2014).

Certain *Angiostrongylus* spp. mitochondrial genes are recognized as a source of genetic diversity. Nine unique *COI* gene haplotypes were identified in a study of Brazilian *A. cantonensis* isolates (Monte *et al.* 2012). Wang and Lv (2014) identified 39 *COI* haplotypes in isolates from China. In studies of Thai isolates, as many as 15 mitochondrial *cytochrome b* gene haplotypes were identified (Dusitsittipon *et al.* 2015; Yong *et al.* 2015b). By comparison, only six 18S rDNA haplotypes are known (Eamsobhana *et al.* 2013a, 2015a).

Nuclear genomes

The *A. cantonensis* nuclear genome is large compared with other nematodes (Zarowiecki and Berriman, 2015). Two *A. cantonensis* nuclear genomes have been sequenced; one (~253 Mb) from a Taiwanese isolate (Morassutti *et al.* 2013b) (GenBank accession: PRJEB493) and the other (~260 Mb) from a Thai isolate (Yong *et al.* 2015a) (GenBank accession: PRJNA260338). Morassutti *et al.* (2013b) detected 28 080 putative open reading frames, 3370 of which possessed homology to protein sequences in public databases. Yong *et al.* (2015a) detected 17 482 genes greater than 300 base pairs, 7737 of which were predicted to encode proteins. Of the proteins with known function, most were kinases and transferases. Genes encoding cellular components, including components of the cytoskeleton, were the most abundant (Yong *et al.* 2015a). While most genes obtained matches *Caenorhabditis* spp. sequences, *A. cantonensis* shares a closer phylogenetic relationship to other bursate nematodes such as *Haemonchus contortus* and *Necator americanus* (Morassutti *et al.* 2013b; Yong *et al.* 2015a).

Morassutti *et al.* (2013b) identified genes encoding immunogenic proteins using previously published mass spectrometry data to interrogate their genome. A cDNA walking strategy was used to obtain the complete coding sequence for these genes, with the intention of generating recombinants that might facilitate diagnostic development. Of the immunogenic proteins of interest, six galectins were identified as candidates for expression (Morassutti *et al.* 2012b). Yong *et al.* (2015a) noted nine galectins amongst 34 excreted-secreted proteins identified in their *A. cantonensis* genome.

Wolbachia spp. are endosymbionts of some arthropods and certain filarial nematodes including *Brugia*, *Wuchereria* and *Onchocerca* spp. (Ferri *et al.* 2011; Slatko *et al.* 2014). In *Brugia malayi* the *Wolbachia* endosymbiont is essential for survival and fertility (Foster *et al.* 2005). Tsai *et al.* (2007b) provided PCR evidence for a *Wolbachia* endosymbiont in *A. cantonensis* which was unusual, given that

Wolbachia had never been reported in a non-filarial nematode (Foster *et al.* 2008). To confirm this, Foster *et al.* (2008) used PCR and immunohistochemistry to examine *A. cantonensis* and *A. costaricensis* but found no evidence of *Wolbachia*. In support of this, Morassutti *et al.* (2013b) and Yong *et al.* (2015a) report no evidence of *Wolbachia* in their genome sequences. While Foster *et al.* (2008) suggest that the findings of Tsai *et al.* (2007b) may be the result of contamination, it should be noted that not all specimens within a filarial nematode species harbour *Wolbachia* even if that species is known to possess a *Wolbachia* endosymbiont (Ferri *et al.* 2011).

Mitochondrial genomes

The mitochondrial (mt) genomes of *A. cantonensis*, *A. mackerrasae*, *A. costaricensis* and *A. vasorum* are similar in size and nucleotide composition. Each circular mt genome encodes 36 genes; 12 proteins, 22 transfer RNAs and two rRNAs (Gasser *et al.* 2012; Lv *et al.* 2012, 2014; Aghazadeh *et al.* 2015c; Yong *et al.* 2015c). The first mt genomes of *A. cantonensis* and *A. costaricensis* to be sequenced were 13 497 and 13 585 bp in length, respectively, and were 81.6% similar at the nucleotide level (Lv *et al.* 2012). The mt genome of the canine pathogen, *A. vasorum* is 13 422 bp (Gasser *et al.* 2012), and that of *A. mackerrasae* is 13 640 bp (Aghazadeh *et al.* 2015c). Pairwise comparisons of translated mt protein coding genes from *A. mackerrasae* to those of *A. cantonensis* revealed only a 2.4% difference, indicating that these species diverged very recently (Aghazadeh *et al.* 2015c). This is in contrast to a 16.8% difference observed between *A. mackerrasae* and *A. costaricensis* (Aghazadeh *et al.* 2015c).

Lv *et al.* (2014) sequenced mt genomes from seven Chinese isolates of *A. cantonensis* which revealed five unique types. These Chinese mt genomes were compared with that of a Taiwanese and Thai isolate (Yong *et al.* 2015c). The Taiwanese and Chinese isolates possess smaller mt genomes than the Thai isolate (Yong *et al.* 2015c), which also possessed a genome that was more distantly related compared with the Chinese/Taiwanese genomes. Eighteen tRNAs encoded on the Thai mt genome lacked a TΨC-arm (Yong *et al.* 2015c); a tRNA structure that interacts with elongation factor Tu during protein synthesis (Watanabe *et al.* 2014). While the absence of a TΨC-arm is not uncommon in tRNAs encoded on nematode mt genomes (Watanabe *et al.* 2014), differences were observed between isolates; 17 and 16 tRNAs lacked the TΨC-arm in the Chinese and Taiwanese mt genomes, respectively (Yong *et al.* 2015c). A second 13 652 bp *A. costaricensis* mt genome was sequenced from a Costa Rican isolate (Yong *et al.* 2015d) and compared to the original *A. costaricensis* mt genome, derived from a

Brazilian parasite (Lv *et al.* 2012). Based on phylogenetic analyses and genetic distances calculated for the mt genes, it was proposed that these parasites should be considered distinct members of a species complex, or sibling species (Yong *et al.* 2015d).

Generally, comparisons of mt genes demonstrate greater resolution for differentiating species/strains within the Angiostrongylidae, compared with the nuclear rDNA. These differences could be exploited to develop molecular tools that differentiate *Angiostrongylus* spp., and may facilitate the identification of new, cryptic species (Gasser *et al.* 2012; Yong *et al.* 2015d).

Transcriptomes, proteomes and virulence factors

The transition between larval and adult stages of *A. cantonensis* is a complex process requiring stringent regulation of gene expression. Several studies have investigated differences in gene expression at the transcript and protein level between stages, to identify mechanisms associated with stage transition and function in *Angiostrongylus* spp. While some genes are expressed constitutively across all stages, some are stage specific. Chang *et al.* (2014a) found that transcripts exclusive to L₅ worms were related to growth, development, sexual differentiation and reproduction, while L₃-specific transcripts were mostly related to metabolism. The classes of proteases expressed between these two stages also differed (Chang *et al.* 2014a). Huang *et al.* (2013) observed differences in galectin abundance in the soluble proteomes of male L₅ & male adult worms, and female adult & male adult worms. Stage- and sex-dependent differences in expression of aspartic protease and ornithine decarboxylase antizyme were also noted (Hwang *et al.* 2010; Chen *et al.* 2013), as were differences in antioxidant capacity (Song *et al.* 2012; Huang *et al.* 2013).

Transcriptomes sequenced from all worm stages suggest that proteases are fundamental to *Angiostrongylus* spp. (Ansell *et al.* 2013; Wang *et al.* 2013b). Proteases are important virulence factors in other pathogenic helminths, essential for tissue penetration and nutrient acquisition (Stack *et al.* 2011; McVeigh *et al.* 2012; Morassutti and Graeff-Teixeira, 2012). Chang *et al.* (2014a) identified transcripts for proteases of various types in the L₃ and L₅ stages; most notably metalloproteases, aspartic proteases & cysteine proteases in the L₃ stage and cysteine, aspartic & serine proteases in the L₅ stage. Serine proteases of *A. cantonensis* react to sera from infected patients (Morassutti *et al.* 2012b), and a recombinant asparaginyl endopeptidase was recognized by serum from infected BALB/c and ICR strain mice and, by serum from human angiostrongyliasis patients (Chang *et al.* 2014b). This supports their role at the host-parasite interface.

High expression of cathepsin B-like proteases 1 and 2 (AC-cathB-1, AC-cathB-2) and haemoglobin-type cysteine protease (AC-haem) occurs in L₃ larvae (Ni *et al.* 2012; Yu *et al.* 2014a). As with other helminths, cathepsin B- and D-like proteases of *A. cantonensis* degrade host proteins, including haemoglobin and IgG (Brindley *et al.* 2001; Baig *et al.* 2002; Beckham *et al.* 2006; Jilkova *et al.* 2011; Cheng *et al.* 2012). Expression of AC-cathB-2 is higher in L₃ larvae than in L₁ larvae and adults (Ni *et al.* 2012), while AC-cathB expression is greater in L₄ and L₅ worms (Han *et al.* 2011). High expression of AC-cathBs occurs in the oesophagus of L₃ larvae, supporting their role in host gut penetration (Yu *et al.* 2014a; Long *et al.* 2015). In L₄ worms, AC-cathB-1 is excreted/secreted and localizes to the intestine (Cheng *et al.* 2012). These studies indicate that in L₃ larvae, AC-cathBs are probably important for host gut penetration, and in L₄ and L₅ worms, are involved in host tissue destruction during migration.

Metalloprotease I, AC-cathB-1 and AC-cathB-2 were amongst the most abundant transcripts in pepsin-HCl activated L₃ larvae (Chang *et al.* 2011). Excreted/secreted products of L₁ and L₃ larvae possess demonstrable proteolytic activity (Lai *et al.* 2005b; Lee and Yen, 2005; Rebello *et al.* 2012). Furthermore, serine protease and metalloprotease inhibitors reduced penetration of L₃ larvae through the intestinal wall of mice (Lee and Yen, 2005). Metalloprotease activity is present in L₁, L₃, L₅ and adult worms (Lai *et al.* 2005b), though metalloprotease transcripts were not detected in L₄ worms, an absence that is not understood (He *et al.* 2009). Metalloproteases of *A. cantonensis* also degrade gelatin and host metalloprotease (Lai *et al.* 2005b; Adisakwattana *et al.* 2012). In adult *A. vasorum* 4-5% of all functionally annotated transcripts classified as proteases by homology, including cysteine proteinase 3, aspartyl protease and cathBs (Ansell *et al.* 2013). In adult worms, these proteases probably play a role in the digestion of haemoglobin and other host proteins (Cheng *et al.* 2012; Ansell *et al.* 2013). A role for AC-cathB-2 in reproduction is also proposed, given its localization to the vas deferens, uterus and oviduct of adult worms (Yu *et al.* 2014a).

Helminth-derived molecules that regulate host immune responses are considered important virulence factors; inducing an immune state that tolerates the parasites' presence. Extracts from *Angiostrongylus* spp. reduced inflammation in a rodent model of asthma (Pitrez *et al.* 2015). Similarly, asthmatic rats treated with *A. cantonensis* cystatin experienced reduced airway inflammation (Ji *et al.* 2015). Transcripts for immunomodulatory proteins were among the most abundant in adult *A. vasorum* (Ansell *et al.* 2013). Homologues of *Ancylostoma* secreted protein (activation-associated

secreted proteins or ASPs) were noted in *A. vasorum*. Helminth ASPs modulate host leucocyte activity and recruit neutrophils by neutrophil receptor binding (Ansell *et al.* 2013). Additionally, ASP was identified in a cDNA library of L₄ *A. cantonensis* larvae and a recombinant version of this protein was recognized by sera from infected mice but not from infected rats (Yang *et al.* 2013b). The ASP gene was highly transcribed in brain stage L₄ larvae, and immunization of mice with recombinant ASP before infection reduced neutrophil infiltration in mouse brains, implicating ASPs as being directly involved in the pathogenesis of AEM (Yang *et al.* 2013b). Ansell *et al.* (2013) also identified homologues of high mobility group box protein in adult *A. vasorum*. Homologues of this protein from filarial nematodes possess an immunomodulatory role, triggering secretion of pro-inflammatory cytokines such as IL-6 and tumour necrosis factor (TNF)- α by mouse peritoneal macrophages (Thirugnanam *et al.* 2012).

Constituents of the cytoskeleton, galectins and heat-shock proteins are among the most abundant proteins expressed by *A. cantonensis*, based on proteomic studies (Leon *et al.* 2007; Song *et al.* 2012; Huang *et al.* 2013; Chen *et al.* 2014a). Galectins participate in a diverse range of processes including cell adhesion and host immune modulation (Morassutti and Graeff-Teixeira, 2012; Huang *et al.* 2013). Galectins also react with serum from angiostrongyliasis patients (Morassutti *et al.* 2012b). Stage-specific expression of galectin 1 has been noted; its expression is elevated 11-fold in male L₅ worms compared with male adults (Huang *et al.* 2013). Expression of some galectins is also higher in female adult worms compared with males and vice versa (Song *et al.* 2012; Huang *et al.* 2013), though the significance of this is unknown.

Nitric oxide is produced by the host immune system in response to infections with helminths and other pathogens, to subject them to conditions of oxidative stress (Muro and Perez-Arellano, 2010). Consequently, helminths have evolved defences to counter this. Expression of *A. cantonensis* galectin 10 increased following exposure to H₂O₂, suggesting its involvement in the prevention of oxidative stress (Liu *et al.* 2013). Glutathione-S-transferase (GST) is also involved in helminth defences against oxidative stress, by detoxifying harmful compounds produced by oxygen free radicals (Vibanco-Perez and Landa-Piedra, 1998; Kampkotter *et al.* 2003). Consequently, GST has long been considered a potential target for anthelmintic chemotherapy (Vibanco-Perez and Landa-Piedra, 1998). GST family proteins are highly expressed in adult female *A. costaricensis* (Leon *et al.* 2007). Antioxidants such as peroxiredoxin and thioredoxin peroxidase are also highly expressed, and represent potential chemotherapeutic targets (Leon *et al.* 2007; Huang

et al. 2013). Female adult *A. cantonensis* possess increased antioxidant capacity compared with female L₅ worms (Huang *et al.* 2013), suggesting that female worms increase defences against oxidative stress as they mature, and possibly in preparation for embryogenesis.

MicroRNAs

MicroRNAs (miRNA) are small non-coding RNAs (~21–23 nt) that regulate a wide range of biological processes in parasitic helminths, including development, metabolism and cell differentiation/proliferation (Chen *et al.* 2011a; Britton *et al.* 2014). Consequently, miRNAs have been investigated for their role in helminth biology and as potential diagnostic biomarkers of helminth infection (Britton *et al.* 2014).

Chen *et al.* (2011a) detected 1 072 876 miRNA species in adult *A. cantonensis*; 122 051 were common to both sexes, 548 720 were female specific and 402 105 were male specific. The miRNA profiles differed between young adult and mature adult worms, though certain miRNA species, such as mirR-1, mirR-71 and miR-44, were expressed highly in both stages (Chang *et al.* 2013). Mir-71 family members are the most abundant miRNA species in young adult and mature adult worms (Chen *et al.* 2011a; Chang *et al.* 2013). Mir-71 is thought to promote longevity and stress resistance and along with aca-miR-1–1, may also affect sexual differentiation and development (Chang *et al.* 2013). In L₃ larvae, mir-200c was the most abundant miRNA identified, and its expression was 352-fold higher in L₃ larvae compared with L₄ larvae (Li *et al.* 2014b). Li *et al.* (2014b) noted higher expression of miR-124 in L₄ worms, and implicate it as an immunomodulator. Mouse microglial cells transfected with miR-124 mimics significantly down-regulated transcription of IL-6, IL-1 β and TNF- α (Li *et al.* 2014b).

The utility of miRNAs as biomarkers for *A. cantonensis* infection is also under investigation. Expression of miR-132-3p in mouse brains, and of aca-miR-146a/aca-miR-146a-5p in rodent sera and brains, is elevated in infected animals compared with uninfected controls (Chen *et al.* 2014b; Li *et al.* 2014b; Yu *et al.* 2014b). Consequently, aca-miR-146a and aca-miR-146a-5p could serve as useful biomarkers for early-stage *A. cantonensis* infection.

TREATMENT

Treatment of angiostrongyliasis involves reducing inflammation, reducing ICP, eradicating the worms and reducing pain. Mild cases often resolve spontaneously without therapy, though rarely, severe cases can lead to permanent neurologic damage or, death

in 2–3% of cases (Eamsobhana and Yong, 2009; Morton *et al.* 2013). Repeated lumbar puncture is performed to reduce ICP, thereby relieving severe headaches and other neurological derangements (Jitpimolmard *et al.* 2007; Graeff-Teixeira *et al.* 2009; Diao *et al.* 2011; Murphy and Johnson, 2013). Analgesics such as ibuprofen can also be administered to relieve headache and fever (Wang *et al.* 2010). Administration of intravenous mannitol or glycerol/fructose can reduce ICP in AEM patients (Li *et al.* 2008; Wang *et al.* 2010). Gamma-globulin has also been administered to AEM patients to support immunity (Li *et al.* 2008).

Ocular angiostrongyliasis though non-fatal, can lead to permanent visual impairment if untreated (Sawanyawisuth *et al.* 2007; Feng *et al.* 2013). Surgical removal of larvae from the eye is often performed, along with administration of topical or oral corticosteroids to reduce inflammation (Patikulasila *et al.* 2003; Kumar *et al.* 2005; Sawanyawisuth *et al.* 2007; Feng *et al.* 2013). Laser killing of larvae in the eye can be performed, and is considered preferable to surgery, provided the worms have caused minimal damage at the time of diagnosis (Feng *et al.* 2013). However, some suggest there is little evidence that laser interventions or surgical removal of larvae improve clinical outcomes (Kumar *et al.* 2005; Sawanyawisuth *et al.* 2007). Improvements have been noted upon administration of anthelmintics (Wang *et al.* 2006b), though some investigators advise against this given the risk that dead worms may activate serious intraocular inflammation (Feng *et al.* 2013).

Anthelmintics such as albendazole, mebendazole, flubendazole and ivermectin have been administered in many AEM cases, though their usefulness in AEM management is controversial (Murphy and Johnson, 2013). Some studies suggest that early administration of anthelmintics in AEM may exacerbate symptoms and hasten disease progression (Slom *et al.* 2002; Hidelaratchi *et al.* 2005; Liu *et al.* 2006; Wang *et al.* 2006a). Transaminitis has also been reported following albendazole treatment in an AEM case (Morton *et al.* 2013). However, a double blind, placebo controlled trial confirmed a reduction in headache duration in AEM patients, with no adverse effects, though with borderline efficacy, using albendazole (Jitpimolmard *et al.* 2007; Murphy and Johnson, 2013). Albendazole alone or in combination with corticosteroids, was efficacious, and caused no adverse reactions in AEM mouse models (Lan *et al.* 2004; Tu and Lai, 2006). Similarly, administration of albendazole and prednisolone to a group of 53 Thai patients with eosinophilic meningitis caused no serious adverse reactions (Chotmongkol *et al.* 2009).

Corticosteroids such as dexamethasone and prednisolone form the basis of AEM treatment, by reducing granulomatous inflammation and headache intensity

(Sawanyawisuth *et al.* 2004; Sawanyawisuth, 2008; Thanaviratnanich and Ngamjarus, 2015). The efficacy of corticosteroid & anthelmintic combination therapy for alleviating AEM symptoms is supported by several independent reports (Wan and Weng, 2004; Chotmongkol *et al.* 2006; Leone *et al.* 2007; Diao *et al.* 2009; Lv *et al.* 2009a; Zhou *et al.* 2009; Ueda *et al.* 2015). Furthermore, experiments using peripheral blood mononuclear cells extracted from AEM patients indicate that dexamethasone & albendazole combination therapy dampens the Th2 cytokine response in favour of a Th1 response (Diao *et al.* 2009). Generally, corticosteroids are administered in doses ranging from 3 mg day⁻¹ to 20 mg four times daily, for 1–2 weeks (Chotmongkol *et al.* 2000, 2006; Li *et al.* 2008; Zhou *et al.* 2009; Wang *et al.* 2010; Murphy and Johnson, 2013). Albendazole or mebendazole are administered at doses of 10–20 mg kg⁻¹ day⁻¹ for 1–2 weeks (Chotmongkol *et al.* 2006; Zhou *et al.* 2009; Wang *et al.* 2010). However, no randomized, double-blind, placebo-controlled trials have been performed to assess the efficacy of these regimens, and some investigators report little or no benefit following their administration (Ali *et al.* 2008; Hochberg *et al.* 2011; Tseng *et al.* 2011).

Chotmongkol *et al.* (2009) found no statistical difference between two study groups of more than 50 eosinophilic meningitis patients, treated with either albendazole and prednisolone combination therapy or with prednisolone alone. Furthermore, in rodent models of AEM, dexamethasone treatment alone reduced brain cell apoptosis, BBB permeability and expression of the 14-3-3 β protein in CSF (Tsai *et al.* 2014b, 2015), casting doubt on the therapeutic benefit of administering anthelmintic drugs for the treatment of AEM.

Alternative therapies are also under investigation in rodents. Tribendimidine, a broad spectrum anthelmintic developed in China in the 1980s (Robertson *et al.* 2015), killed *A. cantonensis* in rodents, possibly by damaging its tegument (Wang *et al.* 2013a; Zeng *et al.* 2013a; Feng *et al.* 2014). Treatment of mice with IL-12 and mebendazole reduces AEM severity and causes a shift towards a Th1 response (Du *et al.* 2003). Lai *et al.* (2005a) used albendazole in combination with GM6001 (a matrix MMP-9 inhibitor) to reduce AEM severity in mice. In combination with albendazole, certain Chinese herbs and their extracts exhibited some beneficial effects in mouse models of *Angiostrongylus* optic neuritis and AEM (Lai, 2006; Feng *et al.* 2015). Ginger extracts possess some activity against *A. cantonensis* (Lin *et al.* 2010). Curcumin (a compound found in turmeric) was not efficacious in AEM mouse models when used alone, though in combination with albendazole, reduced CSF eosinophilia to a greater extent than albendazole mono-therapy (Shyu *et al.* 2012). Similarly, the combination of diammonium glycyrrhizinate (a

liquorice root extract) and albendazole was more effective in a mouse model of AEM than albendazole/dexamethasone co-therapy (Li *et al.* 2013b). The marine fungal extract m2-9 had synergistic effects when used in combination with albendazole for treatment of murine AEM (Li *et al.* 2012). Treatment with albendazole and thalidomide was time-dependent; early treatment produced better outcomes in murine AEM (Chen and Lai, 2007).

To summarize, lumbar puncture is effective at reducing ICP in AEM, relieving patients of the associated headaches (Graeff-Teixeira *et al.* 2009; Murphy and Johnson, 2013). Administration of corticosteroids to reduce inflammation in AEM is also warranted based on multiple reports (Chotmongkol *et al.* 2009; Diao *et al.* 2009; Murphy and Johnson, 2013). Conversely, the value of administering anthelmintics is uncertain (Jitpimolmard *et al.* 2007; Chotmongkol *et al.* 2009; Murphy and Johnson, 2013). As reports of severe adverse events resulting from albendazole use are rare, it should generally be considered safe for AEM treatment, particularly in combination with corticosteroids (Murphy and Johnson, 2013). However, given the borderline efficacy of albendazole monotherapy (Jitpimolmard *et al.* 2007), and the lack of any difference in clinical outcome when comparing corticosteroid monotherapy and corticosteroid/albendazole co-therapy (Chotmongkol *et al.* 2009), the value of administering anthelmintics for AEM treatment is questionable. Given the inconsistencies, these studies represent a useful guide for the treatment of angiostrongyliasis at best. Clearly, further research, including randomized, double-blind, placebo-controlled trials, is required to establish an improved consensus for the treatment of this disease.

PREVENTION AND CONTROL

Given its wide distribution and zoonotic nature, it is unlikely that *A. cantonensis* can be completely eradicated. However, disrupting its transmission to humans through the control of its preferred hosts is plausible. Molluscicides and other methods to control snail and slug numbers have been implemented in Hawaii and China (Hata *et al.* 1997; Hollingsworth *et al.* 2013; Yang *et al.* 2013a). In Jamaica, molluscicides are routinely applied to growing produce but this practice was deemed superficial as molluscs remain in the surrounding vegetation and return presumably when the molluscicides lose their potency (Waugh *et al.* 2005). In China, the collection of *P. canaliculata* egg masses and the use of commercial molluscicides have proved effective, though their use is limited by their laborious nature and potential environmental side effects (Yang *et al.* 2013a). Biological controls, including the introduction of ducks and *Mylopharyngodon piceus* (black carp) to rice paddy fields and ponds drastically

reduced snail numbers (Yang *et al.* 2013a). Hollingsworth *et al.* (2013) suggests drowning of terrestrial slugs or snails for several days in a covered bucket filled with soapy water or a 15% solution of salt water. The salt water treatment is not only lethal to terrestrial molluscs, but is also expected to kill any *A. cantonensis* larvae that may be present (Hollingsworth *et al.* 2013).

Standardized food treatment measures may be effective in some areas, though would be ineffective in locations where food is often home grown or collected from the local environment. Thorough washing of vegetables prior to consumption is a simple measure to reduce *A. cantonensis* transmission (Cowie, 2013b). Washing experiments carried out on grown produce by Yeung *et al.* (2013) showed no difference between simple water washes and washing with other domestic chemicals such as bleach, acetic acid and sodium chloride. While some slug slime residue remained after washes, washing was still considered the most appropriate measure for reducing the transmission of larvae in produce (Yeung *et al.* 2013). As evidenced by Wang *et al.* (2011), thorough roasting of molluscs for more than 20 min is required to ensure that all *A. cantonensis* larvae are destroyed in mollusc tissue.

The major difficulty in controlling the spread of angiostrongyliasis lies in its link with diverse social, economic, cultural and environmental factors. In some endemic areas, awareness of *A. cantonensis* is low in the general community (Li *et al.* 2011). Consequently, educating the public to the dangers of raw mollusc consumption in these regions may be an effective starting measure. Educating laypeople on strategies for controlling the spread of *A. cantonensis* would be helpful, particularly in regions where food is home grown or collected locally. However, public education programmes must be executed with caution, as the public concern generated could be excessive given the rarity of angiostrongyliasis. Despite this, public education campaigns have had favourable results in China (Lv *et al.* 2008). Finally, programmes that raise awareness amongst public health workers and physicians might enable them to provide better advice to patients planning overseas travel, and facilitate a more rapid diagnosis with improved clinical outcomes when future cases of angiostrongyliasis are encountered.

Concluding remarks

While angiostrongyliasis is generally considered rare, its global presence is steadily increasing with both clinical cases and epidemiological surveys confirming *A. cantonensis* in regions where it was previously considered absent. AEM is often overlooked at the outset, particularly in regions where angiostrongyliasis is rarely observed. There is a lack of standardization in the procedures for

diagnosing angiostrongyliasis. Serological tests are commercially available, though are not widely used. Other diagnostic tools are under development. There is no consensus on the appropriate treatment for angiostrongyliasis and some treatments are controversial. A combination of anthelmintics and corticosteroids seems effective in most cases, though there have been no randomized, double-blind, placebo-controlled trials to support any treatment regimen described for angiostrongyliasis. Blocking transmission is the most appropriate method of reducing infections. Implementing simple wash protocols for vegetables, public education on the dangers of raw mollusc consumption and implementation of mollusc and rat control measures may be useful. In developed nations, the prevalence of angiostrongyliasis in companion animals and wildlife is usually greater than in humans. Thus, animals play a role in alerting public health authorities of the risk to humans in locations where animals are affected by angiostrongyliasis. Considering the potentially lethal nature of angiostrongyliasis and the recent reports of its increasing geographical range, it is important that this disease is given due consideration in the differential diagnosis of eosinophilic meningitis, even in areas where *A. cantonensis* is rarely reported. This is essential to improve patient prognosis and reduce human and companion animal suffering.

SUPPLEMENTARY MATERIAL

The supplementary material for this article can be found at <http://dx.doi.org/10.1017/S0031182016000652>.

ACKNOWLEDGEMENTS

We acknowledge the University of Technology Sydney and St. Vincent's Hospital, Sydney in support of Ph.D. student, Douglas Chan (scholarship and infrastructure, respectively).

REFERENCES

- Abo-Madyan, A. A., Morsy, T. A., Motawea, S. M., El Garhy, M. F. and Massoud, A. M. (2005). Spot light survey on fresh-water snails of medical importance in Al Fayoum Governorate. Egypt. *Journal of the Egyptian Society of Parasitology* **35**, 49–58.
- Adisakwattana, P., Nuamtanong, S., Yenchitsomanus, P. T., Komalamisra, C. and Meesuk, L. (2012). Degradation of human matrix metalloprotease-9 by secretory metalloproteases of *Angiostrongylus cantonensis* infective stage. *Southeast Asian Journal of Tropical Medicine and Public Health* **43**, 1105–1113.
- Aghazadeh, M., Jones, M. K., Aland, K. V., Reid, S. A., Traub, R. J., McCarthy, J. S. and Lee, R. (2015a). Emergence of neural angiostrongyliasis in eastern Australia. *Vector-Borne and Zoonotic Diseases* **15**, 184–190.
- Aghazadeh, M., Reid, S. A., Aland, K. V., Restrepo, A. C., Traub, R. J., McCarthy, J. S. and Jones, M. K. (2015b). A survey of *Angiostrongylus* species in definitive hosts in Queensland. *International Journal for Parasitology: Parasites and Wildlife* **4**, 323–328.
- Aghazadeh, M., Traub, R. J., Mohandas, N., Aland, K. V., Reid, S. A., McCarthy, J. S. and Jones, M. K. (2015c). The mitochondrial genome of *Angiostrongylus mackerrasae* as a basis for molecular, epidemiological and population genetic studies. *Parasites and Vectors* **8**, 473.

- Ali, A. B., Van den Enden, E., Van Gompel, A. and Van Esbroeck, M. (2008). Eosinophilic meningitis due to *Angiostrongylus cantonensis* in a Belgian traveller. *Travel Medicine and Infectious Disease* **6**, 41–44.
- Alicata, J. (1965a). Biology and distribution of the rat lungworm, *Angiostrongylus cantonensis*, and its relationship to eosinophilic meningoencephalitis and other neurological disorders of man and animals. *Advances in Parasitology* **3**, 223–248.
- Alicata, J. E. (1965b). Occurrence of *Angiostrongylus cantonensis* in Madagascar, Mauritius, Ceylon, and Sarawak. *Journal of Parasitology* **51**, 937.
- Andersen, E., Gubler, D. J., Sorensen, K., Beddard, J. and Ash, L. R. (1986). First report of *Angiostrongylus cantonensis* in Puerto Rico. *American Journal of Tropical Medicine and Hygiene* **35**, 319–322.
- Ansell, B. R., Schnyder, M., Deplazes, P., Korhonen, P. K., Young, N. D., Hall, R. S., Mangiola, S., Boag, P. R., Hofmann, A., Sternberg, P. W., Jex, A. R. and Gasser, R. B. (2013). Insights into the immuno-molecular biology of *Angiostrongylus vasorum* through transcriptomics—prospects for new interventions. *Biotechnology Advances* **31**, 1486–1500.
- Archer, C. E., Appleton, C. C., Mukaratirwa, S. and Hope, K. J. (2011). The rat lung-worm *Angiostrongylus cantonensis*: a first report in South Africa. *South African Medical Journal* **101**, 174–175.
- Asato, R., Taira, K., Nakamura, M., Kudaka, J., Itokazu, K. and Kawanaka, M. (2004). Changing epidemiology of *Angiostrongyliasis cantonensis* in Okinawa prefecture, Japan. *Japanese Journal of Infectious Diseases* **57**, 184–186.
- Baheti, N. N., Sreedharan, M., Krishnamoorthy, T., Nair, M. D. and Radhakrishnan, K. (2008). Neurological picture. Eosinophilic meningitis and an ocular worm in a patient from Kerala, south India. *Journal of Neurology, Neurosurgery, and Psychiatry* **79**, 271.
- Baig, S., Damian, R. T. and Peterson, D. S. (2002). A novel cathepsin B active site motif is shared by helminth bloodfeeders. *Experimental Parasitology* **101**, 83–89.
- Baird, J. K., Neafie, R. C., Lanoie, L. and Connor, D. H. (1987). Abdominal angiostrongylosis in an African man: case study. *American Journal of Tropical Medicine and Hygiene* **37**, 353–356.
- Barrett, J. L., Carlisle, M. S. and Procriv, P. (2002). Neuro-angiostrongylosis in wild Black and Grey-headed flying foxes (*Pteropus* spp). *Australian Veterinary Journal* **80**, 554–558.
- Barrow, K. O., Rose, A. and Lindo, J. F. (1996). Eosinophilic meningitis. Is *Angiostrongylus cantonensis* endemic in Jamaica? *West Indian Medical Journal* **45**, 70–71.
- Bartschi, E., Bordmann, G., Blum, J. and Rothen, M. (2004). Eosinophilic meningitis due to *Angiostrongylus cantonensis* in Switzerland. *Infection* **32**, 116–118.
- Batmanian, J. J. and O'Neill, J. H. (2004). Eosinophilic meningoencephalitis with permanent neurological sequelae. *Internal Medicine Journal* **34**, 217–218.
- Beaver, P. C. and Rosen, L. (1964). Memorandum on the first report of *Angiostrongylus* in man, by Nomura and Lin, 1945. *American Journal of Tropical Medicine and Hygiene* **13**, 589–590.
- Beckham, S. A., Law, R. H., Smooker, P. M., Quinsey, N. S., Caffrey, C. R., McKerrow, J. H., Pike, R. N. and Spithill, T. W. (2006). Production and processing of a recombinant *Fasciola hepatica* cathepsin B-like enzyme (FheatB1) reveals potential processing mechanisms in the parasite. *Biological Chemistry* **387**, 1053–1061.
- Bhaibulaya, M. (1975). Comparative studies on the life history of *Angiostrongylus mackerrasae* Bhaibulaya, 1968 and *Angiostrongylus cantonensis* (Chen, 1935). *International Journal for Parasitology* **5**, 7–20.
- Blair, N. F., Orr, C. F., Delaney, A. P. and Herkes, G. K. (2013). *Angiostrongylus* meningoencephalitis: survival from minimally conscious state to rehabilitation. *Medical Journal of Australia* **198**, 440–442.
- Brindley, P. J., Kalinna, B. H., Wong, J. Y. M., Bogitsh, B. J., King, L. T., Smyth, D. J., Verity, C. K., Abbenante, G., Brinkworth, R. I., Fairlie, D. P., Smythe, M. L., Milburn, P. J., Bielefeldt-Ohmann, H., Zheng, Y. and McManus, D. P. (2001). Proteolysis of human hemoglobin by schistosome cathepsin D. *Molecular and Biochemical Parasitology* **112**, 103–112.
- Britton, C., Winter, A. D., Gillan, V. and Devaney, E. (2014). microRNAs of parasitic helminths – Identification, characterization and potential as drug targets. *International Journal for Parasitology: Drugs and Drug Resistance* **4**, 85–94.
- Brumpt, V., Audebaud, G., Klein, J. M., Jolly, M., Mazaud, R. and Goube, P. (1968). Incidence of *Angiostrongylus cantonensis* (Chen) in Cambodia. *Bulletin de la Société de Pathologie Exotique et de ses Filiales* **61**, 444–462.
- Burns, R. E., Bicknese, E. J., Qvarnstrom, Y., DeLeon-Carnes, M., Drew, C. P., Gardiner, C. H. and Rideout, B. A. (2014). Cerebral *Angiostrongylus cantonensis* infection in a captive African pygmy falcon (*Polihiherax semitorquatus*) in southern California. *Journal of Veterinary Diagnostic Investigation* **26**, 695–698.
- Caldeira, R. L., Carvalho, O. S., Mendonca, C. L., Graeff-Teixeira, C., Silva, M. C., Ben, R., Maurer, R., Lima, W. S. and Lenzi, H. L. (2003). Molecular differentiation of *Angiostrongylus costaricensis*, *A. cantonensis*, and *A. vasorum* by polymerase chain reaction-restriction fragment length polymorphism. *Memorias do Instituto Oswaldo Cruz* **98**, 1039–1043.
- Caldeira, R. L., Mendonca, C. L., Goveia, C. O., Lenzi, H. L., Graeff-Teixeira, C., Lima, W. S., Mota, E. M., Pecora, I. L., Medeiros, A. M. and Carvalho Odos, S. (2007). First record of molluscs naturally infected with *Angiostrongylus cantonensis* (Chen, 1935) (Nematoda: Metastrongylidae) in Brazil. *Memorias do Instituto Oswaldo Cruz* **102**, 887–889.
- Carlisle, M. S., Procriv, P., Grennan, J., Pass, M. A., Campbell, G. L. and Mudie, A. (1998). Cerebrospinal angiostrongyliasis in five captive tamarins (*Saguinus* spp). *Australian Veterinary Journal* **76**, 167–170.
- Chan, D., Barratt, J., Roberts, T., Lee, R., Shea, M., Marriott, D., Harkness, J., Malik, R., Jones, M., Aghazadeh, M., Ellis, J. and Stark, D. (2015). The prevalence of *Angiostrongylus cantonensis/mackerrasae* complex in molluscs from the Sydney Region. *PLoS ONE* **10**, e0128128.
- Chancellor, A. (2007). A souvenir from the Pacific Islands. *Practical Neurology* **7**, 205.
- Chang, E. E. and Yen, C. M. (2004). Eosinophil chemoattracted by eotaxin from cerebrospinal fluid of mice infected with *Angiostrongylus cantonensis* assayed in a microchamber. *Kaohsiung Journal of Medical Sciences* **20**, 209–215.
- Chang, E. E., Chung, L. Y. and Yen, C. M. (2004). Kinetics of change in the eotaxin concentration in serum and cerebrospinal fluid of mice infected with *Angiostrongylus cantonensis*. *Parasitology Research* **92**, 137–141.
- Chang, S. H., Tang, P. and Wang, L. C. (2011). A transcriptomic study on the pepsin-activated infective larvae of *Angiostrongylus cantonensis*. *Molecular and Biochemical Parasitology* **179**, 47–50.
- Chang, S. H., Tang, P., Lai, C. H., Kuo, M. L. and Wang, L. C. (2013). Identification and characterisation of microRNAs in young adults of *Angiostrongylus cantonensis* via a deep-sequencing approach. *Memorias do Instituto Oswaldo Cruz* **108**, 699–706.
- Chang, S.-H., Tang, P., Yen, C.-M., Chow, K.-P. N. and Wang, L.-C. (2014a). A transcriptomic analysis on gene expressions in the infective third and pathogenic fifth larval stages of *Angiostrongylus cantonensis*. *Parasitology International* **63**, 42–48.
- Chang, S. H., Chen, K. Y. and Wang, L. C. (2014b). Identification and characterization of an asparaginyl endopeptidase from *Angiostrongylus cantonensis*. *Parasitology Research* **113**, 2143–2152.
- Chau, T. T., Thwaites, G. E., Chuong, L. V., Sinh, D. X. and Farrar, J. J. (2003). Headache and confusion: the dangers of a raw snail supper. *Lancet* **361**, 1866.
- Chen, C. X., He, H. F., Yin, Z., Zhou, J. H., Li, S. Q., Li, F. H., Chen, J. M., Zhu, W. J., Zhong, X. M., Yang, K. Y., Liu, G. P., Jia, X., Chen, W. T., Li, X. M., Chen, Y. C., Luo, X. D., Chen, D. X. and Shen, H. X. (2012a). Investigation on snails *Achatina fulica* and *Pomacea canaliculata* infected with *Angiostrongylus cantonensis* in Panyu region of Guangzhou City. *Zhongguo Xue Xi Chong Bing Fang Zhi Za Zhi* **24**, 336–338.
- Chen, H. T. (1935). Un nouveau nématode pulmonaire, *Pulmonema cantonensis* n.g., n.sp., des rats de Canton. *Annales de Parasitologie Humaine et Comparée* **13**, 312–317.
- Chen, J., Liu, Q., Yang, X., Wu, X., Zhang, D., He, A. and Zhan, X. (2013). Characterization and immunolocalization of mutated ornithine decarboxylase antizyme from *Angiostrongylus cantonensis*. *Molecular and Biochemical Parasitology* **190**, 76–81.
- Chen, J. X., Chen, M. X., Ai, L., Chen, J. H., Chen, S. H., Zhang, Y. N., Cai, Y. C., Zhu, X. Q. and Zhou, X. N. (2012b). A protein microarray for the rapid screening of patients suspected of infection with various food-borne helminthiases. *PLoS Neglected Tropical Diseases* **6**, e1899.
- Chen, K. M. and Lai, S. C. (2007). Biochemical and pathological evaluation of albendazole/thalidomide co-therapy against eosinophilic meningitis or meningoencephalitis induced by *Angiostrongylus cantonensis*. *Journal of Antimicrobial Chemotherapy* **59**, 264–276.
- Chen, K. M., Liu, J. Y., Lai, S. C., Hsu, L. S. and Lee, H. H. (2006). Association of plasminogen activators and matrix metalloproteinase-9 proteolytic cascade with blood-CNS barrier damage of angiostrongyliasis. *International Journal of Experimental Pathology* **87**, 113–119.
- Chen, K. M., Lee, H. H., Lai, S. C., Hsu, L. S., Wang, C. J. and Liu, J. Y. (2008). Apoptosis in meningoencephalitis of *Angiostrongylus cantonensis*-infected mice. *Experimental Parasitology* **119**, 385–390.
- Chen, K. Y., Cheng, C. J., Yen, C. M., Tang, P. and Wang, L. C. (2014a). Comparative studies on the proteomic expression patterns in the

- third- and fifth-stage larvae of *Angiostrongylus cantonensis*. *Parasitology Research* **113**, 3591–3600.
- Chen, M. X., Ai, L., Xu, M. J., Zhang, R. L., Chen, S. H., Zhang, Y. N., Guo, J., Cai, Y. C., Tian, L. G., Zhang, L. L., Zhu, X. Q. and Chen, J. X. (2011a). *Angiostrongylus cantonensis*: identification and characterization of microRNAs in male and female adults. *Experimental Parasitology* **128**, 116–120.
- Chen, M. X., Wang, K., Ai, L., Yan, W. H., Peng, L. and Zhang, R. L. (2011b). Development of a double antibody sandwich ELISA assay for the diagnosis of angiostrongyliasis. *Journal of Parasitology* **97**, 721–724.
- Chen, M. X., Zhang, R. L., Ai, L., Chen, J. X., Chen, S. H., Huang, D. N., Gao, S. T., Geng, Y. J., Li, X. H. and Zhu, X. Q. (2011c). Seroprevalence of *Angiostrongylus cantonensis* infection in humans in China. *Journal of Parasitology* **97**, 144–145.
- Chen, R., Tong, Q., Zhang, Y., Lou, D., Kong, Q., Lv, S., Zhuo, M., Wen, L. and Lu, S. (2011d). Loop-mediated isothermal amplification: rapid detection of *Angiostrongylus cantonensis* infection in *Pomacea canaliculata*. *Parasites and Vectors* **4**, 204.
- Chen, X., Li, Z. Y., Maleewong, W., Maleewong, P., Liang, J., Zeng, X., Zheng, H., Wu, Z. D. and Sun, X. (2014b). Serum acamir-146a is a potential biomarker for early diagnosis of *Angiostrongylus cantonensis* infection. *Parasitology Research* **113**, 3221–3227.
- Chen, X. G., Li, H. and Lun, Z. R. (2005). Angiostrongyliasis, Mainland China. *Emerging Infectious Diseases* **11**, 1645–1647.
- Cheng, M., Yang, X., Li, Z., He, H., Qu, Z., He, A., Wu, Z. and Zhan, X. (2012). Cloning and characterization of a novel cathepsin B-like cysteine proteinase from *Angiostrongylus cantonensis*. *Parasitology Research* **110**, 2413–2422.
- Cheng, T. and Alicata, J. (1964). Possible role of water in the transmission of *Angiostrongylus cantonensis* (Nematoda: Metastrongylidae). *Journal of Parasitology* **50**, 39–40.
- Chi, M., Kim, H. J., Haug, S., Vagefi, R. and Kersten, R. C. (2014). Acute eosinophilic meningitis and orbital inflammation from presumed angiostrongyliasis. *Canadian Journal of Ophthalmology* **49**, e65–e67.
- Chikweto, A., Bhaiyat, M. I., Macpherson, C. N., Deallie, C., Pinckney, R. D., Richards, C. and Sharma, R. N. (2009). Existence of *Angiostrongylus cantonensis* in rats (*Rattus norvegicus*) in Grenada, West Indies. *Veterinary Parasitology* **162**, 160–162.
- Chiu, P. S. and Lai, S. C. (2013). Matrix metalloproteinase-9 leads to claudin-5 degradation via the NF- κ B pathway in BALB/c mice with eosinophilic meningoencephalitis caused by *Angiostrongylus cantonensis*. *PLoS ONE* **8**, e53370.
- Chiu, P. S. and Lai, S. C. (2014). Matrix metalloproteinase-9 leads to blood-brain barrier leakage in mice with eosinophilic meningoencephalitis caused by *Angiostrongylus cantonensis*. *Acta Tropica* **140**, 141–150.
- Chotmongkol, V., Sawanyawisuth, K. and Thavornpitak, Y. (2000). Corticosteroid treatment of eosinophilic meningitis. *Clinical Infectious Diseases* **31**, 660–662.
- Chotmongkol, V., Sawadpanitch, K., Sawanyawisuth, K., Louhawilai, S. and Limpawattana, P. (2006). Treatment of eosinophilic meningitis with a combination of prednisolone and mebendazole. *American Journal of Tropical Medicine and Hygiene* **74**, 1122–1124.
- Chotmongkol, V., Kittimongkolma, S., Niwattayakul, K., Intapan, P. M. and Thavornpitak, Y. (2009). Comparison of prednisolone plus albendazole with prednisolone alone for treatment of patients with eosinophilic meningitis. *American Journal of Tropical Medicine and Hygiene* **81**, 443–445.
- Chuang, C. C., Chen, C. W., Huang, Y. T. and Du, W. Y. (2016). Anti-ST2 monoclonal antibody inhibits eosinophil infiltration in *Angiostrongylus cantonensis*-infected mice. *Journal of Microbiology, Immunology, and Infection* **49**, 91–96.
- Chuan-Min, Y. and Eng-Rin, C. (1991). Detection of antibodies to *Angiostrongylus cantonensis* in serum and cerebrospinal fluid of patients with eosinophilic meningitis. *International Journal for Parasitology* **21**, 17–21.
- Chye, S. M., Chang, J. H. and Yen, C. M. (2000). Immunodiagnosis of human eosinophilic meningitis using an antigen of *Angiostrongylus cantonensis* L5 with molecular weight 204 kD. *Acta Tropica* **75**, 9–17.
- Chye, S. M., Lin, S. R., Chen, Y. L., Chung, L. Y. and Yen, C. M. (2004). Immuno-PCR for detection of antigen to *Angiostrongylus cantonensis* circulating fifth-stage worms. *Clinical Chemistry* **50**, 51–57.
- Civeyrel, L. and Simberloff, D. (1996). A tale of two snails: is the cure worse than the disease? *Biodiversity and Conservation* **5**, 1231–1252.
- Constantino-Santos, M. A., Basiao, Z. U., Wade, C. M., Santos, B. S. and Fontanilla, I. K. (2014). Identification of *Angiostrongylus cantonensis* and other nematodes using the SSU rDNA in *Achatina fulica* populations of Metro Manila. *Tropical Biomedicine* **31**, 327–335.
- Cowie, R. H. (2013a). Biology, systematics, life cycle, and distribution of *Angiostrongylus cantonensis*, the cause of rat lungworm disease. *Hawai'i Journal of Medicine and Public Health* **72**, 6–9.
- Cowie, R. H. (2013b). Pathways for transmission of angiostrongyliasis and the risk of disease associated with them. *Hawai'i Journal of Medicine and Public Health* **72**, 70–74.
- Crane, A. M., Weiss, M. and Galor, A. (2013). Anterior chamber angiostrongyliasis. *JAMA Ophthalmology* **131**, 951–952.
- da Silva, A. C., Graeff-Teixeira, C. and Zaha, A. (2003). Diagnosis of abdominal angiostrongyliasis by PCR from sera of patients. *Revista do Instituto de Medicina Tropical de Sao Paulo* **45**, 295–297.
- Deng, Z. H., Zhang, Q. M., Huang, S. Y. and Jones, J. L. (2012). First provincial survey of *Angiostrongylus cantonensis* in Guangdong Province, China. *Tropical Medicine and International Health* **17**, 119–122.
- Deo, S. S., Mistry, K. J., Kakade, A. M. and Niphadkar, P. V. (2010). Role played by Th2 type cytokines in IgE mediated allergy and asthma. *Lung India* **27**, 66–71.
- Diao, Z., Chen, X., Yin, C., Wang, J., Qi, H. and Ji, A. (2009). *Angiostrongylus cantonensis*: effect of combination therapy with albendazole and dexamethasone on Th cytokine gene expression in PBMC from patients with eosinophilic meningitis. *Experimental Parasitology* **123**, 1–5.
- Diao, Z., Jin, E. and Yin, C. (2010). *Angiostrongylus cantonensis*: lesions in brain and spinal cord. *American Journal of Tropical Medicine and Hygiene* **82**, 519.
- Diao, Z., Wang, J., Qi, H., Li, X., Zheng, X. and Yin, C. (2011). Treatment of angiostrongyliasis using a combination of albendazole and dexamethasone: the results of a retrospective and comparative study. *Annals of Tropical Medicine and Parasitology* **105**, 65–69.
- Diaz, J. H. (2009). Recognizing and reducing the risks of helminthic eosinophilic meningitis in travelers: differential diagnosis, disease management, prevention, and control. *Journal of Travel Medicine* **16**, 267–275.
- Dorta-Contreras, A. J., Noris-Garcia, E., Escobar-Perez, X. and Padilla Docal, B. (2005). IgG₁, IgG₂ and IgE intrathecal synthesis in *Angiostrongylus cantonensis* meningoencephalitis. *Journal of the Neurological Sciences* **238**, 65–70.
- Dorta-Contreras, A. J., Magraner-Tarrau, M. E. and Sanchez-Zulueta, E. (2009). Angiostrongyliasis in the Americas. *Emerging Infectious Diseases* **15**, 991.
- Dorta-Contreras, A. J., Padilla-Docal, B., Moreira, J. M., Robles, L. M., Arco, J. M., Alarcon, F. and Bu-Coifu-Fanego, R. (2011). Neuroimmunological findings of *Angiostrongylus cantonensis* meningitis in Ecuadorian patients. *Arquivos de Neuro-Psiquiatria* **69**, 466–469.
- Dougherty, E. C. (1946). The genus *Aelurostrongylus* Cameron, 1927 (Nematoda: Metastrongylidae), and its relatives; with descriptions of *Parafilaroides*, gen. nov., and *Angiostrongylus gubernaculatus*, sp. nov. *Proceedings of the Helminthological Society of Washington* **13**, 16–26.
- Du, W. Y., Liao, J. W., Fan, C. K. and Su, K. E. (2003). Combined treatment with interleukin-12 and mebendazole lessens the severity of experimental eosinophilic meningitis caused by *Angiostrongylus cantonensis* in ICR mice. *Infection and Immunity* **71**, 3947–3953.
- Du, W. Y., Chen, C. W., Lin, F. K. and Chuang, C. C. (2013). IL-33 mediates the expressions of IL-5 and IL-13 in *Angiostrongylus cantonensis*-infected mice. *Experimental Parasitology* **135**, 587–594.
- Duffy, M. S., Miller, C. L., Kinsella, J. M. and de Lahunta, A. (2004). *Parastrostrongylus cantonensis* in a nonhuman primate, Florida. *Emerging Infectious Diseases* **10**, 2207–2210.
- Dusitsittipon, S., Thaenkham, U., Watthanakulpanich, D., Adisakwattana, P. and Komalamisra, C. (2015). Genetic differences in the rat lungworm, *Angiostrongylus cantonensis* (Nematoda: Angiostrongylidae), in Thailand. *Journal of Helminthology* **89**, 545–551.
- Eamsobhana, P. (2013). Angiostrongyliasis in Thailand: epidemiology and laboratory investigations. *Hawai'i Journal of Medicine and Public Health* **72**, 28–32.
- Eamsobhana, P. (2014). Eosinophilic meningitis caused by *Angiostrongylus cantonensis* – a neglected disease with escalating importance. *Tropical Biomedicine* **31**, 569–578.
- Eamsobhana, P. and Yong, H. S. (2009). Immunological diagnosis of human angiostrongyliasis due to *Angiostrongylus cantonensis* (Nematoda: Angiostrongylidae). *International Journal of Infectious Diseases* **13**, 425–431.
- Eamsobhana, P., Yoolek, A. and Kreethapon, N. (2003). Blinded multi-laboratory evaluation of an in-house dot-blot ELISA kit for diagnosis of human parastrongyliasis. *Southeast Asian Journal of Tropical Medicine and Public Health* **34**, 1–6.
- Eamsobhana, P., Yoolek, A., Punthuprapasa, P. and Suvoutho, S. (2004). A dot-blot ELISA comparable to immunoblot for the specific diagnosis of human parastrongyliasis. *Journal of Helminthology* **78**, 287–291.
- Eamsobhana, P., Ongrotchanakun, J., Yoolek, A., Punthuprapasa, P., Monkong, N. and Dekumyoy, P. (2006). Multi-immunodot for rapid differential diagnosis of eosinophilic meningitis due to parasitic infections. *Journal of Helminthology* **80**, 249–254.

- Eamsobhana, P., Yoolek, A. and Yong, H. S. (2009). Meiotic chromosomes and sex determination mechanism in Thailand and Hawaii isolates of *Angiostrongylus cantonensis* (Nematoda: Angiostrongylidae). *Tropical Biomedicine* **26**, 346–351.
- Eamsobhana, P., Yoolek, A. and Yong, H. S. (2010). Effect of Thai 'koi-hoi' food flavoring on the viability and infectivity of the third-stage larvae of *Angiostrongylus cantonensis* (Nematoda: Angiostrongylidae). *Acta Tropica* **113**, 245–247.
- Eamsobhana, P., Lim, P. E. and Yong, H. S. (2013a). Genetic diversity of the rat lungworm, *Angiostrongylus cantonensis*, the major cause of eosinophilic meningitis. *Hawai'i Journal of Medicine and Public Health* **72**, 15–17.
- Eamsobhana, P., Wanachiwanawin, D., Dechkum, N., Parsartvit, A. and Yong, H. S. (2013b). Molecular diagnosis of eosinophilic meningitis due to *Angiostrongylus cantonensis* (Nematoda: Metastrongyloidea) by polymerase chain reaction-DNA sequencing of cerebrospinal fluids of patients. *Memorias do Instituto Oswaldo Cruz* **108**, 116–118.
- Eamsobhana, P., Gan, X. X., Ma, A., Wang, Y., Wanachiwanawin, D. and Yong, H. S. (2014). Dot immunogold filtration assay (DIGFA) for the rapid detection of specific antibodies against the rat lungworm *Angiostrongylus cantonensis* (Nematoda: Metastrongyloidea) using purified 31-kDa antigen. *Journal of Helminthology* **88**, 396–401.
- Eamsobhana, P., Lim, P. E. and Yong, H. S. (2015a). Phylogenetics and systematics of *Angiostrongylus* lungworms and related taxa (Nematoda: Metastrongyloidea) inferred from the nuclear small subunit (SSU) ribosomal DNA sequences. *Journal of Helminthology* **89**, 317–325.
- Eamsobhana, P., Prasartvit, A., Gan, X. X. and Yong, H. S. (2015b). Evaluation of dot immunogold filtration assay (DIGFA) for rapid serodiagnosis of eosinophilic meningitis due to *Angiostrongylus cantonensis* (Nematoda: Metastrongyloidea). *Tropical Biomedicine* **32**, 121–125.
- Emerson, J. A., Walden, H. S., Peters, R. K., Farina, L. L., Fredholm, D. V., Qvarnstrom, Y., Xayavong, M., Bishop, H., Slapcinsky, J., McIntosh, A. and Wellehan, J. F., Jr. (2013). Eosinophilic meningoencephalomyelitis in an orangutan (*Pongo pygmaeus*) caused by *Angiostrongylus cantonensis*. *Veterinary Quarterly* **33**, 191–194.
- Evans-Gilbert, T., Lindo, J. F., Henry, S., Brown, P. and Christie, C. D. (2014). Severe eosinophilic meningitis owing to *Angiostrongylus cantonensis* in young Jamaican children: case report and literature review. *Paediatrics and International Child Health* **34**, 148–152.
- Fang, W., Wang, J., Liu, J., Xu, C., Cai, W. and Luo, D. (2012). PCR assay for the cell-free copro-DNA detection of *Angiostrongylus cantonensis* in rat faeces. *Veterinary Parasitology* **183**, 299–304.
- Feng, F., Feng, Y., Liu, Z., Li, W. H., Wang, W. C., Wu, Z. D. and Lv, Z. (2015). Effects of albendazole combined with TSII-A (a Chinese herb compound) on optic neuritis caused by *Angiostrongylus cantonensis* in BALB/c mice. *Parasites and Vectors* **8**, 606.
- Feng, Y., Nawa, Y., Sawanyavisuth, K., Lv, Z. and Wu, Z. D. (2013). Comprehensive review of ocular angiostrongyliasis with special reference to optic neuritis. *Korean Journal of Parasitology* **51**, 613–619.
- Feng, Y., Zeng, X., Li, W. H., Wang, W. C., Chen, W., Ou-Yang, L. S., Sun, X., Feng, F. and Wu, Z. D. (2014). The pathogenesis of optic neuritis caused by *Angiostrongylus cantonensis* in BALB/c mice. *Parasites and Vectors* **7**, 339.
- Ferri, E., Bain, O., Barbuto, M., Martin, C., Lo, N., Uni, S., Landmann, F., Baccei, S. G., Guerrero, R., de Souza Lima, S., Bandi, C., Wanji, S., Diagne, M. and Casiraghi, M. (2011). New insights into the evolution of *Wolbachia* infections in filarial nematodes inferred from a large range of screened species. *PLoS ONE* **6**, e20843.
- Fontanilla, I. K. and Wade, C. M. (2012). First report of *Angiostrongylus cantonensis* in the giant African land snail *Achatina fulica* in French Polynesia detected using the SSU rRNA gene. *Tropical Biomedicine* **29**, 642–645.
- Foronda, P., Lopez-Gonzalez, M., Miquel, J., Torres, J., Segovia, M., Abreu-Acosta, N., Casanova, J. C., Valladares, B., Mas-Coma, S., Bargues, M. D. and Feliu, C. (2010). Finding of *Parastrongylus cantonensis* (Chen, 1935) in *Rattus rattus* in Tenerife, Canary Islands (Spain). *Acta Tropica* **114**, 123–127.
- Foster, J., Ganatra, M., Kamal, I., Ware, J., Makarova, K., Ivanova, N., Bhattacharyya, A., Kapatral, V., Kumar, S., Posfai, J., Vincze, T., Ingram, J., Moran, L., Lapidus, A., Omelchenko, M., Kyrpides, N., Ghedin, E., Wang, S., Goltsman, E., Joukov, V., Ostrovskaya, O., Tsukerman, K., Mazur, M., Comb, D., Koonin, E. and Slatko, B. (2005). The *Wolbachia* genome of *Brugia malayi*: endosymbiont evolution within a human pathogenic nematode. *PLoS Biology* **3**, e121.
- Foster, J. M., Kumar, S., Ford, L., Johnston, K. L., Ben, R., Graeff-Teixeira, C. and Taylor, M. J. (2008). Absence of *Wolbachia* endobacteria in the non-filariid nematodes *Angiostrongylus cantonensis* and *A. costaricensis*. *Parasites and Vectors* **1**, 31.
- Furugen, M., Yamashiro, S., Tamayose, M., Naha, Y., Miyagi, K., Nakasone, C., Uchihara, T., Haranaga, S., Azuma, M., Yara, S., Shinzato, T., Higa, F., Toma, H., Tateyama, M. and Fujita, J. (2006). Elsberg syndrome with eosinophilic meningoencephalitis caused by *Angiostrongylus cantonensis*. *Internal Medicine* **45**, 1333–1336.
- Galor, A. and Eberhard, M. L. (2014). Follow-up on anterior chamber angiostrongyliasis. *JAMA Ophthalmology* **132**, 1029–1030.
- Gasser, R. B., Jabbar, A., Mohandas, N., Schnyder, M., Deplazes, P., Littlewood, D. T. and Jex, A. R. (2012). Mitochondrial genome of *Angiostrongylus vasorum*: comparison with congeners and implications for studying the population genetics and epidemiology of this parasite. *Infection, Genetics and Evolution* **12**, 1884–1891.
- Gebreselassie, N. G., Moorhead, A. R., Fabre, V., Gagliardo, L. F., Lee, N. A., Lee, J. J. and Appleton, J. A. (2012). Eosinophils preserve parasitic nematode larvae by regulating local immunity. *Journal of Immunology* **188**, 417–425.
- Giannelli, A., Colella, V., Abramo, F., do Nascimento Ramos, R. A., Falson, L., Brianti, E., Varcasia, A., Dantas-Torres, F., Knaus, M., Fox, M. T. and Otranto, D. (2015). Release of lungworm larvae from snails in the environment: potential for alternative transmission pathways. *PLoS Neglected Tropical Diseases* **9**, e0003722.
- Gosnell, W. L. and Kramer, K. J. (2013). The role of eosinophils in angiostrongyliasis: multiple roles for a versatile cell? *Hawai'i Journal of Medicine and Public Health* **72**, 49–51.
- Graeff-Teixeira, C., de Avila-Pires, F. D., Machado Rde, C., Camillo-Coura, L. and Lenzi, H. L. (1990). Identification of wild rodents as hosts of *Angiostrongylus costaricensis* in southern Brazil. *Revista do Instituto de Medicina Tropical de Sao Paulo* **32**, 147–150.
- Graeff-Teixeira, C., da Silva, A. C. and Yoshimura, K. (2009). Update on eosinophilic meningoencephalitis and its clinical relevance. *Clinical Microbiology Reviews* **22**, 322–348.
- Gutteridge, B., Bhaibulaya, M. and Findlater, C. (1972). Human larval meningitis possibly following lettuce ingestion in Brisbane. *Pathology* **4**, 63–64.
- Han, Y. P., Li, Z. Y., Li, B. C., Sun, X., Zhu, C. C., Ling, X. T., Zheng, H. Q., Wu, Z. D. and Lv, Z. Y. (2011). Molecular cloning and characterization of a cathepsin B from *Angiostrongylus cantonensis*. *Parasitology Research* **109**, 369–378.
- Harinasuta, C. (1983). Parasitic diseases in the South (Developing World). In *Parasitology: A Global Perspective* (ed. Warren, K. S. and Bowers, J. Z.), pp. 19–44. Springer-Verlag New York Inc., New York, USA.
- Hata, T. Y., Hara, A. H. and Hu, B. K.-S. (1997). Molluscicides and mechanical barriers against slugs, *Vaginula plebeia* Fischer and *Veronicella cubensis* (Pfeiffer) (Stylommatophora: Veronicellidae). *Crop Protection* **16**, 501–506.
- He, H., Cheng, M., Yang, X., Meng, J., He, A., Zheng, X., Li, Z., Guo, P., Pan, Z. and Zhan, X. (2009). Preliminary molecular characterization of the human pathogen *Angiostrongylus cantonensis*. *BMC Molecular Biology* **10**, 97.
- Hidelaratchi, M. D., Riffsy, M. T. and Wijesekera, J. C. (2005). A case of eosinophilic meningitis following monitor lizard meat consumption, exacerbated by anthelmintics. *Ceylon Medical Journal* **50**, 84–86.
- Higgins, D. P., Carlisle-Nowak, M. S. and Mackie, J. (1997). Neural angiostrongyliosis in three captive rufous bettongs (*Aepyprymnus rufescens*). *Australian Veterinary Journal* **75**, 564–566.
- Hochberg, N. S., Park, S. Y., Blackburn, B. G., Sejvar, J. J., Gaynor, K., Chung, H., Leniek, K., Herwaldt, B. L. and Effler, P. V. (2007). Distribution of eosinophilic meningitis cases attributable to *Angiostrongylus cantonensis*, Hawaii. *Emerging Infectious Diseases* **13**, 1675–1680.
- Hochberg, N. S., Blackburn, B. G., Park, S. Y., Sejvar, J. J., Effler, P. V. and Herwaldt, B. L. (2011). Eosinophilic meningitis attributable to *Angiostrongylus cantonensis* infection in Hawaii: clinical characteristics and potential exposures. *American Journal of Tropical Medicine and Hygiene* **85**, 685–690.
- Hollingsworth, R. G., Howe, K. and Jarvi, S. I. (2013). Control measures for slug and snail hosts of *Angiostrongylus cantonensis*, with special reference to the semi-slug *Parmarion martensi*. *Hawai'i Journal of Medicine and Public Health* **72**, 75–80.
- Hou, R. F., Tu, W. C., Lee, H. H., Chen, K. M., Chou, H. L. and Lai, S. C. (2004). Elevation of plasminogen activators in cerebrospinal fluid of mice with eosinophilic meningitis caused by *Angiostrongylus cantonensis*. *International Journal for Parasitology* **34**, 1355–1364.
- Howe, K. (2013). A severe case of rat lungworm disease in Hawaii. *Hawai'i Journal of Medicine and Public Health* **72**, 46–48.
- Hsu, C. C. and Lai, S. C. (2007). Matrix metalloproteinase-2, -9 and -13 are involved in fibronectin degradation of rat lung granulomatous fibrosis

- caused by *Angiostrongylus cantonensis*. *International Journal of Experimental Pathology* **88**, 437–443.
- Hsu, J. J., Chung, S. H., Chen, C. H. and Huang, M. H. (2009). Sacral myeloradiculitis (Elsberg syndrome) secondary to eosinophilic meningitis caused by *Angiostrongylus cantonensis*. *BMJ Case Reports* **2009**. doi: 10.1136/bcr.10.2008.1075.
- Hu, X., Du, J., Tong, C., Wang, S., Liu, J., Li, Y. and He, C. (2011). Epidemic status of *Angiostrongylus cantonensis* in Hainan Island, China. *Asian Pacific Journal of Tropical Medicine* **4**, 275–277.
- Huang, D. N., Chen, M. X., Geng, Y. J., Li, X. H., Gao, S. T. and Zhang, R. L. (2010). Detection of circulating antigen of *Angiostrongylus cantonensis* by 12D5 and 21B7 monoclonal antibodies. *Zhonghua Liu Xing Bing Xue Za Zhi* **31**, 79–82.
- Huang, H. C., Yao, L. L., Song, Z. M., Li, X. P., Hua, Q. Q., Li, Q., Pan, C. W. and Xia, C. M. (2013). Development-specific differences in the proteomics of *Angiostrongylus cantonensis*. *PLoS ONE* **8**, e76982.
- Hwang, K. P. and Chen, E. R. (1991). Clinical studies on angiostrongyliasis cantonensis among children in Taiwan. *Southeast Asian Journal of Tropical Medicine and Public Health* **22**(Suppl.), 194–199.
- Hwang, K. P., Chang, S. H. and Wang, L. C. (2010). Alterations in the expression level of a putative aspartic protease in the development of *Angiostrongylus cantonensis*. *Acta Tropica* **113**, 289–294.
- Ibrahim, M. M. (2007). Prevalence and intensity of *Angiostrongylus cantonensis* in freshwater snails in relation to some ecological and biological factors. *Parasite* **14**, 61–70.
- Incani, R. N., Caleiras, E., Martin, M. and Gonzalez, C. (2007). Human infection by *Angiostrongylus costaricensis* in Venezuela: first report of a confirmed case. *Revista do Instituto de Medicina Tropical de Sao Paulo* **49**, 197–200.
- Intapan, P. M., Maleewong, W., Polsan, Y., Sawanyawisuth, K. and Chotmongkol, V. (2002). Specific IgG antibody subclasses to *Angiostrongylus cantonensis* in patients with angiostrongyliasis. *Asian Pacific Journal of Allergy and Immunology* **20**, 235–240.
- Intapan, P. M., Maleewong, W., Sawanyawisuth, K. and Chotmongkol, V. (2003). Evaluation of human IgG subclass antibodies in the serodiagnosis of angiostrongyliasis. *Parasitology Research* **89**, 425–429.
- Intapan, P. M., Niwattayakul, K., Sawanyawisuth, K., Chotmongkol, V. and Maleewong, W. (2007). Cerebrospinal fluid eotaxin and eotaxin-2 levels in human eosinophilic meningitis associated with angiostrongyliasis. *Cytokine* **39**, 138–141.
- Intapan, P. M., Kittimongkolma, S., Niwattayakul, K., Sawanyawisuth, K. and Maleewong, W. (2008). Cerebrospinal fluid cytokine responses in human eosinophilic meningitis associated with angiostrongyliasis. *Journal of the Neurological Sciences* **267**, 17–21.
- Iwanowicz, D. D., Sanders, L. R., Schill, W. B., Xayavong, M. V., da Silva, A. J., Qvarnstrom, Y. and Smith, T. (2015). Spread of the rat lungworm (*Angiostrongylus cantonensis*) in giant African land snails (*Lissachatina fulica*) in Florida, USA. *Journal of Wildlife Diseases* **51**, 749–753.
- Jarvi, S. I., Farias, M. E., Howe, K., Jacquier, S., Hollingsworth, R. and Pitt, W. (2012). Quantitative PCR estimates *Angiostrongylus cantonensis* (rat lungworm) infection levels in semi-slugs (*Parmarion martensi*). *Molecular and Biochemical Parasitology* **185**, 174–176.
- Jarvi, S. I., Pitt, W. C., Farias, M. E., Shiels, L., Severino, M. G., Howe, K. M., Jacquier, S. H., Shiels, A. B., Amano, K. K., Luiz, B. C., Maher, D. E., Allison, M. L., Holtquist, Z. C. and Scheibelhut, N. T. (2015). Detection of *Angiostrongylus cantonensis* in the blood and peripheral tissues of wild hawaiian rats (*Rattus rattus*) by a quantitative PCR (qPCR) Assay. *PLoS ONE* **10**, e0123064.
- Ji, P., Hu, H., Yang, X., Wei, X., Zhu, C., Liu, J., Feng, Y., Yang, F., Okanurak, K., Li, N., Zeng, X., Zheng, H., Wu, Z. and Lv, Z. (2015). AcCystatin, an immunoregulatory molecule from *Angiostrongylus cantonensis*, ameliorates the asthmatic response in an aluminium hydroxide/ovalbumin-induced rat model of asthma. *Parasitology Research* **114**, 613–624.
- Jilkova, A., Rezacova, P., Lepsik, M., Horn, M., Vachova, J., Fanfrik, J., Brynda, J., McKerrow, J. H., Caffrey, C. R. and Mares, M. (2011). Structural basis for inhibition of cathepsin B drug target from the human blood fluke, *Schistosoma mansoni*. *Journal of Biological Chemistry* **286**, 35770–35781.
- Jin, E., Ma, D., Liang, Y., Ji, A. and Gan, S. (2005). MRI findings of eosinophilic myelomeningoencephalitis due to *Angiostrongylus cantonensis*. *Clinical Radiology* **60**, 242–250.
- Jin, E. H., Ma, Q., Ma, D. Q., He, W., Ji, A. P. and Yin, C. H. (2008). Magnetic resonance imaging of eosinophilic meningoencephalitis caused by *Angiostrongylus cantonensis* following eating freshwater snails. *Chinese Medical Journal* **121**, 67–72.
- Jitpimolmard, S., Sawanyawisuth, K., Morakote, N., Vejajiva, A., Puntumetakul, M., Sanchaisuriya, K., Tassaneeyakul, W., Tassaneeyakul, W. and Korwanich, N. (2007). Albendazole therapy for eosinophilic meningitis caused by *Angiostrongylus cantonensis*. *Parasitology Research* **100**, 1293–1296.
- Kampkotter, A., Volkmann, T. E., de Castro, S. H., Leiers, B., Klotz, L. O., Johnson, T. E., Link, C. D. and Henkle-Duhrsen, K. (2003). Functional analysis of the glutathione S-transferase 3 from *Onchocerca volvulus* (Ov-GST-3): a parasite GST confers increased resistance to oxidative stress in *Caenorhabditis elegans*. *Journal of Molecular Biology* **325**, 25–37.
- Kanpittaya, J., Sawanyawisuth, K., Intapan, P. M., Khotsri, P., Chotmongkol, V. and Maleewong, W. (2012). A comparative study of neuroimaging features between human neuro-gnathostomiasis and angiostrongyliasis. *Neurological Sciences* **33**, 893–898.
- Ke, Y. K., Huang, Y., Ge, X. Q., Peng, M. X., Wang, Q., Huang, L. R., Luo, G. Q., Zhong, L. X., Kan, S. F., Zhang, Y. X. and Zhai, J. X. (2014). Detection of *Angiostrongylus cantonensis* larvae with a high resolution melt PCR assay. *Zhongguo Ji Sheng Chong Xue Yu Ji Sheng Chong Bing Za Zhi* **32**, 452–454.
- Kim, J. R., Hayes, K. A., Yeung, N. W. and Cowie, R. H. (2014). Diverse gastropod hosts of *Angiostrongylus cantonensis*, the rat lungworm, globally and with a focus on the Hawaiian Islands. *PLoS ONE* **9**, e94969.
- Kittimongkolma, S., Intapan, P. M., Laemviteevanich, K., Kanpittaya, J., Sawanyawisuth, K. and Maleewong, W. (2007). Eosinophilic meningitis associated with angiostrongyliasis: clinical features, laboratory investigations and specific diagnostic IgG and IgG subclass antibodies in cerebrospinal fluid. *Southeast Asian Journal of Tropical Medicine and Public Health* **38**, 24–31.
- Kliks, M. M. and Palumbo, N. E. (1992). Eosinophilic meningitis beyond the Pacific Basin: the global dispersal of a peridomestic zoonosis caused by *Angiostrongylus cantonensis*, the nematode lungworm of rats. *Social Science and Medicine* **34**, 199–212.
- Klion, A. D. and Nutman, T. B. (2004). The role of eosinophils in host defense against helminth parasites. *Journal of Allergy and Clinical Immunology* **113**, 30–37.
- Kottwitz, J. J., Perry, K. K., Rose, H. H. and Hendrix, C. M. (2014). *Angiostrongylus cantonensis* infection in captive Geoffroy's tamarins (*Saguinus geoffroyi*). *Journal of the American Veterinary Medical Association* **245**, 821–827.
- Kumar, V., Kyprianou, I. and Keenan, J. M. (2005). Ocular *Angiostrongylus cantonensis*: removal of a live nematode from the anterior chamber. *Eye* **19**, 229–230.
- Kwon, E., Ferguson, T. M., Park, S. Y., Manuzak, A., Qvarnstrom, Y., Morgan, S., Ciminera, P. and Murphy, G. S. (2013). A severe case of *Angiostrongylus cantonensis* infection with encephalitis and neurologic sequelae in Hawai'i. *Hawaii Journal of Medicine and Public Health* **72**, 41–45.
- Lai, S. C. (2006). Chinese herbal medicine Yin-Chen-Extract as an adjunct to anthelmintic albendazole used against *Angiostrongylus cantonensis*-induced eosinophilic meningitis or meningoencephalitis. *American Journal of Tropical Medicine and Hygiene* **75**, 556–562.
- Lai, S. C., Twu, J. J., Jiang, S. T., Hsu, J. D., Chen, K. M., Chiaing, H. C., Wang, C. J., Tseng, C. K., Shyu, L. Y. and Lee, H. H. (2004). Induction of matrix metalloproteinase-9 in murine eosinophilic meningitis caused by *Angiostrongylus cantonensis*. *Annals of Tropical Medicine and Parasitology* **98**, 715–724.
- Lai, S. C., Jiang, S. T., Chen, K. M., Hsu, J. D., Shyu, L. Y. and Lee, H. H. (2005a). Efficacy of albendazole-GM6001 co-therapy against *Angiostrongylus cantonensis*-induced meningitis in BALB/c mice. *Acta Tropica* **93**, 267–273.
- Lai, S. C., Jiang, S. T., Chen, K. M. and Lee, H. H. (2005b). Matrix metalloproteinases activity demonstrated in the infective stage of the nematodes, *Angiostrongylus cantonensis*. *Parasitology Research* **97**, 466–471.
- Lammers, A. J., Goorhuis, A., van de Beek, D., Grobusch, M. P., Bart, A., van Gool, T. and van Vugt, M. (2015). Eosinophilia a deus: a brain nagging souvenir from the Philippines. *Infection* **43**, 615–617.
- Lan, K. P., Wang, C. J., Lai, S. C., Chen, K. M., Lee, S. S., Hsu, J. D. and Lee, H. H. (2004). The efficacy of therapy with albendazole in mice with parasitic meningitis caused by *Angiostrongylus cantonensis*. *Parasitology Research* **93**, 311–317.
- Lee, J. D. and Yen, C. M. (2005). Protease secreted by the infective larvae of *Angiostrongylus cantonensis* and its role in the penetration of mouse intestine. *American Journal of Tropical Medicine and Hygiene* **72**, 831–836.
- Lee, J. D., Chung, L. Y., Wang, L. C., Lin, R. J., Wang, J. J., Tu, H. P., Wu, Z. D. and Yen, C. M. (2014). Sequence analysis in partial genes of five isolates of *Angiostrongylus cantonensis* from Taiwan and biological comparison in infectivity and pathogenicity between two strains. *Acta Tropica* **133**, 26–34.
- Leon, I. R., Neves-Ferreira, A. G., Valente, R. H., Mota, E. M., Lenzi, H. L. and Perales, J. (2007). Improved protein identification

- efficiency by mass spectrometry using N-terminal chemical derivatization of peptides from *Angiostrongylus costaricensis*, a nematode with unknown genome. *Journal of Mass Spectrometry* **42**, 781–792.
- Leone, S., De Marco, M., Ghirga, P., Nicastrì, E., Esposito, M. and Narciso, P. (2007). Eosinophilic meningitis in a returned traveler from Santo Domingo: case report and review. *Journal of Travel Medicine* **14**, 407–410.
- Levett, P. N., Douglas, K. O., Waugh, C. A., Robinson, R. D. and Lindo, J. F. (2004). Failure to detect *Angiostrongylus cantonensis* in rats in Barbados. *West Indian Medical Journal* **53**, 58.
- Li, D. N., He, A., Wang, Y., Liang, Y., Li, Z. Y., Men, J. X. and Zhan, X. M. (2001). Three lethal cases of *Angiostrongylus cantonensis* infected children. *Zhongguo Ji Sheng Chong Xue Yu Ji Sheng Chong Bing Za Zhi* **19**, 310–311.
- Li, H., Xu, F., Gu, J. B. and Chen, X. G. (2008). A severe eosinophilic meningoencephalitis caused by infection of *Angiostrongylus cantonensis*. *American Journal of Tropical Medicine and Hygiene* **79**, 568–570.
- Li, L. S., Zhang, R. Y., Fang, Y. Y., Ouyang, R., Xie, H. G., Jiang, D. W., Xie, X. L., Chen, Z. Y. and Zheng, G. B. (2013a). Prevalence of *Angiostrongylus cantonensis* infection in snails for sale in Fuzhou and Xiamen. *Zhongguo Ji Sheng Chong Xue Yu Ji Sheng Chong Bing Za Zhi* **31**, 464–466.
- Li, S., Yang, F., Ji, P., Zeng, X., Wu, X., Wei, J., Ouyang, L., Liang, J., Zheng, H., Wu, Z. and Lv, Z. (2014a). Eosinophil chemotactic chemokine profilings of the brain from permissive and non-permissive hosts infected with *Angiostrongylus cantonensis*. *Parasitology Research* **113**, 517–525.
- Li, Y., Tang, J. P., Chen, D. R., Fu, C. Y., Wang, P., Li, Z., Wei, W., Li, H. and Dong, W. Q. (2013b). The use of albendazole and diammonium glycyrrhizinate in the treatment of eosinophilic meningitis in mice infected with *Angiostrongylus cantonensis*. *Journal of Helminthology* **87**, 1–11.
- Li, Y. C., Hu, X. M., Tong, C. J., Liu, J., Li, M. T. and Wang, S. Q. (2011). Investigation on serology, risk factor and awareness of *Angiostrongylus cantonensis* in Hainan province. *Zhongguo Ji Sheng Chong Xue Yu Ji Sheng Chong Bing Za Zhi* **29**, 74–75.
- Li, Z., Chen, X., Zen, X., Liang, J., Wei, J., Lv, Z., Sun, X. and Wu, Z. D. (2014b). MicroRNA expression profile in the third- and fourth-stage larvae of *Angiostrongylus cantonensis*. *Parasitology Research* **113**, 1883–1896.
- Li, Z. Y., Sun, R., Li, J., Song, Y. X., Lin, Y. C., Zeng, X., He, H. J., Wei, J., Yang, F., Zheng, H. Q., Lv, Z. Y. and Wu, Z. D. (2012). Efficacy of albendazole combined with a marine fungal extract (m2-9) against *Angiostrongylus cantonensis*-induced meningitis in mice. *Journal of Helminthology* **86**, 410–417.
- Liang, S. H., Huang, H. C., Pan, C. W. and Tan, F. (2005). Detection of *Angiostrongylus cantonensis* circulating antigen by monoclonal antibodies. *Zhonghua Yi Xue Za Zhi* **85**, 3057–3061.
- Lilic, N. and Addison, B. (2013). *Angiostrongylus* meningitis associated with intraparenchymal cerebral haemorrhage. *New Zealand Medical Journal* **126**, 98–102.
- Lin, J. X., Jie, H. Y. and Li, L. S. (2005). An inspiration from the outbreaks of *Angiostrongylus cantonensis*. *Zhongguo Ji Sheng Chong Xue Yu Ji Sheng Chong Bing Za Zhi* **23**, 341–343.
- Lin, R. J., Chen, C. Y., Chung, L. Y. and Yen, C. M. (2010). Larvicidal activities of ginger (*Zingiber officinale*) against *Angiostrongylus cantonensis*. *Acta Tropica* **115**, 69–76.
- Lindo, J. F., Waugh, C., Hall, J., Cunningham-Myrie, C., Ashley, D., Eberhard, M. L., Sullivan, J. J., Bishop, H. S., Robinson, D. G., Holtz, T. and Robinson, R. D. (2002). Enzootic *Angiostrongylus cantonensis* in rats and snails after an outbreak of human eosinophilic meningitis, Jamaica. *Emerging Infectious Diseases* **8**, 324–326.
- Lindo, J. F., Escoffery, C. T., Reid, B., Codrington, G., Cunningham-Myrie, C. and Eberhard, M. L. (2004). Fatal autochthonous eosinophilic meningitis in a Jamaican child caused by *Angiostrongylus cantonensis*. *American Journal of Tropical Medicine and Hygiene* **70**, 425–428.
- Liu, C. Y., Song, H. Q., Zhang, R. L., Chen, M. X., Xu, M. J., Ai, L., Chen, X. G., Zhan, X. M., Liang, S. H., Yuan, Z. G., Lin, R. Q. and Zhu, X. Q. (2011). Specific detection of *Angiostrongylus cantonensis* in the snail *Achatina fulica* using a loop-mediated isothermal amplification (LAMP) assay. *Molecular and Cellular Probes* **25**, 164–167.
- Liu, I. H., Chung, Y. M., Chen, S. J. and Cho, W. L. (2006). Necrotizing retinitis induced by *Angiostrongylus cantonensis*. *American Journal of Ophthalmology* **141**, 577–579.
- Liu, L. H., He, H. J., Lv, Z. Y., Wei, J., Zeng, X., Liang, J. Y., Zheng, H. Q., Yu, X. B., Sun, X. and Wu, Z. D. (2013). The mRNA level of the galectin-10 of *Angiostrongylus cantonensis* induced by reactive oxygen stress. *Parasitology Research* **112**, 933–943.
- Long, Y., Cao, B., Yu, L., Tukayo, M., Feng, C., Wang, Y. and Luo, D. (2015). *Angiostrongylus cantonensis* cathepsin B-like protease (Ac-cathB-1) is involved in host gut penetration. *Parasite* **22**, 37.
- Luessi, F., Sollors, J., Torzewski, M., Muller, H. D., Siegel, E., Blum, J., Sommer, C., Vogt, T. and Thomke, F. (2009). Eosinophilic meningitis due to *Angiostrongylus cantonensis* in Germany. *Journal of Travel Medicine* **16**, 292–294.
- Lunn, J., Lee, R., Smaller, J., MacKay, B., King, T., Hunt, G., Martin, P., Krockenberger, M., Spielman, D. and Malik, R. (2012). Twenty two cases of canine neural angiostrongylosis in eastern Australia (2002–2005) and a review of the literature. *Parasites and Vectors* **5**, 70.
- Lv, S., Zhou, X. N., Zhang, Y., Liu, H. X., Zhu, D., Yin, W. G., Steinmann, P., Wang, X. H. and Jia, T. W. (2006). The effect of temperature on the development of *Angiostrongylus cantonensis* (Chen 1935) in *Pomacea canaliculata* (Lamarck 1822). *Parasitology Research* **99**, 583–587.
- Lv, S., Zhang, Y., Steinmann, P. and Zhou, X. N. (2008). Emerging angiostrongyliasis in Mainland China. *Emerging Infectious Diseases* **14**, 161–164.
- Lv, S., Zhang, Y., Chen, S. R., Wang, L. B., Fang, W., Chen, F., Jiang, J. Y., Li, Y. L., Du, Z. W. and Zhou, X. N. (2009a). Human angiostrongyliasis outbreak in Dali, China. *PLoS Neglected Tropical Diseases* **3**, e520.
- Lv, S., Zhang, Y., Liu, H. X., Hu, L., Yang, K., Steinmann, P., Chen, Z., Wang, L. Y., Utzinger, J. and Zhou, X. N. (2009b). Invasive snails and an emerging infectious disease: results from the first national survey on *Angiostrongylus cantonensis* in China. *PLoS Neglected Tropical Diseases* **3**, e368.
- Lv, S., Zhang, Y., Liu, H. X., Zhang, C. W., Steinmann, P., Zhou, X. N. and Utzinger, J. (2009c). *Angiostrongylus cantonensis*: morphological and behavioral investigation within the freshwater snail *Pomacea canaliculata*. *Parasitology Research* **104**, 1351–1359.
- Lv, S., Zhang, Y., Zhang, L., Liu, Q., Liu, H. X., Hu, L., Wei, F. R., Steinmann, P., Graeff-Teixeira, C., Zhou, X. N. and Utzinger, J. (2012). The complete mitochondrial genome of the rodent intra-arterial nematodes *Angiostrongylus cantonensis* and *Angiostrongylus costaricensis*. *Parasitology Research* **111**, 115–123.
- Lv, S., Zhang, Y., Guo, Y. H., Liu, H. X., Zhou, Z. B., Jiang, M. and Gu, W. B. (2014). Comparative analysis of mitochondrial genomes of *Angiostrongylus cantonensis*. *Zhongguo Xue Xi Chong Bing Fang Zhi Za Zhi* **26**, 175–179.
- Ma, G., Dennis, M., Rose, K., Spratt, D. and Spielman, D. (2013). Tawny frogmouths and brushtail possums as sentinels for *Angiostrongylus cantonensis*, the rat lungworm. *Veterinary Parasitology* **192**, 158–165.
- Mackerras, M. and Sandars, D. (1955). The life history of the rat lungworm, *Angiostrongylus cantonensis* (Chen) (Nematoda: Metastrongylidae). *Australian Journal of Zoology* **3**, 1–25.
- Mackie, J. T., Lacasse, C. and Spratt, D. M. (2013). Patent *Angiostrongylus mackerrasae* infection in a black flying fox (*Pteropus alecto*). *Australian Veterinary Journal* **91**, 366–367.
- Maleewong, W., Sombatsawat, P., Intapan, P. M., Wongkham, C. and Chotmongkol, V. (2001). Immunoblot evaluation of the specificity of the 29-kDa antigen from young adult female worms *Angiostrongylus cantonensis* for immunodiagnosis of human angiostrongyliasis. *Asian Pacific Journal of Allergy and Immunology* **19**, 267–273.
- Malhotra, S., Mehta, D. K., Arora, R., Chauhan, D., Ray, S. and Jain, M. (2006). Ocular angiostrongyliasis in a child – first case report from India. *Journal of Tropical Pediatrics* **52**, 223–225.
- Malvy, D., Ezzedine, K., Receveur, M. C., Pistone, T., Crevon, L., Lemardeley, P. and Josse, R. (2008). Cluster of eosinophilic meningitis attributable to *Angiostrongylus cantonensis* infection in French policemen troop returning from the Pacific Islands. *Travel Medicine and Infectious Disease* **6**, 301–304.
- Maretic, T., Perovic, M., Vince, A., Lukas, D., Dekumyoy, P. and Begovac, J. (2009). Meningitis and radiculomyelitis caused by *Angiostrongylus cantonensis*. *Emerging Infectious Diseases* **15**, 996–998.
- Martin-Alonso, A., Foronda, P., Quispe-Ricalde, M. A., Feliu, C. and Valladares, B. (2011). Seroprevalence of *Angiostrongylus cantonensis* in wild rodents from the Canary Islands. *PLoS ONE* **6**, e27747.
- Martin-Alonso, A., Abreu-Yanes, E., Feliu, C., Mas-Coma, S., Bargas, M. D., Valladares, B. and Foronda, P. (2015). Intermediate hosts of *Angiostrongylus cantonensis* in Tenerife, Spain. *PLoS ONE* **10**, e0120686.
- Martins, Y. C., Tanowitz, H. B. and Kazacos, K. R. (2015). Central nervous system manifestations of *Angiostrongylus cantonensis* infection. *Acta Tropica* **141**, 46–53.
- Mattis, A., Mowatt, L., Lue, A., Lindo, J. and Vaughan, H. (2009). Ocular angiostrongyliasis—first case report from Jamaica. *West Indian Medical Journal* **58**, 383–385.
- McKenzie, R. A., Green, P. E. and Wood, A. D. (1978). *Angiostrongylus cantonensis* infection of the brain of a captive Bennett's wallaby (*Macropus rufogriseus*). *Australian Veterinary Journal* **54**, 86–88.

- McVeigh, P., Maule, A. G., Dalton, J. P. and Robinson, M. W. (2012). *Fasciola hepatica* virulence-associated cysteine peptidases: a systems biology perspective. *Microbes and Infection* **14**, 301–310.
- Mendonça, C. L., Carvalho, O. S., Mota, E. M., Pelajo-Machado, M., Caputo, L. F. and Lenzi, H. L. (1999). Penetration sites and migratory routes of *Angiostrongylus costaricensis* in the experimental intermediate host (*Sarasimula marginata*). *Memorias do Instituto Oswaldo Cruz* **94**, 549–556.
- Monks, D. J., Carlisle, M. S., Carrigan, M., Rose, K., Spratt, D., Gallagher, A. and Procriv, P. (2005). *Angiostrongylus cantonensis* as a cause of cerebrospinal disease in a yellow-tailed black cockatoo (*Calyptorhynchus funereus*) and two tawny frogmouths (*Podargus strigoideus*). *Journal of Avian Medicine and Surgery* **19**, 289–293.
- Monte, T. C., Simoes, R. O., Oliveira, A. P., Novaes, C. F., Thiengo, S. C., Silva, A. J., Estrela, P. C. and Maldonado, A., Jr. (2012). Phylogenetic relationship of the Brazilian isolates of the rat lungworm *Angiostrongylus cantonensis* (Nematoda: Metastrongylidae) employing mitochondrial COI gene sequence data. *Parasites and Vectors* **5**, 248.
- Morassutti, A. L. and Graeff-Teixeira, C. (2012). Interface molecules of *Angiostrongylus cantonensis*: their role in parasite survival and modulation of host defenses. *International Journal of Inflammation* **2012**, 512097.
- Morassutti, A. L., Levert, K., Perelygin, A., da Silva, A. J., Wilkins, P. and Graeff-Teixeira, C. (2012a). The 31-kDa antigen of *Angiostrongylus cantonensis* comprises distinct antigenic glycoproteins. *Vector Borne and Zoonotic Diseases* **12**, 961–968.
- Morassutti, A. L., Levert, K., Pinto, P. M., da Silva, A. J., Wilkins, P. and Graeff-Teixeira, C. (2012b). Characterization of *Angiostrongylus cantonensis* excretory–secretory proteins as potential diagnostic targets. *Experimental Parasitology* **130**, 26–31.
- Morassutti, A. L., Perelygin, A., Levert, K., Lin, S. C., Lee, Y. M., da Silva, A. J., Wilkins, P. P. and Graeff-Teixeira, C. (2013a). Expression of recombinant antigenic proteins from *Angiostrongylus cantonensis*: a brief report. *Hawaii Journal of Medicine and Public Health* **72**, 58–62.
- Morassutti, A. L., Perelygin, A., Carvalho Marcos, O. D. E., Lemos, L. N., Pinto, P. M., Frace, M., Wilkins, P. P., Graeff-Teixeira, C. and Da Silva, A. J. (2013b). High throughput sequencing of the *Angiostrongylus cantonensis* genome: a parasite spreading worldwide. *Parasitology* **140**, 1304–1309.
- Morassutti, A. L., Thiengo, S. C., Fernandez, M., Sawanyawisuth, K. and Graeff-Teixeira, C. (2014). Eosinophilic meningitis caused by *Angiostrongylus cantonensis*: an emergent disease in Brazil. *Memorias do Instituto Oswaldo Cruz* **109**, 399–407.
- Morera, P. (1973). Life history and redescription of *Angiostrongylus costaricensis* Morera and Cespedes, 1971. *American Journal of Tropical Medicine and Hygiene* **22**, 613–621.
- Morera, P. and Cespedes, R. (1971). *Angiostrongylus costaricensis* n. sp. (Nematoda: Metastrongyloidea), a new lungworm occurring in man in Costa Rica. *Revista de Biologia Tropical* **18**, 173–185.
- Morera, P., Lazo, R., Urquiza, J. and Llaguno, M. (1983). First record of *Angiostrongylus costaricensis* Morera and Cespedes, 1971 in Ecuador. *American Journal of Tropical Medicine and Hygiene* **32**, 1460–1461.
- Morley, N. J. (2010). Aquatic molluscs as auxiliary hosts for terrestrial nematode parasites: implications for pathogen transmission in a changing climate. *Parasitology* **137**, 1041–1056.
- Morton, N. J., Britton, P., Palasanthiran, P., Bye, A., Sugo, E., Kesson, A., Ardern-Holmes, S. and Snelling, T. L. (2013). Severe hemorrhagic meningoencephalitis due to *Angiostrongylus cantonensis* among young children in Sydney, Australia. *Clinical Infectious Diseases* **57**, 1158–1161.
- Mota, E. M. and Lenzi, H. L. (2005). *Angiostrongylus costaricensis*: complete redescription of the migratory pathways based on experimental *Sigmodon hispidus* infection. *Memorias do Instituto Oswaldo Cruz* **100**, 407–420.
- Muro, A. and Perez-Arellano, J. L. (2010). Nitric oxide and respiratory helminthic diseases. *Journal of Biomedicine and Biotechnology* **2010**, 958108.
- Murphy, G. S. and Johnson, S. (2013). Clinical aspects of eosinophilic meningitis and meningoencephalitis caused by *Angiostrongylus cantonensis*, the rat lungworm. *Hawaii Journal of Medicine and Public Health* **72**, 35–40.
- Nalini, A., Ramakrishna, A., Dekumoy, P., Kumar, R. R., Pakdee, W., Saini, J. and Hegde, V. S. (2013). Severe form of radiculomyelo – neuropathy with meningo – encephalitis secondary to *Angiostrongylus cantonensis* infection: unusual corpus callosum lesions and serial magnetic resonance imaging findings. *Neurology India* **61**, 414–418.
- Neuhauss, E., Fitarelli, M., Romanzini, J. and Graeff-Teixeira, C. (2007). Low susceptibility of *Achatina fulica* from Brazil to infection with *Angiostrongylus costaricensis* and *A. cantonensis*. *Memorias do Instituto Oswaldo Cruz* **102**, 49–52.
- Ni, F., Wang, Y., Zhang, J., Yu, L., Fang, W. and Luo, D. (2012). Cathepsin B-like and hemoglobin-type cysteine proteases: stage-specific gene expression in *Angiostrongylus cantonensis*. *Experimental Parasitology* **131**, 433–441.
- Odermatt, P., Lv, S. and Sayasone, S. (2010). Less common parasitic infections in Southeast Asia that can produce outbreaks. *Advances in Parasitology* **72**, 409–435.
- Oehler, E., Ghawche, F., Delattre, A., Berberian, A., Levy, M. and Valour, F. (2014). *Angiostrongylus cantonensis* eosinophilic meningitis: a clinical study of 42 consecutive cases in French Polynesia. *Parasitology International* **63**, 544–549.
- Okano, T., Haga, A., Mizuno, E., Onuma, M., Nakaya, Y. and Nagamine, T. (2014). *Angiostrongylus cantonensis* (Nematoda: Metastrongylidae) in the Ryukyu Islands tree rat (*Diplothrix legata*). *Journal of Wildlife Diseases* **50**, 322–325.
- Oliveira, A. P., Gentile, R., Maldonado Junior, A., Lopes Torres, E. J. and Thiengo, S. C. (2015). *Angiostrongylus cantonensis* infection in molluscs in the municipality of Sao Goncalo, a metropolitan area of Rio de Janeiro, Brazil: role of the invasive species *Achatina fulica* in parasite transmission dynamics. *Memorias do Instituto Oswaldo Cruz* **110**, 739–744.
- Owen, I. L. (2005). Parasitic zoonoses in Papua New Guinea. *Journal of Helminthology* **79**, 1–14.
- Padilla-Docal, B., Dorta-Contreras, A. J., Bu-Coifú-Fanego, R., Hernandez, H. F., Barroso, J. C. and Sanchez-Martinez, C. (2008). Intrathecal synthesis of IgE in children with eosinophilic meningoencephalitis caused by *Angiostrongylus cantonensis*. *Cerebrospinal Fluid Research* **5**, 18.
- Pai, S., Madi, D., Achappa, B., Mahalingam, S. and Kendambadi, R. (2013). An interesting case of eosinophilic meningitis. *Journal of Clinical and Diagnostic Research* **7**, 734–735.
- Palominos, P. E., Gasnier, R., Rodriguez, R., Agostini, A. A. and Graeff-Teixeira, C. (2008). Individual serological follow-up of patients with suspected or confirmed abdominal angiostrongyliasis. *Memorias do Instituto Oswaldo Cruz* **103**, 93–97.
- Pan, B., Wu, J., Ruan, C. W., Liang, W. J., Deng, Z. H., Zhang, Q. M., Huang, S. Y., Lin, R. X. and Pei, F. Q. (2011). Investigation on *Angiostrongylus cantonensis* infection in rodents in Guangdong Province. *Zhongguo Ji Sheng Chong Xue Yu Ji Sheng Chong Bing Za Zhi* **29**, 289–292.
- Patikulasila, D., Ittipunkul, N. and Theerakittikul, B. (2003). Intravital angiostrongyliasis: report of 2 cases. *Journal of the Medical Association of Thailand* **86**, 981–985.
- Peng, H., Sun, R., Zhang, Q., Zhao, J., Wei, J., Zeng, X., Zheng, H. and Wu, Z. (2013). Interleukin 33 mediates type 2 immunity and inflammation in the central nervous system of mice infected with *Angiostrongylus cantonensis*. *Journal of Infectious Diseases* **207**, 860–869.
- Pipitgool, V., Sithithaworn, P., Pongmuttasaya, P. and Hinz, E. (1997). *Angiostrongylus cantonensis* in rats and snails in northeast Thailand. *Southeast Asian Journal of Tropical Medicine and Public Health* **28** (Suppl. 1), 190–193.
- Pitrez, P. M., Gualdi, L. P., Barbosa, G. L., Sudbrack, S., Ponzi, D., Cao, R. G., Silva, A. C., Machado, D. C., Jones, M. H., Stein, R. T. and Graeff-Teixeira, C. (2015). Effect of different helminth extracts on the development of asthma in mice: the influence of early-life exposure and the role of IL-10 response. *Experimental Parasitology* **156**, 95–103.
- Podwall, D., Gupta, R., Furuya, E. Y., Sevigny, J. and Resor, S. R. (2004). *Angiostrongylus cantonensis* meningitis presenting with facial nerve palsy. *Journal of Neurology* **251**, 1280–1281.
- Procriv, P. and Carlisle, M. S. (2001). The spread of *Angiostrongylus cantonensis* in Australia. *Southeast Asian Journal of Tropical Medicine and Public Health* **32**(Suppl. 2), 126–128.
- Procriv, P., Spratt, D. M. and Carlisle, M. S. (2000). Neuro-angiostrongyliasis: unresolved issues. *International Journal for Parasitology* **30**, 1295–1303.
- Qi, H., Diao, Z. and Yin, C. (2009). A case of optic nerve compression caused by *Angiostrongylus cantonensis*. *American Journal of Tropical Medicine and Hygiene* **81**, 4.
- Qvarnstrom, Y., Sullivan, J. J., Bishop, H. S., Hollingsworth, R. and da Silva, A. J. (2007). PCR-based detection of *Angiostrongylus cantonensis* in tissue and mucus secretions from molluscan hosts. *Applied and Environmental Microbiology* **73**, 1415–1419.
- Qvarnstrom, Y., da Silva, A. C., Teem, J. L., Hollingsworth, R., Bishop, H., Graeff-Teixeira, C. and da Silva, A. J. (2010). Improved molecular detection of *Angiostrongylus cantonensis* in mollusks and other environmental samples with a species-specific internal transcribed spacer 1-based TaqMan assay. *Applied and Environmental Microbiology* **76**, 5287–5289.
- Qvarnstrom, Y., Bishop, H. S. and da Silva, A. J. (2013). Detection of rat lungworm in intermediate, definitive, and paratenic hosts obtained

- from environmental sources. *Hawai'i Journal of Medicine and Public Health* **72**, 63–69.
- Qvarnstrom, Y., Xayavong, M., da Silva, A. C., Park, S. Y., Whelen, A. C., Calimlim, P. S., Sciuilli, R. H., Honda, S. A., Higa, K., Kitsutani, P., Chea, N., Heng, S., Johnson, S., Graeff-Teixeira, C., Fox, L. M. and da Silva, A. J. (2016). Real-time polymerase chain reaction detection of *Angiostrongylus cantonensis* DNA in cerebrospinal fluid from patients with eosinophilic meningitis. *American Journal of Tropical Medicine and Hygiene* **94**, 176–181.
- Rai, S., Madi, D., Pai, S. and Baliga, S. (2014). Unusual larva in the CSF and unique MRI findings in a case of eosinophilic meningitis. *Journal of Clinical Imaging Science* **4**, 76.
- Rebello, K. M., Siqueira, C. R., Ribeiro, E. L., Valente, R. H., Mota, E. M., Perales, J., Neves-Ferreira, A. G. and Lenzi, H. L. (2012). Proteolytic activity in the adult and larval stages of the human roundworm parasite *Angiostrongylus costaricensis*. *Memorias do Instituto Oswaldo Cruz* **107**, 752–759.
- Reece, R. L., Perry, R. A. and Spratt, D. M. (2013). Neuroangiostrongyliasis due to *Angiostrongylus cantonensis* in gang-gang cockatoos (*Callocephalon fimbriatum*). *Australian Veterinary Journal* **91**, 477–481.
- Robertson, A. P., Puttachary, S., Buxton, S. K. and Martin, R. J. (2015). Tribendimidine: mode of action and nAChR subtype selectivity in *Ascaris* and *Oesophagostomum*. *PLoS Neglected Tropical Diseases* **9**, e0003495.
- Rodriguez, R., da Silva, A. C., Muller, C. A., Alves, S. L., Graeff-Teixeira, C. and Fornari, F. (2014). PCR for the diagnosis of abdominal angiostrongyliasis in formalin-fixed paraffin-embedded human tissue. *PLoS ONE* **9**, e93658.
- Saluja, R., Khan, M., Church, M. K. and Maurer, M. (2015). The role of IL-33 and mast cells in allergy and inflammation. *Clinical and Translational Allergy* **5**, 33.
- Sawanyawisuth, K. (2008). Treatment of angiostrongyliasis. *Transactions of the Royal Society of Tropical Medicine and Hygiene* **102**, 990–996.
- Sawanyawisuth, K. and Chotmongkol, V. (2013). Eosinophilic meningitis. *Handbook of Clinical Neurology* **114**, 207–215.
- Sawanyawisuth, K. and Kitthaweesin, K. (2008). Optic neuritis caused by intraocular angiostrongyliasis. *Southeast Asian Journal of Tropical Medicine and Public Health* **39**, 1005–1007.
- Sawanyawisuth, K., Limpawattana, P., Busaracome, P., Ninpaitoon, B., Chotmongkol, V., Intapan, P. M. and Tanawirattananit, S. (2004). A 1-week course of corticosteroids in the treatment of eosinophilic meningitis. *American Journal of Medicine* **117**, 802–803.
- Sawanyawisuth, K., Kitthaweesin, K., Limpawattana, P., Intapan, P. M., Tiampakao, S., Jitpimolmard, S. and Chotmongkol, V. (2007). Intraocular angiostrongyliasis: clinical findings, treatments and outcomes. *Transactions of the Royal Society of Tropical Medicine and Hygiene* **101**, 497–501.
- Sawanyawisuth, K., Takahashi, K., Hoshuyama, T., Senthong, V., Limpawattana, P., Intapan, P. M., Wilson, D., Tiampakao, S., Jitpimolmard, S. and Chotmongkol, V. (2009). Clinical factors predictive of encephalitis caused by *Angiostrongylus cantonensis*. *American Journal of Tropical Medicine and Hygiene* **81**, 698–701.
- Sawanyawisuth, K., Sawanyawisuth, K., Intapan, P. M., Khotsri, P., Kanpittaya, J., Chotmongkol, V. and Maleewong, W. (2011). Specificity of immunoblotting analyses in eosinophilic meningitis. *Memorias do Instituto Oswaldo Cruz* **106**, 570–572.
- Sawanyawisuth, K., Chindaprasit, J., Senthong, V., Limpawattana, P., Auvichayapat, N., Tassniyom, S., Chotmongkol, V., Maleewong, W. and Intapan, P. M. (2013). Clinical manifestations of Eosinophilic meningitis due to infection with *Angiostrongylus cantonensis* in children. *Korean Journal of Parasitology* **51**, 735–738.
- Schmutzhard, E., Boongird, P. and Vejjajiva, A. (1988). Eosinophilic meningitis and radiculomyelitis in Thailand, caused by CNS invasion of *Gnathostoma spinigerum* and *Angiostrongylus cantonensis*. *Journal of Neurology, Neurosurgery, and Psychiatry* **51**, 80–87.
- Schulte, C., Krebs, B., Jelinek, T., Nothdurft, H. D., von Sonnenburg, F. and Loscher, T. (2002). Diagnostic significance of blood eosinophilia in returning travelers. *Clinical Infectious Diseases* **34**, 407–411.
- Senanayake, S. N., Pryor, D. S., Walker, J. and Konecny, P. (2003). First report of human angiostrongyliasis acquired in Sydney. *Medical Journal of Australia* **179**, 430–431.
- Senthong, V., Chindaprasit, J. and Sawanyawisuth, K. (2013). Differential diagnosis of CNS angiostrongyliasis: a short review. *Hawai'i Journal of Medicine and Public Health* **72**, 52–54.
- Shen, H. X., Chen, K., Pan, S. H., Huang, L. T., Wang, C., Lu, Y. Y., Ma, C. L., Chen, D. X., Chen, X. Y., Li, X. M., Zhu, W. Y. and Ren, W. F. (2012). Investigation on spread and hazards of human parasites in Qushui Village, Suixi County, Zhanjiang City. *Zhongguo Xue Xi Chong Bing Fang Zhi Za Zhi* **24**, 461–463.
- Shyu, L. Y., Chang, H. H., Hsu, J. D., Lin, D. P., Teng, Y. H. and Lee, H. H. (2012). Curcumin alleviates eosinophilic meningitis through reduction of eosinophil count following albendazole treatment against *Angiostrongylus cantonensis* in mice. *Parasitology* **139**, 358–365.
- Simoes, R. O., Monteiro, F. A., Sanchez, E., Thiengo, S. C., Garcia, J. S., Costa-Neto, S. F., Luque, J. L. and Maldonado, A. (2011). Endemic angiostrongyliasis, Rio de Janeiro, Brazil. *Emerging Infectious Diseases* **17**, 1331–1333.
- Simoes, R. O., Maldonado Junior, A., Olifiers, N., Garcia, J. S., Bertolino, A. V. and Luque, J. L. (2014). A longitudinal study of *Angiostrongylus cantonensis* in an urban population of *Rattus norvegicus* in Brazil: the influences of seasonality and host features on the pattern of infection. *Parasites and Vectors* **7**, 100.
- Sinawat, S. and Yospaiboon, Y. (2013). Subretinal angiostrongyliasis-induced optic neuritis. *Clinical Ophthalmology* **7**, 977–979.
- Sinawat, S., Sanguansak, T., Angkawinijwong, T., Ratanapakorn, T., Intapan, P. M. and Yospaiboon, Y. (2008). Ocular angiostrongyliasis: clinical study of three cases. *Eye* **22**, 1446–1448.
- Slatko, B. E., Luck, A. N., Dobson, S. L. and Foster, J. M. (2014). *Wolbachia* endosymbionts and human disease control. *Molecular and Biochemical Parasitology* **195**, 88–95.
- Slom, T. J., Cortese, M. M., Gerber, S. I., Jones, R. C., Holtz, T. H., Lopez, A. S., Zambrano, C. H., Sufit, R. L., Sakolvaree, Y., Chaicumpa, W., Herwaldt, B. L. and Johnson, S. (2002). An outbreak of eosinophilic meningitis caused by *Angiostrongylus cantonensis* in travelers returning from the Caribbean. *New England Journal of Medicine* **346**, 668–675.
- Song, L., Wang, X., Yang, Z., Lv, Z. and Wu, Z. (2016). *Angiostrongylus cantonensis* in the vector snails *Pomacea canaliculata* and *Achatina fulica* in China: a meta-analysis. *Parasitology Research* **115**, 913–923.
- Song, Z., Huang, H., Tan, F., Zhang, E., Hu, J. and Pan, C. (2012). Differential proteomics analysis of female and male adults of *Angiostrongylus cantonensis*. *Experimental Parasitology* **131**, 169–174.
- Spencer, L. A. and Weller, P. F. (2010). Eosinophils and Th2 immunity: contemporary insights. *Immunology and Cell Biology* **88**, 250–256.
- Spratt, D. M. (2015). Species of *Angiostrongylus* (Nematoda: Metastrongyloidea) in wildlife: a review. *International Journal for Parasitology. Parasites and Wildlife* **4**, 178–189.
- Stack, C., Dalton, J. P. and Robinson, M. W. (2011). The phylogeny, structure and function of trematode cysteine proteases, with particular emphasis on the *Fasciola hepatica* cathepsin L family. *Advances in Experimental Medicine and Biology* **712**, 116–135.
- Stockdale-Walden, H. D., Slapcinsky, J., Qvarnstrom, Y., McIntosh, A., Bishop, H. S. and Rosseland, B. (2015). *Angiostrongylus cantonensis* in introduced gastropods in Southern Florida. *Journal of Parasitology* **101**, 156–159.
- Suankratay, C., Wilde, H. and Berger, S. (2001). Thailand: country survey of infectious diseases. *Journal of Travel Medicine* **8**, 192–203.
- Teem, J. L., Qvarnstrom, Y., Bishop, H. S., da Silva, A. J., Carter, J., White-McLean, J. and Smith, T. (2013). The occurrence of the rat lungworm, *Angiostrongylus cantonensis*, in nonindigenous snails in the Gulf of Mexico region of the United States. *Hawai'i Journal of Medicine and Public Health* **72**, 11–14.
- Tesana, S., Srisawangwong, T., Sithithaworn, P. and Laha, T. (2008). *Angiostrongylus cantonensis*: experimental study on the susceptibility of apple snails, *Pomacea canaliculata* compared to *Pila polita*. *Experimental Parasitology* **118**, 531–535.
- Tesana, S., Srisawangwong, T., Sithithaworn, P., Laha, T. and Andrews, R. (2009). Prevalence and intensity of infection with third stage larvae of *Angiostrongylus cantonensis* in mollusks from Northeast Thailand. *American Journal of Tropical Medicine and Hygiene* **80**, 983–987.
- Thanaviratananich, S. and Ngamjarus, C. (2015). Corticosteroids for parasitic eosinophilic meningitis. *Cochrane Database of Systematic Reviews* **2**, CD009088.
- Thiengo, S. C., Maldonado, A., Mota, E. M., Torres, E. J., Caldeira, R., Carvalho, O. S., Oliveira, A. P., Simoes, R. O., Fernandez, M. A. and Lanfredi, R. M. (2010). The giant African snail *Achatina fulica* as natural intermediate host of *Angiostrongylus cantonensis* in Pernambuco, northeast Brazil. *Acta Tropica* **115**, 194–199.
- Thiengo, S. C., Simoes Rde, O., Fernandez, M. A. and Maldonado, A., Jr. (2013). *Angiostrongylus cantonensis* and rat lungworm disease in Brazil. *Hawai'i Journal of Medicine and Public Health* **72**, 18–22.
- Thirugnanam, S., Munirathinam, G., Veerapathran, A., Dakshinamoorthy, G., Reddy, M. V. and Ramaswamy, K. (2012). Cloning and characterization of high mobility group box protein 1

- (HMGB1) of *Wuchereria bancrofti* and *Brugia malayi*. *Parasitology Research* **111**, 619–627.
- Thyssen, A., Mitchell, M., Qvarnstrom, Y., Rao, S., Benke, T. A. and Glode, M. P. (2013). Eosinophilic meningitis in a previously healthy 13-year-old child. *Pediatric Infectious Disease Journal* **32**, 194, 198.
- Tokiwa, T., Harunari, T., Tanikawa, T., Komatsu, N., Koizumi, N., Tung, K. C., Suzuki, J., Kadosaka, T., Takada, N., Kumagai, T., Akao, N. and Ohta, N. (2012). Phylogenetic relationships of rat lungworm, *Angiostrongylus cantonensis*, isolated from different geographical regions revealed widespread multiple lineages. *Parasitology International* **61**, 431–436.
- Tokiwa, T., Hashimoto, T., Yabe, T., Komatsu, N., Akao, N. and Ohta, N. (2013). First report of *Angiostrongylus cantonensis* (Nematoda: Angiostrongylidae) infections in invasive rodents from five islands of the Ogasawara Archipelago, Japan. *PLoS ONE* **8**, e70729.
- Tomanakan, K., Srisurach, N., Sae-tung, S. and Pengpinich, C. (2008). Detection of circulating antibody of *Parastrongylus cantonensis* in sera with eosinophilic meningitis by dot-blot ELISA. *Journal of the Medical Association of Thailand* **91**, 1082–1086.
- Tsai, H. C., Liu, Y. C., Kunin, C. M., Lai, P. H., Lee, S. S., Chen, Y. S., Wann, S. R., Lin, W. R., Huang, C. K., Ger, L. P., Lin, H. H. and Yen, M. Y. (2003). Eosinophilic meningitis caused by *Angiostrongylus cantonensis* associated with eating raw snails: correlation of brain magnetic resonance imaging scans with clinical findings. *American Journal of Tropical Medicine and Hygiene* **68**, 281–285.
- Tsai, H. C., Lee, S. S., Huang, C. K., Yen, C. M., Chen, E. R. and Liu, Y. C. (2004). Outbreak of eosinophilic meningitis associated with drinking raw vegetable juice in southern Taiwan. *American Journal of Tropical Medicine and Hygiene* **71**, 222–226.
- Tsai, H. C., Liu, Y. C., Lee, S. S., Chen, E. R. and Yen, C. M. (2007a). Vascular endothelial growth factor is associated with blood brain barrier dysfunction in eosinophilic meningitis caused by *Angiostrongylus cantonensis* infection. *American Journal of Tropical Medicine and Hygiene* **76**, 592–595.
- Tsai, H. C., Chung, L. Y., Chen, E. R., Liu, Y. C., Lee, S. S., Chen, Y. S., Sy, C. L., Wann, S. R. and Yen, C. M. (2008). Association of matrix metalloproteinase-9 and tissue inhibitors of metalloproteinase-4 in cerebrospinal fluid with blood–brain barrier dysfunction in patients with eosinophilic meningitis caused by *Angiostrongylus cantonensis*. *American Journal of Tropical Medicine and Hygiene* **78**, 20–27.
- Tsai, H. C., Lai, P. H., Sy, C. L., Lee, S. S., Yen, C. M., Wann, S. R. and Chen, Y. S. (2011). Encephalitis caused by *Angiostrongylus cantonensis* after eating raw frogs mixed with wine as a health supplement. *Internal Medicine* **50**, 771–774.
- Tsai, H. C., Tseng, Y. T., Yen, C. M., Chen, E. R., Sy, C. L., Lee, S. S., Wann, S. R. and Chen, Y. S. (2012). Brain magnetic resonance imaging abnormalities in eosinophilic meningitis caused by *Angiostrongylus cantonensis* infection. *Vector Borne and Zoonotic Diseases* **12**, 161–166.
- Tsai, H. C., Chen, Y. S. and Yen, C. M. (2013). Human parasitic meningitis caused by *Angiostrongylus cantonensis* infection in Taiwan. *Hawai'i Journal of Medicine and Public Health* **72**, 26–27.
- Tsai, H. C., Huang, Y. L., Chen, Y. S., Yen, C. M., Tsai, R., Lee, S. S. and Tai, M. H. (2014a). 14-3-3beta protein expression in eosinophilic meningitis caused by *Angiostrongylus cantonensis* infection. *BMC Research Notes* **7**, 97.
- Tsai, H. C., Lee, B. Y., Yen, C. M., Wann, S. R., Lee, S. S., Chen, Y. S. and Tai, M. H. (2014b). Dexamethasone downregulated the expression of CSF 14-3-3beta protein in mice with eosinophilic meningitis caused by *Angiostrongylus cantonensis* infection. *Acta Tropica* **131**, 98–103.
- Tsai, H. C., Lee, B. Y., Yen, C. M., Wann, S. R., Lee, S. S. and Chen, Y. S. (2015). Dexamethasone inhibits brain apoptosis in mice with eosinophilic meningitis caused by *Angiostrongylus cantonensis* infection. *Parasites and Vectors* **8**, 200.
- Tsai, K. H., Huang, C. G., Wang, L. C., Yu, Y. W., Wu, W. J. and Chen, W. J. (2007b). Molecular evidence for the endosymbiont *Wolbachia* in a non-filaroid nematode, *Angiostrongylus cantonensis*. *Journal of Biomedical Science* **14**, 607–615.
- Tseng, Y. K., Tu, W. C., Lee, H. H., Chen, K. M., Chou, H. L. and Lai, S. C. (2004). Ultrastructural localization of matrix metalloproteinase-9 in eosinophils from the cerebrospinal fluid of mice with eosinophilic meningitis caused by *Angiostrongylus cantonensis*. *Annals of Tropical Medicine and Parasitology* **98**, 831–841.
- Tseng, Y. T., Tsai, H. C., Sy, C. L., Lee, S. S., Wann, S. R., Wang, Y. H., Chen, J. K., Wu, K. S. and Chen, Y. S. (2011). Clinical manifestations of eosinophilic meningitis caused by *Angiostrongylus cantonensis*: 18 years' experience in a medical center in southern Taiwan. *Journal of Microbiology Immunology and Infection* **44**, 382–389.
- Tu, W. C. and Lai, S. C. (2006). *Angiostrongylus cantonensis*: efficacy of albendazole-dexamethasone co-therapy against infection-induced plasminogen activators and eosinophilic meningitis. *Experimental Parasitology* **113**, 8–15.
- Tunholi-Alves, V. M., Tunholi, V. M., Lustrino, D., Amaral, L. S., Thiengo, S. C. and Pinheiro, J. (2011). Changes in the reproductive biology of *Biomphalaria glabrata* experimentally infected with the nematode *Angiostrongylus cantonensis*. *Journal of Invertebrate Pathology* **108**, 220–223.
- Tunholi-Alves, V. M., Tunholi, V. M., Pinheiro, J. and Thiengo, S. C. (2012). Effects of infection by larvae of *Angiostrongylus cantonensis* (Nematoda, Metastrongylidae) on the metabolism of the experimental intermediate host *Biomphalaria glabrata*. *Experimental Parasitology* **131**, 143–147.
- Tunholi-Alves, V. M., Tunholi, V. M., Castro, R. N., Sant'Ana, L. D., Santos-Amaral, L., de Oliveira, A. P., Garcia, J., Thiengo, S. C., Pinheiro, J. and Maldonado, A., Jr. (2014). Activation of anaerobic metabolism in *Biomphalaria glabrata* (Mollusca: Gastropoda) experimentally infected by *Angiostrongylus cantonensis* (Nematoda, Metastrongylidae) by high-performance liquid chromatography. *Parasitology International* **63**, 64–68.
- Tunholi-Alves, V. M., Tunholi, V. M., Amaral, L. S., Mota, E. M., Maldonado Junior, A., Pinheiro, J. and Garcia, J. (2015). Biochemical profile of *Achatina fulica* (Mollusca: Gastropoda) after infection by different parasitic loads of *Angiostrongylus cantonensis* (Nematoda, Metastrongylidae). *Journal of Invertebrate Pathology* **124**, 1–5.
- Ubelaker, J. E. (1986). Systematics of species referred to the genus *Angiostrongylus*. *Journal of Parasitology* **72**, 237–244.
- Ueda, M., Takeuchi, Y., Ochiai, J., Mabuchi, C. and Niwa, J. (2015). A case of myelitis with eosinophilia of the cerebrospinal fluid. *Rinsho Shinkeigaku* **55**, 651–653.
- Uga, S., Ono, K., Kataoka, N. and Hasan, H. (1996). Seroepidemiology of five major zoonotic parasite infections in inhabitants of Sidoarjo, East Java, Indonesia. *Southeast Asian Journal of Tropical Medicine and Public Health* **27**, 556–561.
- Vafadari, B., Salamian, A. and Kaczmarek, L. (2016). MMP-9 in translation: from molecule to brain physiology, pathology and therapy. *Journal of Neurochemistry*. doi:10.1111/jnc.13415.
- Vibanco-Perez, N. and Landa-Piedra, A. (1998). Glutathione S-transferase in helminth parasites. *Revista Latinoamericana de Microbiologia* **40**, 73–85.
- Vitta, A., Polseela, R., Nateworanart, S. and Tattiyapong, M. (2011). Survey of *Angiostrongylus cantonensis* in rats and giant African land snails in Phitsanulok province, Thailand. *Asian Pacific Journal of Tropical Medicine* **4**, 597–599.
- Vitta, A., Polsut, W., Fukruksa, C., Yimthin, T., Thanwisai, A. and Dekumyoy, P. (2016). Levels of infection with the lungworm *Angiostrongylus cantonensis* in terrestrial snails from Thailand, with *Cryptozona siamensis* as a new intermediate host. *Journal of Helminthology*, 1–5. doi:10.1017/S0022149X15001042.
- Walker, A. G., Spielman, D., Malik, R., Graham, K., Ralph, E., Linton, M. and Ward, M. P. (2015). Canine neural angiostrongylosis: a case-control study in Sydney dogs. *Australian Veterinary Journal* **93**, 195–199.
- Wallace, G. D. (2013). The discovery of humans in Hawai'i infected with *Angiostrongylus cantonensis*, and early epidemiological findings. *Hawai'i Journal of Medicine and Public Health* **72**, 5.
- Wan, K. S. and Weng, W. C. (2004). Eosinophilic meningitis in a child raising snails as pets. *Acta Tropica* **90**, 51–53.
- Wang, J., Qi, H., Diao, Z., Zheng, X., Li, X., Ma, S., Ji, A. and Yin, C. (2010). An outbreak of *Angiostrongylus cantonensis* in Beijing. *Journal of Parasitology* **96**, 377–381.
- Wang, J., Wei, J., Zeng, X., Liang, J. Y., Wu, F., Li, Z. Y., Zheng, H. Q., He, H. J. and Wu, Z. D. (2013a). Efficacy of tribendimidine against *Angiostrongylus cantonensis* infection in the mice. *Parasitology Research* **112**, 1039–1046.
- Wang, J. J., Chung, L. Y., Lin, R. J., Lee, J. D., Lin, C. W. and Yen, C. M. (2011). Eosinophilic meningitis risk associated with raw *Ampullarium canaliculatus* snails consumption. *Kaohsiung Journal of Medical Sciences* **27**, 184–189.
- Wang, L. C., Jung, S. M., Chen, C. C., Wong, H. F., Wan, D. P. and Wan, Y. L. (2006a). Pathological changes in the brains of rabbits experimentally infected with *Angiostrongylus cantonensis* after albendazole treatment: histopathological and magnetic resonance imaging studies. *Journal of Antimicrobial Chemotherapy* **57**, 294–300.
- Wang, L. C., Wang, I. H. and Jou, J. R. (2006b). Optic neuritis secondary to *Angiostrongylus cantonensis* infection. *Ocular Immunology and Inflammation* **14**, 189–191.
- Wang, L. C., Chen, K. Y., Chang, S. H., Chung, L. Y., Gan, R. C., Cheng, C. J. and Tang, P. (2013b). Transcriptome profiling of the fifth-stage larvae of *Angiostrongylus cantonensis* by next-generation sequencing. *Parasitology Research* **112**, 3193–3202.
- Wang, M. and Lv, S. (2014). Substitution saturation analysis of mitochondrial cytochrome C oxidase subunit I (COX1) gene of *Angiostrongylus*

- cantonensis*. *Zhongguo Ji Sheng Chong Xue Yu Ji Sheng Chong Bing Za Zhi* **32**, 205–209.
- Wang, Q. P., Chen, X. G. and Lun, Z. R. (2007). Invasive freshwater snail, China. *Emerging Infectious Diseases* **13**, 1119–1120.
- Wang, Q.-P., Lai, D.-H., Zhu, X.-Q., Chen, X.-G. and Lun, Z.-R. (2008). Human angiostrongyliasis. *Lancet Infectious Diseases* **8**, 621–630.
- Wang, Q. P., Wu, Z. D., Wei, J., Owen, R. L. and Lun, Z. R. (2012). Human *Angiostrongylus cantonensis*: an update. *European Journal of Clinical Microbiology and Infectious Diseases* **31**, 389–395.
- Watanabe, Y., Suematsu, T. and Ohtsuki, T. (2014). Losing the stem-loop structure from metazoan mitochondrial tRNAs and co-evolution of interacting factors. *Frontiers in Genetics* **5**, 109.
- Waugh, C. A., Shafir, S., Wise, M., Robinson, R. D., Eberhard, M. L. and Lindo, J. F. (2005). Human *Angiostrongylus cantonensis*, Jamaica. *Emerging Infectious Diseases* **11**, 1977–1978.
- Wei, F. R., Liu, H. X., Lv, S., Hu, L. and Zhang, Y. (2010). Multiplex PCR assay for the detection of *Angiostrongylus cantonensis* larvae in *Pomacea canaliculata*. *Zhongguo Ji Sheng Chong Xue Yu Ji Sheng Chong Bing Za Zhi* **28**, 355–358.
- Wei, J., Wu, F., He, A., Zeng, X., Ouyang, L. S., Liu, M. S., Zheng, H. Q., Lei, W. L., Wu, Z. D. and Lv, Z. Y. (2015). Microglia activation: one of the checkpoints in the CNS inflammation caused by *Angiostrongylus cantonensis* infection in rodent model. *Parasitology Research* **114**, 3247–3254.
- Wei, P. C., Tsai, C. H., Chiu, P. S. and Lai, S. C. (2011). Matrix metalloproteinase-12 leads to elastin degradation in BALB/c mice with eosinophilic meningitis caused by *Angiostrongylus cantonensis*. *International Journal for Parasitology* **41**, 1175–1183.
- Wei, Y., Hong, Q., Chen, D., Liang, C., Liu, H., Luo, X. and Zhu, X. (2014). Permissibility of Mongolian gerbil for *Angiostrongylus cantonensis* infection and utility of this animal model for anthelmintic studies. *Parasitology Research* **113**, 1687–1693.
- Wilkins, P. P., Qvarnstrom, Y., Whelen, A. C., Saucier, C., da Silva, A. J. and Eamsobhana, P. (2013). The current status of laboratory diagnosis of *Angiostrongylus cantonensis* infections in humans using serologic and molecular methods. *Hawaii Journal of Medicine and Public Health* **72**, 55–57.
- Wright, J. D., Kelly, W. R., Waddell, A. H. and Hamilton, J. (1991). Equine neural angiostrongylosis. *Australian Veterinary Journal* **68**, 58–60.
- Yang, T. B., Wu, Z. D. and Lun, Z. R. (2013a). The apple snail *Pomacea canaliculata*, a novel vector of the rat lungworm, *Angiostrongylus cantonensis*: its introduction, spread, and control in China. *Hawaii Journal of Medicine and Public Health* **72**, 23–25.
- Yang, X., Qu, Z., He, H., Zheng, X., He, A., Wu, Y., Liu, Q., Zhang, D., Wu, Z., Li, Z. and Zhan, X. (2012). Enzootic angiostrongyliasis in Guangzhou, China, 2008–2010. *American Journal of Tropical Medicine and Hygiene* **86**, 846–849.
- Yang, X., Li, Z., He, H., Cheng, M., Liu, Q., Zhang, D., Chen, J., Wu, X., He, A., Zheng, X., Wu, Y., Wu, Z. and Zhan, X. (2013b). Molecular cloning, expression, and characterization of a putative activation-associated secreted protein from *Angiostrongylus cantonensis*. *Parasitology Research* **112**, 781–788.
- Yeung, N. W., Hayes, K. A. and Cowie, R. H. (2013). Effects of washing produce contaminated with the snail and slug hosts of *Angiostrongylus cantonensis* with three common household solutions. *Hawaii Journal of Medicine and Public Health* **72**, 83–86.
- Yii, C. Y. (1976). Clinical observations on eosinophilic meningitis and meningoencephalitis caused by *Angiostrongylus cantonensis* on Taiwan. *American Journal of Tropical Medicine and Hygiene* **25**, 233–249.
- Yii, C. Y., Chen, C. Y., Chen, E. R., Hsieh, H. C. and Shih, C. C. (1975). Epidemiologic studies of eosinophilic meningitis in southern Taiwan. *American Journal of Tropical Medicine and Hygiene* **24**, 447–454.
- Yong, H. S. and Eamsobhana, P. (2013). Definitive rodent hosts of the rat lungworm *Angiostrongylus cantonensis*. *Raffles Bulletin of Zoology Suppl.* **29**, 111–115.
- Yong, H. S., Eamsobhana, P., Lim, P. E., Razali, R., Aziz, F. A., Rosli, N. S., Poole-Johnson, J. and Anwar, A. (2015a). Draft genome of neurotropic nematode parasite *Angiostrongylus cantonensis*, causative agent of human eosinophilic meningitis. *Acta Tropica* **148**, 51–57.
- Yong, H. S., Eamsobhana, P., Song, S. L., Prasartvit, A. and Lim, P. E. (2015b). Molecular phylogeography of *Angiostrongylus cantonensis* (Nematoda: Angiostrongylidae) and genetic relationships with congeners using *cytochrome b* gene marker. *Acta Tropica* **148**, 66–71.
- Yong, H. S., Song, S. L., Eamsobhana, P., Goh, S. Y. and Lim, P. E. (2015c). Complete mitochondrial genome reveals genetic diversity of *Angiostrongylus cantonensis* (Nematoda: Angiostrongylidae). *Acta Tropica* **152**, 157–164.
- Yong, H. S., Song, S. L., Eamsobhana, P., Goh, S. Y., Lim, P. E., Chow, W. L., Chan, K. G. and Abrahams-Sandi, E. (2015d). Mitochondrial genome supports sibling species of *Angiostrongylus costaricensis* (Nematoda: Angiostrongylidae). *PLoS ONE* **10**, e0134581.
- York, E. M., Butler, C. J. and Lord, W. D. (2014). Global decline in suitable habitat for *Angiostrongylus* (= *Parastrostrongylus*) *cantonensis*: the role of climate change. *PLoS ONE* **9**, e103831.
- York, E. M., Creecy, J. P., Lord, W. D. and Caire, W. (2015). Geographic range expansion for rat lungworm in North America. *Emerging Infectious Diseases* **21**, 1234–1236.
- Yousif, F. and Lammler, G. (1975a). The effect of some biological and physical factors on infection of *Biomphalaria glabrata* with *Angiostrongylus cantonensis*. *Zeitschrift für Parasitenkunde* **47**, 191–201.
- Yousif, F. and Lammler, G. (1975b). The suitability of several aquatic snails as intermediate hosts for *Angiostrongylus cantonensis*. *Zeitschrift für Parasitenkunde* **47**, 203–210.
- Yousif, F., Blahser, S. and Lammler, G. (1980). The cellular responses in *Marisa cornuarietis* experimentally infected with *Angiostrongylus cantonensis*. *Zeitschrift für Parasitenkunde* **62**, 179–190.
- Yu, C., Wang, Y., Zhang, J., Fang, W. and Luo, D. (2014a). Immunolocalization and developmental expression patterns of two cathepsin B proteases (AC-cathB-1, -2) of *Angiostrongylus cantonensis*. *Experimental Parasitology* **144**, 27–33.
- Yu, L., Liao, Q., Chen, X., Xu, L., Zeng, X., Lv, Z., Sun, X., Zhen, H. and Wu, Z. (2014b). Dynamic expression of miR-132, miR-212, and miR-146 in the brain of different hosts infected with *Angiostrongylus cantonensis*. *Parasitology Research* **113**, 91–99.
- Yu, L., Wu, X., Wei, J., Liao, Q., Xu, L., Luo, S., Zeng, X., Zhao, Y., Lv, Z. and Wu, Z. (2015). Preliminary expression profile of cytokines in brain tissue of BALB/c mice with *Angiostrongylus cantonensis* infection. *Parasites and Vectors* **8**, 328.
- Zarowiecki, M. and Berriman, M. (2015). What helminth genomes have taught us about parasite evolution. *Parasitology* **142**(Suppl. 1), S85–S97.
- Zeng, X., Wang, J., Wei, J., Wu, F., Fung, F., Wu, X., Sun, X., Zheng, H., Lv, Z. and Wu, Z. (2013a). *Angiostrongylus cantonensis*: tegumental and hypodermic alterations of the fourth-stage larvae following administration of tribendimidine *in vivo* and *in vitro*. *Parasitology Research* **112**, 3035–3040.
- Zeng, X., Wei, J., Wang, J., Wu, F., Fung, F., Wu, X., Sun, X., Zheng, H., Lv, Z. and Wu, Z. (2013b). *Angiostrongylus cantonensis*: scanning electron microscopic observations on the cuticle of moulting larvae. *Korean Journal of Parasitology* **51**, 633–636.
- Zhang, R. L., Chen, M. X., Gao, S. T., Geng, Y. J., Huang, D. N., Liu, J. P., Wu, Y. L. and Zhu, X. Q. (2008a). Enzootic angiostrongyliasis in Shenzhen, China. *Emerging Infectious Diseases* **14**, 1955–1956.
- Zhang, Y., Huang, D., Tan, Q. M., Chen, D. X. and Zhan, X. M. (2008b). Epidemiological investigation of *Angiostrongylus cantonensis* in Jiangmen of Guangdong Province. *Zhongguo Ji Sheng Chong Xue Yu Ji Sheng Chong Bing Za Zhi* **26**, 370–373.
- Zhao, J., Lv, Z., Wang, F., Wei, J., Zhang, Q., Li, S., Yang, F., Zeng, X., Wu, X. and Wu, Z. (2013). Ym1, an eosinophilic chemotactic factor, participates in the brain inflammation induced by *Angiostrongylus cantonensis* in mice. *Parasitology Research* **112**, 2689–2695.
- Zhou, Z., Barennes, H., Zhou, N., Ding, L., Zhu, Y. H. and Strobel, M. (2009). Two outbreaks of eosinophilic meningitis in Yunann (China) clinical, epidemiological and therapeutical issues. *Bulletin de la Societe de Pathologie Exotique* **102**, 75–80.