



COMPOSITIO MATHEMATICA

Big principal series, p -adic families and \mathcal{L} -invariants

Lennart Gehrmann and Giovanni Rosso

Compositio Math. **158** (2022), 409–436.

[doi:10.1112/S0010437X2200731X](https://doi.org/10.1112/S0010437X2200731X)



FOUNDATION
COMPOSITIO
MATHEMATICA



LONDON
MATHEMATICAL
SOCIETY
EST. 1865





Big principal series, p -adic families and \mathcal{L} -invariants

Lennart Gehrmann and Giovanni Rosso

ABSTRACT

In earlier work, the first named author generalized the construction of Darmon-style \mathcal{L} -invariants to cuspidal automorphic representations of semisimple groups of higher rank, which are cohomological with respect to the trivial coefficient system and Steinberg at a fixed prime. In this paper, assuming that the Archimedean component of the group has discrete series we show that these automorphic \mathcal{L} -invariants can be computed in terms of derivatives of Hecke eigenvalues in p -adic families. Our proof is novel even in the case of modular forms, which was established by Bertolini, Darmon and Iovita. The main new technical ingredient is the Koszul resolution of locally analytic principal series representations by Kohlhaase and Schraen. As an application of our results we settle a conjecture of Spieß: we show that automorphic \mathcal{L} -invariants of Hilbert modular forms of parallel weight 2 are independent of the sign character used to define them. Moreover, we show that they are invariant under Jacquet–Langlands transfer and, in fact, equal to the Fontaine–Mazur \mathcal{L} -invariant of the associated Galois representation. Under mild assumptions, we also prove the equality of automorphic and Fontaine–Mazur \mathcal{L} -invariants for representations of definite unitary groups of arbitrary rank. Finally, we study the case of Bianchi modular forms to show how our methods, given precise results on eigenvarieties, can also work in the absence of discrete series representations.

Introduction

Let $f = \sum_{n=1}^{\infty} a_n q^n$ be a normalized newform of weight 2 and level $\Gamma_0(M)$ such that $M = pN$ with p prime, $p \nmid N$ and $a_p = 1$. Inspired by Teitelbaum’s work (cf. [Tei90]) on \mathcal{L} -invariants for automorphic forms on definite quaternion groups, Darmon in [Dar01] constructed the automorphic \mathcal{L} -invariant $\mathcal{L}(f)^{\pm}$ of f , depending *a priori* on a choice of sign at infinity. Let $a_p(k)$ be the U_p -eigenvalue of the p -adic family passing through f . In [BDI10] Bertolini, Darmon and Iovita (see also [Das05]) prove the following formula:

$$\mathcal{L}^{\pm}(f) = -2 \frac{da_p}{dk} \Big|_{k=2}. \quad (0.1)$$

In particular, the automorphic \mathcal{L} -invariant is independent of the sign at infinity. They also prove a similar formula for Orton’s \mathcal{L} -invariant (cf. [Ort04]) in higher weight. Moreover, if f admits a

Received 25 June 2020, accepted in final form 11 October 2021, published online 25 April 2022.

2020 Mathematics Subject Classification 11F55 (primary), 11F33, 11F70, 11F75, 11F80 (secondary).

Keywords: automorphic forms, Galois representations, \mathcal{L} -invariants, p -adic families.

While working on this paper the first named author was visiting McGill University, supported by Deutsche Forschungsgemeinschaft, and he would like to thank these institutions. The second named author was partly funded by FRQNT grant 2019-NC-254031 and NSERC grant RGPIN-2018-04392. In addition, the authors would like to thank Henri Darmon, Michael Lipnowski, Vytautas Paškūnas and Chris Williams for several intense and stimulating discussions. We also thank the referee for their careful reading of the paper and many suggestions.

© 2022 The Author(s). The publishing rights in this article are licensed to Foundation Compositio Mathematica under an exclusive licence.

Jacquet–Langlands transfer $JL(f)$ to a definite quaternion group, which is split at p , they show the analogous formula

$$\mathcal{L}_T(JL(f)) = -2 \frac{da_p}{dk} \Big|_{k=2}$$

for Teitelbaum’s \mathcal{L} -invariant, thus proving that automorphic \mathcal{L} -invariants are preserved under Jacquet–Langlands transfers. This result was extended to Jacquet–Langlands transfers to indefinite quaternion groups over the rationals, which are split at p , by Dasgupta and Greenberg [DG12], Longo, Rotger and Vigni [LRV12] and Seveso [Sev13].

Over the last few years the construction of automorphic \mathcal{L} -invariants was generalized to various settings, e.g. to Hilbert modular cusp forms of parallel weight 2 by Spieß (see [Spi14]) and to Bianchi modular cusp forms of even weight by Barrera and Williams (see [BW19]). These generalizations have been defined as automorphic \mathcal{L} -invariants appear naturally in many context, from proving exceptional zero conjecture formulae to constructing and studying Stark–Heegner points.

Most recently, the first named author defined automorphic \mathcal{L} -invariants for certain cuspidal automorphic representations of higher rank semi-simple groups over a number field F , which are split at a fixed prime \mathfrak{p} of F (see [Geh21]). The most crucial assumptions on the representation π are that it is cohomological with respect to the trivial coefficient system and that the local factor $\pi_{\mathfrak{p}}$ of π at \mathfrak{p} is the Steinberg representation. Our main aim is to prove the analogue of equation (0.1) for these \mathcal{L} -invariants. Previous works used explicit computations with cocycles and it seems unlikely that one can generalize these to higher rank groups; instead, we give a new, more conceptual approach, which is novel even in the already known cases.

In §1 we recall the definition of automorphic \mathcal{L} -invariants. Just as in the case of modular forms these \mathcal{L} -invariants depend on a choice of sign character at infinity. They also depend on a choice of degree of cohomology, in which the representation occurs. For this introduction, we suppress it because we are mostly interested in the case that there is only one interesting degree. Given a simple root i of the group and a sign character ϵ the space of \mathcal{L} -invariants

$$\mathcal{L}_i(\pi, \mathfrak{p})^\epsilon \subset \text{Hom}^{\text{ct}}(F_{\mathfrak{p}}^*, E)$$

is a subspace of codimension at least one. Here E denotes a large enough p -adic field. If strong multiplicity one holds, its codimension is exactly one. Whether a character of $F_{\mathfrak{p}}^*$ belongs to the space of \mathcal{L} -invariants is decided by certain maps between cohomology groups of \mathfrak{p} -arithmetic subgroups with values in duals of (locally analytic) generalized Steinberg representations. These maps are induced by cup products with one-extensions of the smooth generalized Steinberg representation corresponding to the simple root i with the locally analytic Steinberg representations (see §1.1 for a description of these extensions due to Ding, cf. [Din19]).

As a first step, in §2.2 we show that one can replace generalized Steinberg representations by locally analytic principal series representations and the extension classes by infinitesimal deformations of these principal series. As a consequence, we prove an automorphic analogue of the Colmez–Greenberg–Stevens formula (see Proposition 2.4). Let us remind ourselves that the Colmez–Greenberg–Stevens formula (see, for example, Theorem 3.4 of [Din19]) states that one can compute Fontaine–Mazur \mathcal{L} -invariants of Galois representations by deforming them in p -adic families. Our analogue states that one can compute automorphic \mathcal{L} -invariants from the cohomology of \mathfrak{p} -arithmetic groups with values in duals of big principal series representations, i.e. parabolic inductions of characters with values in units of affinoid algebras.

Thus, we reduce the problem to producing classes in these big cohomology groups. Here is where we need to impose further restrictions. First, we assume that the group under consideration is adjoint. Under this assumption Kohlhaase and Schraen constructed a Koszul resolution of

locally analytic principal series representations (cf. [KS12]), which we recall in §3.1. Using that resolution we can lift overconvergent cohomology classes which are common eigenvectors for all $U_{\mathfrak{p}}$ -operators to big cohomology classes. Second, in order to have a nice enough theory of families of overconvergent cohomology classes (in the spirit of [AS08]) we consider only groups whose Archimedean component fulfils the Harish-Chandra condition. This implies that the automorphic representation we study only shows up in the middle degree cohomology of the associated locally symmetric space. We further assume that the map from the eigenvariety to the weight space \mathcal{W} is étale at the point corresponding to the automorphic representation π . Étaleness is implied by a suitable strong multiplicity one assumption and, thus, holds for example for Hilbert modular forms. Under these hypotheses we can show that we can lift the cohomology class corresponding to π to a big cohomology class valued in functions on an open affinoid neighbourhood of the trivial character in weight space (see Theorem 3.12). This allows us give the generalization of (0.1) in Theorem 3.16. In particular, we see that automorphic \mathcal{L} -invariants are codimension one subspaces under our étaleness assumption.

As a first application, in §4.1 we prove a conjecture of Spieß (cf. [Spi14, Conjecture 6.4]): we show that the \mathcal{L} -invariants of Hilbert modular forms of parallel weight 2 are independent of the sign character used to define them. We further show that in this situation automorphic \mathcal{L} -invariants are invariant under Jacquet–Langlands transfers to quaternion groups which are split at \mathfrak{p} . In fact, we show that all these \mathcal{L} -invariants agree with the Fontaine–Mazur \mathcal{L} -invariant of the associated Galois representation.¹ In particular, one can remove the assumptions in the main theorem of [Spi14] (see Theorem 6.10(b) there). Furthermore, the equality of automorphic and Fontaine–Mazur \mathcal{L} -invariants makes the construction of Stark–Heegner points for modular elliptic curves over totally real fields unconditional. Similarly, we prove the equality of automorphic and Fontaine–Mazur \mathcal{L} -invariants for definite unitary groups under mild assumptions in §4.2, i.e. we give an alternative proof of the main result of [Din19] for our global situation.

We end the paper by considering the easiest case of a group that does not fulfil the Harish-Chandra condition, i.e. we study Bianchi modular forms. As we cannot deform the automorphic representation over the whole weight space, in general one cannot compute the \mathcal{L} -invariant completely in terms of derivatives of Hecke-eigenvalues. But at least in case the Bianchi forms is the base change of a modular form, we overcome this problem; we show that the \mathcal{L} -invariants of the base change equal the \mathcal{L} -invariant of the modular form.

The assumption that the coefficient system is trivial is not necessary for the arguments of this article. However, the construction of automorphic \mathcal{L} -invariants relies on the existence of a well-behaved lattice in the Steinberg representation, which is not known for twists of the Steinberg by an algebraic representation in general. In case the existence of such a lattice is known, e.g. if G is a form of PGL_2 by a result of Vignéras (see [Vig08]), one can easily generalize our results to arbitrary weights and non-critical slopes.

Notation

If X and Y are topological spaces, we write $C(X, Y)$ for the set of continuous functions from X to Y . All rings are assumed to be commutative and unital. The group of invertible elements of a ring R will be denoted by R^* . If M is an R -module we denote the q th exterior power of M by $\Lambda_R^q M$. If R is a ring and G a group, we denote the group ring of G over R by $R[G]$. Given topological groups H and G we write $\mathrm{Hom}^{\mathrm{ct}}(H, G)$ for the space of continuous homomorphism from H to G . Let $\chi: G \rightarrow R^*$ be a character. We write $R[\chi]$ for the G -representation, which

¹ The same result has been obtained by Spieß by different methods; see [Spi20].

underlying R -module is R itself and on which G acts via the character χ . The trivial character of G will be denoted by $\mathbb{1}_G$. Let H be an open subgroup of a locally profinite group G and M an R -linear representation M of H . The compact induction $c\text{-ind}_H^G M$ of M from H to G is the space of all functions $f: G \rightarrow M$ which have finite support modulo H and satisfy $f(gh) = h^{-1}.f(g)$ for all $h \in H, g \in G$.

The setup

We fix an algebraic number field F . In addition, we fix a finite place \mathfrak{p} of F lying above the rational prime p and choose embeddings

$$\overline{\mathbb{Q}_p} \hookrightarrow \overline{\mathbb{Q}} \hookrightarrow \mathbb{C}.$$

If v is a place of F , we denote by F_v the completion of F at v . If v is a finite place, we let \mathcal{O}_v denote the valuation ring of F_v and ord_v the additive valuation such that $\text{ord}_v(\varpi) = 1$ for any local uniformizer $\varpi \in \mathcal{O}_v$.

Let \mathbb{A} be the adèle ring of F , i.e. the restricted product over all completions F_v of F . We write \mathbb{A}^∞ (respectively, $\mathbb{A}^{\mathfrak{p},\infty}$) for the restricted product over all completions of F at finite places (respectively, finite places different from \mathfrak{p}). More generally, if S is a finite set of places of F we denote by \mathbb{A}^S the restricted product of all completions F_v with $v \notin S$.

If H is an algebraic group over F and v is a place of F , we write $H_v = H(F_v)$. We put $H_\infty = \prod_{v|\infty} H_v$.

Throughout the article we fix a connected, adjoint, semi-simple algebraic group G over F . We assume that the base change $G_{F_{\mathfrak{p}}}$ of G to $F_{\mathfrak{p}}$ is split. Let $K_\infty \subseteq G_\infty$ denote a fixed maximal compact subgroup. The integers δ and q are defined via

$$\delta = \text{rk } G_\infty - \text{rk } K_\infty$$

and

$$2q + \delta = \dim G_\infty - \dim K_\infty.$$

Finally, we fix a cuspidal automorphic representation $\pi = \otimes_v \pi_v$ of $G(\mathbb{A})$ with the following properties:

- (i) π is cohomological with respect to the trivial coefficient system;
- (ii) π is tempered at ∞ ; and
- (iii) $\pi_{\mathfrak{p}}$ is the (smooth) Steinberg representation $\text{St}_{G_{\mathfrak{p}}}^\infty(\mathbb{C})$ of $G_{\mathfrak{p}}$.

We denote by $\mathbb{Q}_\pi \subseteq \overline{\mathbb{Q}}$ a fixed finite extension of \mathbb{Q} over which π^∞ has a model (see Theorem C of [Jan18] for the existence of such an extension).

HYPOTHESIS (SMO). We assume that the following strong multiplicity one hypothesis on π holds. If π' is an automorphic representation of G such that:

- (i) $\pi'_v \cong \pi_v$ for all finite places $v \neq \mathfrak{p}$;
- (ii) $\pi'_\mathfrak{p}$ has an Iwahori-invariant vector; and
- (iii) π'_∞ has non-vanishing $(\mathfrak{g}, K_\infty^\circ)$ -cohomology;

then π is cuspidal, $\pi'_\mathfrak{p} \cong \pi_\mathfrak{p}$ and π_∞ is tempered.

It is known that this holds for cuspidal representations of GL_n by work of Jacquet and Shalika [JS81b, JS81a] and Piatetski-Shapiro [Pia79] and thus, in particular, for representations of $G = \text{PGL}_n$.

For general groups strong multiplicity one fails, see, for example, [HP83]. Hence, it is harder to find explicit results about representations π for which the hypothesis SMO holds, but nevertheless

it is expected that SMO should hold in many cases; for example, for generic representation of GSp_4 [Sou87]. The same strategy of [Sou87] could apply as long as we have an injective transfer from (a subclass of) representations of G to representation of GL_n , e.g. tempered representations of unitary groups that at each prime are not endoscopic.

1. Automorphic \mathcal{L} -invariants

In the following, we briefly sketch the construction of automorphic \mathcal{L} -invariants. For more details, see [Geh21].

1.1 Extensions

In this section we recall the computation of certain Ext^1 -groups of (locally analytic) generalized Steinberg representations due to Ding (see [Din19]). We fix a finite extension E of \mathbb{Q}_p . If V and W are admissible locally \mathbb{Q}_p -analytic E -representations of $G_{\mathfrak{p}}$, we write $\mathrm{Ext}_{\mathrm{an}}^1(V, W)$ for the group of locally analytic extensions of V by W .

Given an algebraic subgroup $H \subseteq G_{F_{\mathfrak{p}}}$, we denote the group of $F_{\mathfrak{p}}$ -valued points of H also by H . We fix a Borel subgroup B of the split group $G_{F_{\mathfrak{p}}}$ and a maximal split torus $T \subseteq B$ and denote by Δ the associated basis of simple roots. For a subset $I \subseteq \Delta$ we let $P_I \supseteq B$ be the corresponding parabolic containing B .

Suppose M is a smooth representation of P_I over a ring R ; we define its smooth induction to $G_{\mathfrak{p}}$ as

$$i_{P_I}^{\infty}(M) = \{f: G_{\mathfrak{p}} \rightarrow M \text{ locally constant} \mid f(pg) = p.f(g) \forall p \in P_I, g \in G_{\mathfrak{p}}\}.$$

The generalized R -valued (smooth) Steinberg representation associated with $I \subseteq \Delta$ is given by the quotient

$$v_{P_I}^{\infty}(R) = i_{P_I}^{\infty}(R) \Big/ \sum_{I \subset J \subset \Delta, I \neq J} i_{P_J}^{\infty}(R).$$

Likewise, if V is a locally \mathbb{Q}_p -analytic E -representation of P_I , we define its locally analytic induction to $G_{\mathfrak{p}}$ as the space of functions

$$\mathbb{I}_{P_I}^{\mathrm{an}}(V) = \{f: G_{\mathfrak{p}} \rightarrow \tau \text{ locally } \mathbb{Q}_p\text{-analytic} \mid f(pg) = p.f(g) \forall p \in P_I, g \in G_{\mathfrak{p}}\}.$$

We define the locally analytic generalized Steinberg representation with respect to I as the quotient

$$\mathbb{V}_{P_I}^{\mathrm{an}}(E) = \mathbb{I}_{P_I}^{\mathrm{an}}(E) \Big/ \sum_{I \subset J \subset \Delta, I \neq J} \mathbb{I}_{P_J}^{\mathrm{an}}(E).$$

We put $\mathrm{St}_{G_{\mathfrak{p}}}^{\infty}(R) = v_B^{\infty}(R)$ and $\mathrm{St}_{G_{\mathfrak{p}}}^{\mathrm{an}}(E) = \mathbb{V}_B^{\mathrm{an}}(E)$. Similarly, replacing locally analyticity with continuity we define the continuous Steinberg representation $\mathrm{St}_{G_{\mathfrak{p}}}^{\mathrm{ct}}(E)$ and, more generally, $\mathbb{V}_{P_I}^{\mathrm{ct}}(E)$. It is easy to see that $\mathbb{V}_{P_I}^{\mathrm{ct}}(E)$ is the universal unitary completion of both $v_{P_I}^{\infty}(E)$ and $\mathbb{V}_{P_I}^{\mathrm{an}}(E)$.

Let $i \in \Delta$ be a simple root and $\lambda \in \mathrm{Hom}^{\mathrm{ct}}(B, E)$ a continuous homomorphism. Note that λ is automatically locally analytic and is trivial on the unipotent radical of B . Thus, it can be identified with a character on T . We write τ_{λ} for the two-dimensional representation of B given by

$$\tau_{\lambda}(b) = \begin{pmatrix} 1 & \lambda(b) \\ 0 & 1 \end{pmatrix}. \tag{1.1}$$

By the exactness of the parabolic induction functor for locally analytic extensions (see [Koh11, Proposition 5.1 and Remark 5.4]) we have a short exact sequence of the form

$$0 \longrightarrow \mathbb{I}_B^{\text{an}}(E) \longrightarrow \mathbb{I}_B^{\text{an}}(\tau\lambda) \longrightarrow \mathbb{I}_B^{\text{an}}(E) \longrightarrow 0.$$

We write

$$\widehat{\mathcal{E}}^{\text{an}}(\lambda) \in \text{Ext}_{\text{an}}^1(\mathbb{I}_B^{\text{an}}(E), \mathbb{I}_B^{\text{an}}(E)) \tag{1.2}$$

for the associated extension class. Further, we define the class

$$\widetilde{\mathcal{E}}_i^{\text{an}}(\lambda) \in \text{Ext}_{\text{an}}^1(i_{P_i}^\infty(E), \mathbb{I}_B^{\text{an}}(E)) \tag{1.3}$$

as the pullback of this extension along $i_{P_i}^\infty(E) \rightarrow \mathbb{I}_B^{\text{an}}(E)$. Finally, taking pushforward along $\mathbb{I}_B^{\text{an}}(E) \rightarrow \text{St}_{G_p}^{\text{an}}(E)$ yields the extension class

$$\mathcal{E}_i^{\text{an}}(\lambda) \in \text{Ext}_{\text{an}}^1(i_{P_i}^\infty(E), \text{St}_{G_p}^{\text{an}}). \tag{1.4}$$

By an easy calculation we see that the map

$$\text{Hom}^{\text{ct}}(B, E) \longrightarrow \text{Ext}_{\text{an}}^1(i_{P_i}^\infty(E), \text{St}_{G_p}^{\text{an}}(E)), \quad \lambda \longmapsto \mathcal{E}_i^{\text{an}}(\lambda)$$

defines a homomorphism. The inclusion $B \hookrightarrow P_i$ induces an injection

$$\text{Hom}^{\text{ct}}(P_i, E) \longrightarrow \text{Hom}^{\text{ct}}(B, E).$$

The quotient can be identified with the space $\text{Hom}^{\text{ct}}(F_p^*, E)$ via the map

$$\text{Hom}^{\text{ct}}(F_p^*, E) \longrightarrow \text{Hom}^{\text{ct}}(B, E) / \text{Hom}^{\text{ct}}(P_i, E), \quad \lambda \longmapsto \lambda \circ i. \tag{1.5}$$

Alternatively, let i^\vee denote the coroot associated with i . Then, the kernel of the map

$$\text{Hom}^{\text{ct}}(B, E) \longrightarrow \text{Hom}^{\text{ct}}(F_p^*, E), \quad \lambda \longmapsto \lambda_i = \lambda \circ i^\vee \tag{1.6}$$

is equal to $\text{Hom}^{\text{ct}}(P_i, E)$ and, hence, the map induces an isomorphism

$$\text{Hom}^{\text{ct}}(B, E) / \text{Hom}^{\text{ct}}(P_i, E) \longrightarrow \text{Hom}^{\text{ct}}(F_p^*, E),$$

which is the inverse of the isomorphism above up to multiplication by two.

THEOREM 1.1 (Ding). *The following hold.*

- (i) *The map $\text{Hom}^{\text{ct}}(B, E) \rightarrow \text{Ext}_{\text{an}}^1(i_{P_i}^\infty(E), \text{St}_{G_p}^{\text{an}}(E))$, $\lambda \mapsto \mathcal{E}_i^{\text{an}}(\lambda)$ is surjective with kernel $\text{Hom}^{\text{ct}}(P_i, E) \subseteq \text{Hom}^{\text{ct}}(B, E)$.*
- (ii) *The canonical map $\text{Ext}_{\text{an}}^1(v_{P_i}^\infty(E), \text{St}_{G_p}^{\text{an}}(E)) \rightarrow \text{Ext}_{\text{an}}^1(i_{P_i}^\infty(E), \text{St}_{G_p}^{\text{an}}(E))$ is an isomorphism.*
- (iii) *The induced map $\text{Hom}^{\text{ct}}(F_p^*, E) \rightarrow \text{Ext}_{\text{an}}^1(v_{P_i}^\infty(E), \text{St}_{G_p}^{\text{an}}(E))$, $\lambda \mapsto \mathcal{E}_i^{\text{an}}(\lambda \circ i)$ is an isomorphism.*

Proof. The third claim is a direct consequence of the first two. For the proof of the first two claims in the case $G = \text{PGL}_n$, see § 2.2 of [Din19]. The general case is proven in § 2.4 of [Geh21]. □

1.2 Flawless lattices

We recall the notion of flawless smooth representations over a ring R .

DEFINITION 1.2. A smooth R -representation M of G_p is called flawless if:

- (i) M is projective as an R -module; and

(ii) there exists a finite length exact resolution

$$0 \longrightarrow C_m \longrightarrow \cdots \longrightarrow C_0 \longrightarrow M \longrightarrow 0$$

of M by smooth R -representations C_i of G_p , where each C_i is a finite direct sum of modules of the form

$$\text{c-ind}_{K_p}^{G_p}(L)$$

with $K_p \subseteq G_p$ a compact, open subgroup and L a smooth representation of K_p that is finitely generated projective over R .

The following is our main example.

THEOREM 1.3 (Borel–Serre). *The Steinberg representation $\text{St}_{G_p}^\infty(R)$ is flawless for every ring R .*

Proof. It is enough to prove that $\text{St}_{G_p}^\infty(\mathbb{Z})$ is flawless. By [BS76, Theorem 5.6], $\text{St}_{G_p}^\infty(\mathbb{Z})$ can be identified with the cohomology with compact supports of the Bruhat–Tits building of G_p . Thus, its simplicial complex gives a flawless resolution of $\text{St}_{G_p}^\infty(\mathbb{Z})$. \square

1.3 Cohomology of p -arithmetic groups

Let A be a \mathbb{Q}_p -affinoid algebra in the sense of Tate. Given a compact, open subgroup $K^p \subseteq G(\mathbb{A}^{p,\infty})$, an $A[G_p]$ -module V and an $A[G(F)]$ -module W , which is free and of finite rank over A , we define $\mathcal{C}_A(K^p, V; W)$ as the space of all A -linear maps $\Phi: G(\mathbb{A}^{p,\infty})/K^p \times V \rightarrow W$. The A -module $\mathcal{C}_E(K^p, V; W)$ carries a natural $G(F)$ -action given by

$$(\gamma \cdot \Phi)(g, v) = \gamma \cdot (\Phi(\gamma^{-1}g, \gamma^{-1} \cdot v)).$$

Suppose V is a topological A -module equipped with a continuous A -linear G_p -action we put $\mathcal{C}_A^{\text{ct}}(K^p, V; W) = C(G(\mathbb{A}^{p,\infty})/K^p, \text{Hom}_A^{\text{ct}}(V, W))$. Here W is endowed with its canonical topology as a finitely generated free A -module.

Let E be a finite extension of \mathbb{Q}_p with ring of integers \mathcal{O}_E . Suppose that V is a smooth E -representation of G_p that admits a flawless G_p -stable \mathcal{O}_E -lattice M . As M is finitely generated as an $\mathcal{O}_E[G_p]$ -module, the completion of V with respect to M is the universal unitary completion V^{un} of V . The following automatic continuity statement holds (see [Geh21, Proposition 3.11]).

PROPOSITION 1.4. *Suppose that V is a smooth E -representation of G_p that admits a flawless G_p -stable \mathcal{O}_E -lattice M . Then the canonical map*

$$\text{H}^d(G(F), \mathcal{C}_E^{\text{ct}}(K^p, V^{\text{un}}; E(\epsilon))) \longrightarrow \text{H}^d(G(F), \mathcal{C}_E(K^p, V; E(\epsilon)))$$

is an isomorphism for every character $\epsilon: \pi_0(G_\infty) \rightarrow \{\pm 1\}$, every compact, open subgroup $K^p \subseteq G(\mathbb{A}^{p,\infty})$ and every degree $d \geq 0$.

This proposition combined with Theorem 1.3 implies the following.

COROLLARY 1.5. *The canonical map*

$$\text{H}^d(G(F), \mathcal{C}_E^{\text{ct}}(K^p, \text{St}_{G_p}^{\text{ct}}(E); E(\epsilon))) \longrightarrow \text{H}^d(G(F), \mathcal{C}_E(K^p, \text{St}_{G_p}^\infty(E); E(\epsilon)))$$

is an isomorphism.

For a compact, open subset $K_p \subseteq G_p$ and a character $\epsilon: \pi_0(G_\infty) \rightarrow \{\pm 1\}$ we put

$$\text{H}^d(X_{K^p \times K_p}, W)^\epsilon = \text{H}^d(G(F), \mathcal{C}_E(K^p, E[G_p/K_p]; W(\epsilon))).$$

If the level $K^p \times K_p$ is neat, this group is naturally isomorphic to (the epsilon component of) the singular cohomology with coefficients in W of the locally symmetric space of level $K^p \times K_p$

associated with G . More generally, let V be a topological A -module, which is locally convex as a \mathbb{Q}_p -vector space, with a continuous A -linear $K_{\mathfrak{p}}$ -action. We consider $\text{c-ind}_{K_{\mathfrak{p}}}^{G_{\mathfrak{p}}} V$ with the locally convex inductive limit topology. This is a continuous A -module with a continuous G_p -action. We put

$$\mathrm{H}^d(X_{K^{\mathfrak{p}} \times K_{\mathfrak{p}}}, \mathrm{Hom}_A^{\mathrm{ct}}(V, W))^{\epsilon} = \mathrm{H}^d(G(F), \mathcal{C}_A^{\mathrm{ct}}(K^{\mathfrak{p}}, \text{c-ind}_{K_{\mathfrak{p}}}^{G_{\mathfrak{p}}} V; W(\epsilon))).$$

Again, this space can be identified with the cohomology of the corresponding locally symmetric space with values in the sheaf associated with $\mathrm{Hom}_A^{\mathrm{ct}}(V, A)$.

1.4 Automorphic \mathcal{L} -invariants

For the remainder of the article we fix a finite extension E of \mathbb{Q}_p that contains \mathbb{Q}_{π} and a compact, open subgroup $K^{\mathfrak{p}} = \prod_{v|\mathfrak{p}\infty} K_v \subseteq G(\mathbb{A}^{\mathfrak{p},\infty})$ such that $(\pi^{\mathfrak{p},\infty})^{K^{\mathfrak{p}}} \neq 0$. We may assume that K_v is hyperspecial for every finite place v such that π_v is spherical.

As $v_{P_I}^{\infty}(E) = v_I^{\infty}(\mathbb{Z}) \otimes E$ we have a canonical isomorphism

$$\mathcal{C}_{\mathbb{Z}}(K^{\mathfrak{p}}, v_{P_I}^{\infty}(\mathbb{Z}); W) \cong \mathcal{C}_E(K^{\mathfrak{p}}, v_{P_I}^{\infty}(E); W)$$

for any E -vector space W . Hence, we abbreviate this space by $\mathcal{C}(K^{\mathfrak{p}}, v_{P_I}^{\infty}; W)$ (and similarly for $\mathrm{St}_{G_{\mathfrak{p}}}^{\infty}$ in place of $v_{P_I}^{\infty}$).

Let $I_{\mathfrak{p}} \subseteq G_{\mathfrak{p}}$ be an Iwahori subgroup. By Frobenius reciprocity the choice of an Iwahori-fixed vector yields a $G_{\mathfrak{p}}$ -equivariant homomorphism

$$\text{c-ind}_{I_{\mathfrak{p}}}^{G_{\mathfrak{p}}} E \longrightarrow \mathrm{St}_{G_{\mathfrak{p}}}^{\infty}, \tag{1.7}$$

which, in turn, induces a Hecke-equivariant map

$$\mathrm{ev}^{(d)}: \mathrm{H}^d(G(F), \mathcal{C}(K^{\mathfrak{p}}, \mathrm{St}_{G_{\mathfrak{p}}}^{\infty}; E(\epsilon))) \longrightarrow \mathrm{H}^d(X_{K^{\mathfrak{p}} \times I_{\mathfrak{p}}}, E)^{\epsilon}. \tag{1.8}$$

Let

$$\mathbb{T} = \mathbb{T}(K^{\mathfrak{p}} \times I_{\mathfrak{p}})_E = C_c(K^{\mathfrak{p}} \times I_{\mathfrak{p}} \backslash G(\mathbb{A}^{\infty}) / K^{\mathfrak{p}} \times I_{\mathfrak{p}}, E)$$

be the E -valued Hecke algebra of level $K^{\mathfrak{p}} \times I_{\mathfrak{p}}$. By abuse of notation, we denote the model of π^{∞} over E also by π^{∞} . If V is a \mathbb{T} -module, we put

$$V[\pi] = \sum_f \mathrm{im}(f)$$

where we sum over all \mathbb{T} -homomorphisms $f: (\pi^{\infty})^{K^{\mathfrak{p}} \times I_{\mathfrak{p}}} \rightarrow V$. Similarly, we define

$$\mathbb{T}^{\mathfrak{p}} = \mathbb{T}(K^{\mathfrak{p}})_E = C_c(K^{\mathfrak{p}} \backslash G(\mathbb{A}^{\mathfrak{p},\infty}) / K^{\mathfrak{p}}, E)$$

to be the Hecke algebra away from \mathfrak{p} and, given a $\mathbb{T}^{\mathfrak{p}}$ -module V , we put

$$V[\pi^{\mathfrak{p}}] = \sum_f \mathrm{im}(f)$$

where we sum over all $\mathbb{T}^{\mathfrak{p}}$ -homomorphisms $f: (\pi^{\mathfrak{p},\infty})^{K^{\mathfrak{p}}} \rightarrow V$.

For the proof of the following proposition that crucially relies on the hypothesis (SMO) we refer to [Geh21, Proposition 3.6].

PROPOSITION 1.6. *The map $\mathrm{ev}^{(d)}$ induces an isomorphism*

$$\mathrm{H}^d(G(F), \mathcal{C}(K^{\mathfrak{p}}, \mathrm{St}_{G_{\mathfrak{p}}}^{\infty}; E(\epsilon)))[\pi^{\mathfrak{p}}] \xrightarrow{\cong} \mathrm{H}^d(X_{K^{\mathfrak{p}} \times I_{\mathfrak{p}}}, E)^{\epsilon}[\pi].$$

on isotypic components. There exists an integer $m_{\pi} \geq 0$ such that

$$\dim_E \mathrm{H}^d(X_{K^{\mathfrak{p}} \times I_{\mathfrak{p}}}, E)^{\epsilon}[\pi] = m_{\pi} \cdot \dim(\pi^{\infty})^{K^{\mathfrak{p}} \times I_{\mathfrak{p}}} \cdot \binom{\delta}{d-q} \forall d \geq 0$$

for each sign character ϵ .

By Corollary 1.5, we have a canonical isomorphism

$$H^d(G(F), \mathcal{C}(K^{\mathfrak{p}}, \text{St}_{G_{\mathfrak{p}}}^{\infty}; E(\epsilon))) \cong H^d(G(F), \mathcal{C}_E^{\text{ct}}(K^{\mathfrak{p}}, \text{St}_{G_{\mathfrak{p}}}^{\text{ct}}; E(\epsilon))).$$

Composing with the homomorphism coming from dualizing the continuous inclusion $\text{St}_{G_{\mathfrak{p}}}^{\text{an}} \hookrightarrow \text{St}_{G_{\mathfrak{p}}}^{\text{ct}}$ yields the map

$$H^d(G(F), \mathcal{C}(K^{\mathfrak{p}}, \text{St}_{G_{\mathfrak{p}}}^{\infty}; E(\epsilon))) \longrightarrow H^d(G(F), \mathcal{C}_E^{\text{ct}}(K^{\mathfrak{p}}, \text{St}_{G_{\mathfrak{p}}}^{\text{an}}; E(\epsilon)))$$

in cohomology. Thus, for every $i \in \Delta$ we get a well-defined cup-product pairing

$$\begin{aligned} H^d(G(F), \mathcal{C}(K^{\mathfrak{p}}, \text{St}_{G_{\mathfrak{p}}}^{\infty}; E(\epsilon))) \times \text{Ext}_{\text{an}}^1(v_{P_i}^{\infty}(E), \text{St}_{G_{\mathfrak{p}}}^{\text{an}}(E)) \\ \longrightarrow H^{d+1}(G(F), \mathcal{C}(K^{\mathfrak{p}}, v_{P_i}^{\infty}; E(\epsilon))), \end{aligned}$$

which commutes with the action of the Hecke algebra $\mathbb{T}^{\mathfrak{p}}$. By Theorem 1.1 we have a canonical isomorphism $\text{Hom}^{\text{ct}}(F_{\mathfrak{p}}^*, E) \cong \text{Ext}_{\text{an}}^1(v_{P_i}^{\infty}(E), \text{St}_{G_{\mathfrak{p}}}^{\text{an}}(E))$. Hence, taking cup product with the extension $\mathcal{E}_i^{\text{an}}(\lambda \circ i)$ associated with a homomorphism $\lambda \in \text{Hom}^{\text{ct}}(F_{\mathfrak{p}}^*, E)$ in (1.4) yields a map

$$c_i^{(d)}(\lambda)^{\epsilon}: H^d(G(F), \mathcal{C}(K^{\mathfrak{p}}, \text{St}_{G_{\mathfrak{p}}}^{\infty}; E(\epsilon)))[\pi^{\mathfrak{p}}] \longrightarrow H^{d+1}(G(F), \mathcal{C}(K^{\mathfrak{p}}, v_{P_i}^{\infty}; E(\epsilon)))[\pi^{\mathfrak{p}}]$$

on $\pi^{\mathfrak{p}, \infty}$ -isotypic parts.

DEFINITION 1.7. Given a character $\epsilon: \pi_0(G_{\infty}) \rightarrow \{\pm 1\}$, an integer $d \in \mathbb{Z}$ with $0 \leq d \leq \delta$ and a root $i \in \Delta$ we define

$$\mathcal{L}_i^{(d)}(\pi, \mathfrak{p})^{\epsilon} \subseteq \text{Hom}^{\text{ct}}(F_{\mathfrak{p}}^*, E)$$

as the kernel of the map $\lambda \mapsto c_i^{(q+d)}(\lambda)^{\epsilon}$.

Alternatively, we can consider cup products with extensions associated with homomorphisms $\lambda: B \rightarrow E$ and define the \mathcal{L} -invariant as a subspace of

$$\text{Hom}^{\text{ct}}(B, E) / \text{Hom}^{\text{ct}}(P_i, E).$$

This subspace is mapped to the \mathcal{L} -invariant defined above via the map (1.6) induced by the coroot i^{\vee} associated with i .

PROPOSITION 1.8. For all sign characters ϵ , every degree $d \in [0, \delta] \cap \mathbb{Z}$ and every root $i \in \Delta$ the \mathcal{L} -invariant

$$\mathcal{L}_i^{(d)}(\pi, \mathfrak{p})^{\epsilon} \subseteq \text{Hom}^{\text{ct}}(F_{\mathfrak{p}}^*, E)$$

is a subspace of codimension at least one, which does not contain the space of smooth homomorphisms.

Suppose $m_{\pi} = 1$. Then, in the extremal cases $d = 0$ and $d = \delta$ the codimension is exactly one.

Proof. This is Proposition 3.14 of [Geh21]. □

Remark 1.9. Let $\log_p: E^* \rightarrow E$ denote the branch of the p -adic logarithm such that $\log_p(p) = 0$. Let us assume that E contains the images of all embeddings $\sigma: F_{\mathfrak{p}} \rightarrow \overline{\mathbb{Q}_p}$. We put $\log_{p, \sigma} = \log_p \circ \sigma: F_{\mathfrak{p}}^* \rightarrow E$. The set

$$\{\text{ord}_{\mathfrak{p}}\} \cup \{\log_{p, \sigma} \mid \sigma: F_{\mathfrak{p}} \rightarrow E\}$$

is a basis of $\text{Hom}^{\text{ct}}(F_{\mathfrak{p}}^*, E)$. Suppose $\mathcal{L} \subseteq \text{Hom}^{\text{ct}}(F_{\mathfrak{p}}^*, E)$ is a subspace of codimension one that does not contain $\text{ord}_{\mathfrak{p}}$. Then for each embedding σ there exists a unique element $\mathcal{L}^{\sigma} \in E$ such that $\log_{p, \sigma} - \mathcal{L}^{\sigma} \text{ord}_{\mathfrak{p}} \in \mathcal{L}$ and these elements clearly form a basis of \mathcal{L} .

2. \mathcal{L} -invariants and big principal series

By the Colmez–Greenberg–Stevens formula (see [Din19, Theorem 3.4]) one can calculate Fontaine–Mazur \mathcal{L} -invariants of certain local Galois representations by deforming them (or rather the attached (φ, Γ) -modules) in rigid analytic families. The goal of this section is to prove an automorphic version of that formula.

This implies that \mathcal{L} -invariants can be related to the existence of cohomology classes with values in (duals of) families of principal series.

2.1 Linear algebra over the dual numbers

Let R be a ring and $R[\varepsilon] = R[X]/X^2$ the ring of dual numbers over R . Let M be an $R[\varepsilon]$ -module. We define several maps on dual spaces associated with M . First, multiplication with ε induces the map $\mu: M/\varepsilon M \rightarrow \varepsilon M \hookrightarrow M$. We denote by

$$\mu_\varepsilon^*: \text{Hom}_R(M, R) \longrightarrow \text{Hom}_R(M/\varepsilon M, R)$$

its R -dual. Second, reducing modulo ε yields the map

$$\text{red}: \text{Hom}_{R[\varepsilon]}(M, R[\varepsilon]) \longrightarrow \text{Hom}_R(M/\varepsilon M, R).$$

Finally, the map $\text{add}: R[\varepsilon] \rightarrow R, a + b\varepsilon \mapsto a + b$ induces the map

$$\text{add}_*: \text{Hom}_{R[\varepsilon]}(M, R[\varepsilon]) \longrightarrow \text{Hom}_R(M, R).$$

The following easy computation is left to the reader.

LEMMA 2.1. *The maps μ_ε^* , red and add are functorial in M and the following equality holds: $\text{red} = \mu_\varepsilon^* \circ \text{add}_*$.*

2.2 Infinitesimal deformations of principal series

Let $T \subseteq B \subset G_{F_p}$ be the maximal torus respectively the Borel subgroup chosen in § 1.1. The map

$$\tau: \text{Hom}^{\text{ct}}(B, E) \longrightarrow \text{Hom}^{\text{ct}}(B, E[\varepsilon]^*), \quad \lambda \longmapsto [x \mapsto 1 + \lambda(x)\varepsilon]$$

defines an injective group homomorphism. Its image is the set of all continuous characters $\chi: B \rightarrow E[\varepsilon]^*$ such that $\chi \equiv 1 \pmod{\varepsilon}$. The underlying E -representation of $\tau(\lambda)$ is the two-dimensional representation τ_λ defined in (1.1). Thus, we can view $\mathbb{I}_B^{\text{an}}(\tau_\lambda) = \mathbb{I}_B^{\text{an}}(\tau(\lambda))$ as an $E[\varepsilon]$ -representation of G_p . Reducing modulo ε induces the map

$$\text{red}_\lambda^{d,\varepsilon}: \text{H}^d(G(F), \mathcal{C}_{E[\varepsilon]}^{\text{ct}}(K^{\mathfrak{p}}, \mathbb{I}_B^{\text{an}}(\tau_\lambda); E[\varepsilon](\epsilon))) \longrightarrow \text{H}^d(G(F), \mathcal{C}_E^{\text{ct}}(K^{\mathfrak{p}}, \mathbb{I}_B^{\text{an}}(E); E(\epsilon)))$$

in cohomology.

Let i^\vee be the coroot associated with i . Given a character $\lambda: B \rightarrow E$ we put $\lambda_i = \lambda \circ i^\vee$.

LEMMA 2.2. *Let $\lambda: B \rightarrow E$ be a continuous character. If the isotypic component*

$$\text{H}^{q+d}(G(F), \mathcal{C}_E^{\text{ct}}(K^{\mathfrak{p}}, \mathbb{I}_B^{\text{an}}(E); E(\epsilon)))[\pi^{\mathfrak{p}}]$$

is contained in the image of $\text{red}_\lambda^{q+d,\varepsilon}$, then the homomorphism λ_i belongs to $\mathcal{L}_i^{(d)}(\pi, \mathfrak{p})^\epsilon$.

Proof. The inclusion $\mathbb{I}_B^{\text{an}}(E) \hookrightarrow \mathbb{I}_B^{\text{an}}(\tau_\lambda)$ induces the map

$$\text{H}^{q+d}(G(F), \mathcal{C}_E^{\text{ct}}(K^{\mathfrak{p}}, \mathbb{I}_B^{\text{an}}(\tau_\lambda); E(\epsilon))) \longrightarrow \text{H}^{q+d}(G(F), \mathcal{C}_E^{\text{ct}}(K^{\mathfrak{p}}, \mathbb{I}_B^{\text{an}}(E); E(\epsilon)))$$

in cohomology. Its image is the kernel of the cup product with the extension class $\widehat{\mathcal{E}}^{\text{an}}(\lambda)$ defined in (1.2).

Thus, by Lemma 2.1 our assumption implies that the π -isotypic component $H^{q+d}(G(F), \mathcal{C}_E^{\text{ct}}(K^{\mathfrak{p}}, \mathbb{I}_B^{\text{an}}(E); E(\epsilon)))[\pi^{\mathfrak{p}}]$ is contained in the kernel of the cup product with $\widehat{\mathcal{E}}^{\text{an}}(\lambda)$. We have the following commutative diagram.

$$\begin{CD}
 H^{q+d}(G(F), \mathcal{C}_E^{\text{ct}}(K^{\mathfrak{p}}, \mathbb{I}_B^{\text{an}}(E); E(\epsilon))) @>\cup \widehat{\mathcal{E}}^{\text{an}}(\lambda)>> H^{q+d+1}(G(F), \mathcal{C}_E^{\text{ct}}(K^{\mathfrak{p}}, \mathbb{I}_B^{\text{an}}(E); E(\epsilon))) \\
 @VV=V @VVV \\
 H^{q+d}(G(F), \mathcal{C}_E^{\text{ct}}(K^{\mathfrak{p}}, \mathbb{I}_B^{\text{an}}(E); E(\epsilon))) @>\cup \widetilde{\mathcal{E}}^{\text{an}}(\lambda)>> H^{q+d+1}(G(F), \mathcal{C}_E(K^{\mathfrak{p}}, i_{P_i}^{\infty}(E); E(\epsilon))) \\
 @V\uparrow VV @V\uparrow VV \\
 H^{q+d}(G(F), \mathcal{C}_E^{\text{ct}}(K^{\mathfrak{p}}, \text{St}_{G_{\mathfrak{p}}}^{\text{ct}}(E); E(\epsilon))) @>\cup \mathcal{E}_i^{\text{an}}(\lambda)>> H^{q+d+1}(G(F), \mathcal{C}_E(K^{\mathfrak{p}}, v_{P_i}^{\infty}(E); E(\epsilon)))
 \end{CD}$$

The claim now follows from the first part of the next lemma and a simple diagram chase. □

LEMMA 2.3. *Let $J \subseteq \Delta$ be a subset.*

(i) *The canonical map*

$$H^d(G(F), \mathcal{C}_E(K^{\mathfrak{p}}, v_{P_J}^{\infty}(E); E(\epsilon)))[\pi^{\mathfrak{p}}] \longrightarrow H^d(G(F), \mathcal{C}_E(K^{\mathfrak{p}}, i_{P_J}^{\infty}(E); E(\epsilon)))[\pi^{\mathfrak{p}}]$$

is injective for all d .

(ii) *It is an isomorphism in degree $d = q + |J|$.*

Proof. The Jordan–Hölder decomposition of $i_{P_J}^{\infty}(E)$ consists of all generalized Steinberg representations $v_{P_I}^{\infty}(E)$ with $J \subseteq I$, each occurring with multiplicity one. Thus, the second claim follows from [Geh21, Proposition 3.9].

Via the well-known resolution (see, for example, [Orl05, Proposition 11])

$$0 \rightarrow i_{P_{\Delta}}^{\infty}(E) \rightarrow \bigoplus_{\substack{I \subseteq J \subseteq \Delta \\ |\Delta \setminus I|=1}} i_{P_I}^{\infty}(E) \rightarrow \cdots \rightarrow \bigoplus_{\substack{J \subseteq I \subseteq \Delta \\ |I \setminus J|=1}} i_{P_I}^{\infty}(E) \rightarrow i_{P_J}^{\infty}(E) \rightarrow v_{P_J}^{\infty}(E) \rightarrow 0$$

one can reduce the first claim to the following statement: Let $\mathcal{E}_{J,I}$ be any smooth extension of $v_{P_J}^{\infty}(E)$ by $v_{P_I}^{\infty}(E)$, where $J \subseteq I \subseteq \Delta$ with $|I| = |J| + 1$. Then the map

$$H^d(G(F), \mathcal{C}_E(K^{\mathfrak{p}}, v_{P_J}^{\infty}(E); E(\epsilon)))[\pi^{\mathfrak{p}}] \longrightarrow H^d(G(F), \mathcal{C}_E(K^{\mathfrak{p}}, \mathcal{E}_{J,I}; E(\epsilon)))[\pi^{\mathfrak{p}}]$$

is injective for all d . Equivalently, it is enough to show that the cup product

$$H^d(G(F), \mathcal{C}_E(K^{\mathfrak{p}}, v_{P_I}^{\infty}(E); E(\epsilon)))[\pi^{\mathfrak{p}}] \xrightarrow{\cup \mathcal{E}_{J,I}} H^{d+1}(G(F), \mathcal{C}_E(K^{\mathfrak{p}}, v_{P_J}^{\infty}(E); E(\epsilon)))[\pi^{\mathfrak{p}}]$$

is the zero map for all d . Let $\mathcal{E}_{I,J}$ be the unique up to scalar non-split smooth extension of $v_{P_I}^{\infty}(E)$ by $v_{P_J}^{\infty}(E)$. By [Geh21, Corollary 3.8], the cup product

$$H^{d+1}(G(F), \mathcal{C}_E(K^{\mathfrak{p}}, v_{P_J}^{\infty}(E); E(\epsilon)))[\pi^{\mathfrak{p}}] \xrightarrow{\cup \mathcal{E}_{I,J}} H^{d+2}(G(F), \mathcal{C}_E(K^{\mathfrak{p}}, v_{P_I}^{\infty}(E); E(\epsilon)))[\pi^{\mathfrak{p}}]$$

is an isomorphism. Therefore, it is enough to prove that taking the cup product with $\mathcal{E}_{J,I} \cup \mathcal{E}_{I,J}$ induces the zero map on $H^{\bullet}(G(F), \mathcal{C}_E(K^{\mathfrak{p}}, v_{P_I}^{\infty}(E); E(\epsilon)))[\pi^{\mathfrak{p}}]$. This is true because $\mathcal{E}_{J,I} \cup \mathcal{E}_{I,J}$ is a smooth 2-extension of $v_{P_I}^{\infty}(E)$ by itself and the space of all such extensions is zero by [Orl05, Theorem 1]. □

2.3 The automorphic Colmez–Greenberg–Stevens formula

Let A be an E -affinoid algebra and $\chi: B \rightarrow A^*$ a locally analytic character. The parabolic induction $\mathbb{I}_B^{\text{an}}(\chi)$ is naturally an $A[G_{\mathfrak{p}}]$ -module. Given an ideal $\mathfrak{m} \subseteq A$ we let $\chi_{\mathfrak{m}}: B \rightarrow (A/\mathfrak{m})^*$,

$x \mapsto \chi(x) \bmod \mathfrak{m}$ denote the reduction of χ modulo \mathfrak{m} . Similar to Proposition 2.2.1 of [Han17], one can prove that

$$\mathrm{Hom}_A^{\mathrm{ct}}(\mathbb{I}_B^{\mathrm{an}}(\chi), A) \otimes_A A/\mathfrak{m} \cong \mathrm{Hom}_{A/\mathfrak{m}}^{\mathrm{ct}}(\mathbb{I}_B^{\mathrm{an}}(\chi_{\mathfrak{m}}), A/\mathfrak{m}).$$

Let $\mathfrak{m} \in \mathrm{Spm} A$ be an E -rational point such that $\chi_{\mathfrak{m}} = \mathbb{1}_{G_{\mathfrak{p}}}$. Thus, the reduction map induces the maps

$$\mathrm{red}_{\chi}^{d,\epsilon}: \mathrm{H}^d(G(F), \mathcal{C}_A^{\mathrm{ct}}(K^{\mathfrak{p}}, \mathbb{I}_B^{\mathrm{an}}(\chi); A(\epsilon))) \longrightarrow \mathrm{H}^d(G(F), \mathcal{C}_E^{\mathrm{ct}}(K^{\mathfrak{p}}, \mathbb{I}_B^{\mathrm{an}}(E); E(\epsilon)))$$

in cohomology.

Let i^{\vee} be the coroot associated with i . We put $\chi_i = \chi \circ i^{\vee} \in \mathrm{Hom}(F_{\mathfrak{p}}^*, A)$. Suppose $v: \mathrm{Spec} E[\epsilon] \rightarrow \mathrm{Spm} A$ is an element of the tangent space of $\mathrm{Spm} A$ at \mathfrak{m} . The pullback $\chi_{i,v}$ of χ_i along v is of the form $\chi_{i,v} = 1 + (\partial/\partial v)\chi_i \cdot \epsilon$ for a unique homomorphism $(\partial/\partial v)\chi_i: F_{\mathfrak{p}}^* \rightarrow E$.

Lemma 2.2 immediately implies the following.

PROPOSITION 2.4. *Suppose that the image of $\mathrm{red}_{\chi}^{q+d,\epsilon}$ contains the $\pi^{\mathfrak{p},\infty}$ -isotypic component of $\mathrm{H}^{q+d}(G(F), \mathcal{C}_E^{\mathrm{ct}}(K^{\mathfrak{p}}, \mathbb{I}_B^{\mathrm{an}}(E); E(\epsilon)))$. Then the homomorphism $(\partial/\partial v)\chi_i$ belongs to $\mathcal{L}_i^{(d)}(\pi, \mathfrak{p})^{\epsilon}$ for every element v of the tangent space of $\mathrm{Spm} A$ at \mathfrak{m} .*

3. Overconvergent cohomology

After giving a brief overview on Kohlhaase and Schraen’s Koszul resolution of locally analytic principal series (see [KS12]) we recall the control theorem relating overconvergent cohomology to classical cohomology as proven by Ash–Stevens, Urban and Hansen (see [AS08, Urb11, Han17]). Combining the two results allows us to construct classes in the cohomology of (duals of) principal series. If G_{∞} fulfils the Harish-Chandra condition, i.e. if $\delta = 0$, we can lift the construction to families of principal series. This implies our main theorem.

3.1 Koszul complexes

In [KS12], Kohlhaase and Schraen construct a resolution of locally analytic principal series representations via a Koszul complex. We recall their construction in a slightly more general setup: instead of restricting to p -adic fields as coefficient rings we allow affinoid algebras. The proofs of [KS12] carry over verbatim to this more general framework.

Let us fix some notation: we denote the Borel opposite of $B \subseteq G_{F_{\mathfrak{p}}}$ by \bar{B} . Let $\bar{N} \subseteq \bar{B}$ be its unipotent radical. The chosen torus $T \subseteq B \subseteq G_{F_{\mathfrak{p}}}$ determines an apartment in the Bruhat–Tits building of $G_{\mathfrak{p}}$. We chose a chamber C of that apartment and a special vertex v of C as in §3.5. of [Car79]. The stabilizer $G_{\mathfrak{p},0} \subseteq G_{\mathfrak{p}}$ of v is a maximal compact subgroup of $G_{\mathfrak{p}}$ and the stabilizer $I_{\mathfrak{p}} \subseteq G_{\mathfrak{p},0}$ of C is an Iwahori subgroup. Let \mathfrak{G}_0 be the Bruhat–Tits group scheme over $\mathcal{O}_{\mathfrak{p}}$ associated with $G_{\mathfrak{p},0}$. We define

$$I_{\mathfrak{p}}^n = \ker(I_{\mathfrak{p}} \longrightarrow \mathfrak{G}_0(\mathcal{O}_{\mathfrak{p}}/\mathfrak{p}^n)).$$

The open normal subgroups $I_{\mathfrak{p}}^n \subseteq I_{\mathfrak{p}}$ form a system of neighbourhoods of the identity in $I_{\mathfrak{p}}$. The subgroup $T_0 = T \cap G_{\mathfrak{p},0}$ is maximal compact subgroup of T .

Let $X^*(T)$ (respectively, $X_*(T)$) denote the group of $F_{\mathfrak{p}}$ -rational characters (respectively, cocharacters) of T . The natural pairing

$$\langle \cdot, \cdot \rangle: X^*(T) \times X_*(T) \longrightarrow \mathbb{Z}$$

is a perfect pairing. There is a natural isomorphism $T/T_0 \cong X_*(T)$ characterized by

$$\langle \chi, t \rangle = \mathrm{ord}_{\mathfrak{p}}(\chi(t)).$$

We denote by Φ^+ the set of positive roots with respect to B and put

$$T^- = \{t \in T \mid \text{ord}_p(\alpha(t)) \leq 0 \ \forall \alpha \in \Phi^+\}.$$

Let A be a \mathbb{Q}_p -affinoid algebra. Restricting to T_0 gives a bijection between locally analytic characters $\chi: B \cap I_p \rightarrow A^*$ and locally analytic characters $\chi: T_0 \rightarrow A^*$. Given such a character χ we write

$$\mathcal{A}_\chi = \{f: I_p \rightarrow A \text{ locally analytic} \mid f(bk) = \chi(b)f(k) \ \forall b \in B \cap I_p, k \in I_p\}$$

for the locally analytic induction of χ to I_p . It is naturally an $A[I_p]$ -module. Restricting a function $f \in \mathcal{A}_\chi$ to the intersection $I_p \cap \bar{N}$ induces an isomorphism of \mathcal{A}_χ with the space of all locally analytic functions from $I_p \cap \bar{N}$ to A . There exists a minimal integer $n_\chi \geq 1$ such that χ restricted to $B \cap I_p^{n_\chi}$ is rigid analytic. For any $n \geq n_\chi$ we define the $A[I_p]$ -submodule

$$\mathcal{A}_\chi^n = \{f \in \mathcal{A}_\chi \mid f \text{ is rigid analytic on any coset in } I_p/I_p^n\}.$$

For later purposes, we define the dual spaces

$$\mathcal{D}_\chi^n = \text{Hom}_A^{\text{ct}}(\mathcal{A}_\chi^n, A).$$

By Frobenius reciprocity we can identify $\text{End}_{A[G_p]}(\text{c-ind}_{I_p}^{G_p}(\mathcal{A}_\chi^n))$ with the space of all functions $\Psi: G_p \rightarrow \text{End}_A(\mathcal{A}_\chi^n)$ such that:

- (i) Ψ is I_p -biequivariant, i.e. $\Psi(k_1 g k_2) = k_1 \Psi(g) k_2$ for all $k_1, k_2 \in I_p, g \in G_p$; and
- (ii) for every $f \in \mathcal{A}_\chi^n$ the function $G_p \rightarrow \mathcal{A}_\chi^n, g \mapsto \Psi(g)(f)$ is compactly supported.

Let t be an element of T^- and $f \in \mathcal{A}_\chi^n$. The function $I_p \rightarrow A, u \mapsto f(tut^{-1})$ defines an element of \mathcal{A}_χ^n .

LEMMA 3.1. *For every element $t \in T^-$ there exists a unique I_p -biequivariant function $\Psi_t: G_p \rightarrow \text{End}_A(\mathcal{A}_\chi^n)$ such that:*

- (i) $\text{supp}(\Psi_t) = I_p t^{-1} I_p$; and
- (ii) $\Psi_t(t^{-1})(f)(u) = f(tut^{-1})$ for any $f \in \mathcal{A}_\chi^n$ and $u \in I_p \cap \bar{N}$.

Proof. This is a minor generalization of [KS12, Lemma 2.2]. □

For $t \in T^-$ we denote by U_t the endomorphism of $\text{c-ind}_{I_p}^{G_p}(\mathcal{A}_\chi^n)$ corresponding to Ψ_t . The following is a straightforward generalization of [KS12, Lemma 2.3].

LEMMA 3.2. *We have $U_t U_{\tilde{t}} = U_{t\tilde{t}}$ for all $t, \tilde{t} \in T^-$.*

Now let us fix a character $\chi: T \rightarrow A^*$ and let χ_0 be its restriction to T_0 . Given an open subset $C \subseteq G_p$, which is stable under multiplication with B from the left, we denote by

$$\mathbb{I}_B^{\text{an}}(\chi)(C) \subseteq \mathbb{I}_B^{\text{an}}(\chi)$$

the subset of all functions with support in C . Restricting functions to I_p gives an I_p -equivariant A -linear isomorphism

$$\mathbb{I}_B^{\text{an}}(\chi)(BI_p) \xrightarrow{\cong} \mathcal{A}_{\chi_0}.$$

Thus, by Frobenius reciprocity its inverse induces a G_p -equivariant A -linear map

$$\text{aug}_\chi: \text{c-ind}_{I_p}^{G_p}(\mathcal{A}_{\chi_0}^n) \rightarrow \text{c-ind}_{I_p}^{G_p}(\mathcal{A}_{\chi_0}^n) \rightarrow \mathbb{I}_B^{\text{an}}(\chi) \tag{3.1}$$

for any integer $n \geq n_{\chi_0}$.

As the group G_{F_p} is adjoint, there exist elements $t_i \in T^-, i \in \Delta$, such that

$$t_i \in \bigcap_{j \in \Delta \setminus \{i\}} \ker(j)$$

and

$$\text{ord}_{\mathfrak{p}}(i(t_i)) = -1.$$

The element t_i is uniquely determined by the value $i(t_i)^{-1} \in F_{\mathfrak{p}}^*$, which is a uniformizer. Every element $t \in T$ can uniquely be written as $t = t_0 \prod_{i \in \Delta} t_i^{n_i}$ with $t \in T_0$ and integers $n_i \in \mathbb{Z}$. Let us fix a choice of $t_i, i \in \Delta$, and put

$$y_i = U_{t_i} - \chi(t_i).$$

PROPOSITION 3.3. *The $G_{\mathfrak{p}}$ -equivariant A -linear map aug_{χ} is surjective with kernel $\sum_{i \in \Delta} \text{im}(y_i)$.*

Proof. The same proof as for [KS12, Proposition 2.4] works here. □

By Lemma 3.1 the $G_{\mathfrak{p}}$ -representation $\text{c-ind}_{I_{\mathfrak{p}}}^{G_{\mathfrak{p}}}(\mathcal{A}_{\chi_0}^n)$ is a module over the polynomial algebra $A[X_i \mid i \in \Delta]$, where X_i acts through the operator T_i . The Koszul complex of $\text{c-ind}_{I_{\mathfrak{p}}}^{G_{\mathfrak{p}}}(\mathcal{A}_{\chi_0}^n)$ with respect to the endomorphisms $(y_i)_{i \in \Delta}$ is the complex $\Lambda_A^{\bullet}(A^{\Delta}) \otimes \text{c-ind}_{I_{\mathfrak{p}}}^{G_{\mathfrak{p}}}(\mathcal{A}_{\chi_0}^n)$ with boundary maps

$$d_k(e_{i_1} \wedge \cdots \wedge e_{i_k} \otimes f) = \sum_{l=1}^k (-1)^{l+1} e_{i_1} \wedge \cdots \wedge \widehat{e_{i_l}} \wedge \cdots \wedge e_{i_k} \otimes y_{i_l}(f). \tag{3.2}$$

The following is the main technical theorem of Kohlhaase–Schraen (cf. [KS12, Theorem 2.5]) generalized to affinoid coefficient rings.

THEOREM 3.4 (Kohlhaase–Schraen). *For any $n \geq n_{\chi_0}$ the augmented Koszul complex*

$$\Lambda_A^{\bullet}(A^{\Delta}) \otimes_A \text{c-ind}_{I_{\mathfrak{p}}}^{G_{\mathfrak{p}}}(\mathcal{A}_{\chi_0}^n) \longrightarrow \mathbb{I}_B^{\text{an}}(\chi) \longrightarrow 0$$

with boundary maps (3.2) and augmentation map (3.1) is exact.

Remark 3.5. All of the results above remain valid if one replaces $\mathcal{A}_{\chi_0}^n$ by \mathcal{A}_{χ_0} .

Example 3.6. Suppose $G_{\mathfrak{p}} = \text{PGL}_n(F_{\mathfrak{p}})$, T is the torus of diagonal matrices and B is the Borel subgroup of upper triangular matrices. The simple roots of T with respect to B are given by

$$i(\text{diag}(x_1, \dots, x_n)) = x_i x_{i+1}^{-1}$$

for $1 \leq i \leq d - 1$. For each simple root $1 \leq i \leq d - 1$, we might choose for t_i the image of the diagonal matrix

$$\text{diag}(1, \dots, 1, \pi, \dots, \pi),$$

where π is a uniformizer and exactly the first i entries are equal to one. For the Iwahori subgroup $I_{\mathfrak{p}}$, we may choose the image in $G_{\mathfrak{p}}$ of all matrices in $\text{GL}_n(\mathcal{O}_{\mathfrak{p}})$, which are upper triangular modulo \mathfrak{p} . Then $I_{\mathfrak{p}}^n$ consists of all matrices in $I_{\mathfrak{p}}$ which are congruent to the identity modulo \mathfrak{p}^n .

There is also a smooth variant of the above result, which is probably well-known. As we could not find a reference in the literature, we give a proof of said variant in the following.

Let Ω be field of characteristic zero. With the same formula as before, we can define commuting Hecke operators $U_t \in \text{End}_{G_{\mathfrak{p}}}(\text{c-ind}_{I_{\mathfrak{p}}}^{G_{\mathfrak{p}}}(E))$ for $t \in T^-$. Let $\mathbb{1} : T_0 \rightarrow \Omega^{\times}$ denote the constant

character on T_0 . In case $\Omega = E$ we can identify E with the subspace of constant functions in \mathcal{A}_1^n and the induced embedding

$$\text{c-ind}_{I_p}^{G_p}(E) \hookrightarrow \text{c-ind}_{I_p}^{G_p}(\mathcal{A}_1^n) \tag{3.3}$$

is equivariant with respect to the operators $U_t, t \in T^-$.

The operators $U_t \in \text{End}_{G_p}(\text{c-ind}_{I_p}^{G_p}(\Omega))$ are invertible. Moreover, by the Bernstein decomposition the map

$$\Omega[X_i^{\pm 1} | i \in \Delta] \longrightarrow \text{End}_{G_p}(\text{c-ind}_{I_p}^{G_p}(\Omega)), X_i \longmapsto U_{t_i}$$

is injective and $\text{End}_{G_p}(\text{c-ind}_{I_p}^{G_p}(\Omega))$ is a free module of finite rank (and, therefore, flat) over $\Omega[X_i^{\pm 1} | i \in \Delta]$ (see, for example, [Ree92] for more details). As $\text{c-ind}_{I_p}^{G_p}(\Omega)$ is a flat module over its endomorphism algebra by a theorem of Borel (see [Bor76, Theorem 4.10]), it is thus also flat as a $\Omega[X_i^{\pm 1} | i \in \Delta]$ -module.

Therefore, for any choice of elements $a_i \in \Omega^\times$ the Koszul complex

$$\Lambda_\Omega^\bullet(\Omega^\Delta) \otimes_\Omega \text{c-ind}_{I_p}^{G_p}(\Omega)$$

associated with the regular sequence $y_i = U_{t_i} - a_i, i \in \Delta$, is a resolution of the G_p -representation

$$M_{\underline{a}} = \text{coker}\left(\bigoplus_{i \in \Delta} \text{c-ind}_{I_p}^{G_p}(\Omega) \xrightarrow{(y_i)_{i \in \Delta}} \text{c-ind}_{I_p}^{G_p}(\Omega)\right).$$

Let $\chi_{\underline{a}}: T \rightarrow \Omega^*$ be the unique smooth unramified character such that $\chi_{\underline{a}}(t_i) = a_i$. As before, we extend $\chi_{\underline{a}}$ to a character of B . Let $\phi \in i_B^\infty(\chi_{\underline{a}})$ be the unique element such that:

- (i) ϕ is invariant under I_p ;
- (ii) the support of ϕ is BI_p ; and
- (iii) $\phi(1) = 1$.

Note that in case $\chi_{\underline{a}} = \mathbb{1}$ is the trivial character the image of ϕ under the quotient map $i_B^\infty(\Omega) \twoheadrightarrow \text{St}^\infty(\Omega)$ is non-zero and, thus, generates the space of Iwahori invariants of the Steinberg representation.

By Frobenius reciprocity ϕ induces a G_p -equivariant homomorphism

$$\text{c-ind}_{I_p}^{G_p} \Omega \longrightarrow i_B^\infty(\chi_{\underline{a}}). \tag{3.4}$$

One can argue as in the proof of [Oll14, Proposition 4.4] that the map (3.4) induces an isomorphism

$$M_{\underline{a}} \longrightarrow i_B^\infty(\chi_{\underline{a}}).$$

In conclusion, we see that the augmented Koszul complex

$$\Lambda_A^\bullet(A^\Delta) \otimes_A \text{c-ind}_{I_p}^{G_p}(\Omega) \longrightarrow i_B^\infty(\chi_{\underline{a}}) \longrightarrow 0 \tag{3.5}$$

is exact.

3.2 Overconvergent cohomology

We show that the result from the previous section together with the theory of overconvergent cohomology allows us to construct cohomology classes with values in duals of locally analytic representations.

Before sticking to the case which is most relevant for our applications let us consider the general case: let A be a \mathbb{Q}_p -affinoid algebra and $\chi: T \rightarrow A$ a locally analytic character.

Denote by χ_0 its restriction to T_0 and fix elements $t_i \in T^-, i \in \Delta$, as before. For $n \geq n_{\chi_0}$ the continuous dual of the augmentation map (3.1) of the previous section yields the map

$$\text{aug}_\chi^*: H^d(G(F), C_E^{\text{ct}}(K^{\mathfrak{p}}, \mathbb{I}_B^{\text{an}}(\chi); A(\epsilon))) \longrightarrow H^d(X_{K^{\mathfrak{p}} \times I_{\mathfrak{p}}}, \mathcal{D}_{\chi_0}^n)^\epsilon$$

for every sign character ϵ and every integer $d \geq 0$. We denote the operator induced by $U_{t_i}, i \in \Delta$, on the right-hand side also by U_{t_i} and similar for $U_{\tilde{t}}$. Then, Proposition 3.3 implies the following.

COROLLARY 3.7. *We have*

$$\text{im}(\text{aug}_\chi^*) \subseteq \bigcap_{i \in \Delta} \ker(U_{t_i} - \chi(t_i)).$$

For the remainder of the section we study the case of the trivial character $\mathbb{1} = \mathbb{1}_T: T \rightarrow E^*$ of T with values in the units of the p -adic field E . We denote its restriction to T_0 also by $\mathbb{1}$. Let $n \geq 1$ be any integer. The map (3.3) induces a homomorphism

$$H^d(X_{K^{\mathfrak{p}} \times I_{\mathfrak{p}}}, \mathcal{D}_{\mathbb{1}}^n)^\epsilon \longrightarrow H^d(X_{K^{\mathfrak{p}} \times I_{\mathfrak{p}}}, E)^\epsilon \tag{3.6}$$

in cohomology, that is, equivariant with respect to the commuting actions of the Hecke algebra $\mathbb{T}^{\mathfrak{p}}$ and the operators $U_t, t \in T^-$.

In the following, we study the finite slope parts of the above cohomology groups (see, for example, [Han17, § 2.3], for definitions and notation).

THEOREM 3.8 (Ash–Stevens, Urban and Hansen). *Let $\tilde{t} = \prod_{i \in \Delta} t_i$.*

- (i) *The space $H^d(X_{K^{\mathfrak{p}} \times I_{\mathfrak{p}}}, \mathcal{D}_{\mathbb{1}}^n)^\epsilon$ admits a slope decomposition with respect to $U_{\tilde{t}}$ and every rational number h .*
- (ii) *If h is small enough with respect to \tilde{t} and the trivial character $\mathbb{1}$ (see, for example, the first formula on page 1690 of [Urb11] or equation (21) of [AS08]), then the map*

$$H^d(X_{K^{\mathfrak{p}} \times I_{\mathfrak{p}}}, \mathcal{D}_{\mathbb{1}}^n)^{\epsilon, \leq h} \longrightarrow H^d(X_{K^{\mathfrak{p}} \times I_{\mathfrak{p}}}, E)^{\epsilon, \leq h}.$$

is an isomorphism. In particular, this is an isomorphism for $h = 0$.

Proof. For the first claim see § 2.3 of [Han17] and for the second see [Han17, Theorem 3.2.5]. The third claim is [Urb11, Proposition 4.3.10]. Note that in all cases the authors consider all primes lying above p at once. The same proofs work in our partial \mathfrak{p} -adic setup. □

Composing the dual of the augmentation map

$$\text{aug}_{\mathbb{1}}^*: H^d(G(F), C_E^{\text{ct}}(K^{\mathfrak{p}}, \mathbb{I}_B^{\text{an}}(E); E(\epsilon))) \longrightarrow H^d(X_{K^{\mathfrak{p}} \times I_{\mathfrak{p}}}, \mathcal{D}_{\mathbb{1}}^n)^\epsilon \tag{3.7}$$

with the map (3.6) yields the map

$$H^d(G(F), C_E^{\text{ct}}(K^{\mathfrak{p}}, \mathbb{I}_B^{\text{an}}(E); E(\epsilon))) \longrightarrow H^d(X_{K^{\mathfrak{p}} \times I_{\mathfrak{p}}}, E)^\epsilon. \tag{3.8}$$

As the diagram

$$\begin{array}{ccc} \text{c-ind}_{I_{\mathfrak{p}}}^{G_{\mathfrak{p}}} E & \xrightarrow{(3.3)} & \text{c-ind}_{I_{\mathfrak{p}}}^{G_{\mathfrak{p}}} \mathcal{A}_{\mathbb{1}}^n \xrightarrow{\text{aug}_{\mathbb{1}}} \mathbb{I}_B^{\text{an}}(E) \\ (1.7) \downarrow & & \downarrow \\ \text{St}_{G_{\mathfrak{p}}}^{\infty}(E) & \xrightarrow{\hspace{10em}} & \text{St}_{G_{\mathfrak{p}}}^{\text{ct}}(E) \end{array}$$

is commutative up to multiplication with a non-zero constant, which comes from the choice of a Iwahori-fixed vector in (1.7), so is the induced diagram in cohomology as follows.

$$\begin{CD} H^d(G(F), \mathcal{C}_E^{\text{ct}}(K^{\mathfrak{p}}, \mathbb{I}_B^{\text{an}}(E); E(\epsilon))) @>(3.8)>> H^d(X_{K^{\mathfrak{p}} \times I_{\mathfrak{p}}}, E)^{\epsilon} \\ @VVV @VV\text{ev}^{(d)}V \\ H^d(G(F), \mathcal{C}_E^{\text{ct}}(K^{\mathfrak{p}}, \text{St}_{G_{\mathfrak{p}}}^{\text{ct}}(E); E(\epsilon))) @>\cong>> H^d(G(F), \mathcal{C}_E^{\text{ct}}(K^{\mathfrak{p}}, \text{St}_{G_{\mathfrak{p}}}^{\infty}(E); E(\epsilon))) \end{CD}$$

PROPOSITION 3.9. Let $a_i \in E^{\times}$, $i \in \Delta$, be elements with p -adic valuation equal to 0 and $\chi_{\underline{a}}: T \rightarrow E^{\times}$ the unique smooth unramified character such that $\chi_{\underline{a}}(t_i) = a_i$. The natural inclusion $i_B^{\infty}(\chi_{\underline{a}}) \hookrightarrow \mathbb{I}_B^{\text{an}}(\chi_{\underline{a}})$ induces an isomorphism

$$H^d(G(F), \mathcal{C}_E^{\text{ct}}(K^{\mathfrak{p}}, \mathbb{I}_B^{\text{an}}(\chi_{\underline{a}}); E(\epsilon))) \xrightarrow{\cong} H^d(G(F), \mathcal{C}_E(K^{\mathfrak{p}}, i_B^{\infty}(\chi_{\underline{a}}); E(\epsilon)))$$

for all $d \geq 0$. In particular, the map (3.8) induces an isomorphism

$$H^q(G(F), \mathcal{C}_E^{\text{ct}}(K^{\mathfrak{p}}, \mathbb{I}_B^{\text{an}}(E); E(\epsilon)))[\pi^{\mathfrak{p}}] \xrightarrow{\cong} H^q(X_{K^{\mathfrak{p}} \times I_{\mathfrak{p}}}, E)^{\epsilon}[\pi]$$

on π -isotypic components in degree q .

Proof. The second claim is a direct consequence of the first claim and Lemma 2.3.

Given an $I_{\mathfrak{p}}$ -representation M we denote by $\text{Coind}_{I_{\mathfrak{p}}}^{G_{\mathfrak{p}}} M$ the coinduction of M to $G_{\mathfrak{p}}$, i.e. the module of all functions $f: G_{\mathfrak{p}} \rightarrow M$ such that $f(kg) = kf(g)$ for all $k \in I_{\mathfrak{p}}$, $g \in G_{\mathfrak{p}}$. By taking continuous duals the Koszul complex of Theorem 3.4 with $y_i = U_{t_i} - a_i$ yields a quasi-isomorphism

$$\text{Hom}_E^{\text{ct}}(\mathbb{I}_B^{\text{an}}(\chi_{\underline{a}}), E) \xrightarrow{\cong} \Lambda_E^{\bullet}(E^{\Delta}) \otimes_E \text{Coind}_{I_{\mathfrak{p}}}^{G_{\mathfrak{p}}}(\mathcal{D}_{\mathbb{1}}^n)$$

of complexes. Note that taking continuous duals in this case is exact by the Hahn–Banach theorem. Similarly by the smooth Koszul resolution (3.5) we have a quasi-isomorphism

$$\text{Hom}_E(i_B^{\infty}(\chi_{\underline{a}}), E) \xrightarrow{\cong} \Lambda_E^{\bullet}(E^{\Delta}) \otimes_E \text{Coind}_{I_{\mathfrak{p}}}^{G_{\mathfrak{p}}}(E)$$

and the canonical diagram

$$\begin{CD} \text{Hom}_E^{\text{ct}}(\mathbb{I}_B^{\text{an}}(\chi_{\underline{a}}), E) \otimes \Lambda_E^{\bullet}(E^{\Delta}) \otimes_E \text{Coind}_{I_{\mathfrak{p}}}^{G_{\mathfrak{p}}}(\mathcal{D}_{\mathbb{1}}^n) @>>> \Lambda_E^{\bullet}(E^{\Delta}) \otimes_E \text{Coind}_{I_{\mathfrak{p}}}^{G_{\mathfrak{p}}}(\mathcal{D}_{\mathbb{1}}^n) \\ @VVV @VVV \\ \text{Hom}_E(i_B^{\infty}(\chi_{\underline{a}}), E) \otimes \Lambda_E^{\bullet}(E^{\Delta}) \otimes_E \text{Coind}_{I_{\mathfrak{p}}}^{G_{\mathfrak{p}}}(E) @>>> \Lambda_E^{\bullet}(E^{\Delta}) \otimes_E \text{Coind}_{I_{\mathfrak{p}}}^{G_{\mathfrak{p}}}(E) \end{CD}$$

is commutative.

From the associated spectral sequences for double complex, we deduce that it is enough to prove that the map of complexes

$$\Lambda_E^{\bullet}(E^{\Delta}) \otimes_E H^d(X_{K^{\mathfrak{p}} \times I_{\mathfrak{p}}}, \mathcal{D}_{\mathbb{1}}^n) \longrightarrow \Lambda_E^{\bullet}(E^{\Delta}) \otimes_E H^d(X_{K^{\mathfrak{p}} \times I_{\mathfrak{p}}}, E) \tag{3.9}$$

is a quasi-isomorphism for all $d \geq 0$. The modules on the left-hand side admit a slope decomposition for $U_{\tilde{t}}$ by Theorem 3.8(i), whereas the modules on the right-hand side admit a slope decomposition because they are finite-dimensional E -vector spaces.

As the operators y_i commute with $U_{\tilde{t}}$ they respect the slope decomposition on both sides (see [Urb11, Lemma 2.3.2]). On the slope less than or equal to zero part the map (3.9) is even an isomorphism of complexes by Theorem 3.8(ii).

Thus, we are reduced to proving that

$$\Lambda_E^\bullet(E^\Delta) \otimes_E H^d(X_{K^p \times I_p}, \mathcal{D}_1^n)^{>h} \longrightarrow \Lambda_E^\bullet(E^\Delta) \otimes_E H^d(X_{K^p \times I_p}, E)^{>h}$$

is a quasi-isomorphism for all $d \geq 0$.

In fact, both sides are acyclic: standard properties of Koszul complexes imply that the operators y_i act as multiplication by zero on the cohomology of the complexes. By the definition of a slope decomposition the operator $U_{\tilde{t}} - \prod_{i \in \Delta} a_i$ acts via isomorphisms on both complexes and, thus, on their cohomology. However, because $U_{\tilde{t}} - \prod_{i \in \Delta} a_i$ lies in the ideal generated by the operators y_i , it also acts via multiplication by zero, which proves the claim. \square

Remark 3.10. In order to keep the article short and avoid unnecessary notation, we decided to stick to the special case above. However, Proposition 3.9 also holds in a much more general situation, e.g. one can allow arbitrary coefficient systems and arbitrary non-critical principal series representations, given that the slope is small enough with respect to \tilde{t} .

3.3 Overconvergent families

Let \mathcal{W} be the weight space of T , i.e. it is the rigid space over E such that for every E -affinoid algebra A its A -points are given by

$$\mathcal{W}(A) = \text{Hom}^{\text{ct}}(T, A^*).$$

It is smaller than the usual weight space as we only consider characters of the torus in G_{F_p} . For any open affinoid $\mathcal{U} \subset \mathcal{W}$ and we let $\chi_{\mathcal{U}}$ be the corresponding universal weight. As in § 3.1 we define the space of n -analytic functions and distributions $\mathcal{A}_{\chi_{\mathcal{U}}}^n$ and $\mathcal{D}_{\chi_{\mathcal{U}}}^n$ for $n \gg 0$.

We suppose now that the group G_∞ has discrete series or, equivalently, that $\delta = 0$ (see [Kna01, Theorem 12.20] for the equivalence of these two properties). Thus, by Proposition 1.6, the representation π^∞ appears only in the middle degree cohomology $H^q(X_{K^p \times I_p}, E)$. Hence, we put $\mathcal{L}_i(\pi, \mathfrak{p})^\epsilon = \mathcal{L}_i^{(q)}(\pi, \mathfrak{p})^\epsilon$.

Let $\mathbb{T}_{\text{sph}}^{\mathfrak{p}} \subseteq \mathbb{T}^{\mathfrak{p}}$ be the spherical Hecke algebra of level $K^{\mathfrak{p}} \times I_{\mathfrak{p}}$ over E , i.e. the commutative Hecke algebra generated by all Hecke operators at finite places v such that K_v is hyperspecial. We define \mathbb{T}_{sph} to be the commutative algebra generated by $\mathbb{T}_{\text{sph}}^{\mathfrak{p}}$ and all U_t -operators for $t \in T^-$. Let $\mathfrak{m}_\pi \subseteq \mathbb{T}_{\text{sph}}$ (respectively, $\mathfrak{m}_\pi^{\mathfrak{p}} \subseteq \mathbb{T}_{\text{sph}}^{\mathfrak{p}}$) the maximal ideal associated with π . We assume the following weak non-Eisenstein assumption on the maximal ideal \mathfrak{m}_π throughout this section.

HYPOTHESIS (NE). We have

$$H^d(X_{K^p \times I_p}, E)_{\mathfrak{m}_\pi} = 0$$

unless $d = q$.

Example 3.11. If the group G is definite, the hypothesis is automatically true. By strong multiplicity one and the Jacquet–Langlands correspondence, it is also true for inner forms of PGL_2 over totally real number fields.

Let \mathcal{U} be an open affinoid of \mathcal{W} and define $\mathcal{O}_{\mathcal{U}, \mathbb{1}}$ to be the rigid localization of $\mathcal{O}_{\mathcal{W}}(\mathcal{U})$ at the weight $\mathbb{1}$ (which is the cohomological weight of π), i.e.

$$\mathcal{O}_{\mathcal{U}, \mathbb{1}} = \varinjlim_{\mathbb{1} \in \mathcal{U}' \subset \mathcal{U}} \mathcal{O}_{\mathcal{W}}(\mathcal{U}').$$

The following theorem is instrumental in calculating \mathcal{L} -invariants.

THEOREM 3.12. *After localization at the ideal \mathfrak{m}_π of \mathbb{T}_{sph} and restricting to a small enough open affinoid \mathcal{U} containing $\mathbb{1}$, the canonical reduction map*

$$H^q(X_{K^p \times I_p}, \mathcal{D}_{\chi_{\mathcal{U}}}^n)_{\mathfrak{m}_\pi}^\epsilon \rightarrow H^q(X_{K^p \times I_p}, E)_{\mathfrak{m}_\pi}^\epsilon$$

is surjective. Moreover, $H^q(X_{K^p \times I_p}, \mathcal{D}_{\chi_{\mathcal{U}}}^n)_{\mathfrak{m}_\pi}^\epsilon$ is a free $\mathcal{O}_{\mathcal{U}, \mathbb{1}}$ -module of rank equal to the dimension of $H^q(X_{K^p \times I_p}, E)_{\mathfrak{m}_\pi}^\epsilon$.

In addition, we have $H^d(X_{K^p \times I_p}, \mathcal{D}_{\chi_{\mathcal{U}}}^n)_{\mathfrak{m}_\pi}^\epsilon = 0$ for all $d \neq q$.

Proof. The case of PGL_2 over a totally real field is proven in detail in [BDJ21, Theorem 2.14]. The main ingredient in their proof is the vanishing of the cohomology outside middle degree. Thus, the same proof works in our more general setup. \square

DEFINITION 3.13. We say that \mathfrak{m}_π is \mathfrak{p} -étale (with respect to ϵ) if every Hecke operator $h \in \mathbb{T}_{\text{sph}}$ acts on $H^q(X_{K^p \times I_p}, \mathcal{D}_{\chi_{\mathcal{U}}}^n)_{\mathfrak{m}_\pi}^\epsilon$ as multiplication by an element $\alpha_h \in \mathcal{O}_{\mathcal{U}, \mathbb{1}}^*$.

Theorem 3.12 immediately implies the following.

COROLLARY 3.14. *Assume that $\dim_E H^q(X_{K^p \times I_p}, E)_{\mathfrak{m}_\pi}^\epsilon = 1$. Then \mathfrak{m}_π is \mathfrak{p} -étale.*

Example 3.15. Suppose G is an inner form of PGL_2 over a totally real number field. Then the above corollary together with strong multiplicity one implies that \mathfrak{m}_π is \mathfrak{p} -étale.

Let $t_i \in T^-$, $i \in \Delta$, be a choice of elements as in §3.1. Suppose \mathfrak{m}_π is \mathfrak{p} -étale (with respect to ϵ) with associated eigenvalues $\alpha_{U_{t_i}} \in \mathcal{O}_{\mathcal{U}, \mathbb{1}}^*$, $i \in \Delta$. By possibly shrinking \mathcal{U} , we may assume that $\alpha_{U_{t_i}} \in \mathcal{O}_{\mathcal{U}}^*$. Every element $t \in T$ can be written uniquely as a product $t = t_0 \prod_{i \in \Delta} t_i^{n_i}$ with $t_0 \in T_0$, $n_i \in \mathbb{Z}$. Hence, there exists a unique character $\chi_\alpha: T \rightarrow \mathcal{O}_{\mathcal{U}}^*$ such that

$$\chi_\alpha|_{T_0} = \chi_U$$

and

$$\chi_\alpha(t_i) = \alpha_{U_{t_i}}.$$

THEOREM 3.16. *Suppose that \mathfrak{m}_π is \mathfrak{p} -étale (with respect to ϵ) with associated character $\chi_\alpha: T \rightarrow \mathcal{O}_{\mathcal{U}}^*$. Then for every tangent vector v of \mathcal{U} at $\mathbb{1}$ we have:*

$$\frac{\partial}{\partial v} \chi_{\alpha, i} \in \mathcal{L}_i(\pi, \mathfrak{p})^\epsilon$$

where $\chi_{\alpha, i}$ denotes the composition $\chi_\alpha \circ i^\vee$. Moreover, the subspace $\mathcal{L}_i(\pi, \mathfrak{p})^\epsilon \subseteq \text{Hom}^{\text{ct}}(F_{\mathfrak{p}}^*, E)$ has codimension one.

Proof. Let $\mathfrak{m}_{\mathbb{1}} \subseteq \mathcal{O}_{\mathcal{U}}$ be the maximal ideal corresponding to the trivial character. We put $A = \mathcal{O}_{\mathcal{U}}/\mathfrak{m}_{\mathbb{1}}^2$ and $\chi_0 = \chi_{\mathcal{U}} \bmod \mathfrak{m}_{\mathbb{1}}^2$. Let us write $\tilde{\chi}_\alpha = \chi_\alpha \bmod \mathfrak{m}_{\mathbb{1}}^2: T \rightarrow A$. Using Theorem 3.12 and arguments with Koszul complexes as in the proof of Proposition 3.9 one shows that the image of the map

$$H^q(G(F), \mathcal{C}_A^{\text{ct}}(K^p, \mathbb{I}_B^{\text{an}}(\tilde{\chi}_\alpha); A(\epsilon)))_{\mathfrak{m}_\pi^p} \longrightarrow H^q(X_{K^p \times I_p}, E)_{\mathfrak{m}_\pi^p}^\epsilon$$

is the intersection of the kernels of the homomorphisms $U_{t_i} - 1$. Therefore, the first claim follows from Proposition 2.4.

We may assume that E is large enough. We explain how to choose appropriate tangent vectors so that $\mathcal{L}_i(\pi, \mathfrak{p})^\epsilon$ contains elements of the form $\log_{p, \sigma} - \mathcal{L}^\sigma \text{ord}_{\mathfrak{p}}$ for every embedding $\sigma: F_{\mathfrak{p}} \rightarrow E$. This will show that the \mathcal{L} -invariant has codimension at most one.

Let $E\langle k_\sigma \rangle$ be the ring of power series in the k_σ and let $\mathcal{O}_{\mathfrak{p}, \mathbb{1}}^*$ denote the free part of $\mathcal{O}_{\mathfrak{p}}^*$. By smoothness of the weight space, we can embed $\mathcal{O}_{\mathcal{U}}^*$ in $E\langle k_\sigma \rangle$ and the structural morphism of

$E[\mathcal{O}_{\mathfrak{p},1}^*]$ into $\mathcal{O}_{\mathcal{U}}^*$ can be described via the map

$$\mathcal{O}_{\mathfrak{p}}^* \ni \langle u \rangle \mapsto \prod_{\sigma} \sigma(u)^{k_{\sigma}}.$$

We can, hence, identify $\chi_{\alpha,i}$ with a map from $F_{\mathfrak{p}}^*$ to $\mathcal{O}_{\mathcal{U}}^*$. For all $n \in \mathbb{Z}$ and $u \in \mathcal{O}_{\mathfrak{p},1}^*$ we have

$$\chi_{\alpha,i}(\pi^n u) = \alpha_{U_{t_i}}^n \prod_{\sigma} \sigma(u)^{k_{\sigma}}.$$

Note that

$$\frac{d}{dk_{\sigma'}} (\sigma(u)^{k_{\sigma}}) = \delta_{\sigma,\sigma'} \log_p(\sigma(u)) \sigma(u)^{k_{\sigma}},$$

where

$$\delta_{\sigma,\sigma'} = \begin{cases} 1 & \text{if } \sigma = \sigma' \\ 0 & \text{otherwise.} \end{cases}$$

Take for v the direction where only k_{σ} varies, derive $\chi_{\alpha,i}$ along v , and evaluate at $k_{\sigma'} = 0$ for all σ' (corresponding to the weight $\mathbb{1}$) to obtain

$$\frac{\partial}{\partial v} \chi_{\alpha,i}(\pi^n u) = \frac{d}{dk_{\sigma}} \alpha_{U_{t_i}}(\mathbb{1}) \text{ord}_{\mathfrak{p}}(\pi^n) + \alpha_{U_{t_i}}(\mathbb{1}) \log_p(\sigma(u)).$$

By the Steinberg hypothesis, $\alpha_{U_{t_i}}(\mathbb{1})$ is not vanishing and we are done.

By Proposition 1.8, the \mathcal{L} -invariant has codimension at least one and, hence, the second claim follows. □

4. Applications

4.1 Hilbert modular forms

We want to study the case of inner forms of PGL_2 , which are split at \mathfrak{p} , over a totally real number field F in detail. As there is only one simple root we drop it from the notation. If G is equal to PGL_2 , one can attach $2^{[F:\mathbb{Q}]}$ *a priori* different \mathcal{L} -invariants $\mathcal{L}(\pi, p)^{\epsilon}$ to π ; in this case, a conjecture of Spieß (cf. [Spi14, Conjecture 6.4]) states that the definition does not depend on ϵ , i.e.

$$\mathcal{L}(\pi, \mathfrak{p})^{\epsilon} = \mathcal{L}(\pi, \mathfrak{p})^{\epsilon'}$$

for all choices of sign characters ϵ and ϵ' . In the same paper, Remark 6.6(b), Spieß also states that ‘an interesting and difficult problem’ is to show that the \mathcal{L} -invariant is invariant under Jacquet–Langlands transfers. In this section, thanks to Theorem 3.16, we settle both Spieß’ conjecture and his question.

Moreover, let $\rho_{\pi}: \text{Gal}(\overline{\mathbb{Q}}/F) \rightarrow \text{GL}_2(E)$ be the Galois representation associated with π (or rather associated with the Jacquet–Langlands transfer $\text{JL}(\pi)$ of π to PGL_2), which exists by work of Taylor (see [Tay89]). As π is \mathfrak{p} -ordinary, by Theorem 2 of [Wil88] we know that the restriction $\rho_{\pi,\mathfrak{p}}: \text{Gal}(\overline{\mathbb{Q}_p}/F_{\mathfrak{p}})$ of ρ_{π} to a decomposition group at \mathfrak{p} is ordinary. Moreover, because $\pi_{\mathfrak{p}}$ is Steinberg, a result of Saito (see [Sai09]) implies that this restriction is a non-split extension of the trivial character by the cyclotomic character. Therefore, it gives a class $\langle \rho_{\pi,\mathfrak{p}} \rangle \in H^1(\text{Gal}(\overline{\mathbb{Q}_p}/F_{\mathfrak{p}}), E(1))$, where $E(1)$ denotes the Tate twist of E . The Fontaine–Mazur \mathcal{L} -invariant $\mathcal{L}^{\text{FM}}(\rho_{\pi,\mathfrak{p}})$ of $\rho_{\pi,\mathfrak{p}}$ is the orthogonal complement of $\langle \rho_{\pi,\mathfrak{p}} \rangle$ with respect to the local

Tate pairing

$$H^1(\text{Gal}(\overline{\mathbb{Q}_p}/F_{\mathfrak{p}}), E(1)) \times H^1(\text{Gal}(\overline{\mathbb{Q}_p}/F_{\mathfrak{p}}), E) \longrightarrow E.$$

By local class field theory we have a canonical isomorphism

$$H^1(\text{Gal}(\overline{\mathbb{Q}_p}/F_{\mathfrak{p}}), E) \cong \text{Hom}^{\text{ct}}(F_{\mathfrak{p}}^*, E)$$

and, thus, we consider $\mathcal{L}^{\text{FM}}(\rho_{\pi, \mathfrak{p}})$ as a codimension-one subspace of $\text{Hom}^{\text{ct}}(F_{\mathfrak{p}}^*, E)$. We show that automorphic \mathcal{L} -invariants equal the Fontaine–Mazur \mathcal{L} -invariant of the associated Galois representation.

THEOREM 4.1. *Suppose G is an inner form of PGL_2 over a totally real number field F , which is split at the prime \mathfrak{p} of F . Let π be a cuspidal automorphic representation of parallel weight 2 of G that is Steinberg at \mathfrak{p} .*

- (i) *The automorphic \mathcal{L} -invariant of π is independent of ϵ , i.e.*

$$\mathcal{L}(\pi, \mathfrak{p})^\epsilon = \mathcal{L}(\pi, \mathfrak{p})^{\epsilon'}$$

for all choices of ϵ and ϵ' . We therefore put $\mathcal{L}(\pi, \mathfrak{p}) = \mathcal{L}(\pi, \mathfrak{p})^\epsilon$ for any choice of sign character ϵ .

- (ii) *Let $\text{JL}(\pi)$ be the Jacquet–Langlands transfer of π to PGL_2 . The equality*

$$\mathcal{L}(\pi, \mathfrak{p}) = \mathcal{L}(\text{JL}(\pi), \mathfrak{p})$$

holds.

- (iii) *Let ρ_π the Galois representation associated with π . Then*

$$\mathcal{L}(\pi, \mathfrak{p}) = \mathcal{L}^{\text{FM}}(\rho_{\pi, \mathfrak{p}}).$$

Proof. We actually prove that $\mathcal{L}(\pi, \mathfrak{p})^\epsilon = \mathcal{L}^{\text{FM}}(\rho_{\pi, \mathfrak{p}})$ for every ϵ , which implies all other claims. As \mathfrak{m}_π is \mathfrak{p} -étale (with respect to ϵ) we can deform the Hecke eigenvalues of π to a family over an open affinoid subspace of weight space in the following sense: there exists an open affinoid $\mathcal{U} \subseteq \mathcal{W}$ containing the trivial character such that the eigenvalue α_h of each Hecke operator $h \in \mathbb{T}_{\text{sph}}$ acting on $H^q(X_{K^{\mathfrak{p}} \times I_{\mathfrak{p}}}, \mathcal{D}_{\chi_{\mathcal{U}}}^n)^\epsilon$ is an element of $\mathcal{O}_{\mathcal{W}}(\mathcal{U})$. As explained in [BDJ21, Theorem 2.14(ii)], we may shrink \mathcal{U} further such that the common eigenspace associated to the eigenvalues α_h is a free $\mathcal{O}_{\mathcal{W}}(\mathcal{U})$ -submodule of $H^q(X_{K^{\mathfrak{p}} \times I_{\mathfrak{p}}}, \mathcal{D}_{\chi_{\mathcal{U}}}^n)^\epsilon$ of rank equal to the dimension of $H^q(X_{K^{\mathfrak{p}} \times I_{\mathfrak{p}}}, E)_{\mathfrak{m}_\pi}^\epsilon$. By shrinking \mathcal{U} even further we may assume that $\alpha_{t_i} \in \mathcal{O}_{\mathcal{W}}(\mathcal{U})^\times$ has absolute value one for each $i \in \Delta$. Then, by the classicality of small slope overconvergent eigenforms, the specialization of these eigenvalues at a classical point $\lambda \in \mathcal{U}$ are the eigenvalues associated to a cuspidal representation π_λ of weight λ .

Note that the Galois representation ρ_π is irreducible, as the Hilbert modular form associated with π is cuspidal. Thus, by standard arguments (see, for example, Theorem B of [Che14]) there exists (after possibly shrinking \mathcal{U} again) a family of Galois representations $\rho_{\pi, \mathcal{U}}$ over \mathcal{U} passing through ρ_π , i.e. the specialization of $\rho_{\pi, \mathcal{U}}$ at each classical point $\lambda \in \mathcal{U}$ is isomorphic to the Galois representation attached to π_λ . It follows from the results of Saito (see Theorem 1 of [Sai09]) that for every classical point $z \in \mathcal{U}$ the restriction of the Galois representation $\rho_{\pi, z}$ to a decomposition group is determined by the $U_{\mathfrak{p}}$ -eigenvalue. Moreover, the Weil–Deligne module of ρ_π restricted at the decomposition group at \mathfrak{p} is the Weil–Deligne module associated with the Steinberg representation via the local Langlands correspondence. In particular, our representation (with the filtration induced by the ordinarity of the Galois representation) is non-critical special, as defined in § 3.1 of [Din19]; indeed we know that the corresponding (φ, Γ) -module is an extension of special characters, so we just have to check whether the extensions are split or not. If one

extension were split, then the Weil–Deligne representation would also be split, contradicting the fact the monodromy for the Steinberg is maximal.

Comparing the Galois theoretic Colmez–Greenberg–Stevens formula (see [Din19, Theorem 3.4]) with its automorphic counterpart of Theorem 3.16 yields the result. \square

Remark 4.2. (i) In the Hilbert modular form case, the constructions simplify substantially and, thus, one can check that our methods also work for higher weights assuming that the representation is non-critical.

(ii) One can prove the first two claims of Theorem 4.1 without passing to the Galois side. Namely, using methods as in §5 of [Han17] one can show that the eigenvalue of the overconvergent \mathfrak{p} -adic family passing through π is independent of the sign character and stable under the Jacquet–Langlands transfer. Thus, by Theorem 3.16, we can conclude the argument.

(iii) By definition, Fontaine–Mazur \mathcal{L} -invariants are stable under restricting the Galois representation to the absolute Galois group of finite extensions. Thus, we can deduce from Theorem 4.1 that automorphic \mathcal{L} -invariants of Hilbert modular forms are stable under abelian base change with respect to totally real extensions.

(iv) In the case of modular elliptic curves over totally real fields of class number one the equality of automorphic and Fontaine–Mazur \mathcal{L} -invariants was conjectured by Greenberg (see [Gre09, Conjecture 2]). Theorem 4.1 thus implies that the construction of Stark–Heegner points over totally real fields is unconditional (see [GMS15] for a detailed discussion of Stark–Heegner points).

4.2 Unitary groups

Let \tilde{F} be a CM field with totally real subfield F . We assume that the prime \mathfrak{p} of F is split in \tilde{F} . Let U be the unitary group attached to a positive definite hermitian space over \tilde{F} and G the associated adjoint group. By construction, $G_{\mathfrak{p}}$ is isomorphic to $\mathrm{PGL}_n(F_{\mathfrak{p}})$. We can identify the simple roots with respect to the upper triangular Borel with the set $\{1, \dots, n - 1\}$. Let π be an automorphic representation of G such that $\pi_{\infty} = \mathbb{C}$ and $\pi_{\mathfrak{p}}$ is the Steinberg representation of $\mathrm{PGL}_n(F_{\mathfrak{p}})$. In this case the only sign character is the trivial character and, therefore, we drop it from the notation.

By Shin’s appendix to [Gol14] the base change $\mathrm{BC}(\pi)$ of π to PGL_n over \tilde{F} exists and it is Steinberg at both primes of \tilde{F} lying above \mathfrak{p} . By the work of many people (see Theorem 2.1.1 of [BGGT14] for a detailed discussion) we can attach a p -adic Galois representation

$$\rho_{\pi} = \rho_{\mathrm{BC}(\pi)} : \mathrm{Gal}(\overline{\mathbb{Q}}/\tilde{F}) \longrightarrow \mathrm{GL}_n(E)$$

to $\mathrm{BC}(\pi)$. As we consider the trivial coefficient system the Steinberg representation is ordinary (cf. [Ger19, Lemma 5.6]). Therefore, as shown in [Tho15, Theorem 2.4], one deduces from the local–global compatibility theorem of Caraiani (see [Car14]) that the restriction $\rho_{\pi, \mathfrak{p}}$ of ρ_{π} to the decomposition group of a prime above \mathfrak{p} can be brought in the following form: it is upper triangular and the i th diagonal entry is the $(1 - i)$ -power of the cyclotomic character.

Therefore, we have $n - 1$ canonical two-dimensional subquotients $\rho_{\pi, \mathfrak{p}, i}$ which are extensions of $E(-n + i)$ by $E(-n + i + 1)$. We can consider the associated Fontaine–Mazur \mathcal{L} -invariant $\mathcal{L}_i^{\mathrm{FM}}(\rho_{\pi, \mathfrak{p}}) = \mathcal{L}^{\mathrm{FM}}(\rho_{\pi, \mathfrak{p}, i}(n - i))$. Recall that the Weil–Deligne representation associated with the Steinberg has always maximal monodromy so the corresponding (φ, Γ) -module is non-critical split *à la* Ding, see §3.1 of [Din19]. Replacing the local–global compatibility results of Saito by those of Caraiani (see [Car14]), we can apply Theorem 3.4 of [Din19] to the (φ, Γ) -module associated with the Galois representation ρ_{π} and then the same proof as that of Theorem 4.1 yields the following result.

THEOREM 4.3. *Let F be a totally real number field and G the adjoint of a unitary group over F compact at infinity and split at \mathfrak{p} . Let π be an automorphic representation of G such that $\pi_\infty = \mathbb{C}$ and $\pi_{\mathfrak{p}}$ is Steinberg. Suppose π satisfies (SMO), that we can choose the tame level $K^{\mathfrak{p}}$ such that \mathfrak{m}_π is \mathfrak{p} -étale and that the Galois representation ρ_π attached to π is irreducible. Then we have*

$$\mathcal{L}_i^{\text{FM}}(\rho_\pi) = \mathcal{L}_i(\pi, \mathfrak{p})$$

for every $i = 1, \dots, n - 1$.

Remark 4.4. Something can be said also when $F = \mathbb{Q}$ and $G = \text{Sp}_{2g}$. In this case the Galois representations have been constructed by Scholze [Sch15] but local–global compatibility is not known. Still, if one supposes that the $2g + 1$ -dimensional Galois representation $\text{Std}(\rho_\pi)$ associated with π is semistable with maximal monodromy, then the Greenberg–Benois $\mathcal{L}(\text{Std}(\rho_\pi))$ -invariant has been calculated in [Ros15, Theorem 1.3].

In this case the root system of Sp_{2g} can be identified with the set $\{1, \dots, g\}$ via the identification with the root system of GL_g , which is embedded in Sp_{2g} by

$$\text{GL}_g \longrightarrow \text{Sp}_{2g}, \quad A \longmapsto \begin{pmatrix} {}^t A^{-1} & 0 \\ 0 & A \end{pmatrix},$$

and we see that the automorphic \mathcal{L} -invariant $\mathcal{L}_1(\pi, p)$ coincides with that of [Ros15].

The same calculation in §4.2 of [Ros15] gives us that $\mathcal{L}_i(\pi, p)$ is the Greenberg–Benois \mathcal{L} -invariant for $\text{Std}(\rho_\pi)(i - 1)$. (This case has not been treated in [Ros15] as the L -values are not Deligne-critical, but Benois’ definition applies also in this case, see formula (96) in [Ben21].)

If $F_{\mathfrak{p}} \neq \mathbb{Q}_p$, then the comparison is more subtle, as there is only just one Greenberg–Benois \mathcal{L} -invariant per p -adic place, and from the Galois side one needs to consider Galois invariant characters of $F_{\mathfrak{p}}^*$.

5. Beyond discrete series: the Bianchi case

When G_∞ does not fulfil the Harish-Chandra condition, the representation π contributes to several degrees of cohomology of the associated locally symmetric space and the techniques used in Theorem 3.12 break down.

There are two tools to tackle the problem: first, one can use Hansen’s Tor-spectral sequence (see Theorem 3.3.1 of [Han17])

$$\text{Tor}_i^{\mathcal{O}_{\mathcal{W}}}(H^{q+i}(X_{K^{\mathfrak{p}} \times I_{\mathfrak{p}}}, \mathcal{D}_{\chi_{\mathcal{U}}})^{\leq h}, E) \implies H^q(X_{K^{\mathfrak{p}} \times I_{\mathfrak{p}}}, \mathcal{D}_{\chi_{\Sigma}})^{\leq h},$$

where $\mathcal{U} \subseteq \mathcal{W}$ is an open affinoid and $\Sigma \subseteq \mathcal{U}$ is Zariski-closed, to analyse the overconvergent cohomology groups in question; second, one can use cases of Langlands functoriality in p -adic families to reduce to groups, which fulfil the Harish-Chandra condition.

In good situations it should be possible to calculate at least one of the \mathcal{L} -invariants $\mathcal{L}_i^{(d)}(\pi, \mathfrak{p})$ for $0 \leq d \leq \delta$, using Proposition 2.4. The main difficulty is that, in general, most classes do not lift to a big cohomology class as classes in H^{q+1} give lifting obstructions. However, as soon as one can show that at least a class lifts to a family in H^{q+d} (combined with some results on the tangent directions in the eigenvariety) Proposition 2.4 lets us calculate $\mathcal{L}_i^{(d)}(\pi, \mathfrak{p})$. If Venkatesh’s conjecture (stating that the π -isotypic component of the cohomology is generated by the minimal degree cohomology as a module over the derived Hecke algebra) holds, then these \mathcal{L} -invariants $\mathcal{L}_i^{(d)}(\pi, \mathfrak{p})$, for varying d , are essentially all the same by the main result of [Geh19a].

Using unpublished work of Hansen (see also [BW21]) we study the case of Bianchi modular forms. We now fix F to be a quadratic imaginary field where p is unramified and π a cuspidal representation of $\mathrm{PGL}_{2,F}$ of parallel weight 2 such that $\pi_{\mathfrak{q}}$ is Steinberg for all primes \mathfrak{q} lying above p . We put $G = \mathrm{PGL}_{2,F}$ if p is inert and, if p is split, we define G to be the Weil restriction $\mathrm{Res}_{F/\mathbb{Q}} \mathrm{PGL}_{2,F}$. In the first case there is only one simple root and, thus, we drop it from the notation. In the second case, the simple roots can be identified with the two primes above \mathfrak{p} . In both cases, the only sign character is the trivial one and we shall drop it from the notation as well. (The following theorem indicates that in the case of a split prime the partial eigenvarieties we considered before may not be big enough. The case that p is split and π is only Steinberg at one of the primes above p could be handled similarly but one would have to introduce new notation. For the sake of brevity, we do not discuss it further.)

We recall some result on the eigenvariety for Bianchi modular forms due to Hansen and Barrera–Williams (see [BW21, Lemma 4.4] and the proof of Theorem 4.5).

THEOREM 5.1 (Hansen, Barrera–Williams). *Let $\mathcal{U} \subseteq \mathcal{W}$ be an open affinoid neighbourhood of the trivial character.*

- (i) *The system of eigenvalues associated with π appears in $H^d(X_{K^{\mathfrak{p}} \times I_{\mathfrak{p}}}, \mathcal{D}_{\chi_{\mathcal{U}}})$ if and only if $d = 2$.*
- (ii) *There is at least one curve $\mathcal{S} \subseteq \mathcal{U}$ passing through $\mathbb{1}$ such that*

$$H^1(X_{K^{\mathfrak{p}} \times I_{\mathfrak{p}}}, \mathcal{D}_{\chi_{\mathcal{S}}})_{\mathfrak{m}_{\pi}} \neq 0.$$

If such a curve \mathcal{S} is smooth at $\mathbb{1}$, the space is free of rank one over $\mathcal{O}_{\mathcal{S}, \mathbb{1}}$ and the canonical map

$$H^1(X_{K^{\mathfrak{p}} \times I_{\mathfrak{p}}}, \mathcal{D}_{\chi_{\mathcal{S}}})_{\mathfrak{m}_{\pi}} \longrightarrow H^1(X_{K^{\mathfrak{p}} \times I_{\mathfrak{p}}}, E)_{\mathfrak{m}_{\pi}}$$

is surjective.

The following proposition completes the picture by taking into account the cohomology in degree two.

PROPOSITION 5.2. *For every curve \mathcal{S} as in Theorem 5.1(ii) that is smooth at $\mathbb{1}$ we have*

$$H^2(X_{K^{\mathfrak{p}} \times I_{\mathfrak{p}}}, \mathcal{D}_{\chi_{\mathcal{S}}})_{\mathfrak{m}_{\pi}} \neq 0.$$

More precisely, the space is free of rank one over $\mathcal{O}_{\mathcal{S}, \mathbb{1}}$ and the canonical map

$$H^2(X_{K^{\mathfrak{p}} \times I_{\mathfrak{p}}}, \mathcal{D}_{\chi_{\mathcal{S}}})_{\mathfrak{m}_{\pi}} \otimes_{\mathcal{O}_{\mathcal{S}}(\mathcal{S})} E \longrightarrow H^2(X_{K^{\mathfrak{p}} \times I_{\mathfrak{p}}}, E)_{\mathfrak{m}_{\pi}}$$

is an isomorphism. Moreover, $H^2(X_{K^{\mathfrak{p}} \times I_{\mathfrak{p}}}, \mathcal{D}_{\chi_{\mathcal{S}}})_{\mathfrak{m}_{\pi}}$ and $H^1(X_{K^{\mathfrak{p}} \times I_{\mathfrak{p}}}, \mathcal{D}_{\chi_{\mathcal{S}}})_{\mathfrak{m}_{\pi}}$ are isomorphic as modules over the Hecke algebra.

Proof. Let m be a generator of the maximal ideal of the localization. From the short exact sequence

$$0 \rightarrow \mathcal{D}_{\chi_{\mathcal{S}}} \xrightarrow{\times m} \mathcal{D}_{\chi_{\mathcal{S}}} \rightarrow \mathcal{D}_{\mathbb{1}} \rightarrow 0$$

we obtain a long exact sequence in cohomology

$$\begin{aligned} \xrightarrow{\times m} H^1(X_{K^{\mathfrak{p}} \times I_{\mathfrak{p}}}, \mathcal{D}_{\chi_{\mathcal{S}}})_{\mathfrak{m}_{\pi}} &\longrightarrow H^1(X_{K^{\mathfrak{p}} \times I_{\mathfrak{p}}}, E)_{\mathfrak{m}_{\pi}} \xrightarrow{0} H^2(X_{K^{\mathfrak{p}} \times I_{\mathfrak{p}}}, \mathcal{D}_{\chi_{\mathcal{S}}})_{\mathfrak{m}_{\pi}} \\ \xrightarrow{\times m} H^2(X_{K^{\mathfrak{p}} \times I_{\mathfrak{p}}}, \mathcal{D}_{\chi_{\mathcal{S}}})_{\mathfrak{m}_{\pi}} &\longrightarrow H^2(X_{K^{\mathfrak{p}} \times I_{\mathfrak{p}}}, E)_{\mathfrak{m}_{\pi}} \longrightarrow 0. \end{aligned}$$

Here the second map is the zero map as the first map is surjective by the second part of Theorem 5.1, and the last term is zero because the system of eigenvalues for π does not appear in degrees greater by Theorem 5.1(i).

We then get that multiplication by m is injective on $H^2(X_{K^p \times I_p}, \mathcal{D}_{\chi_S})_{\mathfrak{m}_\pi}$ and, therefore, the map

$$H^2(X_{K^p \times I_p}, \mathcal{D}_{\chi_S})_{\mathfrak{m}_\pi} \otimes_{\mathcal{O}_{S,1}} E \longrightarrow H^2(X_{K^p \times I_p}, E)_{\mathfrak{m}_\pi}$$

is an isomorphism. By hypothesis, $H^2(X_{K^p \times I_p}, E)_{\mathfrak{m}_\pi}$ is one-dimensional. Therefore, the $\mathcal{O}_{S,1}$ -module $H^2(X_{K^p \times I_p}, \mathcal{D}_{\chi_S})_{\mathfrak{m}_\pi}$ is cyclic by Nakayama’s Lemma. If it were torsion, then multiplication by m would not be injective on it, which is a contradiction; so it is free.

Let r be a generator of the ideal of $\mathcal{O}_{U,1}$ corresponding to S . The modules $H^1(X_{K^p \times I_p}, \mathcal{D}_{\chi_S})_{\mathfrak{m}_\pi}$ and $H^2(X_{K^p \times I_p}, \mathcal{D}_{\chi_S})_{\mathfrak{m}_\pi}$ are kernel and cokernel, respectively, of the multiplication by r map on $H^2(X_{K^p \times I_p}, \mathcal{D}_{\chi_U})_{\mathfrak{m}_\pi}$. Multiplication by the appropriate power of r yields the sought-after isomorphism. \square

We would like to apply Theorem 3.16 to conclude that \mathcal{L} -invariants are independent of the cohomological degree. There are, however, several problems. First, it is not clear whether one can find a curve \mathcal{S} as in the theorem that is smooth at 1 . Second, if one finds such a curve, the automorphic Colmez–Greenberg–Stevens formula in the inert case only produces one element in the intersection of $\mathcal{L}^{(0)}(f, p)$ and $\mathcal{L}^{(1)}(f, p)$. As these spaces are two-dimensional by Proposition 1.8 this gives us no new information. In the split case the situation is better: one can at least compute the \mathcal{L} -invariant of one of the primes lying above p and, if the tangent space of \mathcal{S} at 1 is generic enough, one can compute both.

COROLLARY 5.3. *Let F be an imaginary quadratic field unramified at p and let π be a cuspidal Bianchi newform of parallel weight 2. Suppose that p splits in F as $p = \mathfrak{p}\bar{\mathfrak{p}}$ and f is Steinberg at both, \mathfrak{p} and $\bar{\mathfrak{p}}$. Then at least one of the equalities*

$$\mathcal{L}^{(0)}(\pi, \mathfrak{p}) = \mathcal{L}^{(1)}(\pi, \mathfrak{p})$$

or

$$\mathcal{L}^{(0)}(\pi, \bar{\mathfrak{p}}) = \mathcal{L}^{(1)}(\pi, \bar{\mathfrak{p}})$$

holds.

Remark 5.4. It was shown in [Geh19a] that the theorem above would also follow from Venkatesh’s conjectures on the action of derived Hecke algebras.

The situation simplifies substantially if π is the base change of a modular form f .

THEOREM 5.5. *Let F be an imaginary quadratic field unramified at p . Let f be a newform of weight 2 that is Steinberg at p and $\text{BC}(f)$ its base change to F . The following hold.*

(i) *If $p = \mathfrak{p}\bar{\mathfrak{p}}$ is split in F , then*

$$\mathcal{L}^{(0)}(\text{BC}(f), \mathfrak{p}) = \mathcal{L}^{(1)}(\text{BC}(f), \mathfrak{p}) = \mathcal{L}^{(0)}(\text{BC}(f), \bar{\mathfrak{p}}) = \mathcal{L}^{(1)}(\text{BC}(f), \bar{\mathfrak{p}}) = \mathcal{L}(f, p).$$

(ii) *If p is inert, we have*

$$\mathcal{L}^{(0)}(\text{BC}(f), p) \cap \text{Hom}^{\text{ct}}(\mathbb{Q}_p^*, E) = \mathcal{L}^{(1)}(\text{BC}(f), p) \cap \text{Hom}^{\text{ct}}(\mathbb{Q}_p^*, E) = \mathcal{L}(f, p).$$

(iii) *If p is inert, we have*

$$\mathcal{L}^{(0)}(\text{BC}(f), p) = \mathcal{L}^{(1)}(\text{BC}(f), p).$$

Proof. In § 5 of [BW19] it is shown that one may take $\mathcal{S} \subseteq \mathcal{W}$ to be the parallel weight curve (see § 5 of [BW19]), which is clearly smooth at 1 . Using p -adic Langlands functoriality as explained in [BW19], the first two claims follow by applying Theorem 3.16 (respectively, its analogue in the Bianchi setting) for the modular form f and its base change to F . The last claim is a

consequence of the second claim and the Galois invariance of automorphic \mathcal{L} -invariants attached to base change representations (see Lemma 3.1 of [Geh19b]). \square

Remark 5.6. In the inert case the equality $\mathcal{L}^{(0)}(\mathrm{BC}(f), p) \cap \mathrm{Hom}^{\mathrm{ct}}(\mathbb{Q}_p^*, E) = \mathcal{L}(f, p)$ was proven in [Geh19b, Lemma 3.3], using Artin formalism for p -adic L -functions and the exceptional zero formula. An analogous result for higher weight forms was proven by Barrera-Salazar and Williams in [BW19, Proposition 10.2].

Our approach can be adapted to higher weights making the construction of Stark–Heegner cycles of [VW21] unconditional in the base change case.

REFERENCES

- AS08 A. Ash and G. Stevens, *p*-adic deformations of arithmetic cohomology, Preprint (2008), <https://math.bu.edu/people/gsh/preprints/Ash-Stevens-02-08.pdf>.
- BGGT14 T. Barnet-Lamb, T. Gee, D. Geraghty and R. Taylor, *Potential automorphy and change of weight*, Ann. of Math. (2) **179** (2014), 501–609.
- BDJ21 D. Barrera Salazar, M. Dimitrov and A. Jorza, *p*-adic L -functions of Hilbert cusp forms and the trivial zero conjecture, J. Eur. Math. Soc. (2021), doi:10.4171/JEMS/1165.
- BW19 D. Barrera Salazar and C. Williams, *Exceptional zeros and \mathcal{L} -invariants of Bianchi modular forms*, Trans. Amer. Math. Soc. **372** (2019), 1–34.
- BW21 D. Barrera Salazar and C. Williams, *Families of Bianchi modular symbols: critical base-change p -adic L -functions and p -adic Artin formalism*, Selecta Math. (N.S.) **27** (2021), 82.
- Ben21 D. Benois, *p*-adic heights and *p*-adic Hodge theory, Mém. Soc. Math. Fr. (N.S.) **167** (2021).
- BDI10 M. Bertolini, H. Darmon and A. Iovita, *Families of automorphic forms on definite quaternion algebras and Teitelbaum’s conjecture*, Astérisque **331** (2010), 29–64.
- Bor76 A. Borel, *Admissible representations of a semi-simple group over a local field with vector fixed under an Iwahori subgroup*, Invent. Math. **35** (1976), 233–260.
- BS76 A. Borel and J.-P. Serre, *Cohomologie d’immeubles et de groupes S -arithmétiques*, Topology **15** (1976), 211–232.
- Car14 A. Caraiani, *Monodromy and local-global compatibility for $l = p$* , Algebra Number Theory **8** (2014), 1597–1646.
- Car79 P. Cartier, *Representations of p -adic groups: a survey*, in *Automorphic forms, representations and L -functions (Proc. Sympos. Pure Math., Oregon State Univ., Corvallis, Ore., 1977), Part 1*, Proceedings of Symposia in Pure Mathematics, vol. XXXIII (American Mathematical Society, Providence, RI, 1979), 111–155.
- Che14 G. Chenevier, *The p -adic analytic space of pseudocharacters of a profinite group and pseudorepresentations over arbitrary rings*, in *Automorphic forms and Galois representations. Vol. 1*, London Mathematical Society Lecture Note Series, vol. 414 (Cambridge University Press, Cambridge, 2014), 221–285.
- Dar01 H. Darmon, *Integration on $H_p \times H$ and arithmetic applications*, Ann. of Math. (2) **154** (2001), 589–639.
- Das05 S. Dasgupta, *Stark–Heegner points on modular jacobians*, Ann. Sci. Éc. Norm. Supér (4) **38** (2005), 427–469.
- DG12 S. Dasgupta and M. Greenberg, *L -invariants and Shimura curves*, Algebra Number Theory **6** (2012), 455–485.
- Din19 Y. Ding, *Simple \mathcal{L} -invariants for GL_n* , Trans. Amer. Math. Soc. **372** (2019), 7993–8042.
- Geh19a L. Gehrmann, *Derived Hecke algebra and automorphic L -invariants*, Trans. Amer. Math. Soc. **372** (2019), 7767–7784.

- Geh19b L. Gehrmann, *Functoriality of automorphic L -invariants and applications*, Doc. Math. **24** (2019), 1225–1243.
- Geh21 L. Gehrmann, *Automorphic \mathcal{L} -invariants for reductive groups*, J. Reine Angew. Math. **779** (2021), 57–103.
- Ger19 D. Geraghty, *Modularity lifting theorems for ordinary Galois representations*, Math. Ann. **373** (2019), 1341–1427.
- Gol14 W. Goldring, *Galois representations associated to holomorphic limits of discrete series*, Compos. Math. **150** (2014), 191–228.
- Gre09 M. Greenberg, *Stark–Heegner points and the cohomology of quaternionic Shimura varieties*, Duke Math. J. **147** (2009), 541–575.
- GMS15 X. Guitart, M. Masdeu and M. H. Sengün, *Darmon points on elliptic curves over number fields of arbitrary signature*, Proc. Lond. Math. Soc. (3) **111** (2015), 484–518.
- Han17 D. Hansen, *Universal eigenvarieties, trianguline Galois representations, and p -adic Langlands functoriality*, J. Reine Angew. Math. **730** (2017), 1–64, with an appendix by James Newton.
- HP83 R. Howe and I. I. Piatetski-Shapiro, *Some examples of automorphic forms on Sp_4* , Duke Math. J. **50** (1983), 55–106.
- JS81a H. Jacquet and J. A. Shalika, *On Euler products and the classification of automorphic forms. II*, Amer. J. Math. **103** (1981), 777–815.
- JS81b H. Jacquet and J. A. Shalika, *On Euler products and the classification of automorphic representations. I*, Amer. J. Math. **103** (1981), 499–558.
- Jan18 F. Januszewski, *Rational structures on automorphic representations*, Math. Ann. **370** (2018), 1805–1881.
- Kna01 A. W. Knap, *Representation theory of semisimple groups*, Princeton Landmarks in Mathematics (Princeton University Press, Princeton, NJ, 2001). An overview based on examples, reprint of the 1986 original.
- Koh11 J. Kohlhaase, *The cohomology of locally analytic representations*, J. Reine Angew. Math. **651** (2011), 187–240.
- KS12 J. Kohlhaase and B. Schraen, *Homological vanishing theorems for locally analytic representations*, Math. Ann. **353** (2012), 219–258.
- LRV12 M. Longo, V. Rotger and S. Vigni, *On rigid analytic uniformization of Jacobians of Shimura curves*, Amer. J. Math. **134** (2012), 1197–1246.
- Oll14 R. Ollivier, *Resolutions for principal series representations of p -adic GL_n* , Münster J. Math. **7** (2014), 225–240.
- Orl05 S. Orlik, *On extensions of generalized Steinberg representations*, J. Algebra **293** (2005), 611–630.
- Ort04 L. Orton, *An elementary proof of a weak exceptional zero conjecture*, Can. J. Math. **56** (2004), 373–405.
- Pia79 I. I. Piatetski-Shapiro, *Multiplicity one theorems*, in *Automorphic forms, representations and L -functions (Proc. Sympos. Pure Math., Oregon State Univ., Corvallis, Ore., 1977), Part 1*, Proceedings of Symposia in Pure Mathematics, vol. XXXIII (American Mathematical Society, Providence, RI, 1979), 209–212.
- Ree92 M. Reeder, *On certain Iwahori invariants in the unramified principal series*, Pacific J. Math. **153** (1992), 313–342.
- Ros15 G. Rosso, *\mathcal{L} -invariant for Siegel–Hilbert forms*, Doc. Math. **20** (2015), 1227–1253.
- Sai09 T. Saito, *Hilbert modular forms and p -adic Hodge theory*, Compos. Math. **145** (2009), 1081–1113.
- Sch15 P. Scholze, *On torsion in the cohomology of locally symmetric varieties*, Ann. of Math. (2) **182** (2015), 945–1066.

- Sev13 M. A. Seveso, *The Teitelbaum conjecture in the indefinite setting*, Amer. J. Math. **135** (2013), 1525–1557.
- Sou87 D. Soudry, *A uniqueness theorem for representations of $\mathrm{GSO}(6)$ and the strong multiplicity one theorem for generic representations of $\mathrm{GSp}(4)$* , Israel J. Math. **58** (1987), 257–287.
- Spi14 M. Spieß, *On special zeros of p -adic L -functions of Hilbert modular forms*, Invent. Math. **196** (2014), 69–138.
- Spi20 M. Spieß, *On \mathcal{L} -invariants associated to Hilbert modular forms*, Preprint (2020), [arXiv:2005.11892](https://arxiv.org/abs/2005.11892).
- Tay89 R. Taylor, *On Galois representations associated to Hilbert modular forms*, Invent. Math. **98** (1989), 265–280.
- Tei90 J. T. Teitelbaum, *Values of p -adic L -functions and a p -adic Poisson kernel*, Invent. Math. **101** (1990), 395–410.
- Tho15 J. A. Thorne, *Automorphy lifting for residually reducible l -adic Galois representations*, J. Amer. Math. Soc. **28** (2015), 785–870.
- Urb11 E. Urban, *Eigenvarieties for reductive groups*, Ann. of Math. (2) **174** (2011), 1685–1784.
- VW21 G. Venkat and C. Williams, *Stark–Heegner cycles attached to Bianchi modular forms*, J. Lond. Math. Soc. **104** (2021), 394–422.
- Vig08 M.-F. Vignéras, *A criterion for integral structures and coefficient systems on the tree of $\mathrm{PGL}(2, F)$* , Pure Appl. Math. Q. **4** (2008), 1291–1316.
- Wil88 A. Wiles, *On ordinary λ -adic representations associated to modular forms*, Invent. Math. **94** (1988), 529–573.

Lennart Gehrmann lennart.gehrmann@uni-due.de

Fakultät für Mathematik, Universität Duisburg-Essen, Thea-Leymann-Straße 9, 45127 Essen, Germany

Giovanni Rosso giovanni.rosso@concordia.ca

Departments of Mathematics and Statistics, Concordia University, 1455 De Maisonneuve Blvd. W., Montréal, Québec, Canada H3G 1M8