

- Michel MF, Wagenvoort JHT. Surveillance of postoperative infections in thoracic surgery. *J Hosp Infect* 1994;27:139-147.
9. Nelson RM, Dries DJ. The economic implications of infection in cardiac surgery. *Ann Thorac Surg* 1986;42:240-246.
 10. Rubinstein E, Green M, Modan M, Amit P. The effects of nosocomial infections on the length and costs of hospital stay. *J Antimicrob Chemother* 1982;9(suppl A):93-100.
 11. Loop FD, Lytle BW, Cosgrove DM, et al. Sternal wound complications after isolated coronary artery bypass grafting: early and late mortality, morbidity, and cost of care. *Ann Thorac Surg* 1990;49:179-187.
 12. Calia FM, Wolinsky E, Mortimer EA Jr, Abrams JS, Rammelkamp CH. Importance of the carrier state as a source of *Staphylococcus aureus* in wound sepsis. *Journal of Hygiene, Cambridge* 1969;67:49-57.
 13. Kluytmans JAJW, Mouton JW, Ijzerman EPF, et al. Nasal carriage of *Staphylococcus aureus* as a major risk factor for wound infections after cardiac surgery. *J Infect Dis* 1995;171:216-219.
 14. Weinstein HJ. The relation between the nasal-staphylococcal-carrier state and the incidence of postoperative complications. *N Engl J Med* 1959;260:1303-1308.
 15. Williams REO, Jevons MP, Shooter RA, et al. Nasal staphylococci and sepsis in hospital patients. *British Medical Journal* 1959;ii:658-663.
 16. Doebbeling BN, Breneman DL, Neu HC, et al. Elimination of *Staphylococcus aureus* nasal carriage in health care workers: analysis of six clinical trials with calcium mupirocin ointment. *Clin Infect Dis* 1993;17:466-474.
 17. Holton DL, Nicolle LE, Diley D, Bernstein K. Efficacy of mupirocin nasal ointment in eradicating *Staphylococcus aureus* nasal carriage in chronic hemodialysis patients. *J Hosp Infect* 1991;17:133-137.
 18. Reagan DR, Doebbeling BN, Pfaller MA, et al. Elimination of coincident *Staphylococcus aureus* nasal and handcarriage with intranasal application of mupirocin calcium ointment. *Ann Intern Med* 1991;114:101-106.
 19. Kluytmans JAJW, Mouton JW, VandenBergh MFQ, et al. Reduction of surgical-site infections in cardiothoracic surgery by elimination of nasal carriage of *Staphylococcus aureus*. *Infect Control Hosp Epidemiol* 1996;17:780-785.
 20. Garner JS, Jarvis WR, Grace Emori T, Horan TC, Hughes JM. CDC definitions for nosocomial infections, 1988. *Am J Infect Control* 1988;16:128-140.
 21. Gertman PM, Restuccia JD. The Appropriateness Evaluation Protocol: a technique for assessing unnecessary days of hospital care. *Med Care* 1981;19:855-871.
 22. Wong ES. Surgical site infections. In: Mayhall CG, ed. *Hospital Epidemiology and Infection Control*. Baltimore, MD: Williams & Wilkins; 1996:154-175.
 23. Haley WH, Schaberg DR, Von Allmen SD, McGowan JE Jr. Estimating the extra charges and prolongation of hospitalization due to nosocomial infections: a comparison of methods. *J Infect Dis* 1980;141:248-257.
 24. Daschner F. Cost-effectiveness on hospital infection control—lessons from the 1990s. *J Hosp Infect* 1989;13:325-336.
 25. Wakefield DS, Pfaller MA, Hammons GT, Massanari RM. Use of the appropriateness evaluation protocol for estimating the incremental costs associated with nosocomial infections. *Med Care* 1987;25:481-488.
 26. Freeman J, McGowan JE Jr. Methodological issues in hospital epidemiology, III: investigating the modifying effects of time and severity of underlying illness on estimates of cost of nosocomial infection. *Reviews of Infectious Diseases* 1984;6:285-300.
 27. Green MS, Rubinstein E, Amit P. Estimating the effects of nosocomial infections on the length of hospitalization. *J Infect Dis* 1982;145:667-672.
 28. Wakefield DS, Pfaller M, Ludke RL, Wenzel RP. Methods for estimating days of hospitalization due to nosocomial infections. *Med Care* 1992;30:373-376.
 29. McGowan JE Jr. Cost and benefit in control of nosocomial infection: methods for analysis. *Reviews of Infectious Diseases* 1981;3:790-797.
 30. Finkler SA. The distinction between costs and charges. *Ann Intern Med* 1982;96:102-109.
 31. Kaiser AB, Petracek MR, Lea JW, et al. Efficacy of cefazolin, cefamandole, and gentamicin as prophylactic agents in cardiac surgery. *Ann Surg* 1987;206:791-797.
 32. Wakefield DS, Helms CM, Massanari RM, Motomi M, Pfaller M. Cost of nosocomial infection: relative contribution of laboratory, antibiotic, and per diem costs in serious *Staphylococcus aureus* infections. *Am J Infect Control* 1988;16:185-192.
 33. Restuccia JD, Gertman PM, Dayno SJ, et al. A comparative analysis of appropriateness of hospital use. *Health Aff* 1984;3:130.
 34. Rishpon S, Lubach S, Epstein LM. Reliability of a method of determining the necessity for hospitalization days in Israel. *Med Care* 1986;24:279-282.
 35. Stevenson RC, Blackmann SC, Willimas CL, Bartzokas CA. Measuring the saving attributable to an antibiotic prescribing policy. *J Hosp Infect* 1988;11:16-25.

Chlamydia Infections Most Common US Notifiable Infectious Disease

Gina Pugliese, RN, MS
Martin S. Favero, PhD

The National Notifiable Disease Surveillance System is a national passive surveillance system comprising 52 diseases designated by the Council of State and Territorial Epidemiologists as reportable to the CDC. Although 1995 was the first year that genital infections with *Chlamydia trachomatis* were nationally notifiable, this condition was the most commonly reported disease for 1995. The 10 most frequently report-

ed nationally notifiable infectious diseases for 1995 were, in descending order, chlamydia, gonorrhea, AIDS, salmonellosis, hepatitis A, shigellosis, tuberculosis, primary and secondary syphilis, Lyme disease, and hepatitis B. The sexually transmitted diseases of chlamydia, gonorrhea, AIDS, syphilis, and hepatitis B accounted for 87% of cases reported. The most commonly reported infectious disease among children aged <5 years was salmonellosis and shigellosis. Gonorrhea remained the most common disease reported among persons

aged 15-24 years. These statistics reflect only those diseases that are reported to state and local health departments, who then report to the CDC. Because resources for conducting surveillance vary widely by disease, patterns of detection may vary widely. As such, the true incidences in the United States probably are underestimated.

FROM: Centers for Disease Control and Prevention. Ten leading notifiable infectious diseases, United States, 1995. *MMWR* 1995;45(41):883-884.