

On the stochastic bifurcations regarding random iterations of polynomials of the form $z^2 + c_n$

TAKAYUKI WATANABE 

*Center for Mathematical Science and Artificial Intelligence,
Chubu University Academy of Emerging Sciences, 1200 Matsumotocho,
Kasugai-shi, Aichi 487-8501, Japan
(e-mail: takawatanabe@isc.chubu.ac.jp)*

(Received 16 July 2022 and accepted in revised form 26 January 2024)

Abstract. In this paper, we consider random iterations of polynomial maps $z^2 + c_n$, where c_n are complex-valued independent random variables following the uniform distribution on the closed disk with center c and radius r . The aim of this paper is twofold. First, we study the (dis)connectedness of random Julia sets. Here, we reveal the relationships between the bifurcation radius and connectedness of random Julia sets. Second, we investigate the bifurcation of our random iterations and give quantitative estimates of bifurcation parameters. In particular, we prove that for the central parameter $c = -1$, almost every random Julia set is totally disconnected with much smaller radial parameters r than expected. We also introduce several open questions worth discussing.

Key words: random dynamical systems, stochastic bifurcation, quadratic polynomials, the Mandelbrot set, mean stability, non-autonomous iterations

2020 Mathematics Subject Classification: 37H20, 37F10 (Primary); 37F12, 37F46, 37H12 (Secondary)

1. Introduction

1.1. Background. In this paper, we discuss the concrete and interesting random holomorphic dynamical systems with parameters $c \in \mathbb{C}$ and $r \geq 0$. More precisely, we will denote $f_c(z) = z^2 + c$ throughout this paper and we consider random iterations of the form $f_{c_n} \circ \dots \circ f_{c_2} \circ f_{c_1}$, where $c_n (n = 1, 2, \dots)$ are complex-valued independent random variables that follow the uniform distribution on the closed disk $\bar{B}(c, r) = \{c' \in \mathbb{C} : |c' - c| \leq r\}$ on the c -plane. The reader is referred to [31, Remark 4.10]. See Setting 4.1 for the rigorous setting.

The aim of this paper is twofold. First, we study the (dis)connectedness of random Julia sets and relate it to the bifurcation, and second, we investigate the bifurcation of our proposed random iterations and give quantitative estimates of bifurcation parameters.

In recent decades, there has been a rapid growth of studies on random dynamical systems. In deterministic autonomous dynamical systems, time-evolution rules are

independent of time and homogeneous; however, they are dependent in random dynamical systems. Motivated by scientific demands, great efforts have been devoted to establishing fundamental results. For this field, refer to the textbook by Arnold [1].

Fornæss and Sibony first studied random dynamical systems of complex analytic maps [12]. They proved that small ‘random perturbations’ of a single map produce very stable dynamics on average. It is then natural to ask what happens when larger noise is added. This question is the central motivation of this paper.

Sumi expanded the results of Fornæss and Sibony by studying minimal sets and established foundations for the study of random holomorphic dynamical systems. He defined the mean stability, and showed that the set of mean stable systems is open and dense in the space of all random dynamical systems having mild conditions [27]. We will quickly review this theory in §2.2. For details, see [28, 29]. See also [30, 31] for the author generalizing the setting from independent and identically distributed (i.i.d.) to non-i.i.d.

These previous studies motivate us to study the bifurcation of (i.i.d.) random quadratic dynamical systems induced by the uniform distribution on $\bar{B}(c, r)$. More precisely, for a fixed parameter $c \in \mathbb{C}$, we can define the bifurcation radius $r_{\text{bif}}(c)$ as the parameter satisfying that the random dynamical system is mean stable if and only if $r \notin \{0, r_{\text{bif}}(c)\}$. Compare this with Theorem 2.16 and Definition 2.17.

The dynamical systems of quadratic polynomials $f_c(z) = z^2 + c$ ($c \in \mathbb{C}$) are extremely important for studying holomorphic dynamical systems. The celebrated Mandelbrot set is an iconic figure on the parameter plane, whose boundary is known as the deterministic bifurcation locus of quadratic polynomials. The Mandelbrot set \mathcal{M} is the set of all parameters whose critical orbit is bounded: $\mathcal{M} = \{c \in \mathbb{C} : f_c^{on}(0) \not\rightarrow \infty (n \rightarrow \infty)\}$, where $f_c^{on} = f_c \circ \dots \circ f_c \circ f_c$ is the (autonomous) n th iterate of a map f_c . The condition $f_c^{on}(0) \not\rightarrow \infty$ is equivalent to the Julia set of f_c being connected. See McMullen’s book [19] and Milnor’s book [21] for details.

Several authors have reported great studies on sample-wise dynamics for random iterations. For a fixed infinite sequence $(c_n)_{n=1}^{\infty}$, we consider the Julia set of compositions $f_{c_n} \circ \dots \circ f_{c_2} \circ f_{c_1}$, which is called a non-autonomous Julia set or a random Julia set.

Brück, Büger, and Reitz investigated the (dis)connectedness of random Julia sets [4]. They found interesting examples that illustrated the difficulty of random iteration compared with deterministic iteration and established several conditions relating the noise amplitude to the connectedness of random Julia sets. See Remark 3.2 and [3, 5].

Brück, Büger, and Reitz showed that if center $c = 0$ and radius $r \leq 1/4$, then every random Julia set of $f_{c_n} \circ \dots \circ f_{c_2} \circ f_{c_1}$ is connected. Later in [17], Lech and Zdunik proved that if $c = 0$ and $r > 1/4$, then almost every random Julia set is *totally* disconnected. For our viewpoint, this is equivalent to $r_{\text{bif}}(0) = 1/4$. One motivation of this paper is to generalize this result for the cases where $c \neq 0$.

1.2. Main results. First, we consider a sufficient condition for the random Julia sets to be totally disconnected almost surely.

Main Result A. (Theorem 3.7) In this statement, the distribution μ of c_n needs not to be the uniform distribution on a disk. Under some assumption, if the critical point 0 tends

to ∞ with probability one, then the random Julia set J_ω is totally disconnected almost surely.

Second, we reveal the relationships between the bifurcation radius r_{bif} and the connectedness of random Julia sets. More precisely, the following holds.

Main Result B. (Theorem 4.6) Suppose that the interior of $\bar{B}(c, r)$ contains a superattracting parameter. Then the following four are equivalent.

- (1) The inequality $r \leq r_{\text{bif}}(c)$ holds.
- (2) The random orbit of $z = 0$ does not tend to ∞ surely.
- (3) The random Julia set is connected for every sample-path.
- (4) The set of all total orbits of $z = 0$ is bounded in \mathbb{C} .

Also, the following four are equivalent.

- (1') The inequality $r > r_{\text{bif}}(c)$ holds.
- (2') The random orbit of $z = 0$ tends to ∞ almost surely.
- (3') The random Julia set is totally disconnected almost surely.
- (4') The set of all total orbits of $z = 0$ is not bounded in \mathbb{C} .

Furthermore, either the former or the latter is valid.

In particular, the following dichotomy holds.

COROLLARY 1.1. Let c be a superattracting parameter. If $r \leq r_{\text{bif}}(c)$, then every random Julia set is connected. If $r > r_{\text{bif}}(c)$, then almost every random Julia set is totally disconnected.

Corollary 1.1 is a generalization of the fact that the autonomous Julia set of a single map f_c is either connected or totally disconnected according to $c \in \mathcal{M}$ or not.

Next, we show some properties and present some quantitative estimates for the bifurcation radius as follows.

Main Result C. (Lemma 4.9, Theorems 4.10, 4.12, 4.14, 4.18, and Examples 4.21 and 4.23) All the following hold.

- (i) For every $c \in \mathbb{C}$, we have $r_{\text{bif}}(c) \leq \text{dist}(c, \partial\mathcal{M})$.
- (ii) For every $c, c' \in \mathbb{C}$, we have $|r_{\text{bif}}(c) - r_{\text{bif}}(c')| \leq |c - c'|$.
- (iii) If $0 \leq c \leq 1/4$, then $r_{\text{bif}}(c) = 1/4 - c$.
- (iv) If $-1/2 \leq c < 0$, then $r_{\text{bif}}(c) \leq 1/4 - c - c^2$.
- (v) We have $0.0386 \dots \leq r_{\text{bif}}(-1) \leq 0.0399 \dots$.
- (vi) For the airplane parameter $\tilde{c}_3 \approx -1.75487766$, we have $r_{\text{bif}}(\tilde{c}_3) \leq 0.0021$.

Here, $\text{dist}(c, \partial\mathcal{M})$ denotes the Euclidean distance from c to the boundary of the Mandelbrot set $\partial\mathcal{M}$. Statement (iii) gives the examples of c satisfying equality $r_{\text{bif}}(c) = \text{dist}(c, \partial\mathcal{M})$. However, statements (iv), (v), and (vi) give the examples of c satisfying strict inequality $r_{\text{bif}}(c) < \text{dist}(c, \partial\mathcal{M})$.

It is worth noting that for $c = -1$ or \tilde{c}_3 , statements (v) and (vi) illustrate a large gap $r_{\text{bif}}(c) \ll \text{dist}(c, \partial\mathcal{M})$. Furthermore, we also have $r_{\text{bif}}(c + \epsilon) \ll \text{dist}(c + \epsilon, \partial\mathcal{M})$ for

a small $\epsilon \in \mathbb{C}$. The proofs of statements (v) and (vi) rely on parabolic implosion in a stochastic context. See Theorem 4.19.

Combining statement (iv) with Corollary 1.1, we give a stronger result compared with the result by Lech and Zdunik. Compare Corollary 4.15 with Theorem 3.5. Moreover, we give a result for the case where c is far from 0.

COROLLARY 1.2. (Corollaries 4.15 and 4.25) *Almost every random Julia set is totally disconnected:*

- if the central parameter $-1/4 \leq c < 0$ and the noise amplitude $r > 1/4 - c - c^2$; or
- if the central parameter $c = -1 + \epsilon$ and the noise amplitude $r > 0.0399 \dots + |\epsilon|$; or
- if the central parameter $c = \tilde{c}_3 + \epsilon$ and the noise amplitude $r > 0.0021 + |\epsilon|$.

This corollary gives new examples which satisfy that the random Julia set is totally disconnected almost surely even if $\bar{B}(c, r) \subset \text{int } \mathcal{M}$.

In addition, Main Result B gives a quasiconformal conjugacy between Julia sets.

Main Result D. (Theorem 4.29 and Corollary 4.30) Suppose that the interior of $\bar{B}(c, r)$ contains a superattracting parameter \tilde{c} and suppose $r < r_{\text{bif}}(c)$. Then for every $(c_n)_{n=1}^\infty, (c'_n)_{n=1}^\infty \in \bar{B}(c, r)^\mathbb{N}$, there exist $K \geq 1$ and a sequence of maps $\{\varphi_n\}_{n \geq 0}$ such that φ_n maps the Julia sets of $(c_n)_{n=1}^\infty$ onto the Julia sets of $(c'_n)_{n=1}^\infty$ K -quasiconformally and $\varphi_{n+1} \circ f_{c_{n+1}} = f_{c'_{n+1}} \circ \varphi_n$ on the iterated Julia set for every $n \geq 0$. In particular, there exists a quasiconformal map φ which maps the autonomous Julia set $J_{\tilde{c}}$ onto the non-autonomous Julia set.

1.3. *Structure of the paper.* In §2, we discuss the elementary properties in the general form and define the bifurcation radius. Moreover, we quickly review Sumi’s theory on minimal sets and mean stability in §2.2. In §3, we establish some tools to decide connectedness and total disconnectedness. These are developed by Brück, Büger, and Reitz [4], and Lech and Zdunik [17]. In addition, we show Main Result A relating bifurcation and connectedness. In §4, we prove Main Results B, C, and D. In §5, we discuss some open problems and generalization to other cases.

2. *Preliminaries*

In this section, we define the bifurcation radius and discuss the elementary properties required in subsequent sections. We consider minimal sets and mean stability, which appear in §2.2. They describe set-valued dynamics and we can apply them to quenched, or averaged, dynamics. The minimal sets provide the basis for our discussion and define the bifurcation radius, which is the main subject in this paper.

2.1. *Setting and notation.* In this section, we consider a slightly general setting.

Setting 2.1. Let μ be a Borel probability measure on the parameter plane \mathbb{C} with compact support. Denote the support by $\text{supp } \mu$. We define the probability measure \mathbb{P}_μ as the one-sided infinite product of μ , which is supported on $\Omega_\mu = \prod_{n=1}^\infty \text{supp } \mu$ with the Borel σ -algebra \mathfrak{B}_μ .

The most simple and important example for μ is the normalized Lebesgue measure on $\bar{B}(c, r) = \{c' \in \mathbb{C} : |c' - c| \leq r\}$, which means the uniform distribution. Another example is the Dirac measure at c , which can be regarded as a deterministic dynamical system. We analyze these examples in §4.

Our random dynamical systems are the stochastic process perturbed by i.i.d. noise in a sense. One can generalize the setting to a non-i.i.d. setting, which we do not pursue here. For details, the reader is referred to [30, 31].

Regarding noise, a natural consideration may be a distribution μ with unbounded support, say the Gaussian distribution. However, in unbounded cases, there are no planar attractors which we intend to investigate. Thus, we assume that $\text{supp } \mu$ is compact. See [5] for the detail and see also [9] for more wild phenomena for the distributions with unbounded supports.

The following is a well-known fact. See, for example, [11, §6].

LEMMA 2.2. *Denote by σ the natural left shift, that is, $\sigma\omega = (c_2, c_3, \dots)$ if $\omega = (c_1, c_2, \dots)$. Then $\sigma : \Omega_\mu \rightarrow \Omega_\mu$ is a measure-preserving ergodic transformation with respect to \mathbb{P}_μ .*

We now define the Julia set for every noise realization ω .

Definition 2.3. For each $n \in \mathbb{N}$ and infinite sequence of parameters $\omega = (c_n)_{n=1}^\infty \in \mathbb{C}^\mathbb{N}$, we denote $f_\omega^{(n)} := f_{c_n} \circ \dots \circ f_{c_2} \circ f_{c_1}$.

Definition 2.4. For each $\omega = (c_n)_{n=1}^\infty \in \mathbb{C}^\mathbb{N}$, we define the non-autonomous Julia set or the random Julia set of ω by

$$J_\omega = \{z \in \mathbb{C} : \{f_\omega^{(n)}\}_{n=1}^\infty \text{ is not normal on any neighborhood of } z\}.$$

If there exists $c_0 \in \mathbb{C}$ such that $c_n = c_0$ for every $n \in \mathbb{N}$, then for $\omega = (c_n)_{n=1}^\infty$, the non-autonomous Julia set J_ω is equal to the usual (autonomous) Julia set of f_{c_0} .

Analogously to the usual case, the non-autonomous Julia set is the common boundary of the basin at infinity and the filled Julia set defined as follows.

Definition 2.5. For each $\omega = (c_n)_{n=1}^\infty \in \mathbb{C}^\mathbb{N}$, we define the non-autonomous basin at infinity of ω by $A_\omega = \{z \in \widehat{\mathbb{C}} : f_\omega^{(n)}(z) \rightarrow \infty \text{ as } n \rightarrow \infty\}$. We define the non-autonomous filled Julia set of ω by $K_\omega = \widehat{\mathbb{C}} \setminus A_\omega$.

Here, $\widehat{\mathbb{C}} = \mathbb{C} \cup \{\infty\}$ denotes the Riemann sphere, which is homeomorphic to the real two-dimensional sphere. We endow $\widehat{\mathbb{C}}$ with the spherical metric d . A polynomial map $f : \mathbb{C} \rightarrow \mathbb{C}$ can be extended analytically to $f : \widehat{\mathbb{C}} \rightarrow \widehat{\mathbb{C}}$ by letting $f(\infty) = \infty$.

For our convenience, we list the elementary properties of non-autonomous Julia sets, basins at infinity, and filled Julia sets. For $R > 0$, we denote by D_R the open disk $D_R = \{z \in \mathbb{C} : |z| < R\}$ on the dynamical plane.

LEMMA 2.6. *Let μ be a Borel probability measure on \mathbb{C} with compact support. Then we have all the following.*

- (1) *There exists $R > 0$ such that $f_\omega^{(n)}(\widehat{\mathbb{C}} \setminus D_R) \subset \widehat{\mathbb{C}} \setminus D_{2R}$ for every $n \in \mathbb{N}$ and $\omega \in \Omega_\mu$.*
- (2) *Take $R > 0$ as above. The non-autonomous basin at infinity is then the union of the increasing sequence of the open subsets; $A_\omega = \bigcup_{n=1}^\infty (f_\omega^{(n)})^{-1}(\widehat{\mathbb{C}} \setminus D_R)$ for every $\omega \in \Omega_\mu$. Hence, A_ω is an open neighborhood of ∞ . Conversely, $K_\omega = \bigcap_{n=1}^\infty (f_\omega^{(n)})^{-1}(\overline{D_R})$ is a non-empty compact subset for every $\omega \in \Omega_\mu$.*
- (3) *For $\omega \in \Omega_\mu$, we have $\partial A_\omega = J_\omega = \partial K_\omega$.*
- (4) *For $\omega \in \Omega_\mu$, we have $A_\omega = f_{c_1}^{-1}(A_{\sigma\omega})$, $K_\omega = f_{c_1}^{-1}(K_{\sigma\omega})$, and $J_\omega = f_{c_1}^{-1}(J_{\sigma\omega})$.*

For the proof, see [4]. Note that the existence of such R is due to the compactness of $\text{supp } \mu$.

2.2. *Minimal sets, mean stability, and bifurcation.* Consider the action of polynomial semigroups whose product is the composition of maps. Semigroup actions are related to random dynamical systems as set-valued dynamics. Here, we review the theory of polynomial semigroups, which enables us to define bifurcations.

Generally, one can consider rational semigroups, not polynomial. However, we focus on polynomial maps since we are interested in quadratic polynomials. The interested reader is referred to [15, 24].

We start with the definition of polynomial semigroups. Denote by Poly the space of all polynomials of degree two or more endowed with the topology of uniform convergence on $\widehat{\mathbb{C}}$. We say that G is a polynomial semigroup if G is a non-empty subset of Poly closed under mapping composition. Let Γ be a subset of Poly. We say that G is generated by Γ if for every $g \in G$, there exist $n \in \mathbb{N}$ and $\gamma_1, \gamma_2, \dots, \gamma_n \in \Gamma$ such that $g = \gamma_n \circ \dots \circ \gamma_2 \circ \gamma_1$.

Definition 2.7. Let μ be a Borel probability measure on \mathbb{C} with compact support. We define the polynomial semigroup G_μ as the semigroup generated by the compact set $\Gamma = \{f_c \in \text{Poly} : c \in \text{supp } \mu\}$. Here $f_c(z) = z^2 + c$ is a quadratic polynomial, as we will assume throughout this paper.

Note that the map $\text{Poly} \times \widehat{\mathbb{C}} \ni (g, z) \mapsto g(z) \in \widehat{\mathbb{C}}$ is continuous. This implies that the map $\Omega \times \widehat{\mathbb{C}} \ni (\omega, z) \mapsto f_\omega^{(n)}(z) \in \widehat{\mathbb{C}}$ is also continuous for every $n \in \mathbb{N}$.

For every polynomial semigroup, we define the Julia set and Fatou set as follows. The following proposition connects the Julia set of polynomial semigroup and the random Julia sets. The idea of the proof can be found in [25].

Definition 2.8. For a polynomial semigroup G , define the Julia set by

$$J(G) = \{z \in \widehat{\mathbb{C}} : G \text{ is not normal on any neighborhood of } z\}.$$

We call the complement $F(G) = \widehat{\mathbb{C}} \setminus J(G)$ the Fatou set of G .

PROPOSITION 2.9. *For a Borel probability measure μ on \mathbb{C} with compact support, we have $J(G_\mu) = \bigcup_{\omega \in \Omega_\mu} J_\omega$.*

Next, we define minimal sets which play a crucial role in studying random holomorphic dynamical systems.

Definition 2.10. Let G be a polynomial semigroup. A subset $L' \subset \widehat{\mathbb{C}}$ is said to be forward invariant under G , or forward G -invariant, if $g(L') \subset L'$ for every $g \in G$. A subset $L \subset \widehat{\mathbb{C}}$ is called a *minimal* set of G if L is a minimal element of the set of all forward G -invariant, non-empty, and compact subsets of $\widehat{\mathbb{C}}$ with respect to the inclusion relation \subset .

We can easily show that a non-empty compact set $L \subset \widehat{\mathbb{C}}$ is minimal if and only if for every $z \in L$, we have $\overline{\bigcup_{g \in G} \{g(z)\}} = L$.

Trivially, the singleton $\{\infty\}$ is a minimal set of G for every polynomial semigroup G . Thus, we are interested mainly in planar minimal sets as defined below.

Definition 2.11. A subset $L \subset \widehat{\mathbb{C}}$ is said to be *planar* if $\infty \notin L$.

For example, if $\mu = \delta_c$ is the Dirac measure at c , then every periodic cycle of f_c is a minimal set of G_{δ_c} . Hence, there is an infinite number of planar minimal sets of G_{δ_c} for every $c \in \mathbb{C}$.

For the number of minimal sets, we have the following property.

LEMMA 2.12. *Let G_1 and G_2 be polynomial semigroups generated by Γ_1 and Γ_2 , respectively. If $\Gamma_1 \subset \Gamma_2$, then every minimal set L_2 of G_2 contains some minimal set L_1 of G_1 .*

Proof. By definition, a minimal set L_2 of G_2 is forward invariant under G_2 . Since $\Gamma_1 \subset \Gamma_2$, the set L_2 is forward invariant also under G_1 . It follows from Zorn’s lemma that there exists a minimal set L_1 of G_1 such that $L_1 \subset L_2$. □

Definition 2.13. Let G be the polynomial semigroup generated by a subset Γ of Poly. A minimal set $L \subset \widehat{\mathbb{C}}$ of G is said to be *attracting* if there exist $N \in \mathbb{N}$ and non-empty open subsets U and V of the Fatou set $F(G)$ such that the following two conditions hold.

- (i) $L \Subset V \Subset U \Subset F(G)$.
- (ii) For every $f_1, f_2, \dots, f_N \in \Gamma$, we have $f_N \circ \dots \circ f_2 \circ f_1(U) \Subset V$.

Here, $A \Subset B$ means that the closure \overline{A} is a relatively compact subset of B .

One can easily show that if the polynomial semigroup G is generated by a compact subset, then the minimal set $\{\infty\}$ is attracting. By using the hyperbolic metric, we have the following characterization of attracting minimal sets. For the proof, see [28, Remark 3.5] or [31, Lemma 2.8].

PROPOSITION 2.14. *Let G be the polynomial semigroup generated by a compact subset Γ of Poly. A minimal set $L \subset F(G)$ of G is attracting if and only if the following holds. There exist $N \in \mathbb{N}$, $0 < \alpha < 1$, $C > 0$, and a neighborhood W of L such that for every $k \in \mathbb{N}$, $f_1, f_2, \dots, f_{kN} \in \Gamma$, and $z, z_0 \in W$, the spherical metric satisfies*

$$d(f_{kN} \circ \dots \circ f_1(z), f_{kN} \circ \dots \circ f_1(z_0)) \leq C\alpha^k d(z, z_0).$$

Now, we present the classification theorem, which states that every minimal set either is attracting, intersects the Julia set, or intersects some rotation domain. Compare the following theorem with the classification theorem for autonomous iteration of a single map. For the latter classical theorem, see [21, Theorem 16.1] or [6, Theorem IV.2.1].

THEOREM 2.15. *Let G be the polynomial semigroup generated by a compact subset Γ of Poly. A minimal set $L \subset \widehat{\mathbb{C}}$ of G is either one of the following three.*

- (i) L is attracting.
- (ii) $L \cap J(G) \neq \emptyset$.
- (iii) $L \subset F(G)$ and there exist $g \in G$ and a connected component U of $F(G)$ with $L \cap U \neq \emptyset$ such that $g(U) \subset U$ and U is a subset of a Siegel disk of autonomous iteration of g .

For the proof, see [28, Lemma 3.8]. The classification theorem leads to the following definitions. We say that a minimal set is *J-touching* if it is of type (ii) in Theorem 2.15. In addition, a minimal set is *sub-rotative* if it is of type (iii) in Theorem 2.15.

However, these two types of minimal sets are uncommon compared with the attracting minimal sets as stated by Theorem 2.16. Denote by \mathcal{H} the set of all parameters $c \in \mathcal{M}$ such that f_c is hyperbolic.

THEOREM 2.16. *For every $c \in \mathbb{C}$ and $r > 0$, define the polynomial semigroup $G_{c,r}$ as the semigroup generated by $\{f_{c'} \in \text{Poly} : |c' - c| \leq r\}$. Here $f_c(z) = z^2 + c$. Then $G_{c,r}$ has an attracting minimal set $\{\infty\}$. Other than $\{\infty\}$, possible planar minimal sets are described as follows.*

- If $c \notin \mathcal{H}$, then $G_{c,r}$ has no planar minimal sets for every $r > 0$.
- If $c \in \mathcal{H}$, then there exists $r_{\text{bif}}(c) > 0$ such that the following hold.
 - If $0 < r < r_{\text{bif}}(c)$, then $G_{c,r}$ has exactly one planar minimal set and it is attracting.
 - If $r = r_{\text{bif}}(c)$, then $G_{c,r}$ has exactly one planar minimal set and it is not attracting.
 - If $r_{\text{bif}}(c) < r$, then $G_{c,r}$ has no planar minimal sets.

Theorem 2.16 motivates us to define the bifurcation radius as follows. Note that our bifurcation occurs at most once because f_c is of degree 2, and in general, our bifurcation occurs at most $d - 1$ times if we work on polynomial maps of degree d .

Definition 2.17. For every $c \in \mathbb{C}$, we define the bifurcation radius $r_{\text{bif}}(c) \geq 0$ according to the following cases. If $c \notin \mathcal{H}$, we define $r_{\text{bif}}(c) = 0$. If $c \in \mathcal{H}$, we define the bifurcation radius $r_{\text{bif}}(c) > 0$ as the unique $r > 0$ such that $G_{c,r}$ has the non-attracting planar minimal set.

Equivalently, the bifurcation radius $r_{\text{bif}}(c)$ is the maximum value for $r \geq 0$ such that $G_{c,r}$ has a planar minimal set, or equivalently, is the infimum value for $r \geq 0$ such that $G_{c,r}$ has no planar minimal sets.

The rest of this subsection is devoted to emphasizing the importance of deciding on the bifurcation parameters by presenting some interesting phenomena. See [27, §3.6] for the details.

Definition 2.18. Let G be the polynomial semigroup generated by a compact subset Γ of Poly. We say that G is mean stable if all the minimal sets of G are attracting.

By Theorem 2.16, for every $c \in \mathbb{C}$ and $r \geq 0$, the polynomial semigroup $G_{c,r}$ is mean stable if and only if $r \notin \{0, r_{\text{bif}}(c)\}$.

Before describing the dynamics of a mean stable system, we define the function representing the probability of random orbits tending to an attractor.

Definition 2.19. Let μ be a Borel probability measure on \mathbb{C} with compact support. For every attracting minimal set L of G_μ , we define the function $T_{L,\mu}: \widehat{\mathbb{C}} \rightarrow [0, 1]$ by

$$T_{L,\mu}(z) := \mathbb{P}_\mu(\{\omega \in \Omega_\mu : d(f_\omega^{(n)}(z), L) \rightarrow 0 \ (n \rightarrow \infty)\})$$

for every point $z \in \widehat{\mathbb{C}}$, where $d(w, L) = \min_{w' \in L} d(w, w')$.

The following theorem reveals an interesting noise-induced order. That is, almost every orbit is attracted by some attracting minimal set and the probability $T_{L,\mu}(z)$ tending to the attractor L depends continuously on initial point z if the system is mean stable. For the details, see [27, Theorem 3.15].

Definition 2.20. For a polynomial semigroup G , define the kernel Julia set by

$$J_{\text{ker}}(G) = \bigcap_{g \in G} g^{-1}(J(G)).$$

THEOREM 2.21. *Let μ be a Borel probability measure on \mathbb{C} with compact support. If $J_{\text{ker}}(G) = \emptyset$, then $T_{\{\infty\},\mu}$ is continuous on $\widehat{\mathbb{C}}$. Moreover, the sum satisfies $\sum_L T_{L,\mu}(z) = 1$ for every $z \in \widehat{\mathbb{C}}$ where the sum runs over all attracting minimal sets L .*

The following gives a useful sufficient condition for $J_{\text{ker}}(G) = \emptyset$.

LEMMA 2.22. *Let μ be a Borel probability measure on \mathbb{C} with compact support. If $\text{int}(\text{supp } \mu) \neq \emptyset$, then $J_{\text{ker}}(G_\mu) = \emptyset$.*

See [27, Lemma 5.34] for the proof. Notably, the continuity of $T_{L,\mu}$ in Theorem 2.21 does not occur in deterministic dynamical systems. Namely, if μ is the Dirac measure at c , then the function $T_{\{\infty\},\mu}$ takes 1 on $A(f_c)$ and 0 on $K(f_c)$, where $A(f_c) = A_\omega$ and $K(f_c) = K_\omega$ for $\omega = (c, c, \dots)$. Thus, $T_{\{\infty\},\mu}$ is discontinuous on the autonomous Julia set $J(f_c) = J_\omega$.

Theorem 2.21 suggests that the dynamics of a mean stable system is similar to that of a hyperbolic map. In the latter case, every orbit on the Fatou set is contracted to an attracting cycle, while the complementary Julia set has zero areas.

The set of all hyperbolic rational maps is conjectured to be open and dense in the space of all rational maps. Even in the quadratic polynomial case, whether or not $\overline{\mathcal{H}} = \mathcal{M}$ remains unsolved. As is well known, this conjecture can be affirmatively solved if one proves that the boundary $\partial\mathcal{M}$ is locally connected. Meanwhile, it is still open whether or not the boundary of the Mandelbrot set has a positive area, although Shishikura showed that it has the full Hausdorff dimension.

Compared with the autonomous case, the following results show that we can solve random versions of the abovementioned conjectures. For the proof, see [28, 31].

THEOREM 2.23. *For every $c \in \mathbb{C}$ and $r \geq 0$, define the polynomial semigroup $G_{c,r}$ as the semigroup generated by $\{f_{c'} \in \text{Poly} : |c' - c| \leq r\}$. Then we have all the following statements.*

- (i) The set $\{(c, r) \in \mathbb{C} \times [0, \infty) : G_{c,r} \text{ is mean stable}\}$ is open and dense in $\mathbb{C} \times [0, \infty)$ with the usual topology.
- (ii) For all but countably many $r \in [0, \infty)$, the set $\{c \in \mathbb{C} : G_{c,r} \text{ is not mean stable}\}$ has zero two-dimensional Lebesgue measure.

We also have the following tameness if noise is added.

THEOREM 2.24. *Let μ be a Borel probability measure on \mathbb{C} with compact support. If $J_{\ker}(G) = \emptyset$, then the random Julia set J_ω has zero two-dimensional Lebesgue measure for \mathbb{P}_μ -almost every $\omega \in \Omega_\mu$.*

One may mistakenly consider that Theorem 2.24 is insignificant since it is difficult to construct autonomous Julia sets with positive areas. However, compared with the autonomous case, constructing the non-autonomous Julia set of positive areas is not very difficult. Jones *et al* found that there exists $\omega = (c_n)_{n=1}^\infty$ with $c_n \in \{0, 1/2, 1/4\}$ for all $n \in \mathbb{N}$ such that J_ω has positive areas. They use the parabolic fixed point $1/2$ of the map $f_{1/4}$. For the proof, see [10, §5.2]. Developing the idea, Comerford proved the existence of measurable invariant sequences of line fields [8].

As presented above, the dynamics of mean stable systems is similar to that of hyperbolic polynomial maps. However, they cause interesting phenomena that cannot occur in deterministic dynamical systems, e.g., continuity of $T_{L,\mu}$, finiteness of the set of all minimal sets, etc. This motivates us to determine the bifurcation radius where the mean stability breaks.

3. Total disconnectedness

The systematic study of random iterations of the quadratic polynomial was done by Brück and Büger. In particular, Brück, Büger, and Reitz investigated the (dis)connectedness of the non-autonomous Julia sets in [4]. Here, we present some conditions for the random Julia set to be connected or totally disconnected.

The following [4, Theorem 1.1] shows that the critical orbits determine connectedness of Julia sets. Compared with the autonomous case, we need to modify the statement suitably.

THEOREM 3.1. *Let $\omega = (c_n)_{n=1}^\infty \in \mathbb{C}^\mathbb{N}$. Then the non-autonomous Julia set J_ω is connected if and only if the orbit $\{f_{\sigma^k \omega}^{(n)}(0)\}_{n=1}^\infty$ is bounded in \mathbb{C} for every $k \in \mathbb{N}$.*

Recall that if $\omega = (c_n)_{n=1}^\infty$, then $f_{\sigma^k \omega}^{(n)} = f_{c_{k+n}} \circ \dots \circ f_{c_{k+2}} \circ f_{c_{k+1}}$ by definition. We remark on the difficulty of the disconnected case compared with the autonomous case.

Remark 3.2. For an autonomous case, the Julia set of a quadratic polynomial is either connected or totally disconnected [21, Problem 9-f]. However, for a non-autonomous case, we can construct a Julia set that has more than one but finitely many connected components. For instance, if $|c_1| > 1$ and $c_n = 0$ for every $n \geq 2$, then $J_\omega = f_{c_1}^{-1}(J_{\sigma \omega})$ has exactly two connected components where $\omega = (c_n)_{n=1}^\infty$. Similarly, we can construct various non-autonomous Julia sets with finitely many connected components. Furthermore, there exists $\omega = (c_n)_{n=1}^\infty$ such that $|c_n| < 2$ for every $n \in \mathbb{N}$ and $f_{\sigma^k \omega}^{(n)}(0) \rightarrow \infty$ as $n \rightarrow \infty$ for every $k \in \mathbb{N}$, but the filled Julia set K_ω contains a segment (see [4, Example 4.4]). Thus, it

is interesting to determine when the non-autonomous Julia set J_ω is connected, has finitely many connected components, is totally disconnected, and so on.

For the case when μ is the normalized Lebesgue measure on the disk $\bar{B}(0, r)$, Brück, Bürger, and Reitz showed that $r = 1/4$ is the critical value where connectedness drastically changes as we see below. If $r \leq 1/4$, then the random Julia set J_ω is connected for every $\omega \in \Omega_\mu$ [4, Remark 1.2]. Brück, Bürger, and Reitz then questioned whether the random Julia set J_ω is *totally* disconnected almost surely if $r > 1/4$.

Lech and Zdunik [17] showed that it is true. We pay attention to their idea for a while.

Definition 3.3. Let μ be a Borel probability measure on \mathbb{C} with compact support. Let $R > 0$ be a large number as in Lemma 2.6(1).

For every $\omega \in \Omega_\mu$, define the escape time $k(z, \omega)$ of z from D_R by

$$k(z, \omega) = \begin{cases} \min\{n \in \mathbb{N} : |f_\omega^{(n)}(z)| > R\} & \text{if } z \in A_\omega, \\ \infty & \text{if } z \in K_\omega. \end{cases}$$

We say that the critical point 0 is *typically fast escaping* if there exists $\gamma > 0$ such that $\mathbb{P}_\mu(\{\omega \in \Omega_\mu : k(0, \omega) > k\}) \leq e^{-\gamma k}$ for every $k \in \mathbb{N}$.

Note that the escape time is related to the value of the Green function

$$G_\omega(z) = \lim_{n \rightarrow \infty} \frac{1}{2^n} \log_+ |f_\omega^{(n)}(z)|,$$

where $\log_+ a = \max\{a, 0\}$ for $a \in \mathbb{R}$. By [17, Proposition 8], there exists $C > 0$ such that for every $\omega \in \Omega_\mu$ and $z \in A_\omega \cap D_R$, we have

$$C^{-1}2^{-k(z,\omega)} \leq G_\omega(z) \leq C2^{-k(z,\omega)}.$$

Hence, typical fast escaping can be defined using the Green functions. For details, see [17]. For other discussions on the Green function, see also [12, 16, 26].

The following theorem gives a sufficient condition of random Julia sets to be totally disconnected almost surely. For the proof, see [17, Theorem 11].

THEOREM 3.4. *Let μ be a Borel probability measure on \mathbb{C} with compact support. If the critical point 0 is typically fast escaping, then the Julia set J_ω is totally disconnected for \mathbb{P}_μ -almost every $\omega \in \Omega_\mu$.*

Using the theorem above, the authors of [17] proved the following theorem.

THEOREM 3.5. *Let μ be a Borel probability measure on \mathbb{C} with compact support. If $\text{supp } \mu \supset \bar{B}(0, 1/4)$ and $\text{supp } \mu \neq \bar{B}(0, 1/4)$, then the random Julia set J_ω is totally disconnected for \mathbb{P}_μ -almost every $\omega \in \Omega_\mu$.*

Consider the case where μ is the normalized Lebesgue measure on the disk $\bar{B}(c, r)$. From above, the connectedness changes at $r = 1/4$ if $c = 0$. Since the value $1/4$ is the distance from $c = 0$ to the boundary of the Mandelbrot set, one may conjecture that the transition from connected to disconnected Julia sets equals the distance from c to the boundary of the Mandelbrot set for any $c \in \mathcal{M}$. However, Theorem 3.5 states that

even if $\text{supp } \mu$ is contained in the closure of the main cardioid of the Mandelbrot set, it can happen that the random Julia set J_ω is totally disconnected almost surely.

Before giving the sufficient condition of typical fast escaping, we show the following lemma.

LEMMA 3.6. *Let μ be a Borel probability measure on \mathbb{C} with compact support. Define $E = \{z \in \widehat{\mathbb{C}} : T_{\{\infty\},\mu}(z) = 1\}$. If $z \in E$ and $k \in \mathbb{N}$, then $f_\omega^{(k)}(z) \in E$ for \mathbb{P}_μ -almost every $\omega \in \Omega_\mu$.*

Proof. Since \mathbb{P}_μ is the product of the probability measure μ , Fubini's theorem implies

$$T_{\{\infty\},\mu}(z) = \int_{\bar{B}(c,r)} T_{\{\infty\},\mu}(f_{c_1}(z)) \, d\mu(c_1)$$

for every $z \in \widehat{\mathbb{C}}$. By induction, we have $T_{\{\infty\},\mu}(z) = \int_{\Omega} T_{\{\infty\},\mu}(f_\omega^{(k)}(z)) \, d\mathbb{P}_\mu(\omega)$. Thus, if $T_{\{\infty\},\mu}(z) = 1$, then $T_{\{\infty\},\mu}(f_\omega^{(k)}(z)) = 1$ with probability one. \square

The following is the new and original result that gives a sufficient condition for typical fast escaping.

THEOREM 3.7. *Let μ be a Borel probability measure on \mathbb{C} with compact support. Suppose that $T_{\{\infty\},\mu}$ is continuous. If $T_{\{\infty\},\mu}(0) = 1$, then the critical point 0 is typically fast escaping, and hence the random Julia set J_ω is totally disconnected for \mathbb{P}_μ -almost every $\omega \in \Omega_\mu$.*

Proof. The latter statement is a consequence of Theorem 3.4. Thus, it suffices to show that the critical point 0 is typically fast escaping. Let $R > 0$ be a large number as in Lemma 2.6(1). Set $E = \{z \in \widehat{\mathbb{C}} : T_{\{\infty\},\mu}(z) = 1\}$, then E is compact since we assume that $T_{\{\infty\},\mu}$ is continuous. (This is the only place where the assumption of continuity is used.)

For every $z \in E$, there exist $n_z \in \mathbb{N}$ and $c_1^z, c_2^z, \dots, c_{n_z}^z \in \text{supp } \mu$ such that $|f_{c_{n_z}^z} \circ \dots \circ f_{c_2^z} \circ f_{c_1^z}(z)| > R$. Since this map $f_{c_{n_z}^z} \circ \dots \circ f_{c_2^z} \circ f_{c_1^z}$ is continuous, there exists an open neighborhood O_z of z such that for every $z' \in O_z$, we have $|f_{c_{n_z}^z} \circ \dots \circ f_{c_2^z} \circ f_{c_1^z}(z')| > R$. We may and do assume that O_z is precompact.

We now consider the open covering $\{O_z\}_{z \in E}$ of E . Since E is compact, there exists a finite subcover. Namely, there exist $\ell \in \mathbb{N}$ and ℓ open sets O_j ($j = 1, \dots, \ell$) such that $E \subset \bigcup_{j=1}^{\ell} O_j$. Moreover, for each $j = 1, \dots, \ell$, there exist $n_j \in \mathbb{N}$ and $c_1^j, c_2^j, \dots, c_{n_j}^j \in \text{supp } \mu$ such that for every $z' \in O_j$, we have $|f_{c_{n_j}^j} \circ \dots \circ f_{c_2^j} \circ f_{c_1^j}(z')| > R$. Let $N = \max_{j=1, \dots, \ell} n_j$. By Lemma 2.6(1), we can assume that $n_j = N$ for every $j = 1, \dots, \ell$. Since $\mathbb{C} \ni c \mapsto f_c \in \text{Poly}$ is continuous, for every $n = 1, 2, \dots, N$, there exists an open neighborhood U_n^j of c_n^j such that for every $c_n \in U_n^j$ and $z' \in O_j$, we have $|f_{c_n} \circ \dots \circ f_{c_2} \circ f_{c_1}(z')| > R$.

Define $\mathfrak{G}^k(z) = \{\omega \in \Omega_\mu : |f_\omega^{(k)}(z)| \leq R\}$ for every $z \in \widehat{\mathbb{C}}$. Then, by the construction above, we have $\mathbb{P}_\mu(\Omega_\mu \setminus \mathfrak{G}^N(z)) \geq \mathbb{P}_\mu(\prod_{n=1}^N U_n^j \times \prod_{N+1}^\infty \bar{B}(c, r))$ for every $j = 1, \dots, \ell$ and $z \in O_j$. Since $c_n^j \in \text{supp } \mu$ and U_n^j is its open neighborhood, we have $\mu(U_n^j) > 0$ for every $j = 1, \dots, \ell$ and $n = 1, 2, \dots, N$. Define $\alpha = \min_{j=1, \dots, \ell} \prod_{n=1}^N \mu(U_n^j) > 0$. Then, for every $z \in E$, we have $\mathbb{P}_\mu(\mathfrak{G}^N(z)) \leq 1 - \alpha$.

It follows from Fubini’s theorem that

$$\mathbb{P}_\mu(\mathfrak{S}^{k+N}(z)) = \int_{\Omega_\mu} \mathbb{P}_\mu(\mathfrak{S}^N(f_\omega^{(k)}(z))) d\mathbb{P}_\mu(\omega)$$

for every $k \in \mathbb{N}$. If $\omega \notin \mathfrak{S}^k(z)$, then $\mathfrak{S}^N(f_\omega^{(k)}(z)) = \emptyset$ by Lemma 2.6. Thus, the integral on the complement of $\mathfrak{S}^k(z)$ is zero; $\mathbb{P}_\mu(\mathfrak{S}^{k+N}(z)) = \int_{\mathfrak{S}^k(z)} \mathbb{P}_\mu(\mathfrak{S}^N(f_\omega^{(k)}(z))) d\mathbb{P}_\mu(\omega)$. By Lemma 3.6, we have $\mathbb{P}_\mu(\mathfrak{S}^{k+N}(z)) \leq (1 - \alpha)\mathbb{P}_\mu(\mathfrak{S}^k(z))$ if $z \in E$.

Repeating this, we have $\mathbb{P}_\mu(\mathfrak{S}^m(0)) \leq (1 - \alpha)^m$ for every $m \in \mathbb{N}$. Thus, we can find a constant $\gamma > 0$ such that

$$\mathbb{P}_\mu(\{\omega \in \Omega_\mu : k(0, \omega) > k\}) \leq \mathbb{P}_\mu(\mathfrak{S}^k(0)) \leq e^{-\gamma k}$$

for every $k \in \mathbb{N}$. This completes the proof. □

COROLLARY 3.8. *Let μ be a Borel probability measure on \mathbb{C} whose support is compact and satisfies $\text{int}(\text{supp } \mu) \neq \emptyset$. If $T_{\{\infty\},\mu}(0) = 1$, then the random Julia set J_ω is totally disconnected for \mathbb{P}_μ -almost every $\omega \in \Omega_\mu$.*

Proof. This is a consequence of Lemma 2.22 and Theorems 2.21 and 3.7. □

COROLLARY 3.9. *Let μ be a Borel probability measure on \mathbb{C} whose support is compact. If G_μ has no planar minimal sets, then the random Julia set J_ω is totally disconnected for \mathbb{P}_μ -almost every $\omega \in \Omega_\mu$.*

Proof. Since G_μ has no planar minimal sets, the function satisfies $T_{\{\infty\},\mu} \equiv 1$ by Theorem 2.21. Since the constant function is continuous, Theorem 3.7 implies the conclusion. □

For instance, let $\mu = \delta_0/2 + \delta_c/2$ be the convex combination of the Dirac measures at 0 and c with $|c| > 1$. If $|z| > 1$, then $f_0^{on}(z) \rightarrow \infty$ as $n \rightarrow \infty$. If $|z| \leq 1$, then $|f_c(z)| \geq -|z^2| + |c| > 1$, and hence $f_0^{on} \circ f_c(z) \rightarrow \infty$ as $n \rightarrow \infty$. This shows that G_μ has no planar minimal sets, which implies that the random Julia set is totally disconnected \mathbb{P}_μ -almost surely.

In the next section, we apply Corollary 3.8 to the case where μ is the normalized Lebesgue measure on the disk $\bar{B}(c, r)$.

4. Main results on uniform noise process

Throughout this section, we consider the case where μ is the normalized Lebesgue measure on the disk $\bar{B}(c, r)$ as in Setting 4.1.

Setting 4.1. For fixed $c \in \mathbb{C}$ and $r \geq 0$, we define the following Borel probability measure $\mu_{c,r}$ on \mathbb{C} with compact support. If $r > 0$, define $\mu_{c,r}$ as the normalized Lebesgue measure on $\bar{B}(c, r) = \{c' \in \mathbb{C} : |c' - c| \leq r\}$. If $r = 0$, define $\mu_{c,r}$ as the Dirac measure at c . We define the probability space $(\Omega_{c,r}, \mathbb{P}_{c,r}) = (\Omega_{\mu_{c,r}}, \mathbb{P}_{\mu_{c,r}})$ as in Setting 2.1. Namely, denote by $\mathbb{P}_{c,r}$ the one-sided infinite product of $\mu_{c,r}$ supported on $\Omega_{c,r} = \prod_{n=1}^\infty \bar{B}(c, r)$ with the Borel σ -algebra. In addition, let the polynomial semigroup $G_{c,r}$ be defined by the semigroup generated by $\{f_{c'} \in \text{Poly} : c' \in \bar{B}(c, r)\}$ and define the function

$T_{c,r}: \widehat{\mathbb{C}} \rightarrow [0, 1]$, abusing notation, by

$$T_{c,r}(z) = \mathbb{P}_{c,r}(\{\omega \in \Omega_{c,r}: f_\omega^{(n)}(z) \rightarrow \infty (n \rightarrow \infty)\}).$$

Remark 4.2. Setting 4.1 can be interpreted as an additive-noise stochastic process in the following way. Fix $c \in \mathbb{C}$ and $r \geq 0$. On a probability space $(\Omega, \mathfrak{B}, \mathbb{P})$, take independent random variables $v_n: \Omega \rightarrow \mathbb{C}$ that follow the uniform distribution on $\bar{B}(0, r)$ for all $n \in \mathbb{N}$. Fix $z_0 \in \mathbb{C}$ and define the random variable $Z_0: \Omega \rightarrow \mathbb{C}$ by a constant map with value z_0 . Define the \mathbb{C} -valued stochastic process $\{Z_n\}_{n \geq 0}$ on $(\Omega, \mathfrak{B}, \mathbb{P})$, inductively by $Z_{n+1} = f_c(Z_n) + v_{n+1}$ for every $n \geq 0$. Then the distribution of Z_n are identical to that of the random variable $f_\bullet^{(n)}(z_0): \Omega_{c,r} \rightarrow \mathbb{C}$ for every $n \in \mathbb{N}$, where $f_\bullet^{(n)}(z_0)$ is the map $\omega \mapsto f_\omega^{(n)}(z_0)$. This stochastic process describes the process driven by adding uniform noise independently to the deterministic dynamical systems of f_c with the initial value z_0 . Here, the parameter r describes the size of noise that satisfies $|v_n| \leq r$ almost surely for every $n \in \mathbb{N}$.

We investigate the bifurcation radius $r_{\text{bif}}(c)$ defined in Definition 2.17. Our final goal is to determine $r_{\text{bif}}(c)$ and we provide some partial results in this paper.

4.1. *Relation between bifurcation and connectedness.* In this subsection, we reveal the relation between bifurcation and connectedness. We start the discussion with the following corollary.

COROLLARY 4.3. *Suppose that $r > r_{\text{bif}}(c)$. Then the random Julia set J_ω is totally disconnected for $\mathbb{P}_{c,r}$ -almost every ω .*

Proof. If $r > r_{\text{bif}}(c)$, then $T_{c,r} \equiv 1$ by Theorems 2.16 and 2.21. It follows from Corollary 3.8 that the random Julia set J_ω is totally disconnected almost surely. □

In the following, suppose that the interior of the support $\bar{B}(c, r)$ contains a superattracting parameter.

LEMMA 4.4. *Suppose that the interior of $\bar{B}(c, r)$ contains a superattracting parameter \tilde{c} . If there exists $\omega \in \Omega_{c,r}$ such that $f_\omega^{(n)}(0) \rightarrow \infty$ as $n \rightarrow \infty$, then $r > r_{\text{bif}}(c)$.*

Proof. We show that for every $z \in \widehat{\mathbb{C}}$, there exist $c_1, c_2, \dots \in \bar{B}(c, r)$ such that $f_{c_n} \circ \dots \circ f_{c_2} \circ f_{c_1}(z) \rightarrow \infty$, which implies that $G_{c,r}$ has no planar minimal sets.

The proof is divided into three cases. Denote by $K(f_{\tilde{c}})$ the autonomous filled Julia set of the superattracting map $f_{\tilde{c}}$. If $z \notin K(f_{\tilde{c}})$, then $f_{\tilde{c}}^{on}(z) \rightarrow \infty$ by definition. If $z \in \partial K(f_{\tilde{c}})$, then there exists $c_1 \in \bar{B}(c, r)$ such that $f_{c_1}(z) \notin K(f_{\tilde{c}})$ and hence $f_{\tilde{c}}^{on} \circ f_{c_1}(z) \rightarrow \infty$. If $z \in \text{int}K(f_{\tilde{c}})$, then by the no wandering domain theorem, the orbit $f_{\tilde{c}}^{on}(z)$ converges to the attracting cycle of 0. By the assumption, we have $0 \in A_\omega$. Since the non-autonomous basin A_ω is open, there exists an open neighborhood D of 0 such that for every $z' \in D$, we have $f_\omega^{(n)}(z') \rightarrow \infty$ as $n \rightarrow \infty$. Take $N \in \mathbb{N}$ so that $f_{\tilde{c}}^{\circ N}(z) \in D$, then $f_\omega^{(n)}(f_{\tilde{c}}^{\circ N}(z)) \rightarrow \infty$ as $n \rightarrow \infty$. This completes the proof. □

LEMMA 4.5. *Suppose that the interior of $\bar{B}(c, r)$ contains a superattracting parameter \tilde{c} . If $L = \bigcup_{g \in G_{c,r}} \{g(0)\}$ is bounded in \mathbb{C} , then L is a planar minimal set of $G_{c,r}$.*

Proof. Fix any $z \in L$. First, we show $\overline{\bigcup_{g \in G_{c,r}} \{g(z)\}} \subset L$. By definition, there exists $g_n \in G_{c,r}$ such that $g_n(0) \rightarrow z$ as $n \rightarrow \infty$. Then, for every $g \in G_{c,r}$, we have $g \circ g_n(0) \rightarrow g(z)$ as $n \rightarrow \infty$. Since $G_{c,r}$ is a semigroup, we have $g \circ g_n \in G_{c,r}$. Thus, we have $g(z) \in L$, which implies $\bigcup_{g \in G_{c,r}} \{g(z)\} \subset L$. Next, we show $\bigcup_{g \in G_{c,r}} \{g(z)\} \supset L$. For every open set U which intersects L , take $g_1 \in G_{c,r}$ such that $g_1(0) \in U$. Since g_1 is continuous, there exists an open set U_0 such that $0 \in U_0$ and $g_1(U_0) \subset U$. Since $\overline{\bigcup_{g \in G_{c,r}} \{g(z)\}} \subset L$, the autonomous orbit $\{f_{\tilde{c}}^{on}(z)\}_{n \in \mathbb{N}}$ is bounded, and hence $z \in K(f_{\tilde{c}})$. If $z \in \partial K(f_{\tilde{c}})$, then there exists $c_1 \in \tilde{B}(c, r)$ such that $f_{c_1}(z) \notin K(f_{\tilde{c}})$, but this contradicts $\overline{\bigcup_{g \in G_{c,r}} \{g(z)\}} \subset L$. Thus, $z \in \text{int}K(f_{\tilde{c}})$. It follows from the no wandering domain theorem that there exists $N \in \mathbb{N}$ such that $f_{\tilde{c}}^{oN}(z) \in U_0$. Thus, $g_1 \circ f_{\tilde{c}}^{oN}(z) \in U$, which implies $\overline{\bigcup_{g \in G_{c,r}} \{g(z)\}} \supset L$. This completes the proof. \square

The following is the main theorem of this paper.

THEOREM 4.6. *Suppose that the interior of $\tilde{B}(c, r)$ contains a superattracting parameter \tilde{c} . Then the following four are equivalent.*

- (1) *The inequality $r \leq r_{\text{bif}}(c)$ holds.*
- (2) *For every $\omega \in \Omega_{c,r}$, the orbit $\{f_{\omega}^{(n)}(0)\}_{n=1}^{\infty}$ is bounded; hence, $T_{c,r}(0) = 0$.*
- (3) *The random Julia set J_{ω} is connected for every $\omega \in \Omega_{c,r}$.*
- (4) *The set $\overline{\bigcup_{g \in G_{c,r}} \{g(0)\}}$ is a planar minimal set of $G_{c,r}$.*

In contraposition to this, the following four are equivalent.

- (1') *The inequality $r > r_{\text{bif}}(c)$ holds.*
- (2') *For $\mathbb{P}_{c,r}$ -almost every $\omega \in \Omega_{c,r}$, the orbit $f_{\omega}^{(n)}(0) \rightarrow \infty$ ($n \rightarrow \infty$): $T_{c,r}(0) = 1$.*
- (3') *The random Julia set J_{ω} is totally disconnected for $\mathbb{P}_{c,r}$ -almost every $\omega \in \Omega_{c,r}$.*
- (4') *The set $\overline{\bigcup_{g \in G_{c,r}} \{g(0)\}}$ is not bounded in \mathbb{C} .*

Furthermore, either the former or the latter is valid.

Proof. It is obvious that item (1) or (1') is valid. First, we show the equivalence of the four from items (1) to (4).

(1) \Rightarrow (2) : If $r \leq r_{\text{bif}}(c)$, then $f_{\omega}^{(n)}(0) \not\rightarrow \infty$ as $n \rightarrow \infty$ for every $\omega \in \Omega_{c,r}$ by Lemma 4.4. Hence, $T_{c,r}(0) = 0$.

(2) \Rightarrow (3) : Take a large $R > 0$ as in Lemma 2.6(1). Suppose that for every $\omega \in \Omega_{c,r}$, we have $\{f_{\omega}^{(n)}(0)\}_{n=1}^{\infty}$ is bounded. Then $|f_{\omega}^{(n)}(0)| < R$ for every $n \in \mathbb{N}$. By Theorem 3.1, the random Julia set J_{ω} is connected for every $\omega \in \Omega_{c,r}$.

(3) \Rightarrow (4) : Suppose that the random Julia set J_{ω} is connected for every $\omega \in \Omega_{c,r}$. For a large $R > 0$ as in Lemma 2.6(1), it follows from Theorem 3.1 that $|f_{\sigma^k \omega}^{(n)}(0)| \leq R$ for every $\omega \in \Omega_{c,r}$, $n \in \mathbb{N}$, and $k \in \mathbb{N}$. Thus, the set $L = \overline{\bigcup_{g \in G_{c,r}} \{g(0)\}}$ is bounded in \mathbb{C} . By Lemma 4.5, the set L is a planar minimal set.

(4) \Rightarrow (1) : This is trivial. See the latter part of Definition 2.17.

The argument above completes the proof of the equivalence from items (1) to (4). Next, we show the equivalence of the four from items (1') to (4').

(1') \Rightarrow (2') : If $r > r_{\text{bif}}(c)$, then $G_{c,r}$ has no planar minimal sets by Theorem 2.16. It follows from Theorem 2.21 that $T_{c,r}(z) = 1$ for every $z \in \hat{\mathbb{C}}$, and hence $T_{c,r}(0) = 1$.

(2') \Rightarrow (3') : Suppose $T_{c,r}(0) = 1$. If $r > 0$, then the random Julia set J_ω is totally disconnected $\mathbb{P}_{c,r}$ -almost surely by Corollary 3.8. If $r = 0$, then it trivially follows that the autonomous Julia set is totally disconnected.

(3') \Rightarrow (4') : If item (3') holds, then there exists $\omega \in \Omega_{c,r}$ such that the random Julia set J_ω is disconnected. By Theorem 3.1, there exists $k \in \mathbb{N}$ such that $f_{\sigma^k \omega}^{(n)}(0) \rightarrow \infty$ as $n \rightarrow \infty$. Since $f_{\sigma^k \omega}^{(n)} \in G_{c,r}$, then $\overline{\bigcup_{g \in G_{c,r}} \{g(0)\}}$ is unbounded.

(4') \Rightarrow (1') : Let $R > 0$ be large, as in Lemma 2.6(1). If $\overline{\bigcup_{g \in G_{c,r}} \{g(0)\}}$ is not bounded, then there exists $g \in G_{c,r}$ such that $|g(0)| > R$. By definition of $G_{c,r}$, there exist $N \in \mathbb{N}$ and $c_1, c_2, \dots, c_N \in \bar{B}(c, r)$ such that $g = f_{c_N} \circ \dots \circ f_{c_2} \circ f_{c_1}$. Letting $c_n = c$ for every $n \geq N + 1$ and $\omega = (c_n)_{n=1}^\infty$, we have $f_\omega^{(n)}(0) \rightarrow \infty$ as $n \rightarrow \infty$ by Lemma 2.6. It follows from Lemma 4.4 that $r > r_{\text{bif}}(c)$. \square

Remark 4.7. The orbit $\{f_\omega^{(n)}(0)\}_{n=1}^\infty$ is bounded for every $\omega \in \Omega_{c,r}$ if and only if $T_{c,r}(0) = 0$.

Proof. It is obvious that the ‘only if’ part is true. Suppose $T_{c,r}(0) = 0$ and we show there does not exist $\omega \in \Omega_{c,r}$ such that $f_\omega^{(n)}(0) \rightarrow \infty$ as $n \rightarrow \infty$ by contradiction. Take a large number $R > 0$ as in Lemma 2.6(1). If there exists $\omega = (c_n)_{n \in \mathbb{N}} \in \Omega_{c,r}$ such that $f_\omega^{(n)}(0) \rightarrow \infty$, then there exists $N \in \mathbb{N}$ such that $|f_{c_N} \circ \dots \circ f_{c_2} \circ f_{c_1}(0)| > R$. Since $\mathbb{C} \ni c \mapsto f_c \in \text{Poly}$ is continuous, there exists an open neighborhood U_n of c_n for every $n = 1, 2, \dots, N$ such that $|f_{c'_N} \circ \dots \circ f_{c'_2} \circ f_{c'_1}(0)| > R$ for every $c'_n \in U_n$ ($n = 1, 2, \dots, N$). Moreover, we have $\mathbb{P}_{c,r}(\mathcal{U}) > 0$ for $\mathcal{U} = \prod_{n=1}^N U_n^j \times \prod_{N+1}^\infty \bar{B}(c, r)$. However, for every $\omega' \in \mathcal{U}$, it follows from Lemma 2.6 that $f_{\omega'}^{(n)}(0) \rightarrow \infty$ as $n \rightarrow \infty$, contradicting the assumption $T_{c,r}(0) = 0$. This shows that for every $\omega \in \Omega_{c,r}$, we have $|f_\omega^{(n)}(0)| \leq R$ for every $n \in \mathbb{N}$. \square

4.2. *General properties.* In this subsection, we give a trivial estimate of the bifurcation radius and show the continuity. The following is due to [4].

LEMMA 4.8. *Suppose $r > 0$. If $\bar{B}(c, r) \cap (\mathbb{C} \setminus \mathcal{M}) \neq \emptyset$, then for every $z \in \mathbb{C}$, there exists $\omega \in \Omega_{c,r}$ such that $f_\omega^{(n)}(z) \rightarrow \infty$ as $n \rightarrow \infty$.*

Proof. Fix $c' \in \bar{B}(c, r) \setminus \mathcal{M}$. Then the autonomous filled Julia set $K(f_{c'})$ of $f_{c'}$ has no interior point. Thus, if $z \in K(f_{c'})$, then there exists $c_1 \in \bar{B}(c, r)$ such that $f_{c_1}(z) \notin K(f_{c'})$. In this case, we have $f_{c_1}^{\circ n} \circ f_{c_1}(z) \rightarrow \infty$ as $n \rightarrow \infty$. Also, if $z \notin K(f_{c'})$, then by definition, $f_{c'}^{\circ n}(z) \rightarrow \infty$ as $n \rightarrow \infty$. This completes the proof. \square

Denote by $\text{dist}(c, \partial\mathcal{M})$ the Euclidean distance from the point c to the compact set $\partial\mathcal{M}$.

LEMMA 4.9. *For every $c \in \mathbb{C}$, we have $r_{\text{bif}}(c) \leq \text{dist}(c, \partial\mathcal{M})$.*

Proof. From Lemma 4.8, if $r > \text{dist}(c, \partial\mathcal{M})$, then the polynomial semigroup $G_{c,r}$ has no planar minimal sets. By Definition 2.17, $r_{\text{bif}}(c)$ is the infimum value for $r \geq 0$ such that $G_{c,r}$ has no planar minimal sets. Hence, $r_{\text{bif}}(c) \leq \text{dist}(c, \partial\mathcal{M})$. \square

The following is a part of Main Result C.

THEOREM 4.10. *As a function, the bifurcation radius $r_{\text{bif}}: \mathbb{C} \rightarrow [0, \infty)$ is 1-Lipschitz. That is, $|r_{\text{bif}}(c) - r_{\text{bif}}(c')| \leq |c - c'|$ for every $c, c' \in \mathbb{C}$.*

Proof. Fix $c, c' \in \mathbb{C}$. Pick any $r > |c - c'|$ and set $\epsilon = r - |c - c'| > 0$. By definition, the polynomial semigroup $G_{c, r_{\text{bif}}(c) + \epsilon}$ has no planar minimal sets. Since $\bar{B}(c, r_{\text{bif}}(c) + \epsilon) \subset \bar{B}(c', r_{\text{bif}}(c) + r)$, it follows from Lemma 2.12 that also $G_{c', r_{\text{bif}}(c) + r}$ has no planar minimal sets. This implies that $r_{\text{bif}}(c') \leq r_{\text{bif}}(c) + r$, and hence $r_{\text{bif}}(c') - r_{\text{bif}}(c) \leq |c - c'|$. Exchanging c and c' with each other, we have also $r_{\text{bif}}(c) - r_{\text{bif}}(c') \leq |c' - c|$, which completes the proof. □

4.3. Inside the main cardioid. In this subsection, we present some estimates of $r_{\text{bif}}(c)$ when c is in the main cardioid. Recall that the main cardioid is the set of all parameters $c \in \mathbb{C}$ for which there exists $\lambda \in \mathbb{C}$ with $|\lambda| < 1$ such that $c = \lambda/2 - \lambda^2/4$, and its boundary is contained in $\partial\mathcal{M}$.

COROLLARY 4.11. *If $c = 0$, then $r_{\text{bif}}(0) = 1/4$.*

Proof. This is essentially due to [4]. By Lemma 4.9, we have $r_{\text{bif}}(0) \leq 1/4$. A straight calculation shows that $\overline{D_{1/2}}$ is forward invariant under the polynomial semigroup $G_{0, 1/4}$, and hence $r_{\text{bif}}(0) \geq 1/4$. □

We can easily verify that $\overline{D_{1/2}}$ is the planar minimal set of $G_{0, 1/4}$. Since the autonomous Julia set of $f_{1/4}$ contains $z = 1/2$ which is the parabolic fixed point of $f_{1/4}$, the Julia set of the semigroup satisfies $1/2 \in J(G_{0, 1/4})$ by Proposition 2.9. Thus, the planar minimal set $\overline{D_{1/2}}$ of $G_{0, 1/4}$ is J -touching at $z = 1/2$.

Corollary 4.11 shows that the equality in Lemma 4.9 may hold. Now, we show using Theorem 4.10 that we can determine the bifurcation radius if $0 \leq c \leq 1/4$. The following is a part of Main Result C.

THEOREM 4.12. *If $0 \leq c \leq 1/4$, then $r_{\text{bif}}(c) = 1/4 - c$.*

Proof. By Lemma 4.9, we have $r_{\text{bif}}(c) \leq 1/4 - c$ if $0 \leq c \leq 1/4$. If $c' = 0$, then $r_{\text{bif}}(c') = 1/4$ from Corollary 4.11. Thus, Theorem 4.10 yields that $|r_{\text{bif}}(c) - 1/4| \leq c$ for $0 \leq c \leq 1/4$, which completes the proof. □

We showed that $r_{\text{bif}}(c) = \text{dist}(c, \partial\mathcal{M})$ if $0 \leq c \leq 1/4$. However, the author does not know whether there exists $c \in \mathbb{C}$ other than these, such that $0 < r_{\text{bif}}(c) = \text{dist}(c, \partial\mathcal{M})$. For $c < 0$, we will present some examples which satisfy $r_{\text{bif}}(c) < \text{dist}(c, \partial\mathcal{M})$.

LEMMA 4.13. *For $c = 0$ and $0 < r_0 \leq 1/4$, let $\delta = (1 - \sqrt{1 - 4r_0})/2$. Then the following hold.*

- (i) *The equality $\delta^2 + r_0 = \delta$ holds. Thus, the point δ is a fixed point of f_{r_0} .*
- (ii) *The set $\overline{D_\delta}$ is the planar minimal set of G_{0, r_0} .*

Proof. Statement (i) follows from the direct calculation. Now, we look at statement (ii). By statement (i), we have $f_{c_1}(\overline{D_\delta}) \subset \overline{D_\delta}$ for every $c_1 \in \bar{B}(0, r_0)$. Since $\bar{B}(0, r_0)$ is symmetric under rotation, then the union $\bigcup_{c_1 \in \bar{B}(0, r_0)} \{f_{c_1}(0)\}$ is a round disk centered at the origin.

By induction, $\bigcup_{c_1, c_2, \dots, c_n \in \bar{B}(0, r_0)} \{f_{c_n} \circ \dots \circ f_{c_2} \circ f_{c_1}(0)\}$ is a round disk centered at the origin for every $n \in \mathbb{N}$, and also $\overline{\bigcup_{g \in G_{0, r_0}} \{g(0)\}}$ is a closed round disk contained in \bar{D}_δ . If $c_n = r_0$ for every $n \in \mathbb{N}$, then the autonomous iteration $f_{r_0}^{on}(0)$ converges to the fixed point δ of f_{r_0} . This yields that $\overline{\bigcup_{g \in G_{0, r_0}} \{g(0)\}} = \bar{D}_\delta$, and hence \bar{D}_δ is a minimal set of G_{0, r_0} . \square

The following is a part of Main Result C.

THEOREM 4.14. *Suppose that $0 < \epsilon \leq 1/2$. For $c = -\epsilon$, we have $r_{\text{bif}}(-\epsilon) \leq 1/4 + \epsilon - \epsilon^2$.*

Proof. Fix any $r > 1/4 + \epsilon - \epsilon^2$. We show that $G_{-\epsilon, r}$ has no planar minimal sets by contradiction, which implies $r_{\text{bif}}(-\epsilon) < r$. Suppose that there exists a planar minimal set L of $G_{-\epsilon, r}$. Since $\bar{B}(-\epsilon, r) \supset \bar{B}(0, r - \epsilon)$, it follows from Lemma 2.12 that $G_{0, r - \epsilon}$ also has a planar minimal set. By Corollary 4.11, we have $r - \epsilon \leq 1/4$. In addition, we assumed that $r - \epsilon > 1/4 - \epsilon^2 \geq 0$. By Lemmas 2.12 and 4.13(ii), we have $L \supseteq \bar{D}_\delta$ for $\delta = (1 - \sqrt{1 - 4(r - \epsilon)})/2$. Let $z_0 = i\delta$ and $c_1 = -r - \epsilon$, then $z_0 \in L$, $c_1 \in \bar{B}(-\epsilon, r)$, and $f_{c_1}(z_0) = -\delta^2 - r - \epsilon = -\delta - 2\epsilon$. Now, set $R_1 = (1 + \sqrt{1 - 4(r - \epsilon)})/2 > 1/2$, then

$$-f_{c_1}(z_0) - R_1 = 2\epsilon + \frac{1 - \sqrt{1 - 4(r - \epsilon)}}{2} - \frac{1 + \sqrt{1 - 4(r - \epsilon)}}{2} = 2\epsilon - \sqrt{1 - 4(r - \epsilon)} > 0$$

since $r > 1/4 + \epsilon - \epsilon^2$. Set $z_1 = f_{c_1}(z_0)$, then $z_1 < -R_1$. Let $c_n = r - \epsilon \in \bar{B}(-\epsilon, r)$ for every $n \geq 2$. Since $z_1 < -R_1$, the direct calculation shows $f_{c_2}(z_1) > R_1^2 + r - \epsilon = R_1$. Set $z_2 = f_{c_2}(z_1)$ and $\alpha = z_2 - R_1$, then $z_2 = R_1 + \alpha$ and $\alpha > 0$. We can show by induction that $f_{r - \epsilon}^{on}(z_2) > R_1 + \alpha + n\alpha^2$ for every $n \in \mathbb{N}$. Hence, $f_{r - \epsilon}^{on}(z_1)$ diverges to ∞ as $n \rightarrow \infty$. Thus, we have proved that there exist $z_0 \in L$ and $\omega = (c_n)_{n=1}^\infty \in \Omega_{-\epsilon, r}$ such that $f_\omega^{(n)}(z_0) \rightarrow \infty$ as $n \rightarrow \infty$. This contradicts that L is planar and forward invariant under $G_{-\epsilon, r}$. This completes the proof since r is an arbitrary number that satisfies $r > 1/4 + \epsilon - \epsilon^2$. \square

Note that we can show that $\text{dist}(-\epsilon, \partial\mathcal{M}) = 1/4 + \epsilon$ if $0 < \epsilon \leq 1/4$ as follows. Assume $0 < \epsilon \leq 1/4$, then $\text{dist}(-\epsilon, \partial\mathcal{M}) \leq |-\epsilon - 1/4| = 1/4 + \epsilon$ since $1/4 \in \mathcal{M}$. Let $C = \{\lambda/2 - \lambda^2/4 : \lambda = e^{i\theta}, \theta \in \mathbb{R}\}$. Then $C \subset \partial\mathcal{M}$ and the domain bounded by C is contained in the interior of \mathcal{M} , see [6, Theorem VIII.1.3]. Since

$$\begin{aligned} \left| -\frac{1}{4} - \frac{e^{i\theta}}{2} + \frac{e^{2i\theta}}{4} \right|^2 &= \left(-\frac{1}{4} - \frac{e^{i\theta}}{2} + \frac{e^{2i\theta}}{4} \right) \left(-\frac{1}{4} - \frac{e^{-i\theta}}{2} + \frac{e^{-2i\theta}}{4} \right) \\ &= \frac{6 - (e^{2i\theta} + e^{-2i\theta})}{16} = \frac{6 - 2 \cos 2\theta}{16}, \end{aligned}$$

we have $\text{dist}(-1/4, \partial\mathcal{M}) = \min_{\theta \in \mathbb{R}} \sqrt{6 - 2 \cos 2\theta}/4 = 1/2$ and the minimum is attained at $\theta = 0, \pi$ which correspond to $c = 1/4, -3/4 \in \mathcal{M}$, respectively. Since $\text{int}\bar{B}(-\epsilon, 1/4 + \epsilon) \subset \text{int}\bar{B}(-1/4, 1/2) \subset \text{int}\mathcal{M}$, we have $\text{dist}(-\epsilon, \partial\mathcal{M}) \geq 1/4 + \epsilon$. Therefore, Theorem 4.14 reveals that strict inequality $r_{\text{bif}}(c) < \text{dist}(c, \partial\mathcal{M})$ holds if $-1/4 \leq c < 0$.

Combining the theorem above with Corollary 4.3, we have a little better result than Theorem 3.5.

COROLLARY 4.15. *Suppose that $0 < \epsilon \leq 1/2$. If $c = -\epsilon$ and $r > 1/4 + \epsilon - \epsilon^2$, then the random Julia set J_ω is totally disconnected for $\mathbb{P}_{-\epsilon,r}$ -almost every $\omega \in \Omega_{-\epsilon,r}$.*

This shows that the random Julia set J_ω is totally disconnected for almost every ω even if $\bar{B}(c, r) \subset \text{int } \mathcal{M}$ and $\bar{B}(c, r) \not\supset \bar{B}(0, 1/4)$. This is stronger than Theorem 3.5 proved by Lech and Zdunik.

4.4. Outside the main cardioid. Recall that a parameter \tilde{c} is said to be superattracting if there exists $p \in \mathbb{N}$ such that $f_{\tilde{c}}^{\circ p}(0) = 0$. For example, $\tilde{c} = 0$ and -1 are superattracting parameters with $p = 1$ and 2 , respectively. In this subsection, we show the non-trivial estimates of $r_{\text{bif}}(-1)$.

LEMMA 4.16. *Let $c = -1$, $r \geq 0$, and $\delta \geq 0$. Denote $\bar{D}_\delta = \{z \in \mathbb{C} : |z| \leq \delta\}$. Then $f_{c_2} \circ f_{c_1}(\bar{D}_\delta) \subset \bar{D}_\delta$ for every $c_1, c_2 \in \bar{B}(-1, r)$ if and only if*

$$\delta^4 + 2(1+r)\delta^2 + r^2 + 3r \leq \delta. \tag{1}$$

Proof. Assume that $f_{c_2} \circ f_{c_1}(\bar{D}_\delta) \subset \bar{D}_\delta$ for every $c_1, c_2 \in \bar{B}(-1, r)$. For $z = i\delta$, $c_1 = -1 - r$, and $c_2 = -1 + r$, we have $f_{c_1}(z) = -\delta^2 - 1 - r$ and $f_{c_2} \circ f_{c_1}(z) = \delta^4 + 2(1+r)\delta^2 + r^2 + 3r$. Hence, inequality (1) holds.

Conversely, assume that inequality (1) holds. If $z \in \bar{D}_\delta$ and $c_1 \in \bar{B}(-1, r)$, then $|f_{c_1}(z) + 1| \leq |z|^2 + |c_1 + 1| \leq \delta^2 + r$. In addition, if $|z_1 + 1| \leq \delta^2 + r$ and $c_2 \in \bar{B}(-1, r)$, then

$$\begin{aligned} |f_{c_2}(z_1)| &= |(z_1 + 1)^2 - 2z_1 - 2 + 1 + c_2| \\ &\leq |z_1 + 1|^2 + 2|z_1 + 1| + |1 + c_2| \leq \delta^4 + 2(1+r)\delta^2 + r^2 + 3r. \end{aligned}$$

Thus, if inequality (1) holds, then $f_{c_2} \circ f_{c_1}(\bar{D}_\delta) \subset \bar{D}_\delta$ for every $c_1, c_2 \in \bar{B}(-1, r)$. □

LEMMA 4.17. *Denote by $\delta_{\text{max}} = 0.453 \dots$ the (positive real) root of $\delta^3 + 2\delta - 1$. Define the function $\rho: [0, \delta_{\text{max}}] \rightarrow \mathbb{R}$ by*

$$\rho(\delta) = -\frac{2\delta^2 + 3}{2} + \frac{\sqrt{4\delta^2 + 4\delta + 9}}{2}.$$

Then, for every $\delta \in [0, \delta_{\text{max}}]$, we have $\delta^4 + 2(1 + \rho(\delta))\delta^2 + \rho(\delta)^2 + 3\rho(\delta) = \delta$ and $\rho(\delta) \geq 0$.

Proof. The former part $\delta^4 + 2(1 + \rho(\delta))\delta^2 + \rho(\delta)^2 + 3\rho(\delta) = \delta$ follows from a straight calculation. We show $\rho(\delta) \geq 0$. A simple calculation shows that

$$2\rho(\delta) = \frac{-(2\delta^2 + 3)^2 + 4\delta^2 + 4\delta + 9}{(2\delta^2 + 3) + \sqrt{4\delta^2 + 4\delta + 9}} = \frac{-4\delta(\delta^3 + 2\delta - 1)}{(2\delta^2 + 3) + \sqrt{4\delta^2 + 4\delta + 9}}.$$

Define $\tilde{\rho}(\delta) = \delta^3 + 2\delta - 1$, then $\tilde{\rho}(0) = -1 < 0$ and $\tilde{\rho}'(\delta) = 3\delta^2 + 2 > 0$. Thus, $\rho(\delta) \geq 0$ for every $\delta \in [0, \delta_{\text{max}}]$. □

We now give the lower bound of $r_{\text{bif}}(-1)$, which is one of the main results of this paper.

THEOREM 4.18. *Let ρ be defined as in Lemma 4.17. Then ρ takes the maximum value $r_{\text{max}} = 0.0386 \dots$, and hence $r_{\text{max}} \leq r_{\text{bif}}(-1)$.*

Proof. We have that the derivative $\rho'(\delta) = -2\delta + (2\delta + 1)/\sqrt{4\delta^2 + 4\delta + 9}$. Thus, $\rho'(\delta) = 0$ if and only if $16\delta^4 + 16\delta^3 + 32\delta^2 - 4\delta - 1 = 0$. Let $\delta^* = 0.229 \dots$ be the unique positive root of this quartic equation. Then, $\rho(\delta)$ takes the maximum value at $\delta = \delta^*$. A numerical computation shows that the maximum value r_{max} is approximately $0.0386 \dots$.

By Lemmas 4.16 and 4.17, for every $\delta \in [0, \delta_{\text{max}}]$, the set \bar{D}_δ gives a forward invariant set under the polynomial semigroup $G_{-1, \rho(\delta)}$. Thus, we have $\rho(\delta) \leq r_{\text{bif}}(-1)$. In particular, $r_{\text{max}} = \rho(\delta^*) \leq r_{\text{bif}}(-1)$. This completes the proof. \square

Next, we consider the upper bound of $r_{\text{bif}}(-1)$. The following theorem gives upper bounds, whose proof suggests a kind of parabolic implosion in the stochastic sense.

THEOREM 4.19. *Suppose that $\bar{B}(c, r_0)$ contains a superattracting parameter \tilde{c} . If there exist $N \in \mathbb{N}$ and $c_1, c_2, \dots, c_N \in \bar{B}(c, r_0)$ such that $f_{c_N} \circ \dots \circ f_{c_2} \circ f_{c_1}$ has a parabolic periodic point, then $r_{\text{bif}}(c) \leq r_0$.*

Proof. Fix $r > r_0$ and apply Lemma 4.4. Set $g = f_{c_N} \circ \dots \circ f_{c_2} \circ f_{c_1}$ and let z^* denote the parabolic periodic point of g . Then there exists $p \in \mathbb{N}$ such that $g^{\circ p}(z^*) = z^*$, and the multiplier of $g^{\circ p}$ at z^* is 1. Thus, there exists a critical point z_0 of g such that $g^{\circ np}(z_0) \rightarrow z^*$ as $n \rightarrow \infty$. Since $g = f_{c_N} \circ \dots \circ f_{c_2} \circ f_{c_1}$, there exists $0 \leq k \leq N - 1$ such that $f_{c_k} \circ \dots \circ f_{c_2} \circ f_{c_1}(z_0) = 0$ by the chain rule. Here, if $k = 0$, then we define $f_{c_k} \circ \dots \circ f_{c_2} \circ f_{c_1}$ to be the identity map. In any case, we have $g^{\circ np-1} \circ f_{c_N} \circ \dots \circ f_{c_{k+2}} \circ f_{c_{k+1}}(0) \rightarrow z^*$ as $n \rightarrow \infty$. Now, let $\epsilon = r - r_0 > 0$, then there exists $n_0 \in \mathbb{N}$ such that $n_0 \geq 2$ and

$$|g^{\circ n_0 p-1} \circ f_{c_N} \circ \dots \circ f_{c_{k+2}} \circ f_{c_{k+1}}(0) - z^*| < \epsilon.$$

Set $w = g^{\circ n_0 p-1} \circ f_{c_N} \circ \dots \circ f_{c_{k+2}} \circ f_{c_{k+1}}(0)$. Since the parabolic periodic point z^* belongs to the autonomous Julia set $J(g)$ of g and the Julia set is the boundary of the basin at infinity, perturbation of c_N allows us to find a random orbit which escapes to the basin at infinity. Namely, there exists $c' \in \bar{B}(c_N, \epsilon)$ such that

$$w' = f_{c'} \circ f_{c_{N-1}} \circ \dots \circ f_{c_2} \circ f_{c_1} \circ g^{\circ n_0 p-2} \circ f_{c_N} \circ \dots \circ f_{c_{k+2}} \circ f_{c_{k+1}}(0) \in A(g),$$

where $A(g)$ denotes the basin at infinity for the autonomous iteration of g . Note that $c' \in \bar{B}(c, r)$ since $c_N \in \bar{B}(c, r_0)$ and $\epsilon = r - r_0$. By definition, we have $g^{\circ n}(w') \rightarrow \infty$ as $n \rightarrow \infty$. Thus, Lemma 4.4 gives that $r_{\text{bif}}(c) < r$. This completes the proof since r is an arbitrary number that satisfies $r > r_0$. \square

Theorem 4.19 gives an alternative proof of $r_{\text{bif}}(0) \leq 1/4$ since $f_{1/4}$ has a parabolic fixed point at $z = 1/2$.

The following gives an algorithm to find an upper bound of the bifurcation radius.

THEOREM 4.20. *Suppose c is a superattracting parameter. Fix $N \in \mathbb{N}$ and $k \in \mathbb{N}$, and let $p_i \in \{0, 1, \dots, k - 1\}$ for each $i = 1, 2, \dots, N$. Define $\zeta_k = e^{2\pi i/k}$ and $c_i = c + \rho \zeta_k^{p_i}$, where ρ is a complex variable. Let $\Delta(\rho)$ be the discriminant of the algebraic equation $f_{c_N} \circ \dots \circ f_{c_2} \circ f_{c_1}(z) - z = 0$ in the variable z , which is a polynomial in the variable ρ . If $\Delta(\rho_0) = 0$ for some ρ_0 , then $r_{\text{bif}}(c) \leq |\rho_0|$.*

Proof. Under the assumptions, if $\rho = \rho_0$, then $f_{c_N} \circ \dots \circ f_{c_2} \circ f_{c_1}(z) - z = 0$ has a multiple root. This implies that the map $f_{c_N} \circ \dots \circ f_{c_2} \circ f_{c_1}$ has a parabolic fixed point with multiplier 1. Since $c_1, c_2, \dots, c_N \in \bar{B}(c, r_0)$ with $r_0 = |\rho_0|$, we have $r_{\text{bif}}(c) \leq r_0$ by Theorem 4.19. □

We now give upper bounds of $r_{\text{bif}}(-1)$.

Example 4.21. Suppose $c = -1$. Let $N = 4$, $k = 6$, and $(p_1, p_2, p_3, p_4) = (1, 2, 5, 4)$. Then numerical calculation shows the following result. The discriminant $\Delta(\rho)$ of $f_{c_4} \circ f_{c_3} \circ f_{c_2} \circ f_{c_1}(z) - z$ vanishes at $\rho_0 \approx 0.0399 \dots$, and hence $r_{\text{bif}}(-1) \leq 0.0399 \dots$.

We have some comments on the upper bounds of $r_{\text{bif}}(-1)$.

Remark 4.22. In Example 4.21, the numerical errors should be treated carefully because the discriminant of the equation $f_{c_4} \circ f_{c_3} \circ f_{c_2} \circ f_{c_1}(z) - z = 0$ with respect to z is a polynomial $\Delta(\rho)$ of degree 32, and each coefficient is a very large number in absolute value, that is, of the order of approximately $10^{19} - 10^{26}$. The author used validated numerics to estimate ρ_0 , which showed $r_{\text{bif}}(-1) \leq |\rho_0| \leq 0.0399217$. Since validated numerics gives values including mathematically strict error evaluation, this value 0.0399217 is a rigorous upper bound of $r_{\text{bif}}(-1)$. For more details on validated numerics, see [22], for example.

Note also that all the 6^4 candidates $(p_1, p_2, p_3, p_4) \in \{0, 1, 2, 3, 4, 5\}^4$ were (non-rigorously) verified. That is, for every $(p_1, p_2, p_3, p_4) \in \{0, 1, 2, 3, 4, 5\}^4$, the discriminant $\Delta(\rho)$ of the equation $f_{c_4} \circ f_{c_3} \circ f_{c_2} \circ f_{c_1}(z) - z = 0$ was computed using c_1, \dots, c_4 as in Theorem 4.20, and we numerically computed the roots of $\Delta(\rho)$. This computation shows that 24 candidates (p_1, p_2, p_3, p_4) including $(1, 2, 5, 4)$ attain the same minimum absolute value $|\rho_0| \approx 0.0399 \dots$. Also, the author replaced ζ_6 by $\zeta_4 = e^{2\pi i/4} = i$, and computed the roots of $\Delta(\rho)$ in the same manner, which gave less sharp estimates.

We now give the upper bound for another parameter.

Example 4.23. Suppose $\tilde{c}_3 \approx -1.75487766$ is the airplane parameter such that \tilde{c}_3 is a real number and $f_{\tilde{c}_3}$ has a superattracting periodic point with period 3. Let $N = 3$, $k = 6$, and $(p_1, p_2, p_3) = (0, 3, 0)$. Then a similar numerical computation shows that $r_{\text{bif}}(\tilde{c}_3) \leq 0.0021$.

Remark 4.24. Recall that we can calculate the connected component W of $\text{int } \mathcal{M}$ which contains -1 . The component W is the open disk with center -1 and radius $1/4$, thus $\text{dist}(-1, \partial \mathcal{M}) = 1/4$. For $c = -1$, we have $r_{\text{bif}}(-1) \ll \text{dist}(-1, \partial \mathcal{M})$ by the example above. Similarly, $r_{\text{bif}}(\tilde{c}_3) \ll \text{dist}(\tilde{c}_3, \partial \mathcal{M})$ since $\text{dist}(\tilde{c}_3, \partial \mathcal{M}) \approx 0.00487766$ for

the airplane parameter \tilde{c}_3 . See [13] for the explicit parameterization of period-3 hyperbolic components. This strict inequality is in contrast with the equality $r_{\text{bif}}(0) = \text{dist}(0, \partial\mathcal{M})$.

COROLLARY 4.25. *Almost every random Julia set is totally disconnected:*

- if the central parameter $c = -1 + \epsilon$ and the noise amplitude $r > 0.0399 \dots + |\epsilon|$; or
- if the central parameter $c = \tilde{c}_3 + \epsilon$ and the noise amplitude $r > 0.0021 + |\epsilon|$.

Proof. By Theorem 4.6, $\mathbb{P}_{c,r}$ -almost every random Julia set is totally disconnected if $r > r_{\text{bif}}(c)$. In addition, we have $r_{\text{bif}}(c + \epsilon) \leq r_{\text{bif}}(c) + |\epsilon|$ for every $\epsilon \in \mathbb{C}$ by Theorem 4.10. Thus, the conclusion follows from Examples 4.21 and 4.23. \square

4.5. Application to quasiconformal conjugacy. By combining Theorem 4.6 and the theory of non-autonomous holomorphic motions by Comerford, we can deduce that non-autonomous Julia sets are quasiconformally conjugate to the autonomous Julia set. In this subsection, we see an overview of this conjugacy. For a more detailed meaning, the reader is referred to [7].

For $\omega = (c_1, c_2, \dots) \in \mathbb{C}^{\mathbb{N}}$, we denote $\sigma^m \omega = (c_{m+1}, c_{m+2}, \dots)$ for every $m \geq 0$.

Definition 4.26. Let $\delta > 0$ and let $\omega \in \mathbb{C}^{\mathbb{N}}$ be a bounded sequence of parameters. We say that ω has post-critical distance $\geq \delta$ if

$$\inf_{m \geq 0, n \geq m} \text{dist}(f_{\sigma^m \omega}^{(n-m)}(0), J_{\sigma^m \omega}) \geq \delta.$$

Here, $f_{\omega}^{(0)}$ is understood as the identity map for every ω .

The sequence ω has post-critical distance $\geq \delta$ for some $\delta > 0$ if and only if non-autonomous dynamics is uniformly expanding (hyperbolic) if and only if non-autonomous dynamics is uniformly contractive at critical points. See [7, Theorem 3.3] for the details.

We next define hyperbolic components by analogy with the deterministic case. Denote $\ell^\infty(\mathbb{C}) = \{\omega = (c_n)_{n=1}^\infty : \sup_n |c_n| < \infty\}$.

Definition 4.27. Every connected component of

$$\{\omega \in \ell^\infty(\mathbb{C}) : \omega \text{ has post-critical distance } \geq \delta \text{ for some } \delta > 0\}$$

is called a hyperbolic component.

After formulating the concepts above, Comerford proved the following theorem [7, Theorem 6.1].

THEOREM 4.28. *Suppose that $\omega = (c_n)_{n=1}^\infty$ and $\omega' = (c'_n)_{n=1}^\infty$ lie in the same hyperbolic component. Then there exist $K \geq 1$ and a sequence of maps $\{\varphi_n\}_{n \geq 0}$ such that φ_n maps $J_{\sigma^n \omega}$ onto $J_{\sigma^n \omega'}$ K -quasiconformally and $\varphi_{n+1} \circ f_{c_{n+1}} = f_{c'_{n+1}} \circ \varphi_n$ on $J_{\sigma^n \omega}$ for every $n \geq 0$.*

We say that ω and ω' are quasiconformally conjugate on their iterated Julia sets if such a sequence $\{\varphi_n\}_{n \geq 0}$ exists.

More precisely, the sequence $\{\varphi_n\}_{n \geq 0}$ is bi-equicontinuous, as defined in [7, §4]. Using the theorem above, we show the new results.

THEOREM 4.29. *Suppose that the interior of $\bar{B}(c, r)$ contains a superattracting parameter \tilde{c} and suppose $r < r_{\text{bif}}(c)$. Then every two $\omega, \omega' \in \bar{B}(c, r)^{\mathbb{N}}$ are quasiconformally conjugate on their iterated Julia sets.*

Proof. By Definition 2.17 and Theorem 4.6, the polynomial semigroup $G_{c,r}$ has the planar attracting minimal set $L = \overline{\bigcup_{g \in G_{c,r}} \{g(0)\}}$. Define $\delta = \text{dist}(L, J(G_{c,r}))$, which is strictly positive by Definition 2.13 of attracting minimal sets. For every $\omega \in \bar{B}(c, r)^{\mathbb{N}}$, $m \geq 0$, and $n \geq m$, we have $f_{\sigma^m \omega}^{(n-m)}(0) \in L$ since $f_{\sigma^m \omega}^{(n-m)} \in G_{c,r}$. Also, $J_{\sigma^m \omega} \subset J(G_{c,r})$ by Proposition 2.9. Therefore, ω has post-critical distance $\geq \delta$ with uniform $\delta > 0$. Since the ℓ^∞ -ball $\bar{B}(c, r)^{\mathbb{N}}$ is connected, the ball $\bar{B}(c, r)^{\mathbb{N}}$ is contained in a hyperbolic component. Thus, every two $\omega, \omega' \in \bar{B}(c, r)^{\mathbb{N}}$ are quasiconformally conjugate on their iterated Julia sets by Theorem 4.28. □

More precisely, we can show by using [7, Theorem 1.3] that the maximal dilatation K of the quasiconformal conjugacy $\{\varphi_n\}_{n \geq 0}$ depends on c and r but not on ω nor ω' . This is because the δ in the proof is independent of ω . In addition, we can show that the maximal dilatation K tends to 1 as r tends to 0.

Lastly, we give the following corollary.

COROLLARY 4.30. *Suppose that the interior of $\bar{B}(c, r)$ contains a superattracting parameter \tilde{c} . Denote the autonomous Julia set by $J_{\tilde{c}} = J_{\tilde{\omega}}$ where $\tilde{\omega} = (\tilde{c}, \tilde{c}, \dots)$. If $r < r_{\text{bif}}(c)$, then for every $\omega \in \bar{B}(c, r)^{\mathbb{N}}$, there exists a map φ that maps $J_{\tilde{c}}$ onto the non-autonomous Julia set J_ω quasiconformally. If $r > r_{\text{bif}}(c)$, then for $\mathbb{P}_{c,r}$ -almost every ω , the autonomous Julia set $J_{\tilde{c}}$ is not homeomorphic to the non-autonomous Julia set J_ω .*

Proof. First suppose $r < r_{\text{bif}}(c)$. By Theorem 4.29, there exists a quasiconformal conjugacy $\{\varphi_n\}_{n \geq 0}$ between $\tilde{\omega}$ and ω for every $\omega \in \bar{B}(c, r)^{\mathbb{N}}$. By definition, the map φ_0 quasiconformally maps $J_{\tilde{c}}$ onto J_ω . Next suppose $r > r_{\text{bif}}(c)$. Then by Theorem 4.6, the non-autonomous Julia set J_ω is totally disconnected almost surely. Since the autonomous Julia set $J_{\tilde{c}}$ of superattracting parameter \tilde{c} is connected, these $J_{\tilde{c}}$ and J_ω cannot be homeomorphic each other. □

5. Conclusive discussions

In this section, we discuss three perspectives, namely, open problems, other problems about geometric measure theory, and application to the general settings.

5.1. Open problems. We give the examples of c satisfying the equality $r_{\text{bif}}(c) = \text{dist}(c, \partial\mathcal{M})$ and the examples of c satisfying the strict inequality $r_{\text{bif}}(c) < \text{dist}(c, \partial\mathcal{M})$. The author conjectures that the former ones are uncommon. More precisely, the author considers the answer to the following question is yes.

Question 5.1. If c does not belong to the main cardioid, then does $r_{\text{bif}}(c) < \text{dist}(c, \partial\mathcal{M})$ hold? If $c \notin [0, 1/4]$, then does it hold?

We numerically estimated $0.0386 \dots \leq r_{\text{bif}}(-1) \leq 0.0399 \dots$. However, the actual value remains unknown. More generally, for every superattracting parameter \tilde{c} , it is interesting to determine the bifurcation radius.

Question 5.2. What is the exact value of $r_{\text{bif}}(\tilde{c})$ for a superattracting parameter \tilde{c} ? How does it relate to other dynamical quantities?

In addition, it would be meaningful to investigate their asymptotic behavior.

Question 5.3. Let \tilde{c}_p be a superattracting parameter with period p for every $p \in \mathbb{N}$. How does $r_{\text{bif}}(\tilde{c}_p)$ decrease as $p \rightarrow \infty$?

Our strategy to estimate $r_{\text{bif}}(-1)$ from above is to find multiple roots of some algebraic equation. This raises the following question.

Question 5.4. For every superattracting parameter \tilde{c} , is the bifurcation radius $r_{\text{bif}}(\tilde{c})$ an algebraic number?

At least, if $\tilde{c} = 0$, then $r_{\text{bif}}(0) = 1/4$ is algebraic and is related to the parabolic parameter $c = 1/4$.

We are interested also in parameters which are not superattracting. Recall that Theorem 4.6 is concerned with the case where $\bar{B}(c, r)$ contains superattracting parameters.

Question 5.5. Can we generalize Theorem 4.6 when $\bar{B}(c, r)$ does not contain superattracting parameters?

Question 5.6. How does $r_{\text{bif}}(c)$ decrease as $c \rightarrow \partial\mathcal{M}$? For example, within $\epsilon > 0$, how does $r_{\text{bif}}(-3/4 + \epsilon)$ decrease as $\epsilon \rightarrow 0$?

5.2. Other problems about geometric measure theory. One can consider the Hausdorff dimensions and other geometric properties of random Julia sets. For instance, the uniform perfectness and the Johnness was investigated. See [26, Theorems 1.6 and 1.12].

Regarding dimensions, the following facts are known. Suppose $c = 0$ and $0 < r < 1/4$, and let $(\mathbb{P}_{0,r}, \Omega_{0,r})$ as in Setting 4.1. Denote by $\text{HD}(J)$ the Hausdorff dimension of a set J . Then, there exists $1 < d_{0,r} < 2$ such that $\text{HD}(J_\omega) = d_{0,r}$ for $\mathbb{P}_{0,r}$ -almost every $\omega \in \Omega_{0,r}$. See [18, Theorems 8.8 and 8.10] by Mayer, Skorulski, and Urbański. In addition, denote by \mathcal{H}^d the d -dimensional Hausdorff measure and by \mathcal{P}^d the d -dimensional Packing measure, then $\mathcal{H}^{d_{0,r}}(J_\omega) = 0$ and $\mathcal{P}^{d_{0,r}}(J_\omega) = \infty$ for $\mathbb{P}_{0,r}$ -almost every $\omega \in \Omega_{0,r}$. See [18, Theorem 8.16]. Note that this is different from the deterministic case, and hence almost every J_ω is not bi-Lipschitz equivalent to any autonomous hyperbolic Julia sets.

Sumi kindly taught me that for every $c \in \mathbb{C}$ and $r > 0$, there exists a constant $d_{c,r}$ such that $\text{HD}(J_\omega) = d_{c,r}$ for $\mathbb{P}_{c,r}$ -almost every $\omega \in \Omega_{c,r}$. Rugh showed that the dimension $d_{c,r}$ depends real-analytically on $\text{Re}(c)$, $\text{Im}(c)$, and r within $|c| + r < 1/4$. See [23] for the exact statements. This is the motivation for the following question.

Question 5.7. How does the typical dimension $d_{c,r}$ depend on c and r ?

For other geometric properties, Brück showed that the random Julia set J_ω is a quasi-circle for every $\omega \in \Omega_{0,r}$ if $r < 1/4$. See [2, Corollary 4.5].

Theorem 4.29 generalizes this result. However, the following questions are open.

Question 5.8. Suppose $r > r_{\text{bif}}(c)$. Then for $\mathbb{P}_{c,r} \times \mathbb{P}_{c,r}$ -almost every (ω, ω') , are the Julia sets J_ω and $J_{\omega'}$ quasiconformally equivalent? Note that almost every Julia set is homeomorphic to the (usual) Cantor set.

Question 5.9. If $r = r_{\text{bif}}(c)$, then is almost every random Julia set J_ω quasiconformally equivalent to the autonomous Julia set?

In this paper, we considered the measure-theoretic aspect of the typicality problem. For the topological aspect, see [14], for example.

5.3. Application to general settings. We mainly considered random dynamical systems induced by uniform noise on disks as in Setting 4.1. However, random dynamical systems induced by any distribution μ with compact support can be considered. Even for Setting 2.1, we can apply the numerical estimates given in §4. More precisely, the following hold.

THEOREM 5.10. *Let μ be a Borel probability measure on the parameter plane \mathbb{C} with compact support and let $(\mathbb{P}_\mu, \Omega_\mu)$ be defined as in Setting 2.1. If there exists $c \in \mathbb{C}$ such that $\text{supp } \mu \subset \bar{B}(c, r_{\text{bif}}(c))$, then the polynomial semigroup G_μ has a planar minimal set. If there exists $c \in \mathbb{C}$ such that $\text{int}(\text{supp } \mu) \supset \bar{B}(c, r_{\text{bif}}(c))$, then the random Julia set J_ω is totally disconnected for \mathbb{P}_μ -almost every $\omega \in \Omega_\mu$.*

Proof. Suppose that there exists $c \in \mathbb{C}$ such that $\text{supp } \mu \subset \bar{B}(c, r_{\text{bif}}(c))$. By definition, the polynomial semigroup $G_{c, r_{\text{bif}}(c)}$ has a planar minimal set. Thus, Lemma 2.12 gives that G_μ also has a planar minimal set.

Similarly, if there exists $c \in \mathbb{C}$ such that $\text{int}(\text{supp } \mu) \supset \bar{B}(c, r_{\text{bif}}(c))$, then G_μ does not have any planar minimal sets. Thus, G_μ is mean stable, and $T_{\{\infty\}, \mu}$ is identically equal to 1 by Theorem 2.21. Combining this with Theorem 3.7, we have that the random Julia set J_ω is totally disconnected for \mathbb{P}_μ -almost every $\omega \in \Omega_\mu$. \square

Example 5.11. Let μ_s be the normalized Lebesgue measure on \mathbb{C} with $\text{supp } \mu_s = \{c \in \mathbb{C} : |\text{Re } c| \leq s, |\text{Im } c| \leq s\}$ for $s \geq 0$. Then there exists $s^* > 0$ such that G_{μ_s} is mean stable if and only if $s \notin \{0, s^*\}$. Theorem 5.10 and Corollary 4.11 imply $(4\sqrt{2})^{-1} \leq s^* \leq 4^{-1}$.

Generally, we may apply our results to non-quadratic cases. Since ‘the Mandelbrot set is universal’ [20], the structure of the quadratic family exists for any non-trivial family. Thus, our quadratic results apply to random dynamical systems of any non-trivial family. In this case, it is difficult to obtain quantitative estimates of bifurcation parameters.

Acknowledgments. The author thanks Hiroki Sumi for detailed comments and valuable discussions. The author thanks Isaia Nisoli for the instruction to use validated numerics.

The manuscript has been edited for English language by a professional native proofreader at Xtra, Inc. This work was supported by the Research Institute for Mathematical Sciences, an International Joint Usage/Research Center located in Kyoto University. This work is supported by JSPS Grant-in-Aid for Research Activity Start-up Grant Number JP 21K20323 and Grant-in-Aid for Early-Career Scientists Grant Number JP 23K13000.

REFERENCES

- [1] L. Arnold. *Random Dynamical Systems (Springer Monographs in Mathematics)*. Springer-Verlag, Berlin, 1998.
- [2] R. Brück. Geometric properties of Julia sets of the composition of polynomials of the form $z^2 + c_n$. *Pacific J. Math.* **198**(2) (2001), 347–372.
- [3] R. Brück and M. Büger. Generalized iteration. *Comput. Methods Funct. Theory* **3**(1–2) (2003), 201–252.
- [4] R. Brück, M. Büger and S. Reitz. Random iterations of polynomials of the form $z^2 + c_n$: connectedness of Julia sets. *Ergod. Th. & Dynam. Sys.* **19** (1999), 1221–1231.
- [5] M. Büger. On the composition of polynomials of the form $z^2 + c_n$. *Math. Ann.* **310**(4) (1998), 661–683.
- [6] L. Carleson and T. Gamelin. *Complex Dynamics*. Springer-Verlag, New York, 1993.
- [7] M. Comerford. Holomorphic motions of hyperbolic nonautonomous Julia sets. *Complex Var. Elliptic Equ.* **53**(1) (2008), 1–22.
- [8] M. Comerford. Non-autonomous Julia sets with measurable invariant sequences of line fields. *Discrete Contin. Dyn. Syst.* **33**(2) (2013), 629–642.
- [9] M. Comerford, R. Stankewitz and H. Sumi. Hereditarily non uniformly perfect non-autonomous Julia sets. *Discrete Contin. Dyn. Syst.* **40**(1) (2020), 33–46.
- [10] M. D. Comerford. Properties of Julia sets for the arbitrary composition of monic polynomials with uniformly bounded coefficients. *PhD Thesis*, Yale University, 2001.
- [11] R. Durrett. *Probability—Theory and Examples (Cambridge Series in Statistical and Probabilistic Mathematics, 49)*, 5th edn. Cambridge University Press, Cambridge, 2019.
- [12] J. Fornæss and N. Sibony. Random iterations of rational functions. *Ergod. Th. & Dynam. Sys.* **11** (1991), 687–708.
- [13] D. Giarrusso and Y. Fisher. A parameterization of the period 3 hyperbolic components of the Mandelbrot set. *Proc. Amer. Math. Soc.* **123**(12) (1995), 3731–3737.
- [14] Z. Gong, W. Qiu and Y. Li. Connectedness of Julia sets for a quadratic random dynamical system. *Ergod. Th. & Dynam. Sys.* **23** (2003), 1807–1815.
- [15] A. Hinkkanen and G. Martin. The dynamics of semigroups of rational functions I. *Proc. Lond. Math. Soc.* (3) **73**(2) (1996), 358–384.
- [16] M. Jonsson. Ergodic properties of fibered rational maps. *Ark. Mat.* **38**(2) (2000), 281–317.
- [17] K. Lech and A. Zdunik. Total disconnectedness of Julia sets of random quadratic polynomials. *Ergod. Th. & Dynam. Sys.* **42**(5) (2022), 1764–1780.
- [18] V. Mayer, B. Skorulski and M. Urbański. *Distance Expanding Random Mappings, Thermodynamical Formalism, Gibbs Measures and Fractal Geometry (Lecture Notes in Mathematics, 2036)*. Springer, Heidelberg, 2011.
- [19] C. T. McMullen. *Complex Dynamics and Renormalization (Annals of Mathematics Studies, 135)*. Princeton University Press, Princeton, NJ, 1994.
- [20] C. T. McMullen. The Mandelbrot set is universal. *The Mandelbrot Set, Theme and Variations (London Mathematical Society Lecture Note Series, 274)*. Ed. T. Lei. Cambridge University Press, Cambridge, 2000, pp. 1–17.
- [21] J. Milnor. *Dynamics in One Complex Variable (Annals of Mathematical Studies, 160)*, 3rd edn. Princeton University Press, Princeton, NJ, 2006.
- [22] M. S. Petković and L. D. Petković. *Complex Interval Arithmetic and Its Applications (Mathematical Research, 105)*. Wiley-VCH Verlag Berlin GmbH, Berlin, 1998.
- [23] H. H. Rugh. On the dimensions of conformal repellers. Randomness and parameter dependency. *Ann. of Math.* (2) **168**(3) (2008), 695–748.
- [24] R. Stankewitz. Density of repelling fixed points in the Julia set of a rational or entire semigroup. II. *Discrete Contin. Dyn. Syst.* **32**(7) (2012), 2583–2589.

- [25] H. Sumi. Skew product maps related to finitely generated rational semigroups. *Nonlinearity* **13**(4) (2000), 995–1019.
- [26] H. Sumi. Semi-hyperbolic fibered rational maps and rational semigroups. *Ergod. Th. & Dynam. Sys.* **26**(3) (2006), 893–922.
- [27] H. Sumi. Random complex dynamics and semigroups of holomorphic maps. *Proc. Lond. Math. Soc.* (3) **102**(1) (2011), 50–112.
- [28] H. Sumi. Cooperation principle, stability and bifurcation in random complex dynamics. *Adv. Math.* **245** (2013), 137–181.
- [29] H. Sumi. Negativity of Lyapunov exponents and convergence of generic random polynomial dynamical systems and random relaxed Newton's methods. *Comm. Math. Phys.* **384**(3) (2021), 1513–1583.
- [30] H. Sumi and T. Watanabe. Non-i.i.d. random holomorphic dynamical systems and the probability of tending to infinity. *Nonlinearity* **32**(10) (2019), 3742–3771.
- [31] H. Sumi and T. Watanabe. Non-i.i.d. random holomorphic dynamical systems and the generic dichotomy. *Nonlinearity* **35**(4) (2022), 1857–1875.