

Stability of strong travelling waves for a non-local time-delayed reaction–diffusion equation

Ming Mei

Department of Mathematics and Statistics, McGill University,
Montreal, Quebec H3A 2K6, Canada and
Department of Mathematics, Champlain College, Saint-Lambert,
Quebec J4P 3P2, Canada (mei@math.mcgill.ca)

Joseph W.-H. So

Department of Mathematical and Statistical Sciences,
University of Alberta, Edmonton, Alberta T6G 2G1, Canada
(joseph.so@ualberta.ca)

(MS received 13 March 2006; accepted 7 February 2007)

The paper is concerned with a non-local time-delayed reaction–diffusion equation. We prove the (time) asymptotic stability of a travelling wavefront without a smallness assumption on its wavelength, i.e. the so-called strong wavefront, by means of the (technical) weighted energy method, when the initial perturbation around the wave is small. The exponential convergent rate is also given. Selection of a suitable weight plays a key role in the proof.

1. Introduction

For population dynamics with age structure and diffusion, Metz and Diekmann [16] studied the governing model

$$\frac{\partial u}{\partial t} + \frac{\partial u}{\partial z} - D(z) \frac{\partial^2 u}{\partial x^2} + d(z)u = 0, \quad (1.1)$$

where $u(t, z, x)$ denotes the population density of the species under consideration at time $t \geq 0$, age $z \geq 0$ and location $x \in \Omega$, and $D(z)$ and $d(z)$ are the diffusion rate and death rate of population at age z , respectively. We denote by $r > 0$ the maturation time for the species. Ω is the spatial domain where the species live and it can be bounded or unbounded. We take Ω to be $\mathbb{R} = (-\infty, \infty)$ in this paper.

Let $v(t, x)$ be the total mature population at time t and location x

$$v(t, x) = \int_r^\infty u(t, z, x) dz, \quad (1.2)$$

and let $D_m(z) = D(z)$ for $z \in [r, \infty)$ and $D_i(z) = D(z)$ for $z \in [0, r)$ be the diffusion rates for the mature population and the immature population, respectively. In the event that the mature population is more effective in spatial diffusion than the

immature population, it is then reasonable to assume that

$$\min_{z \in [r, \infty)} D_m(z) \geq \max_{z \in [0, r]} D_i(z). \tag{1.3}$$

We also denote by $d_m(z) = d(z)$ for $z \in [r, \infty)$ and $d_i(z) = d(z)$ for $z \in [0, r)$ the death rates for the mature population and the immature population, respectively. At age zero, $u(t, 0, x)$ is the population density of the newborns. Since only matures can reproduce, we have

$$u(t, 0, x) = b(v(t, x)), \tag{1.4}$$

where $b(\cdot)$ is the birth function.

When the diffusion and death rates for the mature population are independent of age, namely,

$$D_m(z) = D_m \quad \text{and} \quad d_m(z) = d_m \tag{1.5}$$

are constants for $z \in [r, \infty)$, So *et al.* [26], following the approach of Smith and Thieme [22] (see also [9, 27]), reduced equation (1.1) into a non-local time-delayed reaction–diffusion equation for $v(t, x)$ by the Fourier transform method,

$$\frac{\partial v}{\partial t} - D_m \frac{\partial^2 v}{\partial x^2} + d_m v = \varepsilon \int_{-\infty}^{\infty} b(v(t-r, x-y)) f_\alpha(y) dy, \tag{1.6}$$

where

$$\varepsilon = \exp\left(-\int_0^r d_i(z) dz\right) \quad \text{and} \quad \alpha = \int_0^r D_i(z) dz \tag{1.7}$$

represent the impact of the death rate of the immature and the effect of the dispersal rate of the immature on the mature population, respectively. By (1.3), (1.5) and (1.7), we have

$$\alpha \leq r D_m. \tag{1.8}$$

In (1.6), $f_\alpha(y)$ is the heat kernel function

$$f_\alpha(y) = \frac{1}{\sqrt{4\pi\alpha}} e^{-y^2/4\alpha} \quad \text{and} \quad \int_{-\infty}^{\infty} f_\alpha(y) dy = 1. \tag{1.9}$$

In particular, when the birth function $b(v)$ is that used for Nicolson’s blowflies [5, 15, 24–26], that is

$$b(v) = p v e^{-av}, \tag{1.10}$$

where $p > 0$ and $a > 0$ are constants, the constant equilibria for equation (1.1) can be found by solving

$$d_m v = \varepsilon p \int_{-\infty}^{\infty} v e^{-av} f_\alpha(y) dy.$$

By (1.9), this equation admits only two roots:

$$v_- = 0, \quad v_+ = \frac{1}{a} \ln \frac{\varepsilon p}{d_m}. \tag{1.11}$$

If $\varepsilon p/d_m > 1$, then $v_- < v_+$. In [26], So *et al.* showed the existence of travelling waves $\phi(x + ct)$ connecting the two equilibria v_\pm with speed c . For other models

with non-local terms, the existence of travelling waves has been shown in [4,9,27,33] (see also the references therein).

Here, we are interested in the asymptotic stability for such waves. For the Cauchy problem for equation (1.6) with the birth function (1.10) and the initial data

$$v(t, x)|_{t=s} = v_0(s, x) \quad \text{for } s \in [-r, 0], \quad x \in \mathbb{R}, \quad (1.12)$$

where

$$v_0(s, x) \rightarrow v_{\pm} \quad \text{for all } s \in [-r, 0] \text{ as } x \rightarrow \pm\infty,$$

we prove that the global solution $v(t, x)$ of (1.6), (1.10), (1.12) converges to the travelling wave $\phi(x + ct)$ asymptotically (in time), when the initial perturbation around the wave, that is, $|v_0(s, x) - \phi(x + cs)|$, $s \in [-r, 0]$, is suitably small. The exponential convergence rate will also be obtained.

The study of the stability of travelling waves is interesting and usually (technically) difficult. For partial differential equations without time delays, including reaction–diffusion equations, travelling waves have been extensively studied; see, for example, the pioneering works [6, 20] and other more recent contributions [1–3, 7–14, 18–20, 30, 31], and the references therein (see also [28] and the survey papers for viscous equations of conservation laws by Matsumura [10] and the reaction–diffusion equations by Xin [32]). However, results for the time-delayed partial differential equations are very limited and incomplete. The first work related to this topic was done by Schaaf [21] on the linearized stability of the time-delayed Fisher–Kolmogorov–Petrovski–Piskunov equation by means of the spectral method. Later, Ogiwara and Matano [17] and Smith and Zhao [23] studied the nonlinear stability by the method of upper and lower solutions. See also [29]. More recently, Mei *et al.* [15] proved the nonlinear wave stability for the local equation with birth function (1.10), where, for the two steady states connected by the travelling wave, one of the equilibria is an unstable node. Such a case is different from the ‘bistable’ nodes studied in [23]. For the non-local case, Liang and Wu [9] studied theoretically the existence of the travelling waves for (1.6) with a different birth function, $b(v)$, and, furthermore, showed the wave approximations numerically.

Following [9, 15], we treat the non-local case here with nonlinearity (1.10), and prove theoretically the stability for the strong travelling waves. A wave is said to be weak if its wavelength is small, that is, $|v_+ - v_-| \ll 1$; otherwise, the wave is said to be strong. As is well known, one may prove the stability for the weak waves in certain cases, but one cannot usually prove it for the strong wave cases. As in [15], the approach adopted in this paper is still the weighted energy method. In the proof, a key role is played by the selection of a suitable weight function; see the key lemma (lemma 3.6), below.

The rest of the paper is organized as follows. In §2, we state the result on the existence of travelling waves as given in [26]. After defining a suitable weight function, we state the theorem on wave stability. Section 3 is devoted to the proof of the stability theorem using the weighted energy method. The key step in the proof is to establish *a priori* estimates.

Before ending this section, we give some notation. Throughout the paper, $C > 0$ denotes a generic constant, while $C_i > 0$, $i = 0, 1, 2, \dots$, represents a specific constant. Let I be an interval; typically $I = \mathbb{R}$. $L^2(I)$ is the space of square integrable

functions on an interval I , and $H^k(I)$, $k \geq 0$, is the Sobolev space of L^2 -functions $f(x)$ defined on the interval I whose derivatives $d^i f/dx^i$, $i = 1, \dots, k$, also belong to $L^2(I)$. $L^2_w(I)$ represents the weighted L^2 -space with weight $w(x) > 0$. Its norm is defined by

$$\|f\|_{L^2_w} = \left(\int_I w(x)f(x)^2 dx \right)^{1/2}.$$

$H^k_w(I)$ is the weighted Sobolev space with the norm

$$\|f\|_{H^k_w} = \left(\sum_{i=0}^k \int_I w(x) \left| \frac{d^i}{dx^i} f(x) \right|^2 dx \right)^{1/2}.$$

Let $T > 0$ and let \mathcal{B} be a Banach space. We denote by $C^0([0, T]; \mathcal{B})$ the space of \mathcal{B} -valued continuous functions on $[0, T]$, and by $L^2([0, T]; \mathcal{B})$ the space of \mathcal{B} -valued L^2 -functions on $[0, T]$. The corresponding spaces of \mathcal{B} -valued functions on $[0, \infty)$ are defined similarly.

2. Stability of strong travelling waves

A travelling wave of the type in equation (1.6) with the birth function (1.10) connecting with two constant steady states v_{\pm} is a special solution of equation (1.6) of the form $\phi(x + ct)$ ($c > 0$ is the wave speed) satisfying the non-local delayed ordinary differential equation

$$\left. \begin{aligned} c\phi'(\xi) - D_m\phi''(\xi) + d_m\phi(\xi) &= \varepsilon p \int_{-\infty}^{\infty} \phi(\xi - cr - y)e^{-a\phi(\xi - cr - y)} f_{\alpha}(y) dy, \\ \phi(\pm\infty) &= v_{\pm}, \end{aligned} \right\} \tag{2.1}$$

where $\xi = x + ct$ and the prime denotes differentiation with respect to ξ . Using the upper and lower solution method, So *et al.* [26] proved the following result on the existence of monotone wavefronts $\phi(\xi)$ with $\phi'(\xi) > 0$.

PROPOSITION 2.1 (So *et al.* [26]). *If $1 < \varepsilon p/d_m \leq e$, then there exists a critical number c^* ,*

$$0 < c^* < 2\sqrt{D_m(\varepsilon p - d_m)}, \tag{2.2}$$

such that, for every $c > c^$, equation (1.6) has a travelling wavefront solution $\phi(\xi)$ connecting v_{\pm} , with $\phi'(\xi) > 0$ and $v_- < \phi(\xi) < v_+$ for all $\xi \in (-\infty, \infty)$.*

As is well known, in order to prove wave stability it is often necessary to restrict the wavelength to be sufficiently small (that is, $|v_+ - v_-| \ll 1$). Such a wave is called a weak wave. Here, we are interested in establishing the stability of a strong wave. For this, throughout the present paper, let us take $\varepsilon p/d_m$ in proposition 2.1 ($1 < \varepsilon p/d_m \leq e$) to be e , that is,

$$\frac{\varepsilon p}{d_m} = e, \tag{2.3}$$

so that

$$v_+ = \frac{1}{a} \ln \frac{\varepsilon p}{d_m} = \frac{1}{a}$$

is maximum, namely, the wave $\phi(\xi)$ connecting the two equilibria, $v_- = 0$ and $v_+ = 1/a$, is the strongest.

Let

$$\bar{\varepsilon} := \frac{(3 - e)d_m}{2(ad_m + aD_m + \frac{3}{2}ed_m)}. \tag{2.4}$$

We first have the following estimates.

LEMMA 2.2. *For a given strongest travelling wave $\phi(\xi)$, $\xi = x + ct$, there exists a number x_* such that, for $\xi > x_*$, the following inequalities hold:*

$$\left. \begin{aligned} \phi(\xi) &> \frac{1}{a} - \bar{\varepsilon}, \\ |\phi''(\xi)| &< \bar{\varepsilon}, \\ 0 &< e^{a\phi(\xi)}(1 - a\phi(\xi)) < \bar{\varepsilon}. \end{aligned} \right\} \tag{2.5}$$

Proof. For the given strongest travelling wave $\phi(\xi)$, it is easy to see that $0 = v_- \leq \phi(\xi) \leq v_+ = 1/a$ because $\phi(\xi)$ is increasing. Since $\lim_{\xi \rightarrow +\infty} \phi(\xi) = v_+ = 1/a$, $\lim_{\xi \rightarrow +\infty} \phi''(\xi) = 0$ and $\lim_{\xi \rightarrow +\infty} (1 - a\phi(\xi)) = 0$, by the definition of limits, for the given $\bar{\varepsilon}$ there exists a number x_* such that when $\xi > x_*$ the following hold:

$$\begin{aligned} |\phi(\xi) - v_+| &= \left| \phi(\xi) - \frac{1}{a} \right| < \bar{\varepsilon}, \\ |\phi''(\xi)| &< \bar{\varepsilon}, \\ 0 &< e^{a\phi(\xi)}(1 - a\phi(\xi)) < \bar{\varepsilon}. \end{aligned}$$

These inequalities immediately imply (2.5). □

Now we define a weight function $w(\xi)$ as

$$w(\xi) = \begin{cases} e^{-\beta(\xi-x_*)} = e^{\beta|\xi-x_*|}, & \xi < x_*, \\ 1, & \xi \geq x_*, \end{cases} \tag{2.6}$$

where

$$\beta = \frac{c}{2D_m}. \tag{2.7}$$

Our main result for the paper is the following.

THEOREM 2.3 (stability). *Consider the given strongest travelling wavefront $\phi(x + ct)$, where the speed $c > c_*$ satisfies*

$$c > 2\sqrt{D_m(3\varepsilon p - 2d_m)}. \tag{2.8}$$

If

$$v_0(s, x) - \phi(x + cs) \in C^0([-r, 0]; H_w^1(\mathbb{R})), \tag{2.9}$$

where $w = w(x + cs)$, $s \in [-r, 0]$, is the weight function given in (2.6), then there exist positive constants δ_0 and μ , which are dependent only on the coefficients $D_m, d_m, \varepsilon, p, a, r$ and the wave speed c , such that when $\|v_0(s, \cdot) - \phi(\cdot + cs)\|_{H_w^1} \leq \delta_0$ for

$s \in [-r, 0]$ the unique solution $v(t, x)$ of the Cauchy problem (1.6), (1.10), (1.12) exists globally, and it satisfies

$$v(t, x) - \phi(x + ct) \in C^0([0, \infty); H_w^1) \cap L^2([0, \infty); H_w^2)$$

and

$$\sup_{x \in \mathbb{R}} |v(t, x) - \phi(x + ct)| \leq C e^{-\mu t}, \quad 0 \leq t < \infty, \tag{2.10}$$

where $C > 0$ is a constant dependent only on the initial perturbation $v_0(x, s) - \phi(x - cs)$.

REMARK 2.4. (i) Note that (2.2), (2.8) and $3\varepsilon p - 2d_m > \varepsilon p - d_m$ imply that

$$c > 2\sqrt{D_m(3\varepsilon p - 2d_m)} > 2\sqrt{D_m(\varepsilon p - d_m)} > c^*.$$

Thus, theorem 2.3 ensures that when the wave speed is not too close to c^* the strongest wave is asymptotically stable (in time). For speed c close to the critical point c^* , and in particular the case when $c = c^*$, the stability problem is still open.

(ii) By the definition of weight function (2.6) and the definition of the weighted Sobolev space $H_w^1(R)$, we have from (2.9) that

$$\sqrt{w(x + cs)}(v_0(s, x) - \phi(x + cs)) \in H^1(R), \quad s \in [-r, 0].$$

Thus, applying Sobolev's inequality, we obtain

$$\sqrt{w(x + cs)}(v_0(s, x) - \phi(x + cs)) \leq C \|v_0 - \phi\|_{H_w^1}, \quad s \in [-r, 0],$$

which in turn implies that

$$|v_0(s, x) - \phi(x + cs)| \sim w^{-1/2}(x + cs) \sim e^{-\beta|x|/2} \quad \text{as } x \rightarrow -\infty.$$

(iii) Since c is large, by a straightforward but tedious computation we find that the convergence of the initial perturbation $|v_0(s, x) - \phi(x + ct)| \sim O(1) \exp\{-c|x|/4D_m\}$ for $x \rightarrow -\infty$ is faster than the decay of the wavefront to $v_- = 0$ in the form $|\phi(x + cs) - v_-| = |\phi(x + cs)| \sim O(1)e^{-\beta_-|x|}$ as $x \rightarrow -\infty$, where β_- is a positive constant satisfying $\beta_- < c/4D_m$. So, as we show in theorem 2.3, it is not surprising that, when $|v_0(s, x) - \phi(x + cs)| \ll 1$, the solution $v(t, x)$ converges to $\phi(x + ct)$ and not to some shifted wave $\phi(x + ct + x_0)$ with a shift x_0 . In fact, by another tedious computation, as shown in [11], we can formally show that $x_0 = 0$.

3. Proof of stability

This section is devoted to the proof of the stability result, theorem 2.3. Our proof relies on the weighted energy method.

Let $v(t, x)$ be the solution of the Cauchy problem (1.6), (1.10), (1.12), and let $\phi(x + ct)$ be the wavefront. Set

$$V(t, \xi) = v(t, x) - \phi(\xi), \quad \xi = x + ct.$$

The original problem (1.6), (1.10) and (1.12) can be reformulated as

$$\left. \begin{aligned} &V_t(t, \xi) + cV_\xi(t, \xi) - D_m V_{\xi\xi}(t, \xi) + d_m V(t, \xi) \\ &\quad - \varepsilon \int_{\mathbb{R}} b'(\phi(\xi - y - cr))V(t - r, \xi - y - cr)f_\alpha(y) \, dy \\ &\quad = \varepsilon \int_{\mathbb{R}} Q(V(t - r, \xi - y - cr))f_\alpha(y) \, dy, \quad (t, \xi) \in \mathbb{R}_+ \times \mathbb{R}, \\ &V(s, \xi) = v_0(s, \xi - cs) - \phi(\xi) =: V_0(s, \xi), \quad (s, \xi) \in [-r, 0] \times \mathbb{R}. \end{aligned} \right\} \tag{3.1}$$

The nonlinear term $Q(V(t - r, \xi - y - cr))$ is given by

$$Q(V) = b(\phi + V) - b(\phi) - b'(\phi)V, \tag{3.2}$$

where $\phi = \phi(\xi - y - cr)$ and $V = V(t - r, \xi - y - cr)$. It suffices to prove the following stability result for equation (3.1).

THEOREM 3.1. *For the given strongest travelling wave $\phi(\xi)$, $\xi = x + ct$, with speed c satisfying (2.8), if $V_0(s, \xi) \in C^0([-r, 0]; H_w^1(\mathbb{R}))$, where $w(\xi)$ is the weight function defined in (2.6), then there exist positive constants δ_0 and μ such that when $\sup_{s \in [-r, 0]} \|V_0(s)\|_{H_w^1} \leq \delta_0$, the solution $V(t, \xi)$ of the Cauchy problem (3.1) exists uniquely and globally, and satisfies*

$$V(t, \xi) \in C^0([0, \infty); H_w^1(\mathbb{R})) \cap L^2([0, \infty); H_w^2(\mathbb{R}))$$

and

$$\sup_{\xi \in \mathbb{R}} |V(t, \xi)| \leq Ce^{-\mu t}, \quad 0 \leq t \leq \infty. \tag{3.3}$$

Note that $\mu > 0$ and $\delta_0 > 0$ are the same as those in theorem 2.3. We will prove theorem 3.1 based on the following two propositions: one local estimate and an *a priori* estimate by the continuity argument (see also [7, 11, 12, 14]).

For given constants $\tau \geq 0$ and $T > 0$, we define the solution space by

$$X(\tau - r, T + \tau) = \{V \mid V(t, \xi) \in C^0([\tau - r, T + \tau]; H_w^1(\mathbb{R})) \cap L^2([\tau - r, T + \tau]; H_w^2(\mathbb{R}))\}$$

and

$$M_\tau(T) := \sup_{t \in [\tau - r, T + \tau]} \|V(t)\|_{H_w^1},$$

in particular, $M(T) := M_0(T)$ for $\tau = 0$. For simplicity, we henceforth define $V(t) = V(t, \cdot)$. First we state the local estimate.

PROPOSITION 3.2 (local existence). *Consider the Cauchy problem with the initial time $\tau \geq 0$*

$$\left. \begin{aligned} &V_t(t, \xi) + cV_\xi(t, \xi) - D_m V_{\xi\xi}(t, \xi) + d_m V(t, \xi) \\ &\quad - \varepsilon \int_{\mathbb{R}} b'(\phi(\xi - y - cr))V(t - r, \xi - y - cr)f_\alpha(y) \, dy \\ &\quad = \varepsilon \int_{\mathbb{R}} Q(V(t - r, \xi - y - cr))f_\alpha(y) \, dy, \quad (t, \xi) \in (\tau, \infty) \times \mathbb{R}, \\ &V(s, \xi) = v_0(s, \xi - cs) - \phi(\xi) =: V_\tau(s, \xi), \quad (s, \xi) \in [\tau - r, \tau] \times \mathbb{R}. \end{aligned} \right\} \tag{3.4}$$

If $V_\tau(s, \xi) \in H_w^1$ and $M_\tau(0) \leq \delta_1$ for a given positive constant δ_1 , then there exists a small $t_0 = t_0(\delta_1) > 0$ such that $V(t, \xi) \in X(\tau - r, \tau + t_0)$ and $M_\tau(t_0) \leq \sqrt{2(1+r)}M_\tau(0)$.

The proof of proposition 3.2 can be given by the elementary energy method. We omit the details. Next, we state the *a priori* estimate.

PROPOSITION 3.3 (*a priori* estimate). Let $V(t, \xi) \in X(-r, T)$ be a local solution of (3.1) for a given constant $T > 0$. Then there exist positive constants μ, δ_2 and $C_1 > 1$ independent of T such that, when $M(T) \leq \delta_2$,

$$e^{2\mu t} \|V(t)\|_{H_w^1}^2 + \int_0^t e^{2\mu s} \|V(s)\|_{H_w^2}^2 ds \leq C_1 \left(\|V_0(0)\|_{H_w^1}^2 + \int_{-r}^0 \|V(s)\|_{H_w^1}^2 ds \right), \quad 0 \leq t \leq T \quad (3.5)$$

and

$$\|V(t)\|_{H_w^1}^2 \leq C_1 \left(\|V_0(0)\|_{H_w^1}^2 + \int_{-r}^0 \|V_0(s)\|_{H_w^1}^2 ds \right) e^{-2\mu t}, \quad 0 \leq t \leq T. \quad (3.6)$$

REMARK 3.4. Positive constants μ, δ_2 and C_1 , which depend only on the coefficients $D_m, d_m, \varepsilon, p, a, r$ and the wave speed c , will be specified in (3.17), (3.40), and (3.43), below.

We postpone the proof of proposition 3.3 to the last part of this section. Now, based on propositions 3.2 and 3.3, we will prove theorem 3.1 using the continuation argument.

Proof of theorem 3.1. Recall that the constants δ_2, μ and C_1 from proposition 3.3 are independent of T . Let

$$\delta_1 = \max\{\sqrt{C_1(1+r)}M(0), \delta_2\}, \quad (3.7)$$

$$\delta_0 = \min\left\{ \frac{\delta_2}{\sqrt{2(1+r)}}, \frac{\delta_2}{\sqrt{2C_1(1+r)}} \right\} \quad (3.8)$$

and

$$M(0) \leq \delta_0 < \delta_1. \quad (3.9)$$

By proposition 3.2, there exists $t_0 = t_0(\delta_1) > 0$ such that $V(t, \xi) \in X(-r, t_0)$ and

$$M(t_0) \leq \sqrt{2(1+r)}M(0) \leq \sqrt{2(1+r)}\delta_0 \leq \delta_2.$$

Thus, applying proposition 3.3 on the interval $[0, t_0]$, we obtain (3.6) for $t \in [0, t_0]$, and

$$\begin{aligned} \sup_{t \in [0, t_0]} \|V(t)\|_{H_w^1} &\leq \sup_{t \in [0, t_0]} \left\{ C_1 \left(\|V_0(0)\|_{H_w^1}^2 + \int_{-r}^0 \|V_0(s)\|_{H_w^1}^2 ds \right) \right\}^{1/2} e^{-\mu t} \\ &\leq \sqrt{C_1(1+r)}M(0) \leq \sqrt{C_1(1+r)}\delta_0 \\ &\leq \frac{\delta_2}{\sqrt{2(1+r)}}. \end{aligned} \quad (3.10)$$

Now consider the Cauchy problem (3.4) at the initial time $\tau = t_0$. Using (3.9), (3.10) and (3.7), we have

$$\begin{aligned} M_{t_0}(0) &= \sup_{s \in [t_0-r, t_0]} \|V(s)\|_{H_w^1} \\ &\leq \max \left\{ \sup_{s \in [-r, 0]} \|V(s)\|_{H_w^1}, \sup_{s \in [0, t_0]} \|V(s)\|_{H_w^1} \right\} \\ &\leq \max \left\{ M(0), \frac{\delta_2}{\sqrt{2(1+r)}} \right\} \leq \delta_1. \end{aligned} \tag{3.11}$$

Applying proposition 3.2 again yields $V(t, \xi) \in X(-r, 2t_0)$ and

$$M_{t_0}(t_0) \leq \sqrt{2(1+r)}M_{t_0}(0).$$

On the other hand,

$$\begin{aligned} M_{t_0}(0) &= \sup_{t \in [t_0-r, t_0]} \|V(s)\|_{H_w^1} \\ &\leq \max \left\{ \sup_{s \in [-r, 0]} \|V(s)\|_{H_w^1}, \sup_{s \in [0, t_0]} \|V(s)\|_{H_w^1} \right\} \\ &\leq \max \left\{ \delta_0, \frac{\delta_2}{\sqrt{2(1+r)}} \right\} \leq \frac{\delta_2}{\sqrt{2(1+r)}}. \end{aligned} \tag{3.12}$$

Furthermore, we have

$$M_{t_0}(t_0) \leq \sqrt{2(1+r)}M_{t_0}(0) \leq \delta_2.$$

Consequently,

$$\begin{aligned} M(2t_0) &= \sup_{s \in [-r, 2t_0]} \|V(s)\|_{H_w^1} \\ &\leq \max \left\{ \sup_{s \in [-r, 0]} \|V(s)\|_{H_w^1}, \sup_{s \in [0, t_0-r]} \|V(s)\|_{H_w^1}, \sup_{s \in [t_0-r, 2t_0]} \|V(s)\|_{H_w^1} \right\} \\ &\leq \max \left\{ \delta_0, \frac{\delta_2}{\sqrt{2(1+r)}}, \delta_2 \right\} \leq \delta_2. \end{aligned} \tag{3.13}$$

We can apply proposition 3.3 to obtain (3.6) for $0 \leq t \leq 2t_0$ and

$$\begin{aligned} \sup_{t \in [0, 2t_0]} \|V(t)\|_{H_w^1} &\leq \sup_{t \in [0, 2t_0]} \left\{ C_1 \left(\|V_0(0)\|_{H_w^1}^2 + \int_{-r}^0 \|V_0(s)\|_{H_w^1}^2 ds \right) \right\}^{1/2} e^{-\mu t} \\ &\leq \sqrt{C_1(1+r)}M(0) \leq \sqrt{C_1(1+r)}\delta_0 \leq \frac{\delta_2}{\sqrt{2(1+r)}}. \end{aligned} \tag{3.14}$$

Repeating the previous procedure, one can prove that $V(t, x) \in X(-r, \infty)$ and (3.6) for all $0 \leq t < \infty$. Also (3.3) follows immediately from (3.6). The proof is complete. \square

REMARK 3.5. The proof of theorem 3.1 above corrected the mistake in the proof of theorem 3.1 in [15, pp. 586–587], where δ_0 and δ_1 were defined incorrectly.

Next, we will prove proposition 3.3. For this, we need the following important lemma.

LEMMA 3.6 (key inequality). *Let $w(\xi)$ be the weight function given in (2.6) and define*

$$\begin{aligned}
 B_\mu(\xi) := & -c \frac{w'(\xi)}{w(\xi)} - D_m \left(\frac{w'(\xi)}{w(\xi)} \right)^2 + 2d_m \\
 & - 2\mu - \varepsilon \int_{\mathbb{R}} b'(\phi(\xi - y - cr)) f_\alpha(y) dy \\
 & - \varepsilon e^{2\mu r} \frac{b'(\phi(\xi))}{w(\xi)} \int_{\mathbb{R}} w(\xi + y + cr) f_\alpha(y) dy.
 \end{aligned} \tag{3.15}$$

If (2.8) holds, then

$$B_\mu(\xi) \geq C_0(\mu) > 0 \quad \text{for all } \xi \in \mathbb{R}, \tag{3.16}$$

where

$$0 < \mu < \min\{\mu_1, \mu_2\} \tag{3.17}$$

and $\mu_1 > 0$ and $\mu_2 > 0$ are, respectively, the unique solutions to the following equations:

$$\frac{c^2}{4D_m} - (3\varepsilon p - 2d_m) - 2\mu_1 - 2ed_m(e^{2\mu_1 r} - 1) = 0, \tag{3.18}$$

$$\frac{1}{2}(3 - e)d_m - 2\mu_2 - \frac{3}{2}ed_m(e^{2\mu_2 r} - 1) = 0, \tag{3.19}$$

and

$$C_0(\mu) := \min\{C_1(\mu), C_2(\mu)\}, \tag{3.20}$$

$$C_1(\mu) := \frac{c^2}{4D_m} - (3\varepsilon p - 2d_m) - 2\mu - 2ed_m(e^{2\mu r} - 1) > 0, \tag{3.21}$$

$$C_2(\mu) := \frac{1}{2}(3 - e)d_m - 2\mu - \frac{3}{2}ed_m(e^{2\mu r} - 1) > 0. \tag{3.22}$$

Proof. We divide this into two cases.

CASE 1 ($\xi < x_*$). Since $\phi(\xi)$ is increasing from $v_- = 0$ to $v_+ = 1/a$, we thus have $\phi'(\xi) > 0$ and $2 - a\phi(\xi) \geq 2 - av_+ = 2 - \ln e = 1$. According to (1.10), i.e. $b(\phi) = p\phi e^{-a\phi}$, we obtain

$$\frac{d}{d\xi} b'(\phi(\xi)) = -pa(2 - a\phi(\xi))e^{-a\phi(\xi)} \phi'(\xi) < 0.$$

Thus, $b'(\phi(\xi))$ is decreasing for $\xi \in (-\infty, \infty)$. This implies that

$$0 = b'(v_+) < b'(\phi(\xi)) < b'(v_-) = p. \tag{3.23}$$

Using (3.23), (1.9), and the facts that $\varepsilon p = d_m e$ from (2.3),

$$w(\xi) = e^{-\beta(\xi - x_*)}, \quad \frac{w'(\xi)}{w(\xi)} = -\beta, \quad e^{\beta(\xi - x_*)} < 1 \quad \text{for } \xi < x_*,$$

and that $cr - \alpha\beta = c(r - (\alpha/2D_m)) > 0$ from (2.7) and (1.8), which implies that $e^{-\beta(cr-\alpha\beta)} < 1$, we obtain

$$\begin{aligned}
 B_\mu(\xi) &= c\beta - D_m\beta^2 + 2d_m - 2\mu - \varepsilon \int_{\mathbb{R}} b'(\phi(\xi - y - cr))f_\alpha(y) \, dy \\
 &\quad - \varepsilon e^{2\mu r} \frac{b'(\phi(\xi))}{e^{-\beta(\xi-x_*)}} \int_{\mathbb{R}} w(\xi + y + cr)f_\alpha(y) \, dy \\
 &\geq c\beta - D_m\beta^2 + 2d_m - 2\mu - \varepsilon p \int_{\mathbb{R}} f_\alpha(y) \, dy \\
 &\quad - \frac{\varepsilon e^{2\mu r} p}{e^{\beta(\xi-x_*)}} \left[\int_{-\infty}^{x_*-\xi-cr} e^{-\beta(\xi+y+cr-x_*)} f_\alpha(y) \, dy + \int_{x_*-\xi-cr}^{\infty} f_\alpha(y) \, dy \right] \\
 &= c\beta - D_m\beta^2 + 2d_m - 2\mu - \varepsilon p \\
 &\quad - \varepsilon e^{2\mu r} p \left[\int_{-\infty}^{x_*-\xi-cr} e^{-\beta(y+cr)} f_\alpha(y) \, dy + \int_{x_*-\xi-cr}^{\infty} e^{\beta(\xi-x_*)} f_\alpha(y) \, dy \right] \\
 &\geq c\beta - D_m\beta^2 + 2d_m - 2\mu - \varepsilon d_m \\
 &\quad - \varepsilon d_m e^{2\mu r} \left[\int_{-\infty}^{x_*-\xi-cr} e^{-\beta(y+cr)} f_\alpha(y) \, dy + \int_{x_*-\xi-cr}^{\infty} f_\alpha(y) \, dy \right] \\
 &= c\beta - D_m\beta^2 + 2d_m - 2\mu - \varepsilon d_m \\
 &\quad - \varepsilon d_m e^{2\mu r} \left[\int_{-\infty}^{x_*-\xi-cr} \frac{1}{\sqrt{4\pi\alpha}} \exp\left(-\frac{y^2}{4\alpha} - \beta(y+cr)\right) \, dy \right. \\
 &\quad \left. + \int_{x_*-\xi-cr}^{\infty} f_\alpha(y) \, dy \right] \\
 &\geq c\beta - D_m\beta^2 + 2d_m - 2\mu - \varepsilon d_m \\
 &\quad - \varepsilon d_m e^{2\mu r} \left[\frac{e^{-\beta(cr-\alpha\beta)}}{\sqrt{4\pi\alpha}} \int_{-\infty}^{\infty} \exp\left(-\frac{(y+2\alpha\beta)^2}{4\alpha}\right) \, dy + \int_{-\infty}^{\infty} f_\alpha(y) \, dy \right] \\
 &= c\beta - D_m\beta^2 + (2 - e) d_m - 2\mu - \varepsilon d_m e^{2\mu r} [e^{-\beta(cr-\alpha\beta)} + 1] \\
 &\geq c\beta - D_m\beta^2 + (2 - e) d_m - 2\mu - 2\varepsilon d_m e^{2\mu r} \\
 &= \frac{c^2}{4D_m} - (3e - 2) d_m - 2\mu - 2\varepsilon d_m (e^{2\mu r} - 1) \\
 &= \frac{c^2}{4D_m} - (3\varepsilon p - 2d_m) - 2\mu - 2\varepsilon d_m (e^{2\mu r} - 1) \\
 &=: C_1(\mu) > 0 \quad \text{for suitably small } \mu > 0, \tag{3.24}
 \end{aligned}$$

where the last inequality follows from our sufficient condition (2.8), and $0 < \mu < \mu_1$, where $\mu_1 > 0$ is the unique solution of

$$\frac{c^2}{4D_m} - (3\varepsilon p - 2d_m) - 2\mu_1 - 2\varepsilon d_m (e^{2\mu_1 r} - 1) = 0,$$

so then we have

$$C_1(\mu) = \frac{c^2}{4D_m} - (3\varepsilon p - 2d_m) - 2\mu - 2\varepsilon d_m (e^{2\mu r} - 1) > 0$$

for $0 < \mu < \mu_1$.

CASE 2 ($\xi \geq x_*$). In this case, $w(\xi) = 1$. Thus,

$$B_\mu(\xi) = 2d_m - 2\mu - \varepsilon \int_{\mathbb{R}} b'(\phi(\xi - y - cr))f_\alpha(y) dy - \varepsilon e^{2\mu r} b'(\phi(\xi)) \int_{\mathbb{R}} w(\xi + y + cr)f_\alpha(y) dy. \quad (3.25)$$

Note that

$$b'(\phi) = p(1 - a\phi)e^{-a\phi} = pe^{-a\phi} - pa\phi e^{-a\phi} = pe^{-a\phi} - ab(\phi),$$

$\phi(\xi) > 0$, $\phi'(\xi) > 0$ and

$$\int_{\mathbb{R}} f_\alpha(y) dy = 1.$$

Applying equation (2.1) and the first and the second inequalities of (2.5), we can reduce the second term of the right-hand side of (3.25) as follows:

$$\begin{aligned} & -\varepsilon \int_{\mathbb{R}} b'(\phi(\xi - y - cr))f_\alpha(y) dy \\ &= -\varepsilon p \int_{\mathbb{R}} e^{-a\phi(\xi - y - cr)} f_\alpha(y) dy + a\varepsilon \int_{\mathbb{R}} b(\phi(\xi - y - cr))f_\alpha(y) dy \\ &= -\varepsilon p \int_{\mathbb{R}} e^{-a\phi(\xi - y - cr)} f_\alpha(y) dy + a[c\phi'(\xi) - D_m\phi''(\xi) + d_m\phi(\xi)] \\ &\geq -\varepsilon p \int_{\mathbb{R}} f_\alpha(y) dy + ad_m\phi(\xi) - aD_m\phi''(\xi) \\ &= -\varepsilon d_m \int_{\mathbb{R}} f_\alpha(y) dy + ad_m\phi(\xi) - aD_m\phi''(\xi) \\ &= -\varepsilon d_m + ad_m\phi(\xi) - aD_m\phi''(\xi) \\ &\geq -\varepsilon d_m + d_m - ad_m\bar{\varepsilon} - aD_m\bar{\varepsilon} \\ &= -(e - 1)d_m - (ad_m + aD_m)\bar{\varepsilon}. \end{aligned} \quad (3.26)$$

On the other hand, applying the third inequality of (2.5) and noting that $w(\xi) = 1$ and $e^{-\beta(\xi - x_*)} < 1$, we may furthermore estimate the third term of the right-hand side of (3.25) as follows:

$$\begin{aligned} & -\varepsilon e^{2\mu r} b'(\phi(\xi)) \int_{\mathbb{R}} w(\xi + y + cr)f_\alpha(y) dy \\ &= -\varepsilon p e^{2\mu r} e^{a\phi(\xi)} (1 - a\phi(\xi)) \int_{\mathbb{R}} w(\xi + y + cr)f_\alpha(y) dy \\ &= -\varepsilon d_m e^{2\mu r} e^{a\phi(\xi)} (1 - a\phi(\xi)) \int_{\mathbb{R}} w(\xi + y + cr)f_\alpha(y) dy \\ &\geq -\varepsilon d_m \bar{\varepsilon} e^{2\mu r} \left(\int_{-\infty}^{x_* - \xi - cr} + \int_{x_* - \xi - cr}^{+\infty} \right) w(\xi + y + cr)f_\alpha(y) dy \\ &\geq -\varepsilon d_m \bar{\varepsilon} e^{2\mu r} \left[\int_{-\infty}^{x_* - \xi - cr} e^{-\beta(\xi + y + cr - x_*)} f_\alpha(y) dy + \int_{x_* - \xi - cr}^{\infty} f_\alpha(y) dy \right] \end{aligned}$$

$$\begin{aligned}
 &= -ed_m \bar{\varepsilon} e^{2\mu r} \left[\frac{e^{-\beta(cr-\alpha\beta)} e^{-\beta(\xi-x_*)}}{\sqrt{4\pi\alpha}} \right. \\
 &\quad \left. \times \int_{-\infty}^{x_*-\xi-cr} e^{-(y+2\alpha\beta)^2/4\alpha} dy + \int_{x_*-\xi-cr}^{\infty} f_\alpha(y) dy \right] \\
 &\geq -ed_m \bar{\varepsilon} e^{2\mu r} \left[\frac{e^{-\beta(cr-\alpha\beta)}}{\sqrt{4\pi\alpha}} \int_{-\infty}^0 e^{-(y+2\alpha\beta)^2/4\alpha} dy + \int_{-\infty}^{\infty} f_\alpha(y) dy \right] \\
 &= -ed_m \bar{\varepsilon} e^{2\mu r} \left[\frac{1}{2} e^{-\beta(cr-\alpha\beta)} + 1 \right] \\
 &\geq -ed_m \bar{\varepsilon} e^{2\mu r} \left[\frac{1}{2} + 1 \right] \\
 &= -\frac{3}{2} ed_m \bar{\varepsilon} e^{2\mu r}. \tag{3.27}
 \end{aligned}$$

Substituting (3.26) and (3.27) into (3.25), and noting (2.4), we obtain

$$\begin{aligned}
 B_\mu(\xi) &\geq 2d_m - 2\mu - (e - 1) d_m - (ad_m + aD_m)\bar{\varepsilon} - \frac{3}{2} ed_m \bar{\varepsilon} e^{2\mu r} \\
 &= (3 - e)d_m - (ad_m + aD_m + \frac{3}{2} ed_m)\bar{\varepsilon} - 2\mu - \frac{3}{2} ed_m (e^{2\mu r} - 1) \\
 &= \frac{1}{2}(3 - e)d_m - 2\mu - \frac{3}{2} ed_m (e^{2\mu r} - 1) \\
 &=: C_2(\mu) > 0 \quad \text{for some suitably small } \mu > 0, \tag{3.28}
 \end{aligned}$$

where we select μ such that $0 < \mu < \mu_2$. Here $\mu_2 > 0$ is the unique solution to the following equation

$$\frac{1}{2}(3 - e)d_m - 2\mu_2 - \frac{3}{2} ed_m (e^{2\mu_2 r} - 1) = 0,$$

and we always have

$$C_2(\mu) = \frac{1}{2}(3 - e)d_m - 2\mu - \frac{3}{2} ed_m (e^{2\mu r} - 1) > 0$$

for $\mu < \mu_2$.

Now, we take

$$0 < \mu < \min\{\mu_1, \mu_2\}$$

then we have that both (3.24) and (3.28) hold, which leads to (3.16). The proof is complete. \square

Finally, we prove proposition 3.3.

Proof of proposition 3.3. Let $w(\xi)$ be a weight function which will be specified later. Multiplying equation (3.1) by $e^{2\mu t} w(\xi) V(t, \xi)$ for $0 < \mu \leq \mu_0$, we have

$$\begin{aligned}
 &\left\{ \frac{1}{2} e^{2\mu t} w V^2 \right\}_t + \left\{ \left(\frac{1}{2} c w V^2 - D_m w V V_\xi \right) e^{2\mu t} \right\}_\xi + D_m e^{2\mu t} w V_\xi^2 \\
 &\quad + D_m e^{2\mu t} w' V_\xi V + \left\{ -\frac{c}{2} \frac{w'}{w} + d_m - \mu \right\} e^{2\mu t} w V^2 \\
 &\quad - \varepsilon e^{2\mu t} w V \int_{\mathbb{R}} b'(\phi(\xi - y - cr)) V(t - r, \xi - y - cr) f_\alpha(y) dy \\
 &\quad = \varepsilon e^{2\mu t} w V \int_{\mathbb{R}} Q(V(t - r, \xi - y - cr)) f_\alpha(y) dy, \tag{3.29}
 \end{aligned}$$

where $w = w(\xi)$, $V = V(t, \xi)$. Using the Cauchy–Schwarz inequality $ab \leq \frac{1}{2}a^2 + \frac{1}{2}b^2$, we obtain

$$|D_m e^{2\mu t} w'(\xi) V_\xi(t, \xi) V(t, \xi)| \leq \frac{D_m}{2} e^{2\mu t} w V_\xi^2 + \frac{D_m}{2} \left(\frac{w'}{w}\right)^2 e^{2\mu t} w V^2.$$

Substituting this into (3.29) and integrating the resulting inequality over $[0, t] \times \mathbb{R}$, we have

$$\begin{aligned} & e^{2\mu t} \|V(t)\|_{L_w^2}^2 + D_m \int_0^t e^{2\mu s} \|V_\xi(s)\|_{L_w^2}^2 ds \\ & + \int_0^t \int_{\mathbb{R}} \left\{ -c \frac{w'(\xi)}{w(\xi)} + 2d_m - 2\mu - D_m \left(\frac{w'(\xi)}{w(\xi)}\right)^2 \right\} e^{2\mu s} w(\xi) V(s, \xi)^2 d\xi ds \\ & - 2\varepsilon \int_0^t \int_{\mathbb{R}} \int_{\mathbb{R}} e^{2\mu s} b'(\phi(\xi - y - cr)) w(\xi) \\ & \quad \times V(s, \xi) V(s - r, \xi - y - cr) f_\alpha(y) dy d\xi ds \\ & \leq \|v_0(0)\|_{L_w^2}^2 + 2\varepsilon \int_0^t \int_{\mathbb{R}} \int_{\mathbb{R}} e^{2\mu s} w(\xi) \\ & \quad \times V(s, \xi) Q(V(s - r, \xi - y - cr)) f_\alpha(y) dy d\xi ds. \end{aligned} \tag{3.30}$$

Now, using the change of variables $y \mapsto y$, $\xi - y - cr \mapsto \xi$, $s - r \mapsto s$, we get

$$\begin{aligned} & \int_0^t \int_{\mathbb{R}} \int_{\mathbb{R}} e^{2\mu s} w(\xi) b'(\phi(\xi - y - cr)) V^2(s - r, \xi - y - cr) f_\alpha(y) dy d\xi ds \\ & = \int_{-r}^{t-r} \int_{\mathbb{R}} \int_{\mathbb{R}} e^{2\mu(s+r)} w(\xi + y + cr) b'(\phi(\xi)) V^2(s, \xi) f_\alpha(y) dy d\xi ds \\ & = e^{2\mu r} \int_{-r}^{t-r} \int_{\mathbb{R}} e^{2\mu s} \left[\frac{b'(\phi(\xi))}{w(\xi)} \int_{\mathbb{R}} w(\xi + y + cr) f_\alpha(y) dy \right] w(\xi) V^2(s, \xi) d\xi ds. \end{aligned} \tag{3.31}$$

Once again, using the Cauchy–Schwarz inequality, and noting (3.31) and (3.23), then we can estimate the delay term on the left-hand side of (3.30) as follows:

$$\begin{aligned} & 2\varepsilon \left| \int_0^t \int_{\mathbb{R}} \int_{\mathbb{R}} e^{2\mu s} b'(\phi(\xi - y - cr)) w(\xi) V(s, \xi) V(s - r, \xi - y - cr) f_\alpha(y) dy d\xi ds \right| \\ & \leq \varepsilon \int_0^t \int_{\mathbb{R}} \int_{\mathbb{R}} e^{2\mu s} w(\xi) b'(\phi(\xi - y - cr)) \\ & \quad \times [V^2(s, \xi) + V^2(s - r, \xi - y - cr)] f_\alpha(y) dy d\xi ds \\ & = \varepsilon \int_0^t \int_{\mathbb{R}} e^{2\mu s} w(\xi) \left[\int_{\mathbb{R}} b'(\phi(\xi - y - cr)) f_\alpha(y) dy \right] V^2(s, \xi) d\xi ds \\ & \quad + \varepsilon e^{2\mu r} \int_{-r}^{t-r} \int_{\mathbb{R}} e^{2\mu s} \left[\frac{b'(\phi(\xi))}{w(\xi)} \int_{\mathbb{R}} w(\xi + y + cr) f_\alpha(y) dy \right] w(\xi) V^2(s, \xi) d\xi ds \\ & \leq \varepsilon \int_0^t \int_{\mathbb{R}} e^{2\mu s} \left[\int_{\mathbb{R}} b'(\phi(\xi - y - cr)) f_\alpha(y) dy \right] w(\xi) V^2(s, \xi) d\xi ds \end{aligned}$$

$$\begin{aligned}
 &+ \varepsilon e^{2\mu r} \int_0^t \int_{\mathbb{R}} e^{2\mu s} \left[\frac{b'(\phi(\xi))}{w(\xi)} \int_{\mathbb{R}} w(\xi + y + cr) f_\alpha(y) dy \right] w(\xi) V^2(s, \xi) d\xi ds \\
 &+ \varepsilon e^{2\mu r} \int_{-r}^0 \int_{\mathbb{R}} e^{2\mu s} \left[\frac{b'(\phi(\xi))}{w(\xi)} \int_{\mathbb{R}} w(\xi + y + cr) f_\alpha(y) dy \right] w(\xi) V_0^2(s, \xi) d\xi ds.
 \end{aligned}
 \tag{3.32}$$

Substituting (3.32) into (3.30) yields

$$\begin{aligned}
 &e^{2\mu t} \|V(t)\|_{L_w^2}^2 + D_m \int_0^t e^{2\mu s} \|V_\xi(s)\|_{L_w^2}^2 ds + \int_0^t \int_{-\infty}^\infty B_\mu(\xi) e^{2\mu s} w(\xi) V^2(s, \xi) d\xi ds \\
 &\leq \|V_0(0)\|_{L_w^2}^2 \\
 &\quad + \varepsilon e^{2\mu r} \int_{-r}^0 \int_{\mathbb{R}} e^{2\mu s} \left[\frac{b'(\phi(\xi))}{w(\xi)} \int_{\mathbb{R}} w(\xi + y + cr) f_\alpha(y) dy \right] w(\xi) V_0^2(s, \xi) d\xi ds \\
 &\quad + 2\varepsilon \int_0^t \int_{\mathbb{R}} \int_{\mathbb{R}} w(\xi) e^{2\mu s} V(s, \xi) Q(V(s - r, \xi - y - cr)) f_\alpha(y) dy d\xi ds,
 \end{aligned}
 \tag{3.33}$$

where $B_\mu(\xi)$ is defined in (3.15). We need to select a suitable weight function, $w(\xi)$, so that $B_\mu(\xi) > 0$ for all $\xi \in \mathbb{R}$. The choice of $w(\xi)$ is, of course, not unique. One possibility is

$$w(\xi) = \begin{cases} e^{-\beta(\xi - x_*)}, & \xi < x_*, \\ 1, & \xi \geq x_*, \end{cases}$$

as in (2.6) with $\beta = c/2D_m$. According to lemma 3.6, $B_\mu(\xi) \geq C_0(\mu) > 0$ for $0 < \mu \leq \min\{\mu_1, \mu_2\}$, where the positive constants μ_1, μ_2 and $C_0(\mu)$ are defined in (3.18), (3.19) and (3.20), respectively. Then, by using (3.23) and

$$e^{2\mu r} \int_{\mathbb{R}} \frac{w(\xi + y + cr)}{w(\xi)} f_\alpha(y) dy \leq C \quad \text{for all } \xi \in \mathbb{R},
 \tag{3.34}$$

which can be proved analogously to lemma 3.6, we can reduce (3.33) to

$$\begin{aligned}
 &e^{2\mu t} \|V(t)\|_{L_w^2}^2 + D_m \int_0^t e^{2\mu s} \|V_\xi(s)\|_{L_w^2}^2 ds + C_0(\mu) \int_0^t e^{2\mu s} \|V(s)\|_{L_w^2}^2 ds \\
 &\leq \|V_0(0)\|_{L_w^2}^2 + \varepsilon C \int_{-r}^0 \|V_0(s)\|_{L_w^2}^2 ds \\
 &\quad + 2\varepsilon \int_0^t \int_{\mathbb{R}} \int_{\mathbb{R}} e^{2\mu s} w(\xi) V(s, \xi) Q(V(s - r, \xi - y - cr)) f_\alpha(y) dy d\xi ds.
 \end{aligned}
 \tag{3.35}$$

Next, we will estimate the nonlinear term on the right-hand side of (3.35). From (3.2), by Taylor’s formula, we first have $Q(V) = O(V^2)$ as $V \rightarrow 0$, i.e.

$$|Q(V(t - r, \xi - y - cr))| \sim C_5 |V(t - r, \xi - y - cr)|^2
 \tag{3.36}$$

for some positive constant C_5 . Then, by the standard Sobolev embedding inequality $H^1(\mathbb{R}) \hookrightarrow C^0(\mathbb{R})$ and the modified embedding inequality $H_w^1(\mathbb{R}) \hookrightarrow H^1(\mathbb{R})$ for

$w(\xi) > 0$ defined as in (2.6) (see [13]), we obtain

$$|V(t, \xi)| \leq \sup_{\xi \in \mathbb{R}} |V(t, \xi)| \leq C_6 \|V(t, \cdot)\|_{H^1} \leq C_6 \|V(t, \cdot)\|_{H^1_w} \leq C_6 M(t), \tag{3.37}$$

where $C_6 > 0$ is the embedding constant. Applying (3.34), (3.36) and (3.37) and making the change of variables $y \mapsto y, \xi - y - cr \mapsto \xi, s - cr \mapsto s$, we then obtain

$$\begin{aligned} & \int_0^t \int_{\mathbb{R}} \int_{\mathbb{R}} e^{2\mu s} w(\xi) V(s, \xi) Q(V(s-r, \xi-y-cr)) f_\alpha(y) dy d\xi ds \\ & \leq C_7 M(t) \int_0^t \int_{\mathbb{R}} \int_{\mathbb{R}} e^{2\mu s} w(\xi) |V(s-r, \xi-y-cr)|^2 f_\alpha(y) dy d\xi ds \\ & = C_7 M(t) \int_{-r}^{t-r} \int_{\mathbb{R}} \int_{\mathbb{R}} e^{2\mu(s+r)} w(\xi+y+cr) |V(s, \xi)|^2 f_\alpha(y) dy d\xi ds \\ & \leq C_7 M(t) \left\{ \int_0^t \int_{\mathbb{R}} e^{2\mu(s+r)} w(\xi) |V(s, \xi)|^2 \left(\int_{\mathbb{R}} \frac{w(\xi+y+cr)}{w(\xi)} f_\alpha(y) dy \right) d\xi ds \right. \\ & \quad \left. + \int_{-r}^0 \int_{\mathbb{R}} e^{2\mu(s+r)} w(\xi) |V_0(s, \xi)|^2 \left(\int_{\mathbb{R}} \frac{w(\xi+y+cr)}{w(\xi)} f_\alpha(y) dy \right) d\xi ds \right\} \\ & \leq C_8 M(t) \left\{ \int_0^t e^{2\mu s} \|V(s)\|_{L^2_w}^2 ds + \int_{-r}^0 e^{2\mu s} \|V_0(s)\|_{L^2_w}^2 ds \right\} \end{aligned} \tag{3.38}$$

for some positive constants C_7 and C_8 . Substituting (3.38) into (3.35), we finally obtain

$$\begin{aligned} e^{2\mu t} \|V(t)\|_{L^2_w}^2 + D_m \int_0^t e^{2\mu s} \|V_\xi(s)\|_{L^2_w}^2 ds + [C_0(\mu) - C_8 M(t)] \int_0^t e^{2\mu s} \|V(s)\|_{L^2_w}^2 ds \\ \leq \|V_0(0)\|_{L^2_w}^2 + C_9 [1 + M(t)] \int_{-r}^0 \|V_0(s)\|_{L^2_w}^2 ds \end{aligned} \tag{3.39}$$

for some constant $C_9 > 0$.

One can find a positive constant δ_2 such that

$$C_0(\mu) - C_8 \delta_2 > 0 \text{ or, equivalently, } \delta_2 < \frac{C_0(\mu)}{C_8}. \tag{3.40}$$

Clearly, δ_2 depends only on the coefficients $D_m, d_m, \varepsilon, p, a, r$ and the wave speed c , because μ depends on these parameters (see (3.17)–(3.19)). When $M(T) \leq \delta_2$, i.e.

$$C_0(\mu) - C_8 M(T) \geq C_0(\mu) - C_8 \delta_2 > 0,$$

we have

$$e^{2\mu t} \|V(t)\|_{L^2_w}^2 + D_m \int_0^t e^{2\mu s} \|V_\xi(s)\|_{L^2_w}^2 ds \leq \|V_0(0)\|_{L^2_w}^2 + C_{10} \int_{-r}^0 \|V_0(s)\|_{L^2_w}^2 ds \tag{3.41}$$

for some positive constant C_{10} .

Similarly, by differentiating (3.1) with respect to ξ , multiplying the resultant equation by $e^{2\mu t} w(\xi) V_\xi(t, \xi)$, and then integrating it over $[0, t] \times \mathbb{R}$ for $t \leq T$, using

the basic energy estimate (3.41), we finally have

$$e^{2\mu t} \|V_\xi(t)\|_{L^2_w}^2 + D_m \int_0^t e^{2\mu s} \|V_{\xi\xi}(s)\|_{L^2_w}^2 ds \leq C_{11} \left(\|V_0(0)\|_{H^1_w}^2 + \int_{-r}^0 \|V_0(s)\|_{H^1_w}^2 ds \right) \tag{3.42}$$

for some positive constant C_{11} , provided that $M(T) \leq \delta_2$. We omit the detail.

Combining (3.41) and (3.42), we obtain

$$e^{2\mu t} \|V_\xi(t)\|_{H^1_w}^2 + D_m \int_0^t e^{2\mu s} \|V_\xi(s)\|_{H^1_w}^2 ds \leq C_1 \left(\|V_0(0)\|_{H^1_w}^2 + \int_{-r}^0 \|V_0(s)\|_{H^1_w}^2 ds \right) \tag{3.43}$$

for some absolute constant $C_1 = \max\{1 + C_{11}, C_{10} + C_{11}\} > 0$ that is independent of T and $V(t, x)$. Finally, from (3.43), we automatically reach

$$\|V(t)\|_{H^1_w}^2 \leq C_1 \left(\|V_0(0)\|_{H^1_w}^2 + \int_{-r}^0 \|V_0(s)\|_{H^1_w}^2 ds \right) e^{-2\mu t}, \quad 0 \leq t \leq T.$$

The proof is complete. □

Acknowledgments

The authors thank Professors Martin Gander and Georg Schmidt for valuable discussion, and the referee for helpful comments.

References

- 1 J. Bricmont and A. Kupiainen. Stability of moving fronts in the Ginzburg–Landau equations. *Commun. Math. Phys.* **159** (1994), 287–318.
- 2 J.-P. Eckmann and C. E. Wayne. The nonlinear stability of front solutions for parabolic partial differential equations. *Commun. Math. Phys.* **161** (1994), 323–334.
- 3 T. Gallay. Local stability of critical fronts in nonlinear parabolic partial differential equations. *Nonlinearity* **7** (1994), 741–764.
- 4 S. A. Gourley. Travelling front solutions of a nonlocal Fisher equation. *J. Math. Biol.* **41** (2000), 272–284.
- 5 S. A. Gourley and S. Ruan. Dynamics of the diffusive Nicholson’s blowflies equation with distributed delay. *Proc. R. Soc. Edinb. A* **130** (2000), 1275–1291.
- 6 A. M. Il’in and O. A. Oleinik. Asymptotic behavior of solutions of the Cauchy problem for certain quasilinear equations for large time. *Mat. Sb.* **51** (1960), 191–216. (In Russian.)
- 7 S. Kawashima and A. Matsumura. Stability of shock profiles in viscoelasticity with non-convex constitutive relations. *Commun. Pure Appl. Math.* **47** (1994), 1547–1569.
- 8 K. Kirchgässner. On the nonlinear dynamics of travelling fronts. *J. Diff. Eqns* **96** (1992), 256–278.
- 9 D. Liang and J. Wu. Travelling waves and numerical approximations in a reaction–advection–diffusion equation with nonlocal delayed effects. *J. Nonlin. Sci.* **13** (2003), 289–310.
- 10 A. Matsumura. Asymptotic stability of traveling wave solutions for the one-dimensional viscous conservation laws. *Sugaku Expos.* **11** (1998), 215–234.
- 11 A. Matsumura and M. Mei. Convergence to travelling fronts of solutions of the p -system with viscosity in the presence of a boundary. *Arch. Ration. Mech. Analysis* **146** (1999), 1–22.
- 12 A. Matsumura and K. Nishihara. Asymptotic stability of traveling waves for scalar viscous conservation laws with non-convex nonlinearity. *Commun. Math. Phys.* **165** (1994), 83–96.
- 13 M. Mei and K. Nishihara. Nonlinear stability of travelling waves for one-dimensional viscoelastic materials with non-convex nonlinearity. *Tokyo J. Math.* **20** (1997), 241–264.

- 14 M. Mei and B. Rubino. Convergence to traveling waves with decay rates for solutions of the initial boundary problem to a nonconvex relaxation model. *J. Diff. Eqns* **158** (1999), 138–185.
- 15 M. Mei, J. W.-H. So, M. Y. Li and S. S. P. Shen. Asymptotic stability of traveling waves for Nicholson’s blowflies equation with diffusion. *Proc. R. Soc. Edinb. A* **134** (2004), 579–594.
- 16 J. A. J. Metz and O. Diekmann (eds). *The dynamics of physiologically structured populations* (Springer, 1986).
- 17 T. Ogiwara and H. Matano. Monotonicity and convergence results in order-preserving systems in the presence of symmetry. *Discrete Contin. Dynam. Syst.* **5** (1999), 1–34.
- 18 J.-M. Roquejoffre. Stability of travelling fronts in a model for flame propagation. II. Nonlinear stability. *Arch. Ration. Mech. Analysis* **117** (1992), 119–153.
- 19 J.-M. Roquejoffre. Convergence to travelling waves for solutions of a class of semilinear parabolic equations. *J. Diff. Eqns* **108** (1994), 262–295.
- 20 D. H. Sattinger. On the stability of waves of nonlinear parabolic systems. *Adv. Math.* **22** (1976), 312–355.
- 21 K. W. Schaaf. Asymptotic behavior and traveling wave solutions for parabolic functional differential equations. *Trans. Am. Math. Soc.* **302** (1987), 587–615.
- 22 H. L. Smith and H. R. Thieme. Strongly order preserving semiflows generated by functional differential equations. *J. Diff. Eqns* **93** (1991), 332–363.
- 23 H. L. Smith and X.-Q. Zhao. Global asymptotic stability of traveling waves in delayed reaction–diffusion equations. *SIAM J. Math. Analysis* **31** (2000), 514–534.
- 24 J. W.-H. So and Y. Yang. Dirichlet problem for the diffusive Nicholson’s blowflies equation. *J. Diff. Eqns* **150** (1998), 317–348.
- 25 J. W.-H. So, J. Wu and Y. Yang. Numerical steady state and Hopf bifurcation analysis on the diffusive Nicholson’s blowflies equation. *Appl. Math. Computat.* **111** (2000), 53–69.
- 26 J. W.-H. So, J. Wu and X. Zou. A reaction–diffusion model for a single species with age structure. I. Travelling wavefronts on unbounded domains. *Proc. R. Soc. Lond. A* **457** (2001), 1841–1853.
- 27 H. Thieme and X.-Q. Zhao. Asymptotic speeds of spread and traveling waves for integral equations and delayed reaction–diffusion models. *J. Diff. Eqns* **195** (2003), 430–470.
- 28 A. I. Volpert, Vi. A. Volpert and Vl. A. Volpert. *Traveling wave solutions of parabolic systems*. Translations of Mathematical Monographs, vol. 140 (Providence, RI: American Mathematical Society, 1994).
- 29 J.-H. Wu. *Theory and applications of partial functional differential equations*, Applied Mathematical Sciences, vol. 119 (Springer, 1996).
- 30 J. Xin. Existence and stability of traveling waves in periodic media governed by a bistable nonlinearity. *J. Dynam. Diff. Eqns* **3** (1991), 541–573.
- 31 J. Xin. Stability of traveling waves in a solute transport equation. *J. Diff. Eqns* **135** (1997), 269–298.
- 32 J. Xin. Front propagation in heterogeneous media. *SIAM Rev.* **42** (2000), 161–230.
- 33 X.-Q. Zhao. *Dynamical systems in population biology*. Communications in Mathematical Physics, vol. 16 (Springer, 2003).

(Issued 13 June 2008)