

The effect of winter length on duration of dormancy and survival of specialized herbivorous *Rhagoletis* fruit flies from high elevation environments with acyclic climatic variability

J. Rull^{1*}, E. Tadeo², R. Lasa² and M. Aluja²

¹PROIMI Biotecnología-CONICET, LIEMEN-División Control Biológico de Plagas, Av. Belgrano y Pje. Caseros, T4001MVB San Miguel de Tucumán, Tucumán, Argentina; ²Red de Manejo Biorracional de Plagas y Vectores, Instituto de Ecología, A.C., Xalapa, Veracruz 91070, México

Abstract

Dormancy can be defined as a state of suppressed development allowing insects to cope with adverse conditions and plant phenology. Among specialized herbivorous insects exploiting seasonal resources, diapause frequently evolves as a strategy to adjust to predictable plant seasonal cycles. To cope with acyclic and unpredictable climatic events, it has been found for some insects that a proportion of the population undergoes prolonged dormancy. We compared the response of three species in the *Rhagoletis cingulata* species group exploiting plants differing in fruiting phenology from environments varying in frequency and timing of acyclic climatic catastrophic events (frost during flowering and fruit set) and varying also in the time of the onset of the rainy season. Small proportions (<2%) of Central and Northeastern Mexico *R. cingulata* and *Rhagoletis turpiniae*, and a few associated adult parasitoids, emerged without becoming dormant. Pupae exposed to 5°C for different periods of time (0–16 weeks) became dormant and emerged as adults in proportions <30%, compared with other studied species in the group from the USA (up to 80%). Large proportions (19–55%) of exposed pupae entered prolonged dormancy (>10 months), and large proportions of pupae died without emerging as adults. The number of days elapsed from the end of artificial winter and adult eclosion was longer for *R. cingulata* exploiting late fruiting *Prunus serotina* in Northeastern Mexico than for flies recovered from earlier fruiting plants in the central Altiplano. *Rhagoletis turpiniae* and northeastern *R. cingulata* pupae suffered high proportions of parasitism. Large proportions of *R. cingulata* from central Mexico engaging in prolonged dormancy may be explained by the fact that flowering and fruit set for its host, *P. serotina* var *capuli*, driven by the timing of maximum precipitation, matches a period of highest probability of frost often resulting in large areas with fruitless trees at unpredictable time intervals. As a consequence of differences in host plant fruiting phenology, central and northeastern Mexican *R. cingulata* were found to be allochronically isolated. Prolonged dormancy may have resulted in escape from parasitism.

Keywords: diapause, allochronic isolation, host plant phenology, Tephritidae, prolonged dormancy

(Accepted 25 August 2017; First published online 19 September 2017)

*Author for correspondence:
Tel: +54 381 434 4888
Fax: +54 381 434 4887
Email: pomonella@gmail.com

Introduction

Phenology is a major structuring element in all areas of ecology and evolution (Forrest & Miller-Rushing, 2010). In the case of specialized herbivorous insects, seasonal synchronization with suitable stages of host plants is critical for survival, reproduction, and population growth (Tauber *et al.*, 1986; Bale *et al.*, 2002). Evolution of different herbivore life-history strategies for host–plant synchronization is strongly influenced by predictability and duration of seasonal fluctuations (Tauber *et al.*, 1986). Some changes recur on a regular cyclic basis, while others do not provide advance warning allowing physiological or behavioral responses.

Dormancy, defined as a state of suppressed development, encompasses both quiescence and diapause (Košťál, 2006), and allows herbivorous insects to cope with adverse conditions and plant phenology. Diapause is an endogenously mediated interruption of morphogenesis that precedes the advent of adverse conditions and whose end does not necessarily coincide with the end of adversity (Košťál, 2006). While diapause frequently evolves among insects exploiting plants that occur regularly and predictably at a particular time of the year, to cope with acyclic, yet frequent catastrophic events, insects can exhibit spreading of adult emergence over several years through prolonged diapause (Menu, 1993; Menu & Debouzie, 1993). Under such conditions, no single phenotype is well suited for all conditions, and a bet-hedging strategy consist in producing a variety of phenotypes as a way of managing the risk of widespread events such as masting, drought, frost, and floods (Donaldson-Matasci *et al.*, 2013).

All studied species of Tephritid fruit flies in the genus *Rhagoletis* are specialized stenophagous herbivores with adults that meet and mate on the host fruit where the larvae develop after egg laying (Prokopy & Papaj, 2000). Most species are univoltine, exploit groups of plants with discrete yearly phenology, and synchronize adult emergence with fruiting through pupal diapause (Bush, 1966). In *Rhagoletis* the majority of individuals in the population emerge as adults the following season, a small proportion forgoes diapause and emerges within a month of pupation (Dambroski & Feder, 2007; Rull *et al.*, 2016), and other individuals engage after multi-year (2–5 years) diapause (Boyce, 1931; Dean, 1973; Boller & Prokopy, 1976; Moraiti *et al.*, 2012; Neven & Yee, 2017). For *Rhagoletis pomonella*, the most studied species in the genus, it has been found that diapause regulation is a complex gene \times environment interaction and diapause depth and timing of diapause termination or rates of post-diapause development are under differential selection and genetically uncoupled (Feder & Filchak, 1999; Dambroski & Feder, 2007). It has also been found that diapause variation evolved in central Mexican *R. pomonella* populations associated to an inversion polymorphism that introgressed into US populations (Feder *et al.*, 2003; 2005). Subsequently, inversions coupled with host discrimination traits and ecologically different plant niches fostered shifts to novel plants and speciation of the *R. pomonella* species group in the USA (Xie *et al.*, 2007; 2008). Importantly, each sibling species in the *R. pomonella* group infests a different, non-overlapping set of host plants (Bush, 1966; Berlocher, 2000). These species differ in diapause traits differentially adapting them to variation in host fruiting phenology (Berlocher, 2000).

In the case of *Rhagoletis* associated to plants in the genus *Prunus*, Baker & Miller (1978) documented differences in post-diapause eclosion among four European populations of

Rhagoletis cerasi, with an Italian population displaying longer periods in the pupal stage than three populations from colder climates. Moraiti *et al.* (2014), when comparing Greek and German populations, concluded that diapause duration (annual dormancy) is an adaptive response to local climatic conditions produced by phenological differences between local host plant cultivars, while plastic responses to inter annual climatic variability resulted in two types of long life cycles: prolonged and facultative dormancy as response to insufficient and extended chilling, respectively. Long life cycles are expressed as a part of dormancy bet-hedging strategies of *R. cerasi* populations. Similarly, *Rhagoletis indifferens* pupae exposed to constant temperature (26°C) and a 20-week long artificial 5°C winter produced large proportions (56–75%) of live unclosed individuals after 40 weeks (Neven & Yee, 2017). As for *R. cerasi*, these unclosed pupae were most likely engaging in prolonged dormancy.

The *cingulata* species group in the genus *Rhagoletis* is currently composed of five species infesting plants in three families, among which *R. cingulata* and *R. indifferens* are specialized on plants in the genus *Prunus* (Smith & Bush, 2000). Due to their pest status of cherries in the USA, some aspects related to dormancy and phenology have been examined for *R. indifferens* (Frick *et al.*, 1954; Van Kirk & AliNiazee, 1981; 1982; Stark & AliNiazee, 1982; Dowell & Penrose, 2012; Neven & Yee, 2017) and *R. cingulata* (Teixeira *et al.*, 2007). In the case of *R. cingulata*, two Mexican populations exploiting different varieties of black cherries, *Prunus serotina* var *capuli* and *P. serotina* var *virens*, differing in fruiting phenology, have been discovered (Padilla, 1964; Rull *et al.*, 2011). Mexican ‘populations’ exhibit some degree of reproductive isolation with eastern US populations and appear to be allochronically isolated (Tadeo *et al.*, 2015). *Rhagoletis turpiniae* is a species of no economic importance associated to plants in the genus *Turpinia* in central Mexico that has been included in a comparative study examining dormancy among sympatric species of *Rhagoletis* exploiting different host plants (Rull *et al.*, 2016). The remaining species of the group, *Rhagoletis cionanthi* and *Rhagoletis osmanthi*, are very poorly known (Bush, 1966).

Here, we compare the effect of winter length on survival and duration of dormancy among two *R. cingulata* diverging populations that exploit two black cherry varieties differing in fruiting phenology and exposed to different, regular, and catastrophic (frost) environmental fluctuations in highland semi-arid habitats, and *R. turpiniae*, a recently derived species in the *cingulata* species group that exploits host plants in moister, warmer, and more stable environments. Results of this comparative study, contrasted with existing literature on the *pomonella* species group and *R. cerasi*, may contribute in establishing the existence and extent of allochronic isolation between diverging *R. cingulata* populations, the effect of host plant phenology on duration of dormancy, and the effect of catastrophic unpredictable environmental fluctuations on the frequency of individuals undergoing prolonged dormancy. Overall, we aimed to gain insights on how insects inhabiting high elevation temperate climates in the Neotropics adapt to cope with environmental fluctuation.

Materials and methods

Biological material

Infested *P. serotina* var *capuli* was collected under the canopy of several trees in the locality of Humantla, Tlaxcala



Fig. 1. Geographical location of infested fruit collection sites for central Mexican (Huamantla, Tlaxcala) and northeastern (Los Lirios, Coahuila) *Rhagoletis cingulata* and *Rhagoletis turpiniae* (Xalapa, Veracruz).

(19.18.46N, 97.55.18W; 2497 m) (fig. 1) on 13 August 2015 at the edge of the Mexican Altiplano Central. The area is characterized by dry mild weather with an average yearly temperature of 15°C, an average rainfall of 682 mm concentrated from May to September, and a high probability (80%) of frost from early November to late February (Ortiz *et al.*, 1992). *Prunus serotina* var *capuli* flowers from January to March (McVaugh, 1951).

Prunus serotina var *virens* was collected at the locality of Los Lirios (25.23.31N; 100.36.40W; 2244 m), Coahuila on 28 August 2015, on the eastern side of the Sierra Madre Oriental in northeastern Mexico. The area is characterized by dry mild weather with an average mean temperature of 18.2°C and 375 mm of annual rainfall occurring from June to September (<http://es.climate-data.org/>), the probability of frosting events from November to February is high (>50%) in southern Coahuila (Martínez & Ruiz, 2005). *Prunus serotina* var *virens* begins flowering in March, with late spring frost damaging flowers and causing large numbers of newly set fruit to fall (Hough, 1965).

Infested *Turpinia insignis* fruits were collected on the grounds of the Instituto de Ecología A.C. (INECOL), and the Clavijero Botanical garden (CBG), on the old Xalapa-Coatepec road (19.30.45, N: 96.56.40W; 1334 m) from 10–17 August 2015, on the eastern side of the Sierra Madre Oriental in the state of Veracruz in central Mexico. Xalapa has an average mean temperature of 18.2°C, and 1587 mm of annual precipitation peaking from May to October and is virtually free of frost. *Turpinia insignis* can be found from 600 to 2000 m, on moist eastern side of the Sierra Madre Oriental, and flowers from January to April (dry season) (Sosa, 1988).

In all cases, fruits were immediately taken to the laboratory and handled as described by Rull *et al.* (2006) to recover pupae. Pupae were removed from trays every other day, placed in transparent plastic 200 ml cups containing 2 cm of moist vermiculite (such pupae were in consequence 0–2 days old), capped with a mesh covered lid, and taken to a laboratory

under controlled environmental conditions at 24°C, 65% RH, and 13/11 h L/D photoperiod. As a consequence, regardless of collection date, all pupae were aged 1–2 days when submitted to controlled temperature treatments. In all cases, vermiculite was moistened with a 2% (wt./vol.) sodium benzoate solution twice a week during the entire experimental period. Pupae remained under pre-winter conditions for exactly 40 days and were transferred to a conventional refrigerator at 5 ± 1°C for durations described below. The duration of the pre-winter period was established based on criteria detailed by Rull *et al.* (2016) to record proportions of non-dormant flies. During this pre-winter period, date of emergence and sex of individual flies and parasitoids were recorded. Adults recovered <40 days after pupation were considered to be non-dormant.

Temperature regimes

In the case of *R. cingulata* populations from Tlaxcala, 1400 pupae were recovered and separated in individual 200 ml plastic containers (cups) in sets of 100 pupae per cup. Cups were subjected to periods of different lengths at 5°C. Two cups were left at constant 24°C temperature for the entire experimental period and the remaining sets (three cups per temperature regime) were placed at artificial 5°C ‘winter’ periods for 4, 8, 10, and 16 weeks after which they were returned to laboratory conditions as described for the pre-winter period. Winter length treatments were chosen based on previous results with Mexican populations of four species of *Rhagoletis* (Rull *et al.*, 2016). The low-temperature periods (4–16 weeks) selected for this experiment encompass regimes that have resulted in maximum percent post-dormancy adult eclosion, while the longest cold period (16 weeks) produced statistically similar percent eclosion than longer ‘winter’ periods (18–20 weeks). Adult fly and parasitoid emergence were recorded for all treatments until emergence of all adults (30 June 2016). In sum, except for the constant temperature treatment

(two cups, 200 pupae) there were 300 *R. cingulata* pupae from Tlaxcala (placed in three cups with 100 pupae each) for each one of the 10, 5°C 'winter', periods.

For *R. cingulata* from Coahuila, a total of 812 pupae were recovered and subjected to controlled temperature regimes as follows: two cups (one containing 100 and one 112 pupae) were held at constant temperature (24°C), two 100-pupae cups were placed at 5°C for 4 and 8 weeks after the 40-day pre-winter period, and one 100-pupae cup was placed at 5°C for 10 and 16 weeks after the 40-day pre-winter period.

For *R. turpiniae* from Veracruz, a total of 1372 pupae were recovered and subjected to controlled temperature regimes as follows: three cups (two containing 124 and one with 100 pupae) were subjected to constant temperature (24°C) for the entire observational period. Two 100-pupae cups were subjected to a 4-week 5°C winter period, three cups with 100 pupae were subjected to a 8-week 5°C winter period, three cups (two with 100 pupae and one with 194 pupae) were subjected to a 10-week 5°C winter period, and two cups (one with 80 pupae and one with 74) were subjected to a 16-week 5°C winter period.

At the end of the adult emergence recording period (30 June 2016), unclosed pupae for all treatments and replicates were inspected under a dissecting microscope to establish the proportion of pupae having entered a 'long' diapause (live pupae) and the proportion of unclosed pupae having suffered mortality.

Statistical analyses

Cumulative percent post-winter adult eclosion, the proportion of pupae undergoing long diapause (>10 months), and pupal mortality after 10 months, were compared among treatments within each species by means of χ^2 tests of observed proportions vs. expected proportions according to a uniform distribution of percent eclosion, survival, and mortality respectively. Overall percent adult eclosion after dormancy, the overall proportion of pupae undergoing long diapause, percent pupal mortality, and overall parasitism were compared among species (across sites) by means of χ^2 tests of observed vs. expected proportions according to a uniform distribution. The length of time from pupation to adult eclosion (corrected for winter length) was compared among treatments and sex within each fly species, and between male and female flies (pooled together) of each species and their parasitoids by means of general linear models (GLM). The length of time from pupation to adult eclosion (corrected for winter length) was compared across populations/species by building a GLM with species (site), sex, and treatment as dependent variables. In the case of across site comparisons, only the 8-, 10-, and 16-week artificial winter treatments were included in the model, given that no adult eclosion was recorded for pupae of all populations/species when exposed to constant temperatures and short winters (4 weeks).

Results

Non-dormant flies

A very small proportion (1.4 ± 1.1%) of central Altiplano (Tlaxcala) *R. cingulata* adults emerged during the 40-day pre-winter period without becoming dormant (13 males and six females). The developmental period for such adults lasted from pupation to adult eclosion an average (±SD) of 35.2 ± 1.9 days

for males, and 36.2 ± 1.5 days for females. Only one adult of a braconid parasitoid emerged without becoming dormant. The developmental period for this non-dormant adult lasted 35 days.

For northeastern (Coahuila) *R. cingulata*, only one braconid parasitoid emerged without becoming dormant and did so 34 days after pupation.

In the case of *R. turpiniae* (Veracruz) 1.82 ± 1.86% adults emerged during the 40-day pre-winter period without becoming dormant (ten males and 14 females). The developmental period for such adults lasted from pupation to adult eclosion an average (±SD) of 39.4 ± 1.4 days for males, and 39.4 ± 2.6 days for females. Thirty-five adult braconid parasitoids emerged without becoming dormant, and did so after an average (±SD) of 38.1 ± 3.8 days after pupation.

Dormant flies

Percent eclosion according to winter length

There were significant differences in the proportion Tlaxcala *R. cingulata* adults emerging after artificial winters of different duration ($\chi^2 = 99.76$; $df = 4$; $P < 0.01$). Pupae exposed to winter periods of 16 weeks eclosed in greater proportions (34.6%) than pupae exposed to shorter winter periods or constant temperatures (fig. 2a). A large proportion of pupae entered prolonged dormancy periods (>10 months). Pupae exposed to artificial winters of <16 weeks or constant temperatures remained dormant for more than 10 months in greater proportion than those exposed to 5°C for 16 weeks ($\chi^2 = 43.14$; $df = 4$; $P < 0.01$). Pupae exposed to 16-week artificial winters experienced greater mortality than those exposed to artificial winters of shorter duration or constant temperatures ($\chi^2 = 176.4$; $df = 4$; $P < 0.01$).

In the case of northeastern *R. cingulata* from Coahuila, there were significant differences in the proportion of adults emerging after artificial winters of different duration ($\chi^2 = 20.97$; $df = 4$; $P < 0.01$). Pupae exposed to 5°C for 4, 8, and 10 weeks emerged in greater proportions than expected from a uniform distribution of adult emergence (fig. 2b). There were also significant differences in the proportion of pupae entering prolonged diapause ($\chi^2 = 90.11$; $df = 4$; $P < 0.01$). Pupae exposed to constant temperatures and a 4-week artificial winter entered prolonged diapause in greater proportions than expected from a uniform distribution. Pupae exposed to artificial winter periods of 8, 10, and 16 weeks suffered greater mortality than expected from a uniform distribution ($\chi^2 = 16.87$; $df = 4$; $P < 0.01$).

For *R. turpiniae* (Veracruz), pupae exposed to 10- and 16-week artificial winter periods emerged as adults in greater proportion than expected from a uniform distribution ($\chi^2 = 29.92$; $df = 4$; $P < 0.01$) (fig. 2c). Pupae exposed to constant temperatures and 4 weeks of artificial winter entered prolonged diapause in greater proportion than expected from a uniform distribution ($\chi^2 = 44.37$; $df = 4$; $P < 0.01$). Pupae exposed to winters of 10 and 16 weeks suffered mortality in greater proportion than expected from a uniform distribution ($\chi^2 = 11.32$; $df = 4$; $P = 0.023$).

Overall mean proportion (±SD) of adults eclosing after dormancy, pupae undergoing long dormancy, pupal mortality, and parasitism are summarized in table 1. *Rhagoletis cingulata* pupae from Coahuila emerged as adults in greater proportions than expected from a uniform distribution ($\chi^2 = 12.36$; $df = 2$; $P = 0.002$). A greater proportion of *R. cingulata* pupae from Tlaxcala underwent long diapause than expected from a uniform distribution ($\chi^2 = 17.61$; $df = 2$; $P < 0.001$). There were no




Fig. 2. Proportions of: adult eclosion (black bars), live pupae undergoing prolonged dormancy (dark gray bars), and dead pupae (light gray bars), for two populations of *Rhagoletis cingulata* from central (Tlaxcala) and northeastern (Coahuila) Mexico exploiting early (*Prunus serotina* var *capuli*) and late (*P. serotina* var *capuli*) fruiting black cherries and *Rhagoletis turpiniae* (Veracruz). Pupae were exposed to 5°C artificial winters of different lengths (0, 4, 8, 10 and 16 weeks).

statistical differences in percent mortality among species/populations when compared with a uniform distribution ($\chi^2 = 4.96$; $df = 2$; $P = 0.083$). *Rhagoletis cingulata* pupae from Coahuila and *R. turpiniae* pupae from Veracruz suffered a greater proportion of parasitism than that expected from a uniform distribution ($\chi^2 = 38.00$; $df = 2$; $P < 0.001$).

Duration of post-winter dormancy

Rhagoletis cingulata pupae from Tlaxcala subjected to artificial winter periods of different lengths eclosed at significantly different time intervals (after subtracting winter length) ($F = 806.76$; $df = 4.16$; $P < 0.001$). No adults eclosed during the 10-month observational period from pupae subjected to constant temperatures and 4 weeks at 5°C, pupae subjected to 8 weeks at 5°C, took longer to eclose as adults than pupae subjected to winters of 10 and 16 weeks (table 1). No statistical differences were observed in time to adult eclosion between sexes ($F = 0.40$; $df = 1.16$; $P = 0.53$), and the interaction between sex and winter length was not significant ($F = 1.02$; $df = 4.16$; $P = 0.42$).

Rhagoletis cingulata pupae from Coahuila eclosed as adults at different time intervals ($F = 70.79$; $df = 4.5$; $P < 0.001$). Adults emerging from pupae subjected to constant temperatures took longer to eclose than those recovered from any other temperature treatment (table 1). No statistical differences were observed in time to adult eclosion between sexes ($F = 0.007$; $df = 1.5$; $P = 0.93$), and the interaction between sex and winter length was not significant ($F = 1.03$; $df = 4.5$; $P = 0.47$).

Rhagoletis turpiniae from Veracruz eclosed as adults at different time intervals ($F = 383.9$; $df = 4.15$; $P < 0.001$). No adults

eclosed during the 10-month observational period when pupae were subjected to constant temperatures and 4 weeks at 5°C (table 1). No statistical differences were observed in time to adult eclosion between sexes ($F = 0.47$; $df = 1.15$; $P = 0.49$), and the interaction between sex and winter length was not significant ($F = 1.86$; $df = 4.15$; $P = 0.16$).

There were significant differences across sites in the time in days elapsed from the end of artificial winter to adult fly eclosion ($F_{2, 279} = 90.34$; $P < 0.001$). As revealed by unequal *N* honest significant difference tests, flies from Coahuila emerged significantly later than flies from Veracruz, which in turn, emerged significantly later than flies from Tlaxcala (fig. 3). There were significant differences in the time in days elapsed from the end of artificial winters of different duration to adult fly eclosion ($F_{2, 279} = 19.84$; $P < 0.001$). Pupae exposed to 5°C for 8 weeks took longer to emerge (132.21 ± 2.56 ; mean \pm S.E., days) than pupae exposed for 10 weeks (128.49 ± 1.63 days), which in turn, took longer to emerge than those exposed for 16 weeks (114.06 ± 1.09 days). There was however a significant interaction between site and winter length duration ($F_{4, 279} = 7.11$; $P < 0.001$) because pupae from Veracruz took longer to emerge as adults when exposed to 5°C for 10 weeks. There were no significant differences between sexes in time to adult eclosion ($F_{1, 279} = 0.33$; $P = 0.56$).

Parasitoids

Only one unidentified braconid parasitoid infesting *R. cingulata* from Tlaxcala became dormant; this individual emerged 116 days after exposure to a 10-week artificial winter period at 5°C.

Table 1. Mean proportion (\pm SD) of adults eclosing after dormancy (within 10 months of pupation), pupae undergoing long dormancy (>10 months), and pupal mortality for two populations of *Rhagoletis cingulata* populations (Tlaxcala and Coahuila) and *Rhagoletis turpiniae*.

Population	Mean % (\pm SD) dormant flies (%)	Mean % (\pm SD) long dormancy (%)	Mean % (\pm SD) pupal mortality (%)	Mean % (\pm SD) parasitism (%)
<i>R. cingulate</i> (Tlaxcala)	8.7 \pm 14.7	55.2 \pm 24.4*	36 \pm 9.7	3.3 \pm 9.7
<i>R. cingulata</i> (Coahuila)	25.1 \pm 11.4*	19.1 \pm 20.7	54.2 \pm 12.4	27.2 \pm 19.7*
<i>R. turpiniae</i> (Veracruz)	8.9 \pm 8.6	36.9 \pm 20.6	55.8 \pm 15.3	47.7 \pm 27.2*

Within columns, figures followed by an asterisk (*) denote greater proportions than expected from a uniform distribution across sites according to χ^2 tests.




Fig. 3. Duration of dormancy in days for *Rhagoletis cingulata* exploiting late fruiting black cherries *Prunus serotina* var *virens* (black bar), *Rhagoletis turpiniae* exploiting *Turpinia insignis* (dark gray bar), and *R. cingulata* exploiting early fruiting black cherries *P. serotina* var *capuli* (light gray bar).

In the case of *R. cingulata* from Coahuila, 74 parasitoids became dormant when subjected to artificial winter periods of different duration. A GLM revealed a significant interaction between treatment (winter length) and trophic level (herbivore/parasitoid) ($F = 24.86$; $df = 4.6$; $P < 0.001$). When exposed to constant temperatures and short winters (4 and 8 weeks), parasitoids emerged significantly earlier than their fly (herbivore) hosts, and when exposed to winters of 10 and 16 weeks of duration, parasitoids emerged later than their hosts (table 2).

In the case of *R. turpiniae* (Veracruz), 29 parasitoids became dormant after exposure to artificial winters of 8 and 10 weeks. The time from pupation to adult parasitoid eclosion corrected for winter length was significantly longer than that recorded for their fly hosts ($t = 4.44$; $df = 5$; $P < 0.001$) (table 2).

Discussion

As for other species in the genus *Rhagoletis* inhabiting Mexican high elevation temperate habitats (Rull *et al.*, 2016), very small proportions (<2%) of flies from the three species in the *cingulata* species group examined here and a few associated adult parasitoids emerged without becoming dormant. Pupae exposed to 5°C for different periods of time (0–16 weeks) became dormant and emerged as adults in proportions <30%, in contrast to other species in the *cingulata* species group from the USA, where 80% of *R. indifferens* exposed pupae can emerge as adults after 3 months at 3°C (Frick *et al.*, 1954; Brown & AliNiazee, 1977). Large proportions (19–55%) of exposed pupae entered prolonged dormancy (>10 months), and large proportions of pupae died without emerging as adults. The number of days elapsed from the end of artificial winter to

adult eclosion was longer for *R. cingulata* exploiting late fruiting *P. serotina* in northeastern Mexico (Coahuila) than for flies recovered from earlier fruiting plants in the central Altiplano (Tlaxcala), resulting in significant allochronic isolation between Mexican ‘populations’ of *R. cingulata*. *Rhagoletis turpiniae* and northeastern *R. cingulata* pupae suffered high proportions of parasitism.

Small proportions (<2%) of pupae in our experiment emerged within 40 days of pupation as adults without becoming dormant. Similar findings have been reported for Mexican populations of three species of *Rhagoletis* (Rull *et al.*, 2016). In the case of flies in the *cingulata* species group, Van Kirk & AliNiazee (1982) report 3% adult eclosion between 27 and 77 days after pupation for *R. indifferens* held at 22°C, while Frick *et al.* (1954) mentions that 0.3–1.1% of the population failing to become dormant. These patterns are similar to those we found for the three Mexican populations examined here. In the case of flies in the *pomonella* species group, Dambroski & Feder (2007) found greater proportions of non-diapausing flies at Northern than at Southern latitudes; such a trend does not appear to be evident for species in the *cingulata* species group.

Two North American species of *Rhagoletis* of economic importance have been introduced to Europe in the past 20 years (*R. cingulata* and *Rhagoletis completa*) (Aluja *et al.*, 2011; Johannesen *et al.*, 2013; Verheggen *et al.*, 2017), while others (*R. pomonella* in particular) could potentially become established in several temperate fruit growing regions of the world (Geng *et al.*, 2011; Qin *et al.*, 2015). Because extensive pesticide use is no longer an acceptable option to deal with invasive pest species, environmentally friendly pest management techniques such as the sterile insect technique (SIT) are

Table 2. Average (\pm SE) duration (days) corrected for artificial winter length (by subtracting the total number of days held at 5°C according to treatment) from pupation to adult eclosion of three species of *Rhagoletis* fruit flies and their parasitoids.

Treatment	<i>Rhagoletis cingulate</i> Tlaxcala	Parasitoids	<i>R. cingulate</i> Coahuila	Parasitoids	<i>Rhagoletis turpiniae</i> Veracruz	Parasitoids
Constant	N.D ¹	N.D	283.3 \pm 5.6	88.7 \pm 10.1	N.D	N.D
4 weeks	N.D	N.D	136.8 \pm 15.1	70 \pm 5.4	N.D	N.D
8 weeks	128.1 \pm 6.1	N.D	148.3 \pm 3.4	125.4 \pm 13.7	121.4 \pm 5.3	158.2 \pm 5.5
10 weeks	106.9 \pm 3.4	116	145.3 \pm 1.2	185	127.6 \pm 2.7	178.9 \pm 3.4
16 weeks	104.8 \pm 1.4	N.D	131.7 \pm 1.5	155	118.15 \pm 3.3	N.D

¹N.D. = no data (no post-winter adult eclosion).

being developed for area-wide pest management (Nestel *et al.*, 2016). Cost-effective artificial rearing is a prerequisite for SIT development and rearing of natural enemies (Klassen, 2005). While substantial progress has been made for artificial rearing of *R. cerasi* (Köppler *et al.*, 2009), selection of non-diapausing strains of *Rhagoletis* could further facilitate SIT. Non-diapausing strains of *R. pomonella* have been selected for by exposing larvae and pupae to constant light or long photoperiods (Baerwald & Boush, 1967; Prokopy, 1968). However, Brown (1978) reports that the progeny of crosses of non-diapausing *R. indifferens* reverted to diapausing during selection attempts. Exploring the role of photoperiod and latitudinal variation on the frequency of non-diapausing individuals may contribute to development of non-diapausing strains of *Rhagoletis*.

AliNiasee (1975) found for field collected pupae of *R. indifferens* exposed to subfreezing temperatures for different periods of time, a maximum of 77% adult emergence and a maximum of 30% of pupae undergoing prolonged dormancy. Similarly, Frick *et al.* (1954) exposed *R. indifferens* pupae to 4°C for periods ranging from 0 to 230 days, and obtained 0–80% adult emergence, 6.3–87% of pupae undergoing prolonged dormancy, and pupal mortality ranging from 12 to 23%. In this case, optimal adult recovery (80% emergence) was obtained when holding pupae at 4°C for 150–210 days, while short (<90 days) and long (231 days) winters resulted in high proportions (>40%) of pupae engaging in prolonged dormancy. High proportions of pupae engaging in long dormancy are not limited to flies in the *cingulata* species group, in the case of *R. cerasi*, short (<4 months) winters also result in large proportions of pupae engaging in long dormancy (Vallo *et al.*, 1976). While examining fitness tradeoffs of pupae undergoing annual and prolonged dormancy, Moraiti *et al.* (2012) conclude that prolonged dormancy is likely to be part of a bet-hedging strategy that *R. cerasi* populations follow as an adaptation to the environmental variability of their habitats. Bet hedging has been examined in detail for the chesnut weevil *Curculio elephas*, an insect species that exploits fruit undergoing periodic unpredictable masting events (Menu, 1993). In this case, Gourbière & Menu (2009) found that diapause-duration variability is more likely to have evolved by the spread of a bet-hedging strategy than by the establishment of a genetic polymorphism. Here, we found that higher proportions of the central Altiplano (Tlaxcala) pupae engaged in prolonged dormancy when exposed to artificial winters of different length. Flowering and fruit set of *P. serotina* var *capuli* in Tlaxcala overlaps with periods of high likelihood of frosting events (Ortiz *et al.*, 1992), and large area extensions with fruitless trees occur on a periodic basis without following a predictable pattern (JR personal observation). A similar, yet less acute, pattern has been reported in the case of *P. serotina* var *virens* in southern USA (Hough, 1965). However, in

Coahuila, *P. serotina* var *virens* begins flowering in March, a period that does not overlap with the period (November–February) of maximum likelihood (>50%) of frosting events, while *R. turpiniae* occurs in a frost-free environment. When compared with other species in the *cingulata* species group, and a species exploiting hosts under similar ecological conditions, Mexican populations seem to exhibit a high propensity to engage in prolonged dormancy, require cold periods of shorter duration to break diapause, and suffer greater mortality. Because in all cases, pupae was recovered during the same collection date and assigned randomly to temperature treatments, the outcome of our experiments seems to suggest that the observed variability in diapause length is not the result of a genetic polymorphism but rather an effect of phenotypic plasticity. Understanding of this life-history strategy among Mexican *Rhagoletis* in the *R. cingulata* species group will require however lengthy (2–4 years) and specifically designed experiments.

The high mortality levels observed among pupae in our experiment may be due to a combination of factors. In the case of *R. turpiniae*, temperature may have a higher threshold than 0–5°C, since low temperatures in Xalapa rarely drop below 9°C. In the case of Mexican *R. cingulata* populations, humidity may be an important cue for regulation of dormancy. For northeastern *R. cingulata*, and *R. turpiniae*, parasitism levels were also extremely high and mortality could have been caused by unemerged parasitoids. However, in all cases, there seemed to be a trend for higher mortality for pupae exposed to longer winter lengths, suggesting that energy reserves may have been exhausted more rapidly than in the case of species inhabiting environments at more Northern latitudes.

Allochrony has been found to be an important component for the interruption of gene flow between the hawthorn and the apple host races of the apple maggot fly (Feder *et al.*, 1993; Feder & Filchak, 1999; Filchak *et al.*, 2000). In fact, each of the six recognized sibling species in the *R. pomonella* species complex infests a unique set of host plants that overlap in distribution, but differ in fruiting phenology (Bush, 1966; Berlocher, 2000; Xie *et al.*, 2008). For cherry infesting *R. cerasi*, a parallel has been drawn with *R. pomonella* host races (Schwarz *et al.*, 2003). Flies breeding on *Prunus* and those breeding on *Lonicera* show marked differences in adult emergence adjusted to the fruiting phenology of their respective hosts (Boller & Bush, 1974). Recently, in the *cingulata* species group, Yee *et al.* (2015) found that eclosion times of adult *R. indifferens* from allopatric populations infesting sweet (introduced, early fruiting) and bitter (native, late fruiting) cherry also differ according to the phenology of their respective host plants. Bitter cherry flies and their associated parasitoids eclosed later than sympatric sweet cherry adults (Yee *et al.*, 2015). Rull *et al.* (2011) identified three North American geographically isolated populations of *R. cingulata* exploiting

three subspecies of *P. serotina* differing in fruiting phenology. These 'populations' exhibited significant differences in the time from fruit collection to adult eclosion under uncontrolled environmental conditions in central Mexico (Xalapa), and were found to be reproductively isolated (Tadeo *et al.*, 2015). Here we found that under similar environmental conditions, two of these populations (central and northeastern Mexico) differ in duration of dormancy. Northeastern Mexican *R. cingulata* exploiting late fruiting *P. serotina* var. *virens* eclosed from 20 to 39 days later than *R. cingulata* recovered from *P. serotina* var. *capuli* in central Mexico. In sum we can now conclude that these populations are geographically, reproductively, and allochronically isolated.

In the case of Mexican populations of *R. cingulata*, winter length had a significant effect on the duration of dormancy. Pupae exposed to short winters (8 weeks) took longer to eclose as adults than pupae exposed to long (10 and 16 weeks) winters. Brown & AliNiazee (1977) documented similar patterns for *R. indifferens* from western USA. In this case, longer periods under low temperatures (winter) resulted in faster and more synchronous adult eclosion after re-exposure to warm temperatures. Apparently, *Prunus* infesting flies in the *R. cingulata* species group exhibit plasticity in response to environmental variation. The ecological significance of this plasticity may be linked to the effect of environmental variation (in this case winter length) on plant phenology. Long winters may shorten the window for successful egg laying during fruiting.

Flowering and fruit set seasonal patterns for *P. serotina* var. *capuli* and *P. serotina* var. *virens* may be governed by precipitation, while the peak of annual rainfall occurs during the month of June in Tlaxcala, maximum rainfall occurs in September in Coahuila (<http://es.climate-data.org/>). As a result, maximum fruit availability for egg laying and fruit infestation peak during June–July in Tlaxcala and during September in Coahuila (Rull *et al.*, 2011). In turn, these regional differences in host plant phenology appear to have selected for the longer post-winter adult eclosion patterns observed for *R. cingulata* from Coahuila during our study.

High levels of parasitism were detected for two of the three fly species examined here; in all cases, some individual wasps emerged as adults without becoming dormant and overall those individuals engaging in dormancy emerged as adults 30–50 days later than their fly hosts. In general, parasitoids associated to *Rhagoletis* adjust their life cycle to that of their host (Ovruski *et al.*, 2007; Rull *et al.*, 2009; 2016). Interestingly, Forbes *et al.* (2009) found that the braconid wasp *Diachasma alloeum* has formed new incipient species as a result of specializing on diversifying hosts of *R. pomonella*. Traits that differentially adapt flies to early fruiting apple or late fruiting hawthorn quickly evolved and serve as ecological barriers to reproduction, also isolating wasps. Hood *et al.* (2015) further showed that this may be occurring for three species of parasitoids associated with *Rhagoletis* flies in the *pomonella* species complex. Although our data are scarce to reach similar conclusions for braconids associated to *R. cingulata* populations exploiting hosts with different fruiting phenology, it may highlight the opportunity to investigate if parasitoids are diversifying in cascade as their hosts. In particular, we believe that examining if the high tendency of *R. cingulata* from central Mexico to engage in prolonged dormancy has selected for parasitoids adopting a similar life-history strategy or results in temporal escape from parasitism is an interesting avenue for future research.

Acknowledgements

Funding was provided by the International Atomic Energy Agency (IAEA Research Contract No: 18331/R0), the Mexican Campaña Nacional contra Moscas de la Fruta (SAGARPA-CONACOFI-INECOL) and the Instituto de Ecología A.C.

References

- AliNiazee, M.T. (1975) Susceptibility of diapausing pupae of the western cherry fruit fly (Diptera: Tephritidae) and a parasite (Hymenoptera: Diapriidae) to subfreezing temperatures. *Environmental Entomology* **4**, 1011–1013.
- Aluja, M., Guillén, L., Rull, J., Höhn, H., Frey, J., Graf, B. & Samietz, J. (2011) Is the alpine divide becoming more permeable to biological invasions? – insights on the invasion and establishment of the walnut husk Fly, *Rhagoletis completa* (Diptera: Tephritidae) in Switzerland. *Bulletin of Entomological Research* **101**, 451–465.
- Baerwald, R.J. & Boush, M.G. (1967) Selection of a nondiapausing race of apple maggot. *Journal of Economic Entomology* **60**, 682–684.
- Baker, C.R.B. & Miller, G.W. (1978) The effect of temperature on the post-diapause development of four geographical populations of the European fruit fly (*Rhagoletis cerasi*). *Entomologia Experimentalis et Applicata* **23**, 1–13.
- Bale, J.S., Masters, G.J., Hodkinson, I.D., Awmack, C., Bezemer, T.M., Brown, V.K., Butterfield, J., Buse, A., Coulson, J.C., Farrar, J., Good, J.E.G., Harrington, J., Hartley, S., Hefin Jones, T., Lindroth, R.L., Press, M.C., Symrnioudis, I., Watt, A.D. & Whittaker, J.B. (2002). Herbivory in global climate change research: direct effects of rising temperature on insect herbivores. *Global Change Biology* **8**, 1–16.
- Berlacher, S.H. (2000) Radiation and divergence in the *Rhagoletis pomonella* species group: inferences from allozymes. *Evolution* **54**, 543–557.
- Boller, E.F. & Bush, G.L. (1974) Evidence for genetic variation in populations of the European cherry fruit fly, *Rhagoletis cerasi* (Diptera: Tephritidae) based on physiological parameters and hybridization experiments. *Entomologia Experimentalis et Applicata*, **17**, 279–293.
- Boller, E.F. & Prokopy, R.J. (1976) Bionomics and management of *Rhagoletis*. *Annual Review of Entomology* **21**, 223–246.
- Boyce, A.M. (1931) The diapause phenomenon, in insects, with special reference to *Rhagoletis completa* cress. (Diptera: Trypetidae). *Journal of Economic Entomology* **24**, 1018–1024.
- Bush, G.L. (1966). The taxonomy, cytology, and evolution of the genus *Rhagoletis* in North America (Diptera, Tephritidae). *Bulletin of the Museum of Comparative Zoology* **134**, 431–562.
- Brown, R.D. (1978) *Gamma irradiation of the western cherry fruit fly for the sterile male control technique*. PhD Thesis, Oregon State University, 240p.
- Brown, R.D. & AliNiazee, M.T. (1977) Synchronization of adult emergence of the western cherry fruit fly in the laboratory. *Annals of the Entomological Society of America* **70**, 678–680.
- Dambroski, H.R. & Feder, J.L. (2007) Host plant and latitude-related diapause variation in *Rhagoletis pomonella*: a test for multifaceted life history adaptation on different stages of diapause development. *Journal of Evolutionary Biology* **20**, 2101–2112.
- Dean, R.W. (1973). *Bionomics of the Apple Maggot in Eastern*. New York, Cornell University. Volume **3**, 64p.

- Donaldson-Matasci, M.C., Bergstrom, C.T. & Lachmann, M. (2013) When unreliable cues are good enough. *The American Naturalist* **182**, 313–327.
- Dowell, R.V. & Penrose, R.L. (2012) Distribution and phenology of *Rhagoletis fausta* (Osten Sacken 1877) and *Rhagoletis in-differens* curren 1932 (Diptera: Tephritidae) in California. *The Pan-Pacific Entomologist* **88**, 130–150.
- Feder, J.L. & Filchak, K.E. (1999) It's about time: the evidence for host plant-mediated selection in the apple maggot fly, *Rhagoletis pomonella*, and its implications for fitness trade-offs in phytophagous insects. pp. 211–225 in *Proceedings of the 10th International Symposium on Insect-Plant Relationships*. The Netherlands, Springer.
- Feder, J.L., Hunt, T.A. & Bush, G.L. (1993) The effects of climate, host plant phenology and host fidelity on the genetics of apple and hawthorn infesting races of *Rhagoletis pomonella*. *Entomologia Experimentalis et Applicata* **69**, 117–135.
- Feder, J.L., Berlocher, S.H., Roethele, J.B., Dambroski, H., Smith, J.J., Perry, W.L., Garvrilovic, V., Filchak, K.E., Rull, J. & Aluja, M. (2003) Allopatric genetic origins for sympatric host-plant shifts and race formation in *Rhagoletis*. *Proceedings of the National Academy of Sciences* **100**, 10314–10319.
- Feder, J.L., Xie, X., Rull, J., Velez, S., Forbes, A., Leung, B., Dambroski, H., Filchak, K.E. & Aluja, M. (2005) Mayr, Dobzhansky, and bush and the complexities of sympatric speciation in *Rhagoletis*. *Proceedings of the National Academy of Sciences* **102**, 6573–6580.
- Filchak, K.E., Roethele, J.B. & Feder, J.L. (2000) Natural selection and sympatric divergence in the apple maggot *Rhagoletis pomonella*. *Nature* **407**, 739–742.
- Forbes, A.A., Powell, T.H.Q., Stelinski, L.L., Smith, J.J. & Feder, J.L. (2009) Sequential sympatric speciation across trophic levels. *Science* **323**, 776–779.
- Forrest, J. & Miller-Rushing, A.J. (2010) Toward a synthetic understanding of the role of phenology in ecology and evolution. *Philosophical Transactions of the Royal Society B* **365**, 3101–3112.
- Frick, K.E., Simkover, H.G. & Telford, H.S. (1954) Bionomics of the cherry fruit flies in eastern Washington. *Bionomics of the Cherry Fruit Flies in Eastern Washington* **13**, 66 pp.
- Geng, J., Li, Z.H., Rajotte, E.G., Wan, F.H., Lu, X.Y., & Wang, Z.L. (2011) Potential geographical distribution of *Rhagoletis pomonella* (Diptera: Tephritidae) in China. *Insect Science* **18**, 575–582.
- Gourbière, S. & Menu, F. (2009) Adaptive dynamics of dormancy duration variability: evolutionary trade-off and priority effect lead to suboptimal adaptation. *Evolution* **63**, 1879–1892.
- Hood, G.R., Forbes, A.A., Powell, T.H., Egan, S.P., Hamerlinck, G., Smith, J.J. & Feder, J.L. (2015) Sequential divergence and the multiplicative origin of community diversity. *Proceedings of the National Academy of Sciences USA* **112**, E5980–E5989.
- Hough, A.F. (1965) Black cherry (*Prunus serotina* Ehrh.). pp. 539–545 in H.A. Fowells, (ed.) *Silvics of Forest Trees of the United States*. Washington, DC, Comp. U.S. Department of Agriculture, Agriculture Handbook 271.
- Johannesen, J., Keyghobadi, N., Schuler, H., Stauffer, C. & Vogt, H. (2013) Invasion genetics of American cherry fruit fly in Europe and signals of hybridization with the European cherry fruit fly. *Entomologia Experimentalis et Applicata*, **147**, 61–72.
- Klassen, W. (2005) Area-wide integrated pest management and the sterile insect technique. pp. 39–68 in Dyck, V.A., Hendrichs, J. & Robinson, A.S. (eds) *Sterile Insect Technique*. Netherlands, Springer.
- Köppler, K., Kaffer, T. & Vogt, H. (2009) Substantial progress made in the rearing of the European cherry fruit fly, *Rhagoletis cerasi*. *Entomologia Experimentalis et Applicata* **132**, 283–288.
- Košťál, V. (2006) Eco-physiological phases of insect diapause. *Journal of Insect Physiology* **52**, 113–127.
- Martínez, B.O.U.M. & Ruiz, C. (2005) *Riesgo de heladas para la agricultura en la región sureste del estado de Coahuila*. INIFAP-CIRNE. Campo Experimental Saltillo. Coahuila, México, Publicación Especial Núm. 5, 6.6p.
- McVaugh, R. (1951) A revision of the North American black cherries (*Prunus serotina* Ehrh., and relatives). *Brittonia*, **7**, 279–315.
- Menu, F. (1993) Strategies of emergence in the chestnut weevil *Curculio elephas* (Coleoptera: Curculionidae). *Oecologia* **96**, 383–390.
- Menu, F. & Debouzie, D. (1993) Coin-flipping plasticity and prolonged diapause in insects: example of the chestnut weevil *Curculio elephas* (Coleoptera: Curculionidae). *Oecologia* **93**, 367–373.
- Moraiti, C.A., Nakas, C.T. & Papadopoulos, N.T. (2012) Prolonged pupal dormancy is associated with significant fitness cost for adults of *Rhagoletis cerasi* (Diptera: Tephritidae). *Journal of Insect Physiology* **58**, 1128–1135.
- Moraiti, C.A., Nakas, C.T. & Papadopoulos, N.T. (2014) Diapause termination of *Rhagoletis cerasi* pupae is regulated by local adaptation and phenotypic plasticity: escape in time through bet-hedging strategies. *Journal of Evolutionary Biology* **27**, 43–54.
- Nestel, D., Rempoulakis, P., Yanovski, L., Miranda, M.A. & Papadopoulos, N.T. (2016) The evolution of alternative control strategies in a traditional crop: economy and policy as drivers of olive fly control. pp. 47–76 in Horowitz, A.R. & Ishaaya, I. (eds) *Advances in Insect Control and Resistance Management*. Springer International Publishing.
- Neven, L.G. & Yee, W.L. (2017) Impact of prolonged absence of low temperature on adult eclosion patterns of western cherry fruit fly (Diptera: Tephritidae). *Environmental Entomology* **46**, 708–713.
- Ortiz, W.R., Ruiz, S.G. & Fierro, R.S. (1992) Localización microclimática de zonas óptimas para la introducción del cultivo de la soya en el estado de Tlaxcala. *Atmósfera* **5**, 169–179.
- Ovruski, S.M., Wharton, R.A., Rull, J. & Guillén, L. (2007) *Aganaspis alujai* (Hymenoptera: Figitidae: Eucolilinae), a new species attacking *Rhagoletis* (Diptera: Tephritidae) in the Neotropical region. *Florida Entomologist* **90**, 626–634.
- Padilla, R.C. (1964) Mosca del capulín, una nueva plaga descubierta en la región de Texcoco, México. *Fitofilo* **43**: 9–18.
- Prokopy, R.J. (1968) The influence of photoperiod, temperature and food on the initiation of diapause in the apple maggot. *Canadian Entomologist* **100**, 318–329.
- Prokopy, R.J. & Papaj, D.R. (2000) Behavior of flies of the genera *Rhagoletis*, *Zonosemata*, and *Carpomya* (Trypetinae: Carpomyina). *Fruit Flies*. pp. 219–252 in Aluja, M. & Norrbom, A.L. (eds) *Fruit Flies (Tephritidae): Phylogeny and Evolution of Behavior*. Boca Raton, FL, CRC Press.
- Qin, Y., Paini, D.R., Wang, C., Fang, Y. & Li, Z. (2015) Global establishment risk of economically important fruit fly species (Tephritidae). *PLoS ONE* **10**, e0116424.
- Rull, J., Aluja, M., Feder, J. & Berlocher, S. (2006) Distribution and host range of hawthorn-infesting *Rhagoletis* (Diptera: Tephritidae) in Mexico. *Annals of the Entomological Society of America* **99**, 662–672.
- Rull, J., Wharton, R., Feder, J.L., Guillén, L., Sivinski, J., Forbes, A. & Aluja, M. (2009) Latitudinal variation in parasitoid guild composition and parasitism rates of North American

- hawthorn infesting *Rhagoletis*. *Environmental Entomology* **38**, 588–599.
- Rull, J., Aluja, M., Feder, J.L.** (2011) Distribution and basic biology of black cherry-infesting *Rhagoletis* (Diptera: Tephritidae) in México. *Annals of the Entomological Society of America* **104**, 202–211.
- Rull, J., Tadeo, E., Lasa, R. & Aluja, M.** (2016) The effect of winter length on survival and duration of dormancy of four sympatric species of *Rhagoletis* exploiting plants with different fruiting phenology. *Bulletin of Entomological Research* **106**, 818–826.
- Smith, J.J. & Bush, G.L.** (2000) Phylogeny of the subtribe Carpomyina (Trypetinae), emphasizing relationships of the genus *Rhagoletis*. pp. 187–217 in Aluja, M. & Norrbom, A.L. (eds) *Fruit Flies (Tephritidae): Phylogeny and Evolution of Behavior*. Boca Raton, FL, CRC Press.
- Sosa, V.** (1988) *Staphyleaceae*. Flora de Veracruz. Fascículo 57. Xalapa, Veracruz, México.
- Stark, S.B. & AliNiasee, M.T.** (1982) Model of postdiapause development in the western cherry fruit fly. *Environmental Entomology* **11**, 471–474.
- Schwarz, D., McPheron, B.A., Hartl, G.B., Boller, E.F. & Hoffmeister, T.S.** (2003) A second case of genetic host races in *Rhagoletis*?. A population genetic comparison of sympatric host populations in the European cherry fruit fly, *Rhagoletis cerasi*. *Entomologia Experimentalis et Applicata* **108**, 11–17.
- Tadeo, E., Feder, J.L., Egan, S., Schuler, H., Aluja, M., Rull, J.** (2015) Divergence and evolution of reproductive barriers among three allopatric populations of *Rhagoletis cingulata* across eastern North America and México. *Entomología Experimentalis et Applicata* **156**, 301–311.
- Tauber, M.J., Tauber, C.A. & Masaki, S.** (1986) *Seasonal Adaptations of Insects*. New York, NY, Oxford University Press on Demand.
- Teixeira, L.A., Isaacs, R. & Gut, L.J.** (2007) Habitat-specific flight period in the cherry fruit fly *Rhagoletis cingulata* (Loew)(Diptera: Tephritidae). *Environmental Entomology* **36**, 1339–1348.
- Vallo, V., Remund, U. & Boller, E.F.** (1976) Storage conditions of stockpiled diapausing pupae of *Rhagoletis cerasi* for obtaining high emergence rates. *BioControl* **21**, 251–256.
- Van Kirk, J.R. & AliNiasee, M.T.** (1981) Determining low-temperature threshold for pupal development of the western cherry fruit fly for use in phenology models. *Environmental Entomology* **10**, 968–971.
- Van Kirk, J.R. & AliNiasee, M.T.** (1982) Diapause development in the western cherry fruit fly, *Rhagoletis indifferens* Curran (Diptera, Tephritidae). *Journal of Applied Entomology* **93**, 440–445.
- Verheggen, F., Verhaeghe, A., Giordanengo, P., Tassus, X. & Escobar-Gutiérrez, A.** (2017) Walnut husk fly, *Rhagoletis completa* (Diptera: Tephritidae), invades Europe: invasion potential and control strategies. *Applied Entomology and Zoology* **52**, 1–7.
- Xie, X., Rull, J., Michel, A.P., Velez, S., Forbes, A.A., Lobo, N.F., Aluja, M. & Feder, J.L.** (2007) Hawthorn-infesting populations of *Rhagoletis pomonella* in Mexico and speciation mode plurality. *Evolution* **61**, 1091–1105.
- Xie, X., Michel, A.P., Schwarz, D., Rull, J., Velez, S., Forbes, A.A., Aluja, M. & Feder, J.L.** (2008) Radiation and divergence in the *Rhagoletis pomonella* species complex: inferences from DNA sequence data. *Journal of Evolutionary Biology* **21**, 900–913.
- Yee, W.L., Goughnour, R.B., Hood, G.R., Forbes, A.A. & Feder, J.L.** (2015) Chilling and host plant/site-associated eclosion times of Western cherry fruit fly (Diptera: Tephritidae) and a host-specific parasitoid. *Environmental Entomology* **44**, 1029–1042.