

A CALCULATION OF THE PERFECTOIDIZATION OF SEMIPERFECTOID RINGS

RYO ISHIZUKA 

Abstract. We show that perfectoidization can be (almost) calculated by using p -root closure in certain cases, including the semiperfectoid case. To do this, we focus on the universality of perfectoidizations and uniform completions, as well as the p -root closed property of integral perfectoid rings. Through this calculation, we establish a connection between a classical closure operation “ p -root closure” used by Roberts in mixed characteristic commutative algebra and a more recent concept of “perfectoidization” introduced by Bhatt and Scholze in their theory of prismatic cohomology.

§1. Introduction

Let p be a prime number. The method of *perfectoidization*, introduced by Bhatt and Scholze in [8], is an application of the theory of prismatic cohomology to commutative algebra. This yields a universal integral perfectoid ring over a ring, such as a *semiperfectoid ring*, which is a derived p -complete ring that can be written as a quotient of an integral perfectoid ring. This method can be seen as a generalization of the perfect closure of positive characteristic rings. See Section 2 for an explanation of the terminology used in perfectoid theory.

Perfectoidization has various applications to commutative algebra, such as a new theory of almost mathematics (see [8, §10.1]), a general version of the almost purity theorem (see [8, Th. 10.9]), much simpler proof of some previously known theorems in commutative algebra ([8, Rem. 10.13] and [19, Appendix A]), and a mixed characteristic analog of Hilbert–Kunz multiplicity and F -signature (see [10]).

In contrast to these applications, they are not yet widely used in commutative algebra. One problem is that many abstract theories, including homotopy theory, have been used, and therefore perfectoidization has a mysterious ring structure. To the best of the author’s knowledge, perfectoidization has only been explicitly calculated in [22, §2.3.1] and in the proof of [12, Th. 4.4].

1.1 p -root closure

In this paper, we give an explicit description of the perfectoidization of semiperfectoid rings by using p -root closure. Before explaining our first main theorem, we recall the notion of p -root closure.

DEFINITION 1.1 [23]. Let R be a p -torsion-free ring. We say that R is *p -root closed in $R[1/p]$* if $x \in R[1/p]$ satisfies $x^{p^n} \in R$ for some $n \geq 1$, then $x \in R$ holds.

Received May 22, 2023. Revised October 13, 2023. Accepted February 2, 2024.
2020 Mathematics subject classification: Primary 14G45, 46J05.

Keywords: perfectoid rings, perfectoidization, p -root closure, uniform completion.



The p -root closure $C(R)$ of R in $R[1/p]$ is the minimal p -root closed subring of $R[1/p]$ containing R . Focusing on the case of “in $R[1/p]$,” Roberts provided an explicit description of the p -root closure $C(R)$ as follows:

$$C(R) = \{x \in R[1/p] \mid \exists n \geq 1, x^{p^n} \in R\}.$$

The term “in $R[1/p]$ ” is omitted in this paper because only this case is considered.

Here is a brief mention of the history of (p -)root closure. The notion of p -root closure is a special case of *total n -root closure* introduced in commutative algebra by Anderson, Dobbs, and Roitman [2]. Previously, *n -root closedness* was used, for example, by Angermüller [5], Anderson [1], Watkins [26], and Brewer, Costa, and McCrimmon [9]. Furthermore, its origin can be traced back to Sheldon’s definition of *root closedness* in [25].

In the context of commutative algebra in mixed characteristic, Roberts provided the above explicit description of p -root closure and applied this even before perfectoid rings appeared. Most recently, (total) p -root closure has been renamed *p -integral closure* by Česnavičius and Scholze in [11] and is found to be more closely related to the perfectoid theory.

1.2 Main theorem

Under this notation, our main theorems can be stated in the following forms. For simplicity, we only state the theorems in the case of p -torsion-free rings. Readers interested in the general cases may refer to the referenced statements in each theorem, in conjunction with Section 1.5.

THEOREM 1.2 (Theorem 5.7; p -torsion-free case). *Let R be a p -torsion-free ring which satisfies the following conditions:*

1. *The p -adic completion \widehat{R} of R has a map from some integral perfectoid ring.*
2. *The perfectoidization $(\widehat{R})_{\text{perfd}}$ of \widehat{R} is an (honest) integral perfectoid ring.¹*
3. *The p -adic completion $\widehat{C(R)}$ of the p -root closure $C(R)$ is an integral perfectoid ring.*

Then we have an isomorphism

$$(\widehat{R})_{\text{perfd}} \cong \widehat{C(R)}.$$

This result clarifies the ring structure of perfectoidization by using p -root closure, which is a quite explicit closure operation. The left-hand side is constructed abstractly by homotopy theory, but it can be described as a p -root closure followed by a p -adic completion, which are only ring-theoretic operations.

1.3 Applications

Let R be a p -torsion-free ring such that its p -adic completion \widehat{R} is a semiperfectoid ring, that is, \widehat{R} is a quotient of some integral perfectoid ring. In applications of Theorem 1.2, it is crucial that R satisfies the assumptions of the theorem:

1. *The semiperfectoid ring \widehat{R} has a surjective map from some integral perfectoid ring by the definition of semiperfectoid rings.*

¹ As explained in the rest of Section 2, the perfectoidization of an algebra over some integral perfectoid ring is only a complex. So this assumption states that $(\widehat{R})_{\text{perfd}}$ gives an honest ring (see Theorem 2.9).

2. The perfectoidization $(\widehat{R})_{\text{perfd}}$ of \widehat{R} is an (honest) integral perfectoid ring by [8, Cor. 7.3 and Prop. 8.5] or the rest of Section 2.
3. The p -adic completion $\widehat{C(R)}$ of the p -root closure $C(R)$ is an integral perfectoid ring by virtue of [11, Prop. 2.1.8] (see Remark 5.8).

So we can provide an explicit description of the perfectoidization of such R as follows.

THEOREM 1.3 (Corollary 5.9; p -torsion-free case). *Let R be a p -torsion-free ring such that its p -adic completion \widehat{R} becomes a semiperfectoid ring. Then we have an isomorphism $(\widehat{R})_{\text{perfd}} \cong \widehat{C(R)}$. In particular, any p -torsion-free and (classically) p -adically complete semiperfectoid ring S has an isomorphism $S_{\text{perfd}} \cong \widehat{C(S)}$.*

In the study of commutative rings in mixed characteristic, semiperfectoid rings often appear as in [16]. An application of Theorem 1.3 is as follows.

THEOREM 1.4 (Construction 6.1 and Corollary 6.2). *Let (R_0, \mathfrak{m}, k) be a complete Noetherian local domain of mixed characteristic $(0, p)$ with perfect residue field k , and let p, x_2, \dots, x_n be any system of generators of the maximal ideal \mathfrak{m} such that p, x_2, \dots, x_n forms a system of parameters of R_0 . Choose compatible sequences of p -power roots*

$$\{p^{1/p^j}\}_{j \geq 0}, \{x_2^{1/p^j}\}_{j \geq 0}, \dots, \{x_n^{1/p^j}\}_{j \geq 0}$$

inside the absolute integral closure R_0^+ , the integral closure of R_0 in the algebraic closure of the fraction field of R_0 . Set a subring R_∞ of R_0^+ as

$$R_\infty := \bigcup_{j \geq 0} R_0[p^{1/p^j}, x_2^{1/p^j}, \dots, x_n^{1/p^j}] \subseteq R_0^+.$$

Then the p -adic completion \widehat{R}_∞ of R_∞ becomes a p -torsion-free semiperfectoid ring by Cohen's structure theorem (see Construction 6.1 for details). Then $(\widehat{R}_\infty)_{\text{perfd}}$ is isomorphic to the p -adic completion $\widehat{C(R_\infty)}$ of the p -root closure $C(R_\infty)$.

1.4 Strategy of proof

Let us comment on the strategy of the proof of Theorem 1.2. Our proof is attributed to some universalities and the following principle of “rigidity lemma” that makes sense in certain situations (for example, Lemma 5.2).

LEMMA. *Let $f: R \rightarrow R$ be an endomorphism of a ring R . Assume that R has a “good” map $S \rightarrow R$ from some ring S . Then if f is a map of S -algebras, f is exactly the identity map.*

In the proof of Theorem 1.2, two maps of rings $\varphi: \widehat{C(R)} \rightarrow (\widehat{R})_{\text{perfd}}$ and $\psi: (\widehat{R})_{\text{perfd}} \rightarrow \widehat{C(R)}$ are obtained through the universality of uniform completions and perfectoidizations, respectively. By using the lemma mentioned above, we can show that $\psi \circ \varphi$ and $\varphi \circ \psi$ are identity maps. The former is a consequence of the universality of uniform completions (Proposition 3.7), while the latter is a consequence of the use of “almost elements,” which is a concept from almost mathematics (Definition 4.3 and Lemma 5.2).

1.5 p -torsion case

While the aforementioned theorems only deal with p -torsion-free rings, we can show that similar statements hold in general.

For this purpose, we introduce the following symbol. Let R be a ring. The p -torsion-free quotient R^{ptf} is defined as the quotient ring

$$R^{\text{ptf}} := R/R[p^\infty] \cong \text{Im}(R \rightarrow R[1/p]),$$

where $R[p^\infty]$ is the ideal of all p^∞ -torsion elements of R , which is the kernel of the canonical map $R \rightarrow R[1/p]$. Then R^{ptf} is a p -torsion-free ring and we have a canonical map $R \rightarrow R^{\text{ptf}} \hookrightarrow R[1/p]$.

The general case of main theorems are obtained by substituting $C(R)$ and $(\widehat{R})_{\text{perfd}}$ for $C(R^{\text{ptf}})$ and $((\widehat{R})_{\text{perfd}})^{\text{ptf}}$, respectively. Furthermore, the symbols $(-)_{\text{perfd}}$ and $(-)^{\text{ptf}}$ are often interchangeably because of Corollary 5.3.

To show the general version of Theorem 1.2 (i.e., Theorem 5.7), we need to pass from the possibly p -torsion case to the p -torsion-free case as in [4]. With this in mind, we show that any integral perfectoid ring can be canonically modified into a p -torsion-free integral perfectoid ring (see Theorem 4.9) by using *pre-perfectoid pairs* as defined in Section 4.

1.6 Notation

A *Tate ring* is a topological ring A which has an open subring $A_0 \subseteq A$ and an element $t \in A_0$ such that the relative topology on A_0 coincides with the t -adic topology and $A = A_0[1/t]$ as abstract rings. Such an open subring A_0 is called a *ring of definition of A* and such an element t is called a *pseudo-uniformizer of A* . This pair $(A_0, (t))$ of a ring and its ideal is called a *pair of definition of A* . Note that a ring (resp., pair) of definition of A is not necessarily unique.

Conversely, for a ring A_0 and an element $t \in A_0$, the ring $A_0[1/t]$ becomes a Tate ring by taking $\{t^n(A_0/A_0[t^\infty])\}_{n \geq 0}$ as a fundamental system of open neighborhoods of 0. The Tate ring $A_0[1/t]$ has a pair of definition $(A_0/A_0[t^\infty], (t))$. When referring to a ring $A_0[1/t]$ as a *Tate ring*, we refer to the Tate ring that arises from the pair $(A_0/A_0[t^\infty], (t))$.

For a Tate ring A , the symbol A° means the set of all power-bounded elements. This gives an open subring of A . A subring $A^+ \subseteq A$ is a *ring of integral elements* if it is open and integrally closed in A and $A^+ \subseteq A^\circ$.

§2. Perfectoid rings

In this section, we recall and fix some definitions of perfectoid objects.

DEFINITION 2.1 [7, Def. 3.5]. Let S be a (non-zero) ring. Then S is an *integral perfectoid ring* if the following conditions hold:

1. There exists an element $\pi \in S$ such that S is π -adically complete and π^p divides p in S .
2. The Frobenius map $F: S/pS \rightarrow S/pS$ is surjective.
3. The kernel of $\theta: \mathbb{A}_{\text{inf}}(S) \rightarrow S$ is principal, where $\mathbb{A}_{\text{inf}}(S) := W(S^b)$.

This $\pi \in S$ is called a *perfectoid element* in this paper. Here, we do not require that π is a non-zero-divisor in S .

Recently, integral perfectoid rings are simply called *perfectoid rings*. To avoid confusion with perfectoid Tate rings defined later in Definition 2.5, we do not use the term perfectoid rings, but only integral perfectoid rings and perfectoid Tate rings.

LEMMA 2.2 (see [7, Lem. 3.9]). *Let S be an integral perfectoid ring, and let $\pi \in S$ be a perfectoid element. Then S has compatible sequences of p -power roots $\{(u\pi)^{1/p^j}\}_{j \geq 0}$ and $\{(v\pi)^{1/p^j}\}_{j \geq 0}$ of $u\pi$ and $v\pi$ where u and v are unit elements in S .*

We fix the element $\varpi := (v\pi)^{1/p}$ of S . Then ϖ becomes a perfectoid element of S . Without loss of generality, we can assume that a perfectoid element π has a compatible sequence of p -power roots $\{\pi^{1/p^j}\}_{j \geq 0}$.

Proof. The first statement follows from [7, Lem. 3.9].

We next check that ϖ is a perfectoid element of S . Note that the conditions (2) and (3) in Definition 2.1 are independent of the choice of a perfectoid element. Since $\varpi^p = v\pi$ divides p in S , it suffices to show that S is p -adically complete and this is also clear (see [11, §2.1.2] or [13, Prop. 2.8]). \square

REMARK 2.3 [15, Th. 3.52]. Let S be a ring, and let π be an element of S . Then S is an integral perfectoid ring with a perfectoid element $\pi \in S$ if and only if $\pi \in S$ satisfies the following:

1. S is π -adically complete and π^p divides p in S .
2. The p th power map $S/\pi S \xrightarrow{a \mapsto a^p} S/\pi^p S$ is an isomorphism of rings.
3. The multiplicative map

$$\begin{aligned} S[\pi^\infty] &\longrightarrow S[\pi^\infty] \\ s &\longmapsto s^p \end{aligned}$$

is bijective, where the symbol $S[\pi^\infty]$ is the ideal of all π^∞ -torsion elements of S , which is defined as

$$S[\pi^\infty] = \{s \in S \mid \exists n \in \mathbb{Z}_{>0}, \pi^n s = 0 \in S\}. \quad (2.1)$$

We recall the definition of semiperfectoid rings.

DEFINITION 2.4 [8, Notation 7.1]. A ring S is a *semiperfectoid ring* if it is a derived p -complete ring that is isomorphic to a quotient of an integral perfectoid ring.

We next explain perfectoid Tate rings. See Notation (Section 1.6) at the end of the Introduction for the basic terminology of Tate rings.

DEFINITION 2.5 [7], [14]. Let A be a complete Tate ring (more generally, let A be a Banach ring). Then A is a *perfectoid Tate ring* if the following conditions hold:

1. A is uniform, that is, the set of all power-bounded elements A° is bounded in A .
2. There exists a pseudo-uniformizer $\pi \in A$ such that π^p divides p in A° and the Frobenius map on $A^\circ/\pi^p A^\circ$ is surjective.

This $\pi \in A$ is again called a *perfectoid element*.

The following lemma establishes the connection between integral perfectoid rings and perfectoid Tate rings.

LEMMA 2.6 (see [7, Lem. 3.20]). *Let A be a Tate ring, and let A_0^+ be a ring of integral elements in A . If A is a perfectoid Tate ring, then A_0^+ is an integral perfectoid ring. In particular, A° is an integral perfectoid ring.*

Conversely, if A_0^+ is an integral perfectoid ring and bounded in A , the Tate ring A is a perfectoid Tate ring.

REMARK 2.7. Let A be a perfectoid Tate ring, and let $\pi \in A$ be a perfectoid element. Then π is a non-zero-divisor in A . In general, an integral perfectoid ring is not necessarily isomorphic to the set of all power-bounded elements in some perfectoid Tate ring.

LEMMA 2.8 (see [7, Lem. 3.21]). *Let S be an integral perfectoid ring, and let $\pi \in S$ be a perfectoid element. Assume that π is a non-zero-divisor in S . Then the Tate ring $S[1/\pi]$ defined by the pair $(S, (\pi))$ is a perfectoid Tate ring. Additionally, $S \subseteq (S[1/\pi])^\circ$, and its cokernel is annihilated by any fractional power of π .*

For convenience, we summarize a brief overview of the properties of perfectoidization (see [8], [10] for more details).

Let S be an integral perfectoid ring, and let R be a derived p -complete S -algebra. The perfectoidization R_{perfd} of R is defined by using the prismatic cohomology $\Delta_{R/S}$. Note that R_{perfd} is typically a commutative algebra object in $D^{\geq 0}(S)$ and has a map $R \rightarrow R_{\text{perfd}}$ in $D(S)$. The complex R_{perfd} is concentrated in degree 0 in the following cases:

- If $\text{char}(S) = p > 0$, R_{perfd} coincides with the usual perfect closure R_{perf} of R .
- If R can be written as a quotient of S , that is, R is a semiperfectoid ring, then R_{perfd} is an integral perfectoid ring and furthermore the map $R \rightarrow R_{\text{perfd}}$ is surjective.
- If $S \rightarrow R$ is an integral map, R_{perfd} is an integral perfectoid ring.

Furthermore, the following property plays an essential role in this paper.

THEOREM 2.9 (see [8, Cor. 8.14]). *If R_{perfd} is concentrated in degree 0, it becomes an integral perfectoid ring. In this case, the map $R \rightarrow R_{\text{perfd}}$ is the universal map to integral perfectoid rings. Namely, every map $R \rightarrow R'$ to an integral perfectoid ring R' uniquely factors through $R \rightarrow R_{\text{perfd}}$.*

§3. Uniform completion

We use the notion of the uniform completion of Tate rings. In this section, we review the uniform completion outlined in [16].

DEFINITION 3.1. A Tate ring A is uniform if the set of all power-bounded elements A° is bounded.

Any Tate ring has the structure of a seminormed ring as follows (see [20, Def. 2.26] for more details).

DEFINITION 3.2. Let $A := A_0[1/t]$ be a Tate ring. Fix a real number $c > 1$. Then, we can define a seminorm $\|\cdot\|_{A_0,t,c}: A \rightarrow \mathbb{R}_{\geq 0}$ by

$$\|f\|_{A_0,t,c} := \inf_{m \in \mathbb{Z}} \{c^m \mid t^m f \in A_0\}.$$

The seminorm defines a seminormed ring $(A, \|\cdot\|_{A_0,t,c})$. The topology of the seminormed ring $(A, \|\cdot\|_{A_0,t,c})$ is equal to the topology of the Tate ring $A = A_0[1/t]$. In particular, the topology induced from the norm $\|\cdot\|_{A_0,t,c}$ does not depend on the choices of A_0 , t , and c . So we write the seminorm $\|\cdot\|_{A_0,t,c}$ as $\|\cdot\|$ for simplicity.

The spectral seminorm attached to this seminorm $\|\cdot\|$ is defined as

$$\|f\|_{\text{sp}} := \lim_{n \rightarrow \infty} \|f^n\|^{1/n}.$$

By Fekete’s subadditivity lemma, we have $\|f\|_{\text{sp}} \leq \|f\|$.

LEMMA 3.3 (see [20, Lem. 2.29]). *Let $A = A_0[1/t]$ be a uniform Tate ring. Then A° is equal to the unit disk $A_{\|\cdot\|_{\text{sp}} \leq 1}$ by the spectral seminorm in A .*

Next, we define the uniformization and uniform completion of Tate rings as in [6, Exer. 7.2.6] and [17, Def. 2.8.13].

DEFINITION 3.4. Let A be a Tate ring. Fix a pair of definition $(A_0, (t))$ of A and a ring of integral elements A_0^+ of A such that $A_0 \subseteq A_0^+$. We define the following terminology.

1. The *uniformization of A with respect to $(A_0, (t))$ and A_0^+* is the Tate ring $A_0^+[1/t]$.
2. The *uniform completion of A with respect to $(A_0, (t))$ and A_0^+* is the completion of the Tate ring $A_0^+[1/t]$. This is in fact the Tate ring $\widehat{A_0^+[1/t]}$, where $\widehat{A_0^+}$ is the t -adic completion of A_0^+ .

REMARK 3.5. At first glance, the above definitions depend on the choices of $(A_0, (t))$ and A_0^+ . The motivation for these definitions is that we wanted to define them “functorially.”

Furthermore, even if we take a different pair of definition $(A'_0, (t'))$ of A and a different ring of integral elements A'^0_+ such that $A_0 \subseteq A'^0_+$, the uniformization of A with respect to $(A'_0, (t'))$ and A'^0_+ is isomorphic to the uniformization of A with respect to $(A_0, (t))$ and A_0^+ . Its isomorphism is obtained by the identity map on the abstract ring $A = A_0^+[1/t] = A'^0_+[1/t]$ (see [16, Lem. 5.5] and [20, Lem. 2.3]). In particular, the same statement is true for the uniform completion. So the next definitions are well-defined.

DEFINITION 3.6. Let A be a Tate ring. The *uniformization A^u of A* (resp., *uniform completion $A^{\widehat{u}}$ of A*) is the uniformization (resp., uniform completion) of A with respect to a pair of definition $(A_0, (t))$ and a ring of integral elements A_0^+ of A such that $A_0 \subseteq A_0^+$. By the above Remark 3.5, these definitions are independent of the choices of $(A_0, (t))$ and A_0^+ .

Recall that the canonical map of Tate rings $i: A \rightarrow A^u \rightarrow A^{\widehat{u}}$ has the following universal property.

PROPOSITION 3.7 (see [16, Prop. 5.6]). *Let A be a Tate ring. Then the uniform completion $A^{\widehat{u}}$ is a uniform complete Tate ring. Furthermore, the canonical map $i: A \rightarrow A^{\widehat{u}}$ is the universal map to uniform complete Tate rings. That is, every map of Tate rings $h: A \rightarrow B$, where B is a uniform complete Tate ring uniquely factors through $i: A \rightarrow A^{\widehat{u}}$.*

We record some lemmas as follows.

LEMMA 3.8. *Let A be a Tate ring. Then, the completion $\widehat{A^{u^\circ}}$ of the set of all power-bounded elements A^{u° of A^u is isomorphic to the set of all power-bounded elements $(A^{\widehat{u}})^\circ$ of $A^{\widehat{u}}$ as topological ring.*

Proof. Fix a pair of definition $(A_0, (t))$ of A and a ring of integral elements A_0^+ of A such that $A_0 \subseteq A_0^+$. By [20, Lem. 2.3], we have an inclusion $t(A_0^+)_A^* \subseteq A_0^+$, where $(A_0^+)_A^*$ is the complete integral closure of A_0^+ in A . By [20, Prop. 2.4], the canonical map $A^{\widehat{u}} = \widehat{A_0^+[1/t]} \rightarrow \widehat{(A_0^+)_A^*[1/t]}$ is an isomorphism of Tate rings and this map induces an isomorphism of topological rings $(\widehat{A_0^+})_{A^{\widehat{u}}}^* \rightarrow (\widehat{(A_0^+)_A^*})_{A^u}^*$. Since $A^{\widehat{u}}$ (resp., A^u) has a ring of definition $\widehat{A_0^+}$ (resp., A_0^+) and the complete integral closure is equal to the set of all power-bounded elements by [20, Lem. 2.13], we have an isomorphism of topological rings $(A^{\widehat{u}})^\circ \xrightarrow{\cong} \widehat{A^{u^\circ}}$. \square

LEMMA 3.9 (see [16, Prop. 5.6]). *Assume that a Tate ring $A = A_0[1/t]$ has a compatible sequence of p -power roots $\{t^{1/p^j}\}_{j \geq 0}$ of t . The inclusion map $(A^{\widehat{u}})^\circ \hookrightarrow ((A^{\widehat{u}})^\circ)_* := t^{-1/p^\infty}((A^{\widehat{u}})^\circ)$ is an isomorphism of rings.*

Proof. For any $f \in ((A^{\widehat{u}})^\circ)_*$, we have $t^{1/p^n} f \in (A^{\widehat{u}})^\circ$ and then $tf^{p^n} \in (A^{\widehat{u}})^\circ$ for any $n \in \mathbb{Z}_{>0}$. Since $(A^{\widehat{u}})^\circ$ is a ring of definition of $A^{\widehat{u}}$, we have $\|f^{p^n}\| \leq c$ for a fixed $c > 1$ by Definition 3.2. In particular, $\|f^{p^n}\|^{1/p^n} \leq c^{1/p^n}$ for any $n \in \mathbb{Z}_{>0}$. Taking the limit $n \rightarrow \infty$, we have $\|f\|_{\text{sp}} \leq \lim_{n \rightarrow \infty} c^{1/p^n} = 1$. This shows the inclusion $((A^{\widehat{u}})^\circ)_* \subseteq (A^{\widehat{u}})_{\|\cdot\|_{\text{sp}} \leq 1} = (A^{\widehat{u}})^\circ$ by Lemma 3.3. □

§4. Some ring-theoretic properties of pre-perfectoid pairs

Let R be an integral perfectoid ring, and let $\pi \in R$ be a perfectoid element. Our goal in this section is to convert a situation where π is a zero-divisor into a situation where it is a non-zero-divisor, following the approach of [4, §2.3.2]. Our argument is based on [3], [4] and is similar to [6], [24]. If it is sufficient to consider only π -torsion-free rings (resp., p -torsion-free rings), the symbol $(-)^{\pi\text{tf}}$ defined in Definition 4.1 (resp., $(-)^{p\text{tf}}$) can be removed.

For the sake of generality, we define pre-perfectoid pairs as follows.

DEFINITION 4.1. Let (S, π) be a pair such that S is a ring and π is an element of S which has a compatible sequence of p -power roots $\{\pi^{1/p^j}\}_{j \geq 0}$ in S and π^p divides p in S . If the p th power map $S/\pi S \xrightarrow{a \mapsto a^p} S/\pi^p S$ is isomorphism, we call such a pair (S, π) *pre-perfectoid pair*.

For a pre-perfectoid pair (S, π) , the π -torsion-free quotient $S^{\pi\text{tf}}$ of S is defined as the quotient ring

$$S^{\pi\text{tf}} := S/S[\pi^\infty] \cong \text{Im}(S \rightarrow S[1/\pi]),$$

where $S[\pi^\infty]$ is the ideal of all π^∞ -torsion elements of S . Note that $S \twoheadrightarrow S^{\pi\text{tf}}$ is an isomorphism if and only if π is a non-zero-divisor of S . In the case of $S[1/\pi] = S[1/p]$, $S^{\pi\text{tf}}$ is equal to the p -torsion-free quotient $S^{p\text{tf}}$ of S defined in Section 1.5.

For example, an integral perfectoid ring R and a perfectoid element π of R form a pre-perfectoid pair (R, π) because of Remark 2.3(2).

DEFINITION 4.2. Let (S, π) be a pre-perfectoid pair. An S -module M is called $(\pi)^{1/p^\infty}$ -almost zero if $\pi^{1/p^n} \cdot M = 0$ for any $n \in \mathbb{Z}_{>0}$. A map of S -modules $N \rightarrow M$ is called $(\pi)^{1/p^\infty}$ -almost injective (resp., surjective) if its kernel (resp., cokernel) is $(\pi)^{1/p^\infty}$ -almost zero. If a map of S -modules $N \rightarrow M$ is $(\pi)^{1/p^\infty}$ -almost injective and surjective, we call the map a $(\pi)^{1/p^\infty}$ -almost isomorphism.

DEFINITION 4.3. Let (S, π) be a pre-perfectoid pair. We define *the set of almost elements of S* (see, for example, [24, Lem. 5.3]):

$$S_* := \pi^{-1/p^\infty} S := \pi^{-1/p^\infty} S^{\pi\text{tf}} := \{s \in S[1/\pi] \mid \forall n \in \mathbb{Z}_{>0}, \pi^{1/p^n} s \in S^{\pi\text{tf}} \subseteq S[1/\pi]\}. \tag{4.1}$$

For any S -module M , *the set of almost elements of M* is defined as

$$M_* := \pi^{1/p^\infty} M := \{m \in M[1/\pi] \mid \forall n \in \mathbb{Z}_{>0}, \pi^{1/p^n} m \in M^{\pi\text{tf}} \subseteq M[1/\pi]\}, \tag{4.2}$$

where $M^{\pi\text{tf}}$ is the π -torsion-free quotient of M , that is, the image of M in $M[1/\pi]$.

REMARK 4.4. Let (S, π) be a pre-perfectoid pair. Note that $S[1/\pi]$ and $S_*[1/\pi]$ are isomorphic as abstract rings. Furthermore, as Tate rings, $S[1/\pi]$ has a pair of definition $(S^{\pi\text{tf}}, (\pi))$ and $S_*[1/\pi]$ has a pair of definition $(S_*, (\pi))$. Because of $\pi S_* \subseteq S^{\pi\text{tf}} \subseteq S_*$, these are isomorphic as topological rings.

LEMMA 4.5. *Let (S, π) be a pre-perfectoid pair. Assume that S is reduced (e.g., S is an integral perfectoid ring, see [11, §2.1.3] or [13, Prop. 2.20]). Then, the canonical map $S \rightarrow S_*$ is a $(\pi)^{1/p^\infty}$ -almost isomorphism.*

Proof. The kernel of $S \rightarrow S[1/\pi]$ is isomorphic to the ideal $S[\pi^\infty]$ of all π^∞ -torsion elements of S . Since S is reduced, $S[\pi^\infty]$ is $(\pi)^{1/p^\infty}$ -almost zero. In particular, $S \rightarrow S^{\pi\text{tf}}$ is (usual) surjective and $(\pi)^{1/p^\infty}$ -almost injective. Furthermore, the inclusion $S^{\pi\text{tf}} \subseteq \pi^{-1/p^\infty} S$ in $S[1/\pi]$ is (usual) injective and $(\pi)^{1/p^\infty}$ -almost surjective by definition. This completes the proof. \square

LEMMA 4.6. *Let (S, π) a pre-perfectoid pair. Then the p -th power map $S_*/\pi S_* \xrightarrow{a \mapsto a^p} S_*/\pi^p S_*$ is injective and $(\pi)^{1/p^\infty}$ -almost surjective (later in Corollary 4.8, this will become a (usual) surjective map). In particular, S_* is p -root closed in $S_*[1/\pi]$. Namely, if $x \in S_*[1/\pi]$ satisfies $x^{p^n} \in S_*$ for some $n \in \mathbb{Z}_{>0}$, then $x \in S_*$.*

Proof. If π is a non-zero-divisor of S , this lemma is similar to [13, Prop. 2.16(b)]. We only have to make the same proof as [24, Lem. 5.6], being careful that S is not necessarily π -torsion-free in our case.

First, we show that $S_*/\pi S_* \xrightarrow{a \mapsto a^p} S_*/\pi^p S_*$ is injective. Let $t \in S_*$ be an element in the kernel of the p -th power map. There exists some $t' \in S_*$ such that $t^p = \pi^p t'$ in $S_* \subseteq S[1/\pi]$. By definition of S_* , multiplying π^{1/p^n} by the equation for each $n \in \mathbb{Z}_{>0}$, we have

$$(\pi^{1/p^{n+1}} t)^p = \pi^{1/p^n} t^p = \pi^p (\pi^{1/p^n} t') \in \pi^p S^{\pi\text{tf}}. \tag{4.3}$$

Moreover, $\pi^{1/p^{n+1}} t$ and $\pi^{1/p^n} t'$ are elements of $S^{\pi\text{tf}}$. Then, there exist some elements s_{n+1} and s'_n in S such that $s_{n+1}/1 = \pi^{1/p^{n+1}} t$ and $s'_n/1 = \pi^{1/p^n} t'$ in $S^{\pi\text{tf}} \subseteq S[1/\pi]$. By the above equation (4.3), we have $s_{n+1}^p/1 = \pi^p s'_n/1$ in $S^{\pi\text{tf}} \subseteq S[1/\pi]$ and thus, $s_{n+1}^p - \pi^p s'_n$ is in $S[\pi^\infty] \subseteq S$, which is $(\pi)^{1/p^\infty}$ -almost zero. For any $m \in \mathbb{Z}_{>0}$, we have

$$S \ni 0 = \pi^{1/p^m} (s_{n+1}^p - \pi^p s'_n) = (\pi^{1/p^{m+1}} s_{n+1})^p - \pi^p (\pi^{1/p^m} s'_n).$$

In particular, $\pi^{1/p^{m+1}} s_{n+1}$ is in the kernel of the p -th power map $S/\pi S \rightarrow S/\pi^p S$, which is zero by assumption of (S, π) , and so $\pi^{1/p^{m+1}} s_{n+1}$ is in πS . Passing to $S^{\pi\text{tf}} \subseteq S[1/\pi]$, we have

$$\pi S^{\pi\text{tf}} \ni \pi^{1/p^{m+1}} s_{n+1}/1 = \pi^{1/p^{m+1}} \pi^{1/p^{n+1}} t$$

for any $n, m \in \mathbb{Z}_{>0}$. Then, t is in $(\pi S^{\pi\text{tf}})_*$, which is defined in Definition 4.3. The next equality (4.4) shows the injectivity of $S_*/\pi S_* \xrightarrow{a \mapsto a^p} S_*/\pi^p S_*$:

$$(\pi S^{\pi\text{tf}})_* = \pi(S^{\pi\text{tf}})_* = \pi S_* \subseteq S[1/\pi]. \tag{4.4}$$

Proof of (4.4). By definition, we have $(S^{\pi\text{tf}})_* = S_*$ and then the second equality is clear. Any element of $\pi(S^{\pi\text{tf}})_*$ can be written as πt by using some element $t \in (S^{\pi\text{tf}})_*$. Because of $\pi^{1/p^n} (\pi t) = \pi(\pi^{1/p^n} t) \in \pi S^{\pi\text{tf}}$ for any $n \in \mathbb{Z}_{>0}$, we have $\pi t \in (\pi S^{\pi\text{tf}})_*$ and so $\pi(S^{\pi\text{tf}})_* \subseteq (\pi S^{\pi\text{tf}})_*$.

Conversely, take any element $x \in (\pi S^{\pi\text{tf}})_* \subseteq S[1/\pi]$. Then, there exists $t_n \in S^{\pi\text{tf}}$ such that $\pi^{1/p^n} x = \pi t_n \in \pi S^{\pi\text{tf}}$ for each $n \in \mathbb{Z}_{>0}$. Since $S[1/\pi]$ is π^{1/p^n} -torsion-free, we have

$$S[1/\pi] \ni \pi^{1/p^n} (x/\pi) = t_n \in S^{\pi\text{tf}} \tag{4.5}$$

for any $n \in \mathbb{Z}_{>0}$. This shows that $x/\pi \in S[1/\pi]$ is in $(S^{\pi\text{tf}})_*$ and thus $x = \pi(x/\pi)$ is in $\pi(S^{\pi\text{tf}})_*$. \square

Second, we show that $S_*/\pi S_* \xrightarrow{a \mapsto a^p} S_*/\pi^p S_*$ is $(\pi)^{1/p^\infty}$ -almost surjective. Take any element $x \in S_* \subseteq S[1/\pi]$. For each $n \in \mathbb{Z}_{>0}$, we have $\pi^{1/p^n} x \in S^{\pi\text{tf}}$ and thus there exists an $s_n \in S$ such that $\pi^{1/p^n} x = s_n/1 \in S^{\pi\text{tf}}$. Since $S/\pi S \xrightarrow{a \mapsto a^p} S/\pi^p S$ is surjective, there exists some $s'_n \in S$ such that $(s'_n)^p - s_n$ is in $\pi^p S$. Then, we have

$$\pi^{1/p^n} x - (s'_n/1)^p = s_n/1 - (s'_n)^p/1 \in \pi^p S^{\pi\text{tf}} \subseteq \pi^p S_* \tag{4.6}$$

and $s'_n/1$ is in $S^{\pi\text{tf}} \subseteq S_*$. This shows that $S_*/\pi S_* \xrightarrow{a \mapsto a^p} S_*/\pi^p S_*$ is $(\pi)^{1/p^\infty}$ -almost surjective.

Finally, since S_* is π -torsion-free and $S_*/\pi S_* \xrightarrow{a \mapsto a^p} S_*/\pi^p S_*$ is injective as above, by [11, (2.1.7.1)], we can show that S_* is p -root closed in $S_*[1/\pi]$.

LEMMA 4.7. *Let (S, π) be a pre-perfectoid pair. Then $S_*[1/\pi]$ is a uniform Tate ring which satisfies $(S_*[1/\pi])^\circ = S_*$. In particular, S_* is isomorphic to the unit disk of $S_*[1/\pi]$ with respect to the spectral seminorm induced from the seminorm of $S_*[1/\pi]$ as defined in Definition 3.2.*

Proof. Proceeding as in [24, Lem. 5.6], the p -root closedness of S_* in $S_*[1/\pi]$ proved in Lemma 4.6 shows that $(S_*[1/\pi])^\circ = S_*$. In fact, any element $x \in (S_*[1/\pi])^\circ$ makes a topologically nilpotent element $\pi^{1/p^n} x$ for each $n \in \mathbb{Z}_{>0}$. Thus there exists $N = N(n) \in \mathbb{Z}_{>0}$ such that $(\pi^{1/p^n} x)^{p^N} \in S_*$. This shows the equality above and thus $S_*[1/\pi]$ is a uniform Tate ring. Moreover, Lemma 3.3 shows that

$$S_* = (S_*[1/\pi])^\circ = (S_*[1/\pi])_{\|\cdot\|_{\text{sp}} \leq 1} \tag{4.7}$$

and we finish the proof. \square

COROLLARY 4.8. *Let (S, π) be a pre-perfectoid pair. Stronger than Lemma 4.6, we can show that the p -th power map $S_*/\pi S_* \xrightarrow{a \mapsto a^p} S_*/\pi^p S_*$ is (usual) surjective.*

Proof. This proof is similar to [16, Lem. 4.2]. Fix an element $y \in S_*$. Since the p -th power map is $(\pi)^{1/p^\infty}$ -almost surjective by Lemma 4.6, there exist elements a and b in S_* such that $\pi y = a^p + \pi^p b \in S_*$. Set $z := a/\pi^{1/p} \in S_*[1/\pi]$. This satisfies

$$z^p = a^p/\pi = y - \pi^{p-1} b \in S_* \tag{4.8}$$

Since S_* is p -root closed in $S_*[1/\pi]$ by Lemma 4.6, we can show that $z \in S_*$. The equality $\pi y = \pi z^p + \pi^p b \in S_*$ and π -torsion-freeness of $S_* \subseteq S_*[1/\pi]$ show that $y = z^p + \pi^{p-1} b \in S_*$. In particular, every element of $S_*/\pi^{p-1} S_*$ and $S_*/\pi S_*$ is a p -th power. The inclusion $\pi S_* \subseteq \pi^{(p-1)/p} S_*$ induces the commutative diagram

$$\begin{array}{ccccccc}
0 & \longrightarrow & S_*/\pi^{1/p}S_* & \xrightarrow{\cdot\pi^{(p-1)/p}} & S_*/\pi S_* & \longrightarrow & S_*/\pi^{(p-1)/p}S_* \longrightarrow 0 \\
& & \downarrow & & \downarrow & & \downarrow \\
0 & \longrightarrow & S_*/\pi S_* & \xrightarrow{\cdot\pi^{p-1}} & S_*/\pi^p S_* & \longrightarrow & S_*/\pi^{p-1}S_* \longrightarrow 0
\end{array}$$

where the rows are exact sequences and the vertical maps are the p -th power maps. Since the right-most map and left-most map are surjective, the middle map is also surjective by the five lemma. \square

Finally, we show the stability of pre-perfectoid pairs under taking $(-)_*$ and the generalization of Lemma 2.8.

THEOREM 4.9. *For any pre-perfectoid pair (S, π) , the induced pair (S_*, π) is again a pre-perfectoid pair.*

In particular, for any integral perfectoid ring R and a perfectoid element $\pi \in R$ (not necessarily a non-zero-divisor), $R_ = \pi^{-1/p^\infty}R$ is a π -torsion-free integral perfectoid ring with a perfectoid element π .*

Proof. By Lemma 4.6 and Corollary 4.8, the pair (S_*, π) is a pre-perfectoid pair.

Assume that R is an integral perfectoid ring. By the above paragraph of [11, (2.1.3.1)], the image $R^{\pi\text{tf}}$ of R in $R[1/\pi]$ is a π -torsion-free integral perfectoid ring. In particular, similar to the proof of [24, Lem. 5.6], $R_* = (R^{\pi\text{tf}})_*$ is also an integral perfectoid ring with a perfectoid element π . \square

REMARK 4.10. By Lemma 2.2, any integral perfectoid ring R has a compatible sequence of p -power roots $\{\varpi^{1/p^j}\}_{j \geq 0}$ of $\varpi \in R$ such that ϖ^p is some unit multiple of p in R . Then R forms a pre-perfectoid pair (R, ϖ) . In particular, this pre-perfectoid pair (R, ϖ) satisfies all the statements in this section.

§5. Calculations of the perfectoidization

We next show the “universality” of uniformizations and uniform completions for pre-perfectoid pairs and deduce our main theorem (Theorem 5.7). As in the previous section, we can get rid of the symbol $(-)^{\text{ptf}}$ if we only consider p -torsion-free rings.

PROPOSITION 5.1. *Let A_0 be a π -adically topological ring for an element $\pi \in A_0$. Set a Tate ring $A := A_0[1/\pi]$. Let S be an A_0 -algebra such that the image of π in S is a non-zero element. Assume that S is an integral perfectoid ring and π is a perfectoid element of S .*

Then, there exists a unique map of topological rings $(A^{\widehat{u}})^\circ \rightarrow S_$ such that the following diagram commutes:*

$$\begin{array}{ccc}
A_0 & \longrightarrow & S \\
\downarrow & & \downarrow \\
(A^{\widehat{u}})^\circ & \xrightarrow{\exists!} & S_*
\end{array}$$

Proof. Since the A_0 -algebra structure $A_0 \rightarrow S$ is a continuous map with respect to the π -adic topology, this can be extended to the map of Tate rings $A \rightarrow S[1/\pi]$. By Remark 4.4 and Lemma 4.7, the Tate ring $S[1/\pi]$ is isomorphic to the complete uniform Tate ring $S_*[1/\pi]$.

The universality of uniform completion (Proposition 3.7) gives a unique map of Tate rings $A^{\widehat{u}} \rightarrow S_*[1/\pi]$ which extends $A \rightarrow S[1/\pi] = S_*[1/\pi]$. Taking the set of all power-bounded elements, we have a map of topological rings $(A^{\widehat{u}})^{\circ} \rightarrow S_*$ such that the above diagram commutes. The uniqueness of this map is clear. \square

To prove our main theorem (Theorem 5.7), we state the next lemma which is in the form of “rigidity lemma” as in Section 1.4.

LEMMA 5.2. *Let A_0 be a derived p -complete algebra over some integral perfectoid ring of characteristic 0.² Assume that $(A_0)_{\text{perfd}}$ is an (honest) integral perfectoid ring. Then any A_0 -algebra map $(A_0)_{\text{perfd},*} \rightarrow (A_0)_{\text{perfd},*}$ ³ is the identity map.*

Proof. Take any map $f: (A_0)_{\text{perfd},*} \rightarrow (A_0)_{\text{perfd},*}$ such that the following diagram commutes

$$\begin{array}{ccc} (A_0)_{\text{perfd}} & \longleftarrow A_0 & \longrightarrow (A_0)_{\text{perfd}} \\ \downarrow c & & \downarrow c \\ (A_0)_{\text{perfd},*} & \xrightarrow{f} & (A_0)_{\text{perfd},*} \end{array}$$

where the vertical maps are canonical ones. Since $(A_0)_{\text{perfd},*}$ is also an integral perfectoid ring by Theorem 4.9 and $(A_0)_{\text{perfd}}$ is a universal integral perfectoid ring over A_0 , the composite $f \circ c$ is nothing but the canonical map $c: (A_0)_{\text{perfd}} \rightarrow (A_0)_{\text{perfd},*}$. For any $x \in (A_0)_{\text{perfd},*}$, there exists an element $a \in (A_0)_{\text{perfd}}$ such that $c(a) = px$ in $(A_0)_{\text{perfd},*}$. Then, we have $pf(x) = f(px) = f(c(a)) = c(a) = px$ in $(A_0)_{\text{perfd},*}$. Since $(A_0)_{\text{perfd},*}$ is p -torsion-free, we have $f(x) = x$ and we are done. \square

As a consequence of this lemma, we can show the following relation between $(-)^{\text{ptf}}$ and $(-)_{\text{perfd}}$.

COROLLARY 5.3. *Let A_0 be a derived p -complete algebra over some integral perfectoid ring. Assume that $(A_0)_{\text{perfd}}$ and $(A_0^{\text{ptf}})_{\text{perfd}}$ are (honest) integral perfectoid rings. Then the canonical map $(A_0)_{\text{perfd}} \rightarrow (A_0^{\text{ptf}})_{\text{perfd}}$ induces the isomorphism*

$$((A_0)_{\text{perfd}})^{\text{ptf}} \xrightarrow{\cong} (A_0^{\text{ptf}})_{\text{perfd}}.$$

Since the above isomorphism prevents confusion when $((A_0)_{\text{perfd}})^{\text{ptf}}$ is written as $(A_0)_{\text{perfd}}^{\text{ptf}}$, we use this symbol $(-)_{\text{perfd}}^{\text{ptf}}$ in the following.

Proof of Corollary 5.3. The canonical surjective map $A_0 \twoheadrightarrow A_0^{\text{ptf}}$ induces a map $(A_0)_{\text{perfd}} \rightarrow (A_0^{\text{ptf}})_{\text{perfd}}$. Since $(A_0^{\text{ptf}})_{\text{perfd}}$ is p -torsion-free by [19, Lem. A.2], we have a unique map $\varphi: ((A_0)_{\text{perfd}})^{\text{ptf}} \rightarrow (A_0^{\text{ptf}})_{\text{perfd}}$ which extends the map $(A_0)_{\text{perfd}} \rightarrow (A_0^{\text{ptf}})_{\text{perfd}}$.

² Note that any integral perfectoid ring does not contain \mathbb{Q} . So a (p -adically complete) integral perfectoid ring R is of characteristic 0 means that R contains \mathbb{Z} as a subring and $pR \neq R$. We do not assume that R is p -torsion-free.

³ Recall that an integral perfectoid ring $(A_0)_{\text{perfd}}$ has a compatible sequence $\{\varpi^{1/p^n}\}_{n \geq 0}$ of p -power roots of ϖ such that ϖ^p is a unit multiple of p in $(A_0)_{\text{perfd}}$ (see Lemma 2.2 and Remark 4.10). So $(A_0)_{\text{perfd},*}$ is the set of almost elements of the pre-perfectoid pair $((A_0)_{\text{perfd}}, \varpi)$. That is, $(A_0)_{\text{perfd},*} = \varpi^{-1/p^\infty} (A_0)_{\text{perfd}} \subseteq (A_0)_{\text{perfd}}[1/\varpi] = (A_0)_{\text{perfd}}[1/p]$.

Conversely, by [13, Prop. 2.19], the p -torsion-free quotient $((A_0)_{\text{perfd}})^{\text{ptf}}$ is also an integral perfectoid ring over A_0 and thus there exists a unique map $A_0^{\text{ptf}} \rightarrow ((A_0)_{\text{perfd}})^{\text{ptf}}$ which extends the structure map $A_0 \rightarrow ((A_0)_{\text{perfd}})^{\text{ptf}}$. The universality of perfectoidization (Theorem 2.9) shows that this map can be extended to a map $\psi: (A_0^{\text{ptf}})_{\text{perfd}} \rightarrow ((A_0)_{\text{perfd}})^{\text{ptf}}$ uniquely.

Then the A_0^{ptf} -algebra map $\varphi \circ \psi$ is the identity map because of the universality of perfectoidizations (Theorem 2.9). On the other hand, the $(A_0)_{\text{perfd}}$ -algebra map $\psi \circ \varphi$ can be extended to the $(A_0)_{\text{perfd}}$ -algebra endomorphism on $((A_0)_{\text{perfd}})^{\text{ptf}}_*$ because of the inclusion $((A_0)_{\text{perfd}})^{\text{ptf}} \subseteq (((A_0)_{\text{perfd}})^{\text{ptf}})_* \subseteq ((A_0)_{\text{perfd}})[1/p]$. Then the above rigidity lemma (Lemma 5.2) shows that $\psi \circ \varphi$ is in fact the identity map on $((A_0)_{\text{perfd}})^{\text{ptf}}$. This completes the proof. \square

In the proof of [12, Th. 4.4], Dine states that a quotient of a perfectoid Tate ring by some ideal has the perfectoidization that is isomorphic to its uniform completion. We reformulate the proof for the situation of integral perfectoid rings as follows.

THEOREM 5.4 (cf. [12]). *Let A_0 be a derived p -complete algebra over some integral perfectoid ring of characteristic 0. Set a Tate ring $A := A_0[1/p]$. Assume that the perfectoidization $(A_0)_{\text{perfd}}$ is an (honest) integral perfectoid ring and the uniform completion $A^{\widehat{u}}$ of A is a perfectoid Tate ring.*

Then $A^{\widehat{u}}$ is isomorphic to $(A_0)_{\text{perfd}}[1/p]$ as a Tate ring. In particular, $(A^{\widehat{u}})^{\circ}$ and $(A_0)_{\text{perfd},}$ are isomorphic as rings.*

Proof. If we know that $A^{\widehat{u}}$ is a perfectoid Tate ring, the same proof of [12, Th. 4.4] induces the isomorphism $A^{\widehat{u}} \cong (A_0)_{\text{perfd}}[1/p]$ by checking the universality of perfectoidizations and uniform completions as in Proposition 5.1. By Lemma 4.7, taking the set of all power-bounded elements induces the isomorphism $(A^{\widehat{u}})^{\circ} \cong (A_0)_{\text{perfd},*}$. \square

REMARK 5.5. Note that $(A^{\widehat{u}})^{\circ}$ is an integral perfectoid ring if and only if $A^{\widehat{u}}$ is a perfectoid Tate ring by Lemma 2.6. So the conditions for $A^{\widehat{u}}$ to be a perfectoid Tate ring are studied in [18, Th. 3.3.18(ii)]. For example, if A_0 is a semiperfectoid ring, then $(A^{\widehat{u}})^{\circ}$ is an integral perfectoid ring as shown in [12, Th. 4.4]. Compare Remark 5.8 below.

We recall that taking the set of all power-bounded elements is not only a topological operation but also an algebraic operation as mentioned in the next lemma. This lemma is used in the proof of Corollary 6.2.

LEMMA 5.6. *Let A_0 be a π -adically topological ring for an element $\pi \in A_0$ and let A be a Tate ring $A_0[1/\pi]$. Assume that A_0 is integral over some Noetherian ring. Then $(A^{\widehat{u}})^{\circ}$ is the same as the π -adic completion $\widehat{(A_0)_A^+}$ of the integral closure $(A_0)_A^+$ of A_0 in A .*

Proof. By Lemma 3.8, we have $(A^{\widehat{u}})^{\circ} \cong \widehat{A^{u^{\circ}}}$. Recall that $(A_0)_A^+$ becomes a ring of definition of A^u and thus $A^{u^{\circ}}$ is the complete integral closure $((A_0)_A^+)_A^*$ of $(A_0)_A^+$ in A^u by [20, Lem. 2.13(1)]. By assumption, $(A_0)_A^+$ is integral over some Noetherian ring and then $A^{u^{\circ}} = (A_0)_A^+$ by [21, Prop. 7.1]. This completes the proof. \square

The main result of this paper is the following.

THEOREM 5.7. *Let R be a ring which contains \mathbb{Z} as a subring and satisfies the following conditions:*

1. *The p -adic completion $A_0 := \widehat{R}$ of R has a map from some integral perfectoid ring.*
2. *The perfectoidization $(\widehat{R})_{\text{perfd}}$ of \widehat{R} is an (honest) integral perfectoid ring.*
3. *The p -adic completion $\widehat{C(R^{\text{ptf}})}$ of the p -root closure $C(R^{\text{ptf}})$ is an integral perfectoid ring.*

Set a Tate ring $A := A_0[1/p]$. Then there exists an isomorphism $\varphi: \widehat{C(R^{\text{ptf}})} \xrightarrow{\cong} (\widehat{R})_{\text{perfd}}^{\text{ptf}}$ which is a restriction of the unique map $(A^{\widehat{\cdot}})^{\circ} \rightarrow (A_0)_{\text{perfd},}$ taken in Proposition 5.1. In particular, $(\widehat{R})_{\text{perfd}}$ is $(p)^{1/p^\infty}$ -almost isomorphic to $\widehat{C(R^{\text{ptf}})}$. If R is p -torsion-free, we have an honest isomorphism $\widehat{C(R)} \cong (\widehat{R})_{\text{perfd}}$.*

Proof. Since $(A_0)_{\text{perfd}}$ is an integral perfectoid ring whose perfectoid element is ϖ , the ring $(A_0)_{\text{perfd}}^{\text{ptf}}$ is also an integral perfectoid ring by [13, Prop. 2.19]. In particular, $(A_0)_{\text{perfd}}^{\text{ptf}}$ is p -root closed in $(A_0)_{\text{perfd}}[1/p]$ because of the injectivity of the p -th power map $(A_0)_{\text{perfd}}^{\text{ptf}}/\varpi(A_0)_{\text{perfd}}^{\text{ptf}} \xrightarrow{a \mapsto a^p} (A_0)_{\text{perfd}}^{\text{ptf}}/p(A_0)_{\text{perfd}}^{\text{ptf}}$ (see [11, (2.1.7.1)]). The map of Tate rings $R[1/p] \rightarrow (A_0)_{\text{perfd}}[1/p]$ induced by $R \rightarrow A_0 \rightarrow (A_0)_{\text{perfd}}$ gives a unique map $\varphi: \widehat{C(R^{\text{ptf}})} \rightarrow (A_0)_{\text{perfd}}^{\text{ptf}}$ such that the following diagram commutes

$$\begin{array}{ccccc}
 R[1/p] & \longrightarrow & A_0[1/p] & \longrightarrow & (A_0)_{\text{perfd}}[1/p] \\
 \uparrow & & \uparrow & & \uparrow \\
 & & \widehat{C(R^{\text{ptf}})} & \xrightarrow{\exists! \varphi} & (A_0)_{\text{perfd}}^{\text{ptf}} \\
 \uparrow & \nearrow & \uparrow & \searrow & \uparrow \\
 C(R^{\text{ptf}}) & \longrightarrow & C(A_0^{\text{ptf}}) & \longrightarrow & (A_0)_{\text{perfd}}^{\text{ptf}} \\
 \uparrow & & \uparrow & & \uparrow \\
 R & \longrightarrow & A_0 & \longrightarrow & (A_0)_{\text{perfd}}.
 \end{array} \tag{5.1}$$

Taking the p -adic completion of $R \rightarrow C(R^{\text{ptf}})$, we have a map $A_0 \rightarrow \widehat{C(R^{\text{ptf}})}$. By assumption, $\widehat{C(R^{\text{ptf}})}$ is an integral perfectoid ring and then, there exists a unique map $(A_0)_{\text{perfd}} \rightarrow \widehat{C(R^{\text{ptf}})}$ which extends $A_0 \rightarrow \widehat{C(R^{\text{ptf}})}$ as follows.

$$\begin{array}{ccc}
 (A_0)_{\text{perfd}} & \xrightarrow{\exists! \psi} & \widehat{C(R^{\text{ptf}})} \\
 \uparrow & \nearrow & \uparrow \\
 A_0 & & \\
 \uparrow & & \\
 R & \longrightarrow & C(R^{\text{ptf}})
 \end{array} \tag{5.2}$$

Since $\widehat{C(R^{\text{ptf}})}$ is p -torsion-free, there exists a unique map $\psi: (A_0)_{\text{perfd}}^{\text{ptf}} \rightarrow \widehat{C(R^{\text{ptf}})}$ which extends $(A_0)_{\text{perfd}} \rightarrow \widehat{C(R^{\text{ptf}})}$.

Combining (5.1) and (5.2), the composite map $(A_0)_{\text{perfd}}^{\text{ptf}} \xrightarrow{\psi} \widehat{C(R^{\text{ptf}})} \xrightarrow{\varphi} (A_0)_{\text{perfd}}^{\text{ptf}}$ is an R -algebra map and thus an $A_0 = \widehat{R}$ -algebra map. Furthermore, this extends to an A_0 -algebra map of perfectoid Tate rings $(A_0)_{\text{perfd}}[1/p] \rightarrow (A_0)_{\text{perfd}}[1/p]$ and thus extends to an A_0 -algebra map $(A_0)_{\text{perfd},*} \rightarrow (A_0)_{\text{perfd},*}$ by Lemma 4.7. This must be the identity map because of Lemma 5.2 above, and then $\psi \circ \varphi$ is also the identity map.

On the other hand, consider the R -algebra map $\widehat{C(R^{\text{ptf}})} \xrightarrow{\varphi} (A_0)_{\text{perfd}}^{\text{ptf}} \xrightarrow{\psi} \widehat{C(R^{\text{ptf}})}$. Inverting p , we have a map of Tate rings $\widehat{C(R^{\text{ptf}})}[1/p] \rightarrow \widehat{C(R^{\text{ptf}})}[1/p]$ over $R[1/p]$ via $R[1/p] \hookrightarrow C(R^{\text{ptf}})[1/p] \rightarrow \widehat{C(R^{\text{ptf}})}[1/p]$. Since $C(R^{\text{ptf}}) \subseteq R[1/p]$ is contained in the integral closure \widetilde{R} of R in $R[1/p]$, the uniform completion $C(R^{\text{ptf}})[1/p]^{\widehat{u}}$ of $C(R^{\text{ptf}})[1/p]$ can be written as a Tate ring $\widehat{\widetilde{R}}[1/p]$ by Definition 3.6. Similarly, the uniform completion $R[1/p]^{\widehat{u}}$ of $R[1/p]$ also coincides with the Tate ring $\widehat{\widetilde{R}}[1/p]$. So these two uniform completions $C(R^{\text{ptf}})[1/p]^{\widehat{u}}$ and $R[1/p]^{\widehat{u}}$ are isomorphic each other via the canonical map of Tate rings $R[1/p] \rightarrow C(R^{\text{ptf}})[1/p]$.⁴ Furthermore, since $\widehat{C(R^{\text{ptf}})}$ is a p -torsion-free integral perfectoid ring, the uniform completion of $\widehat{C(R^{\text{ptf}})}[1/p]$ is isomorphic to itself by Lemma 2.8. These arguments show that


$$R[1/p]^{\widehat{u}} \xrightarrow{\cong} C(R^{\text{ptf}})[1/p]^{\widehat{u}} \xrightarrow{\cong} \widehat{C(R^{\text{ptf}})}[1/p]^{\widehat{u}} \cong \widehat{C(R^{\text{ptf}})}[1/p]. \tag{5.3}$$

This shows that the above $R[1/p]$ -algebra map of Tate rings $\widehat{C(R^{\text{ptf}})}[1/p] \rightarrow \widehat{C(R^{\text{ptf}})}[1/p]$ coincides with an $R[1/p]$ -algebra map of Tate rings $R[1/p]^{\widehat{u}} \rightarrow R[1/p]^{\widehat{u}}$ and this is the identity map by the universality of uniform completion.

We next check that $\varphi: \widehat{C(R^{\text{ptf}})} \rightarrow (A_0)_{\text{perfd}}^{\text{ptf}}$ is a restriction of the unique map $(A^{\widehat{u}})^{\circ} \rightarrow (A_0)_{\text{perfd},*}$ taken in Proposition 5.1. Similarly as above, the uniform completion of $A_0[1/p]$ is

$$A^{\widehat{u}} = A_0[1/p]^{\widehat{u}} = \widehat{R}[1/p]^{\widehat{u}} \cong R[1/p]^{\widehat{u}} \cong \widehat{C(R^{\text{ptf}})}[1/p]^{\widehat{u}}. \tag{5.4}$$

The uniform completion $\widehat{C(R^{\text{ptf}})}[1/p]^{\widehat{u}}$ of $\widehat{C(R^{\text{ptf}})}[1/p]$ has a universality which gives a unique extension $A^{\widehat{u}} \rightarrow (A_0)_{\text{perfd}}[1/p]^{\widehat{u}}$ of φ as follows (see Proposition 3.7):



⁴ Note that the identity map $R[1/p] = C(R^{\text{ptf}})[1/p]$ is only an isomorphism of rings which is not necessarily an isomorphism of topological rings. Therefore, we go back to the construction of uniform completion and prove the isomorphism $R[1/p]^{\widehat{u}} \cong C(R^{\text{ptf}})[1/p]^{\widehat{u}}$ in this way.

Since A_0 is the p -adic completion of R , $\eta: A^{\widehat{u}} \rightarrow (A_0)_{\text{perfd}}[1/p]$ is a unique extension of $A_0 \rightarrow (A_0)_{\text{perfd}}^{\text{ptf}}$. By Lemma 4.7 and the proof of Proposition 5.1, η induces the unique map $(A^{\widehat{u}})^{\circ} \rightarrow (A_0)_{\text{perfd},*}$ taken in Proposition 5.1 and this extends φ .

If R is p -torsion-free, $A_0 = \widehat{R}$ is also p -torsion-free and so is $(A_0)_{\text{perfd}}$ by [19, Lem. A.2]. Then $C(\widehat{R^{\text{ptf}}}) = \widehat{C(R)}$ is isomorphic to $(\widehat{R})_{\text{perfd}}$. □

REMARK 5.8. Let R be a (not necessarily p -adically complete) ring containing a compatible sequence of p -power roots $\{\varpi^{1/p^j}\}_{j \geq 0}$ of $\varpi \in R$ such that ϖ^p is some unit multiple of p in R . If the Frobenius map $R/pR \xrightarrow{F} R/pR$ is surjective, the p -adic completion $\widehat{C(R)}$ is an integral perfectoid ring by [11, Prop. 2.1.8]. Compare Remark 5.5 above.

In particular, any semiperfectoid ring satisfies the assumptions of Theorem 5.7 and so we have the following corollary.

COROLLARY 5.9. *Let R be a ring such that the p -adic completion \widehat{R} of R becomes a semiperfectoid ring which contains \mathbb{Z} as a subring. Then $(\widehat{R})_{\text{perfd}}^{\text{ptf}}$ is isomorphic to the p -adic completion $\widehat{C(R^{\text{ptf}})}$ of $C(R^{\text{ptf}})$. If R is p -torsion-free, we have $(\widehat{R})_{\text{perfd}} \cong \widehat{C(R)}$.*

Proof. First, the semiperfectoid ring \widehat{R} has a surjective map from some integral perfectoid ring by the definition of semiperfectoid rings (Definition 2.4). Second, The perfectoidization $(\widehat{R})_{\text{perfd}}$ of \widehat{R} is an (honest) integral perfectoid ring by [8, Cor. 7.3 and Prop. 8.5]. Finally, the p -adic completion $\widehat{C(R^{\text{ptf}})}$ of the p -root closure $C(R^{\text{ptf}})$ is an integral perfectoid ring by Remark 5.8. So R satisfies all assumptions of Theorem 5.7, and this completes the proof. □

§6. Connections between p -root Closure and Perfectoidization

We recall a mixed characteristic analog of the perfection of rings, which was introduced in [16]. This construction includes an example from [23, §4] which demonstrates a good behavior of p -root closure from the perspective of Fontaine rings.

CONSTRUCTION 6.1. Let (R_0, \mathfrak{m}, k) be a complete Noetherian local domain of mixed characteristic $(0, p)$ with perfect residue field k and let p, x_2, \dots, x_n be any system of generators of the maximal ideal \mathfrak{m} such that p, x_2, \dots, x_d forms a system of parameters of R_0 . Choose compatible sequences of p -power roots

$$\{p^{1/p^j}\}_{j \geq 0}, \{x_2^{1/p^j}\}_{j \geq 0}, \dots, \{x_n^{1/p^j}\}_{j \geq 0} \tag{6.1}$$

inside the absolute integral closure R_0^+ . Set

$$R_\infty := \bigcup_{j \geq 0} R_0[p^{1/p^j}, x_2^{1/p^j}, \dots, x_n^{1/p^j}] \subseteq R_0^+. \tag{6.2}$$

Let \widetilde{R}_∞ be the integral closure of R_∞ in $R_\infty[1/p]$ and let $\widehat{\widetilde{R}}_\infty$ (resp., \widehat{R}_∞) be the p -adic completion of \widetilde{R}_∞ (resp., R_∞).

By Cohen’s structure theorem, there exists a surjective map $S_0 \twoheadrightarrow R_0$ such that S_0 is a complete unramified regular local ring $W(k)[[t_2, \dots, t_n]]$ and t_i maps to x_i respectively. In particular, R_0 is a finite extension of a subring $T_0 := W(k)[[t_2, \dots, t_d]]$ of S_0 . Then, proceeding as in [16], we have a surjective map $S_\infty \twoheadrightarrow R_\infty$ and its p -adic completion $\widehat{S}_\infty \twoheadrightarrow \widehat{R}_\infty$. Remark that \widehat{S}_∞ is an integral perfectoid ring which has a compatible sequence of p -power roots $\{p^{1/p^j}\}_{j \geq 0}$ of p . Therefore, \widehat{R}_∞ is a semiperfectoid ring.

In [16], we show that $\widehat{C(R_\infty)}$ and \widehat{R}_∞ are integral perfectoid rings and are $(pg)^{1/p^\infty}$ -almost flat and $(pg)^{1/p^\infty}$ -almost faithful T_0 -algebra where g is a non-zero element of \widehat{R}_∞ that becomes a non-zero-divisor in \widehat{R}_∞ .

COROLLARY 6.2. *Keep the notation of Construction 6.1. Then, the integral perfectoid ring \widehat{R}_∞ is $(p)^{1/p^\infty}$ -almost isomorphic to the perfectoidization $(\widehat{R}_\infty)_{\text{perfd}}$ and its restriction to $\widehat{C(R_\infty)}$ induces an isomorphism of $\widehat{C(R_\infty)} \xrightarrow{\cong} (\widehat{R}_\infty)_{\text{perfd}}$.*

Proof. As above, \widehat{R}_∞ is a semiperfectoid ring and has a compatible sequence of p -power roots $\{p^{1/p^j}\}_{j \geq 0}$ of p . So the second statement is already proved in Corollary 5.9.

In particular, the perfectoidization $(\widehat{R}_\infty)_{\text{perfd}}$ is an honest integral perfectoid ring by [8, Prop. 8.5]. Also \widehat{R}_∞ is an integral perfectoid ring by [16, Lem. 4.2]. Note that the proof of [16, Prop. 5.9] shows that the uniform completion of $\widehat{R}_\infty[1/p]$ is isomorphic to the uniform completion $R_\infty[1/p]^{\widehat{u}}$ of $R_\infty[1/p]$. Since \widehat{R}_∞ is a semiperfectoid ring, the uniform completion $\widehat{R}_\infty[1/p]^{\widehat{u}} \cong R_\infty[1/p]^{\widehat{u}}$ is a perfectoid Tate ring by Remark 5.5. In particular, we have $(R_\infty[1/p]^{\widehat{u}})^\circ = \widehat{R}_\infty$ by Lemma 5.6 and deduce the first statement by Theorem 5.4. \square

Acknowledgment. I would like to express my gratitude to Kazuma Shimomoto for his time and effort in this research. Many thanks to Shinnosuke Ishiro for his insightful feedback on the paper's interpretation. Additionally, we thank Yves André, Dimitri Dine, Fumiharu Kato, and Kei Nakazato for reading and providing feedback on this paper. Finally, I would like to express my deepest gratitude to the anonymous referee. The benefit of the referee is everywhere. In particular, the referee gave me the idea of focusing on the “rigidity lemma” (Section 1.4) and precise advice which helps the author to make a clear and readable representation in the Introduction.

REFERENCES

- [1] D. F. Anderson, *Root closure in integral domains*, J. Algebra **79** (1982), no. 1, 51–59.
- [2] D. F. Anderson, D. E. Dobbs, and M. Roitman, *Root closure in commutative rings*, Ann. Sci. de l'Université de Clermont, Math. **95** (1990), no. 26, 1–11.
- [3] Y. André, *Le Lemme d'Abhyankar Perfectoïde*, Publ. Math. Inst. Hautes Etudes Sci. **127** (2018), no. 1, 1–70.
- [4] Y. André, *Weak functoriality of Cohen–Macaulay algebras*, J. Amer. Math. Soc. **33** (2020), no. 2, 363–380.
- [5] G. Angermüller, *On the root and integral closure of Noetherian domains of dimension one*, J. Algebra **83** (1983), no. 2, 437–441.
- [6] B. Bhatt, *Lecture notes for a class on perfectoid spaces*, 2017. <http://www-personal.umich.edu/bhattb/teaching/mat679w17/lectures.pdf>.
- [7] B. Bhatt, M. Morrow, and P. Scholze, *Integral p -adic Hodge theory*, Publ. Math. Inst. Hautes Etudes Sci. **128** (2018), no. 1, 219–397.
- [8] B. Bhatt and P. Scholze, *Prisms and prismatic cohomology*, Ann. Math. **196** (2022), no. 3, 1135–1275.
- [9] J. W. Brewer, D. L. Costa, and K. McCrimmon, *Seminormality and root closure in polynomial rings and algebraic curves*, J. Algebra **58** (1979), no. 1, 217–226.
- [10] H. Cai, S. Lee, L. Ma, K. Schwede, and K. Tucker, *Perfectoid signature, perfectoid Hilbert–Kunz multiplicity, and an application to local fundamental groups*, preprint, [arXiv:2209.04046](https://arxiv.org/abs/2209.04046), 2022.
- [11] K. Česnavičius and P. Scholze, *Purity for flat cohomology*, Ann. Math. **199** (2024), no. 1, 51–180.
- [12] D. Dine, *Topological spectrum and perfectoid Tate rings*, Algebra Number Theory **16** (2022), no. 6, 1463–1500.
- [13] G. Dospinescu, *La Conjecture du Facteur Direct (D'après Y. André et B. Bhatt)*, Astérisque **446** (2023), 141–197.

- [14] J.-M. Fontaine, *Perfectoïdes, Presque Pureté et Monodromie-Poids (D'après Peter Scholze)*, *Astérisque* **352** (2013), 509–534.
- [15] S. Ishiro, K. Nakazato, and K. Shimomoto, *Perfectoid towers and their tilts: With an application to the Étale cohomology groups of local log-regular rings*, preprint, [arXiv:2203.16400](https://arxiv.org/abs/2203.16400), 2022.
- [16] R. Ishizuka and K. Shimomoto, *A mixed characteristic analogue of the perfection of rings and its almost Cohen–Macaulay property*, preprint, [arXiv:2303.13872](https://arxiv.org/abs/2303.13872), 2023.
- [17] K. S. Kedlaya and R. Liu, *Relative p -adic Hodge theory: Foundations*, *Astérisque* **371** (2015), 1–245.
- [18] K. S. Kedlaya and R. Liu, *Relative p -adic Hodge theory, II: Imperfect period rings*, preprint, [arXiv:1602.06899](https://arxiv.org/abs/1602.06899), 2016.
- [19] L. Ma, K. Schwede, K. Tucker, J. Waldron, and J. Witaszek, *An analogue of adjoint ideals and PLT singularities in mixed characteristic*, *J. Algebraic Geom.* **31** (2022), no. 3, 497–559.
- [20] K. Nakazato and K. Shimomoto, *Finite Étale extensions of Tate rings and decompletion of perfectoid algebras*, *J. Algebra* **589** (2022), 114–158.
- [21] K. Nakazato and K. Shimomoto, *A variant of perfectoid Abhyankar’s lemma and almost Cohen–Macaulay algebras*, *Nagoya Math. J.* (2023), 1–52.
- [22] E. Reinecke, *Moduli of curves at infinite level*. Thesis, 2020. <http://deepblue.lib.umich.edu/handle/2027.42/163050>.
- [23] P. C. Roberts, *The root closure of a ring of mixed characteristic*, preprint, [arXiv:0810.0215](https://arxiv.org/abs/0810.0215), 2008.
- [24] P. Scholze, *Perfectoid spaces*, *Publ. Math. Inst. Hautes Etudes Sci.* **116** (2012), no. 1, 245–313.
- [25] P. B. Sheldon, *How changing $D[[x]]$ changes its quotient field*, *Trans. Amer. Math. Soc.* **159** (1971), 223–244.
- [26] J. J. Watkins, *Root and integral closure for $R[[X]]$* , *J. Algebra* **75** (1982), no. 1, 43–58.

Ryo Ishizuka

Department of Mathematics

Tokyo Institute of Technology

Meguro-ku, Tokyo 152-8551

Japan

ishizuka.r.ac@m.titech.ac.jp