

Molecular data infers the involvement of a marine aurantiactinomyxon in the life cycle of the myxosporean parasite *Paramyxidium giardi* (Cnidaria, Myxozoa)

Research Article

Cite this article: Rocha S, Alves Â., Antunes C, Azevedo C, Casal G (2019). Molecular data infers the involvement of a marine aurantiactinomyxon in the life cycle of the myxosporean parasite *Paramyxidium giardi* (Cnidaria, Myxozoa). *Parasitology* **146**, 1555–1563. <https://doi.org/10.1017/S0031182019000866>

Received: 4 April 2019

Revised: 27 May 2019

Accepted: 16 June 2019

First published online: 15 July 2019

Key words:

Actinospore; marine oligochaete; Myxosporea; Portugal; River Minho; SSU rDNA gene; *Tubificoides pseudogaster*

Author for correspondence:

S. Rocha, E-mail: sonia.oliveira.rocha@gmail.com

S. Rocha^{1,2}, Â. Alves¹, C. Antunes^{2,3}, C. Azevedo^{1,2} and G. Casal⁴

¹Laboratory of Cell Biology, Institute of Biomedical Sciences Abel Salazar (ICBAS), University of Porto, Rua Jorge Viterbo Ferreira no. 228, 4050-313 Porto, Portugal; ²Laboratory of Animal Pathology, Interdisciplinary Centre of Marine and Environmental Research (CIIMAR), University of Porto, Terminal de Cruzeiros de Leixões, Av. General Norton de Matos s/n, 4450-208 Matosinhos, Portugal; ³Aquamuseu do Rio Minho, Parque do Castelinho, 4920-290 Vila Nova de Cerveira, Portugal and ⁴University Institute of Health Sciences & Institute of Research and Advanced Training in Health Sciences and Technologies, CESPU, Rua Central da Gandra no. 1317, 4585-116 Gandra, Portugal

Abstract

An aurantiactinomyxon type is described from the marine naidid *Tubificoides pseudogaster* (Dahl, 1960), collected from the lower estuary of a Northern Portuguese River. This type constitutes the first of its collective group to be reported from Portugal, and only the fourth described from a marine oligochaete worldwide. Extensive morphological comparisons of new aurantiactinomyxon isolates to all known types without available molecular data are proposed to be unnecessary, given the artificiality of the usage of morphological criteria for actinosporean differentiation and the apparent strict host specificity of the group. Recognition of naidid oligochaetes as the hosts of choice for marine types of aurantiactinomyxon and other collective groups, suggests that the family Naididae played a preponderant role in the myxosporean colonization of estuarine communities. Molecular analyses of the type in study further infer its involvement in the life cycle of *Paramyxidium giardi* (Cépède, 1906) Freeman and Kristmundsson, 2018, a species that infects the kidney of European eel *Anguilla anguilla* (Linnaeus, 1758) and that has been reported globally, including from Portuguese waters. The low intraspecific difference registered in relation to Icelandic isolates of *P. giardi* (0.6%) is hypothesized to result from the emergence of genotypically different subspecies due to geographic isolation.

Introduction

Myxosporeans are parasitic cnidarians that mainly infect aquatic vertebrates and invertebrates. The involvement of annelids in the life cycle of myxosporeans was first revealed by Wolf and Markiw (1984), upon their discovery that the salmonid-infecting species *Myxobolus cerebralis* Hofer, 1903 has a life cycle stage that develops in the freshwater oligochaete *Tubifex tubifex* (Müller, 1774). Since then, several studies have provided direct and indirect evidence for a two-stage life cycle of these parasites, in which a myxosporean stage that develops in a fish alternates with an actinosporean stage that develops in an oligochaete or polychaete (see Eszterbauer *et al.*, 2015). Both life cycle stages are characterized by the production of spores necessary to achieve dissemination and transmission between vertebrate and invertebrate hosts. Spores are multicellular, comprised by two to several external valve cells that form a wall surrounding one to many polar capsules, and one to many sporoplasms. Despite sharing these main morphological features, the myxospores produced in the fish host and the actinospores produced in the annelid host exhibit highly distinct morphotypes (Lom and Dyková, 2006).

Currently, actinospore morphotypes are divided into ca. 20 collective groups; the differentiation and classification of distinct types within these groups being traditionally based on morphological criteria. The latter include the shape and size of the actinospores' body and valvular processes, shape and position of the polar capsules, and number of secondary cells in the sporoplasm (Lom *et al.*, 1997; Lom and Dyková, 2006). Actinospores of the aurantiactinomyxon collective group, specifically, are defined as having three stout, semicircularly curved, leaf-like valvular processes attached to an ellipsoidal body with protruding polar capsules at the apex and containing a sporoplasm with many secondary cells (Lom and Dyková, 2006). In the past few decades, however, the implementation of molecular tools to the study of actinosporeans has revealed that morphological criteria are unreliable for performing differential diagnoses between types (e.g. Hallett *et al.*, 2002, 2004; Eszterbauer *et al.*, 2006; Xi *et al.*, 2015; Rangel *et al.*, 2016a; Zhao *et al.*, 2016). Congruently, actinospores descriptions are currently based on the combined analysis of morphological and molecular data.

Initially, life cycle studies relied solely on data acquired through means of experimental transmission between vertebrate and invertebrate hosts. The complicated logistics and time-consuming nature of this methodology, however, hinders the achievement of results.

Moreover, its reliability has been discredited by the usage of molecular tools, which show that mixed and non-spore forming infections are common and may lead to erroneous associations (Holzer et al., 2004; Atkinson and Bartholomew, 2009; Marton and Eszterbauer, 2011; Székely et al., 2014). Consequently, life cycle studies now mainly rely on sequencing of the SSU rDNA gene for inferring correspondence between myxosporean and actinosporean stages (see Eszterbauer et al., 2015).

To date, near 60 myxosporean life cycles have been elucidated (see Eszterbauer et al., 2015). Most refer to myxobolid species (*Myxobolus*, *Henneguya* and *Thelohanellus*) that infect fish and oligochaetes in freshwater environments (e.g. Styer et al., 1991; Kent et al., 1993; Lin et al., 1999; Eszterbauer et al., 2000, 2006, 2015; Kallert et al., 2005; Marton and Eszterbauer, 2011; Székely et al., 2014), with fewer providing information pertaining to the life cycles of myxosporean genera such as *Ceratomyxa*, *Ceratonova*, *Chloromyxum*, *Ellipsomyxa*, *Gadimyxa*, *Hofferellus*, *Myxidium*, *Myxobilatus*, *Ortholinea*, *Paramyxidium*, *Parvicapsula*, *Sigmomyxa*, *Sphaerospora* and *Zschokkella* (Grossheider and Körting, 1992; Benajiba and Marques, 1993; Yokoyama et al., 1993; Bartholomew et al., 1997, 2006; Holzer et al., 2004, 2006; Koie et al., 2004, 2007, 2008; Atkinson and Bartholomew, 2009; Rangel et al., 2009, 2017; Karlsbakk and Koie, 2012). In marine environments, polychaetes appear to be the hosts of choice (Koie et al., 2004, 2007, 2008; Rangel et al., 2009, 2011, 2016b), with a few known exceptions occurring in estuarine waters, including Portuguese Rivers (Bartholomew et al., 1997, 2006; Rangel et al., 2015, 2017). Overall, few studies have aimed to investigate myxosporean biodiversity in invertebrate communities of this geographic region, with a total of 20 actinosporean types described from the raabeia (1 type), triactinomyxon (2 types), tetractinomyxon (2 types), sphaeractinomyxon (13 types), synactinomyxon (1 type) and unicapsulaticomyxon (1 type) collective groups (Székely et al., 2005; Rangel et al., 2009, 2011, 2015, 2016a, 2016b, 2017; Rocha et al., 2019a, b). Apart from the Raabeia type of Rocha et al. (2019a) and the Synactinomyxon type of Székely et al. (2005), which were described from the freshwater oligochaetes *Ilyodrilus templetoni* (Southern, 1909) and *Tubifex tubifex*, respectively, all the other above-mentioned types were found infecting marine oligochaetes or polychaetes in downstream estuaries. Both the triactinomyxon and tetractinomyxon types have been paired with their respective myxosporean counterparts, through the positive match of SSU rDNA sequences. The triactinomyxon types, one reported from the oligochaete *Limnodriloides agnes* Habrě, 1967 and the other from an undetermined species of the genus *Tectidrilus* Erséus, 1982, were molecularly identified as the species *Ortholinea auratae* Rangel et al., 2015 and *O. labracis* Rangel et al., 2017, which parasitize the urinary bladder of gilthead seabream *Sparus aurata* Linnaeus, 1758 and European seabass *Dicentrarchus labrax* (Linnaeus, 1758), respectively (Rangel et al., 2015, 2017). In turn, both tetractinomyxon types were described from marine polychaetes, *Hediste diversicolor* (Müller, 1776) and an undetermined species of *Capitella* Blainville, 1828, and molecularly shown to be involved in the life cycle of *Ellipsomyxa mugilis* (Sitjà-Bobadilla and Alvarez-Pellitero, 1993) from the gall bladder of grey mullets and *Sphaerospora dicentrarchi* Sitjà-Bobadilla and Alvarez-Pellitero, 1992 from systemic infections in *D. labrax*, respectively (Rangel et al., 2009, 2016b). To our best knowledge, further 17 myxosporean species have been reported from fishes inhabiting Portuguese rivers and estuaries, including *Paramyxidium giardi* (Cépède, 1906) Freeman and Kristmundsson, 2018 (see Eiras, 2016).

The myxosporean species *Paramyxidium giardi* was originally described as a member of the genus *Myxidium*, from infections in the kidney of European eel *Anguilla anguilla* (Linnaeus, 1758) in

France (Cépède, 1906). Over time, numerous reports of this species were performed from infections occurring in multiple organs of anguillid eels worldwide (Ishii, 1915; Fujita, 1927; Hine, 1975, 1978, 1980; Copland, 1981; Treasurer and Cox, 1997), including specimens of *A. anguilla* from Portuguese estuaries (Ventura and Paperna, 1984; Azevedo et al., 1989; Saraiva and Chubb, 1989; Saraiva and Eiras, 1996; Hermida et al., 2008). In 1993, Benajiba and Marques Rocha et al., 2019a identified the actinosporean counterpart of *P. giardi* as belonging to the aurantiactinomyxon collective group. The authors used cysts from the kidney of infected *A. anguilla* to experiment infection on samples of *Tubifex* that were reported to include non-parasitized specimens, having been collected from a previously studied area (presumably in France). Recent phylogenetic analyses corroborate the potential involvement of aurantiactinomyxon, as well as of raabeia and synactinomyxon types, in the life cycle of *Paramyxidium* spp. (Freeman and Kristmundsson, 2018; unpublished results). Nonetheless, molecular data has not been acquired that confirms the association between this genus and any of the above-mentioned collective groups.

In this study, microscopic and molecular tools are used to describe an aurantiactinomyxon type found infecting the intestinal epithelium of the marine oligochaete *Tubificoides pseudogaster* (Dahl, 1960) in the lower estuary of the River Minho, Portugal. This type constitutes the first of its collective group to be reported from Portuguese waters, and only the fourth described from a marine oligochaete. Its involvement in the life cycle of *Paramyxidium giardi* is molecularly inferred by comparison of the newly obtained SSU rDNA sequence to that available in GenBank from Icelandic isolates used in the species re-description.

Materials and methods

Sampling sites and morphological characterization

As part of a two-year myxozoan survey conducted in the River Minho between 2015 and 2016, mud was collected at low tide from the lower estuary, near the village of “Caminha” (41° 52' N, 08° 50' W). In the laboratory, oligochaetes were collected from the mud and placed individually in 12-well plates containing brackish water (15‰ salinity) and kept at 4 °C. Salinity was chosen according to the value range (15 to 40‰) generally registered for this parameter in the lower estuary throughout the year (see Dias et al., 2016). All oligochaetes were individually examined using the light microscope for the detection of actinosporean stages in internal tissues and cavities. Developmental stages and free actinospores were examined and photographed using an Olympus BX50 light microscope (Olympus, Japan). Morphometry was determined from fresh material, in accordance to Lom et al. (1997). Measurements include the mean value ± standard deviation (s.d.), range of variation, and number of measured actinospores (*n*).

DNA extraction, amplification and sequencing

Genomic DNA from infected oligochaetes was extracted using the GenElute™ Mammalian Genomic DNA Miniprep Kit (Sigma-Aldrich, St Louis, USA), following the manufacturer's instructions. The DNA was stored in 50 µL of TE buffer at -20 °C. The SSU rDNA gene of the actinospores was amplified using both universal and myxosporean-specific primers: the 5'-end by pairing the primer 18E (5'-CTG GTT GAT CCT GCC AGT-3') (Hillis and Dixon, 1991) with the primers MyxspecR (5'-CAA CAA GTT GAT AGG GCA GAA-3') (Fiala, 2006) and MYX4R (5'-CTG ACA GAT CAC TCC ACG AAC-3') (Hallett and Diamant, 2001), and the 3'-end by pairing

the primers MyxospecF (5'-TTC TGC CCT ATC AAC TTG TTG-3') (Fiala, 2006) and MYX4F (5'-GTT CGT GGA GTG ATC TGT CAG-3') (Rocha *et al.*, 2015) with the primer 18R (5'-CTA CGG AAA CCT TGT TAC G-3') (Whipps *et al.*, 2003). In turn, the 16S mitochondrial DNA (mtDNA) gene of the oligochaete host was amplified using the universal primers 16sar-L (5'-CGC CTG TTT ATC AAA AAC AT-3') and 16sbr-H (5'-CCG GTC TGA ACT CAG ATC ACG T-3') (Palumbi *et al.*, 2002). All PCRs were performed in 50 μ L reactions using 10 pmol of each primer, 10 nmol of each dNTP, 2.0 mM MgCl₂, 5 μ L 10 \times Taq polymerase buffer, 1.5 units Taq DNA polymerase (NZYTech, Lisbon, Portugal), and 3 μ L (approximately 100–150 ng) of genomic DNA. The reactions were run on a Hybaid PxE Thermocycler (Thermo Electron Corporation, Milford, Massachusetts), with initial denaturation at 95 °C for 3 min, followed by 35 cycles of 94 °C for 45 s, 53 °C for 45 s, and 72 °C for 90 s. The final elongation step was performed at 72 °C for 7 min. Five- μ L aliquots of the PCR products were electrophoresed through a 1% agarose 1 \times tris-acetate-EDTA buffer (TAE) gel stained with ethidium bromide. PCR products were purified using PuramagTM magnetic beads coated with carboxylic acid groups (MCLAB, San Francisco, California, USA). Sequencing reactions were performed with the same primers used for amplification, using a BigDye Terminator v3.1 Cycle Sequencing Kit from AppliedBiosystems (Thermo Fisher Scientific, Waltham, Massachusetts, EUA), and were run on an ABI3700 DNA analyzer from AppliedBiosystems (Thermo Fisher Scientific, Waltham, Massachusetts, EUA).

Sequence assembly and distance estimation analysis

The SSU rDNA sequence of the parasite was assembled in MEGA7 (Kumar *et al.*, 2016), by aligning the partial sequences obtained for each PCR fragment. A dataset was generated according to the highest similarity scores obtained using BLAST search, and included all known *Paramyxidium* spp., as well as closely related sequences of the aurantiactinomyxon, raabeia and synactinomyxon collective groups. All SSU rDNA sequences available for more distantly related aurantiactinomyxon types were also included in the dataset. Sequences were aligned using MAFFT version 7 available online, and distance estimation was calculated in MEGA7, with the p-distance model and all ambiguous positions removed for each sequence pair.

Results

Myxozoan survey, prevalence of infection and host identification

During this study, 577 oligochaetes were isolated from the mud collected in the lower estuary of the River Minho. Myxozoan infections were determined in a total of 9 specimens, in all of which it was possible to observe mature actinospores alongside younger developmental stages. Five specimens displayed infection in the coelomic cavity by members of the sphaeractinomyxon collective group. The remaining 4 specimens displayed actinosporean development in the intestinal epithelium: two by a type of the aurantiactinomyxon collective group, and the other two by a type of the synactinomyxon collective group. The description of the aurantiactinomyxon type is presented here; descriptions of the remaining types will be published elsewhere.

All 9 oligochaete specimens displaying myxozoan infection were identified as belonging to the marine species *Tubificoides pseudogaster* (Dahl, 1960). Identification was based on combined morphological features and molecular data of the 16S mitochondrial DNA (mtDNA). Infected oligochaetes looked alike, having

similar dimensions, bifid setae and a body tegument devoid of papillae. Sequences of the 16S mtDNA were 98–100% similar to those available for the lineage II of this species, as reported by Kvist *et al.* (2010).

Characterization of the novel *Aurantiactinomyxon* type (*Cnidaria*, *Myxozoa*)

Description: Developmental stages of the gametogamy and sporogony phases developing in the intestinal epithelium (Fig. 1). Youngest developmental stages observed were binucleated cells (Fig. 2A). The latter divided twice to produce a set of four cells that form the initial pansporocyst, comprised by two somatic cells surrounding two generative inner cells (Fig. 2B and C). Three successive mitotic divisions and one meiotic division of the generative cells produced a final set of 16 haploid cells within the pansporocysts (Fig. 2D–F). At the end of gametogamy, these cells united two by two to form 8 diploid zygotes (Fig. 2G). Subsequent mitotic divisions of the zygotes originated 8 immature actinospores, each comprised by a sporoplasmic cell connected to three valvogenic cells and three capsulogenic cells. The valvogenic cells surrounded the capsulogenic cells and appeared positioned centrally, while corresponding sporoplasmic cells leaned against the pansporocyst's wall. Sporoplasmic cells increased in size and multiplied the number of secondary cells prior to migrating into the involucre formed by the valvogenic cells (Fig. 2H–I). At the end of sporogony, 8 mature actinospores were observed within each pansporocyst. Mature actinospores subspherical, 14.4 ± 0.6 (13.6–15.9) μ m long ($n = 25$) and 12.7 ± 0.7 (11.3–13.3) μ m wide ($n = 25$), entirely embraced by 3 equally sized valvular processes, leaf-like with rounded tips, 22.4 ± 2.4 (18.1–27.6) μ m long ($n = 47$) and 15.5 ± 0.9 (13.3–17.0) μ m wide at the base ($n = 25$). Three equally sized spherical polar capsules protruding from the anterior end of the actinospores, 2.6 ± 0.3 (1.9–3.5) μ m in diameter ($n = 53$), each containing a polar tubule exhibiting 4 to 5 coils (Figs 2J–L and 3). Sporoplasm containing an undetermined number of secondary cells.

Type host: *Tubificoides pseudogaster* (Dahl, 1960) (Annelida, Oligochaeta).

Type locality: The lower estuary of the River Minho, near "Caminha" (41° 52' N, 08° 50' W), Portugal.

Site of infection: The intestinal epithelium.

Prevalence: 0.3% (2 infected in a total of 577 oligochaetes examined).

Deposition of material: Series of phototypes deposited together with a representative DNA sample in the Type Material Collection of the Laboratory of Animal Pathology, Interdisciplinary Centre of Marine and Environmental Research, Porto, Portugal, reference CIIMAR 2019.44.

Molecular data: One SSU rDNA gene sequence with a total of 2027 bp, deposited in GenBank with the accession no. MK635346. The latter is representative of two identical sequences that were separately obtained from the actinosporean developmental stages in the intestinal epithelium of two infected oligochaetes.

Remarks: Morphometry was determined from mature actinospores observed in both infected hosts. Morphometric variation between individual measurements was not recorded. Comparison to the ca. 39 known aurantiactinomyxon types without available molecular data revealed some morphometric similarity to the actinospores of the *Aurantiactinomyxon* type of McGeorge *et al.* (1997), *Aurantiactinomyxon* type of Xiao and Desser (1998) and *Aurantiactinomyxon* type 2 of Özer *et al.* (2002). Nonetheless, these types parasitize different oligochaete hosts in freshwater habitats of distant geographic locations (Canada and Scotland), further differing from the one in study in specific morphological aspects of the actinospores. Both the




Fig. 1. Light micrographs showing pansporocysts of the new type of aurantiactinomyxon developing in the intestinal epithelium of the marine oligochaete *Tubificoides pseudogaster* in the lower estuary of the River Minho.

actinospores of the Aurantiactinomyxon type of McGeorge *et al.* (1997) and of the Aurantiactinomyxon type 2 of Özer *et al.* (2002) are spherical instead of subspherical, with the first further having thinner valvular processes, and the second being generally bigger. In turn, the Aurantiactinomyxon type of Xiao and Desser (1998) has pyriform polar capsules instead of spherical, and a slightly smaller actinospore body. It should be noted that no gross similarity was determined in relation to the three marine aurantiactinomyxon types reported in the literature (Hallett *et al.*, 1997), all of which are without available molecular data. In turn, morphological comparison to the freshwater Aurantiactinomyxon type of Benajiba and Marques (1993), previously reported to be the life cycle counterpart of *P. giardi*, was strongly hampered by the paucity of morphometric data given in its original description. Molecular comparison could be performed only in relation to 20 freshwater types with SSU rDNA sequences available from either their original descriptions, follow-up studies, or from the myxosporean species determined to be their life cycle counterparts through experimental transmission. The results showed significant genetic differences of the SSU rDNA sequences in all cases (ranging from 1.4% to as high as 24.8%). In turn, a high similarity value (99.4%) was obtained in relation to the sequence of *Paramyxidium giardi* available in GenBank (MH414925) from infections in the kidney of European eel *A. anguilla*. These two sequences differ in a total of 9 nucleotide positions over 2020 bp. Overall, the parasite was more related to other members of the genus *Paramyxidium*, as well as to actinosporean types known to be phylogenetically associated with the latter, namely raabeia, synactinomyxon and aurantiactinomyxon (similarity values ranged from 98.6% to as low as 81.6%). All other myxosporean sequences available in GenBank shared less than 80% similarity with the new Aurantiactinomyxon type.

Discussion

The majority of known actinosporean stages have been described based solely on morphological criteria. The implementation of molecular tools to the study of myxosporeans, however, has revealed that species having morphologically similar actinospores may be only distantly related based on SSU rDNA sequence data. In the other way around, species having actinospores with divergent morphology may be closely related (Hallett *et al.*, 2002, 2004; Eszterbauer *et al.*, 2006; Xi *et al.*, 2015; Rangel *et al.*, 2016a; Zhao *et al.*, 2016). In the case of aurantiactinomyxon, for instance, previous studies showed that distinct morphotypes can, in fact, share the same genotype (see Hallett *et al.*, 2002; Eszterbauer *et al.*, 2006; Zhao *et al.*, 2016). Consequently, the morphological

characterization of actinospores is important but not enough for the description of new types, making the acquisition of molecular data indispensable for the identification of actinosporean stages. Our study strengthens this contention, given that without the use of molecular data, the Aurantiactinomyxon type described here could easily have been misidentified with the Aurantiactinomyxon type 1 of Negredo and Mulcahy (2001), as these types share a strong morphological resemblance (Negredo and Mulcahy, 2001; Negredo *et al.*, 2003). Despite morphological comparison to types without molecular data having revealed significant similarity to the Aurantiactinomyxon type of McGeorge *et al.* (1997), Aurantiactinomyxon type of Xiao and Desser (1998) and Aurantiactinomyxon type 2 of Özer *et al.* (2002), correspondence to any of these types could never be established with confidence, given their different hosts, habitats and geographical origins. In addition to the known unreliability of the usage of morphological criteria for type differentiation (see Hallett *et al.*, 2002; Eszterbauer *et al.*, 2006; Zhao *et al.*, 2016), aurantiactinomyxon appear to have strict host specificity, considering that no type has been molecularly proven to infect more than a single annelid species. Thus, we suggest that extensive morphological comparisons to all types that are without molecular data and have distinct oligochaete hosts are unnecessary. Actinospores of new isolates should only be identified either by DNA match or through a comprehensive morphological and biological comparison to known types sharing the same annelid host, regardless of there being high morphological similarity to other types.

Morphological features and genetic data identified the host species of the new Aurantiactinomyxon type as being *Tubificoides pseudogaster*. This annelid species is known to host several types of sphaeractinomyxon in Portuguese estuaries (this study; Rangel *et al.*, 2016a) and is further reported here to host infection by a member of the synactinomyxon collective group. Despite this significant biodiversity, concomitant infections were never observed in individual specimens. This is congruent with the very low infection rates commonly reported in wild oligochaete populations (e.g. Hallett *et al.*, 2001; Negredo and Mulcahy, 2001; Atkinson and Bartholomew, 2009). In the same manner, the low value of individual prevalence of infection determined in this study (0.3%) agrees with those previously reported for aurantiactinomyxon types (e.g. Xiao and Desser, 1998; Xi *et al.*, 2015; Zhao *et al.*, 2016; Milanin *et al.*, 2017). Actinospore development was further demonstrated to follow the pattern typically reported for members of the aurantiactinomyxon collective group; development within the host was asynchronous, but sporogony within the pansporocysts was synchronous. Multiple layers of pansporocysts were present in the intestinal epithelium,




Fig. 2. Light micrographs of some developmental stages of the new type of aurantiactinomyxon found infecting the marine oligochaete *Tubificoides pseudogaster* in the lower estuary of the River Minho. (A–F) Gametogamy phase: (A) Binucleated cell; (B) Binucleated cell in cellular division; (C) Initial pansporocyst with two involucre cells and two inner cells; (D–F) Pansporocysts with four, 14 and 16 inner cells, respectively. (G–I) Sporogony phase: (G) Pansporocyst with 8 zygotes undergoing cellular divisions; (H) Pansporocyst showing the valvogenic and capsulogenic cells located centrally (*), while the sporoplasmic cells (SC) lean against the pansporocyst wall; (I) Pansporocyst containing 8 immature actinospores. Notice the capsulogenic cells; (J, K) Mature actinospores in apical view, evidencing the 3 spherical polar capsules protruding from the anterior end, and the rounded tips of the 3 equally sized leaf-like valvular processes; (L) Mature actinospore as observed in lateral view.

partially protruding into the coelomic cavity (e.g. Marcucci *et al.*, 2009; Zhao *et al.*, 2017).

Almost all aurantiactinomyxon types reported in the literature infect oligochaete species belonging to the family Naididae Ehrenberg, 1828 [currently includes the members of the former Tubificidae (Erséus *et al.*, 2008)], be it in freshwater or marine environments. Aurantiactinomyxon pavinsis constitutes the only exception, having been widely reported to infect the freshwater lumbricid *Stygodrilus heringianus* Claparède, 1862 (see Marques, 1984; Oumouna *et al.*, 2003; Holzer *et al.*, 2004; Marcucci *et al.*, 2009). A few types have their oligochaete hosts identified only up to the genus- or family-level, but all infect naids (see Marques, 1984; Grossheider and Körting, 1992; Benajiba and Marques, 1993), while the Aurantiactinomyxon types 7 and 11 of El-Mansy *et al.* (1998), Aurantiactinomyxon type 1 of Oumouna *et al.* (2003) and Aurantiactinomyxon type 1 of Hallett *et al.* (2006) constitute the only members of the collective group that lack host information (El-Mansy *et al.*, 1998; Oumouna *et al.*, 2003; Hallett *et al.*, 2006). Up until this study, only the three aurantiactinomyxon types described by Hallett *et al.* (1997) were known to occur in the marine environment, parasitizing the naids *Limnodriloides toloensis* Erséus, 1984, *Pacifidrilus vanus* (Erséus, 1984) and *P. darvelli* (Erséus, 1984).

The species *Tubificoides pseudogaster* is, therefore, identified here as another member of the family Naididae serving as host for the aurantiactinomyxon collective group in marine environments.

To date, few other actinosporean types have been described from marine oligochaetes and all to parasitize naids. These marine oligochaete-infecting types mostly belong to the sphaeractinomyxon collective group, with few representatives belonging to endocapsa and triactinomyxon. The 13 distinct types of sphaeractinomyxon recently reported from Portuguese estuaries are hosted by *T. pseudogaster*, its congener *T. insularis* (Stephenson, 1923), and *Limnodriloides agnes* Hrabě, 1967 (see Rangel *et al.*, 2016b; Rocha *et al.*, 2019b). Worldwide, the remaining known marine sphaeractinomyxon types were also reported from naids belonging to the genera *Ainudrilus* Finogenova, 1982, *Aktedrilus* Knöllner, 1935, *Bathydrilus* Cook, 1970, *Doliadrilus* Erséus, 1984, *Heronidrilus* Erséus and Jamieson, 1981, *Limnodriloides* Pierantoni, 1903 and *Thalassodrilides* Brinkhurst and Baker, 1979 (Hallett *et al.*, 1997, 1998, 2001; Hallett and Lester, 1999). Similarly, the three marine types included within the endocapsa collective group were reported from naids of the genera *Heronidrilus*, *Heterodrilus* Pierantoni, 1902 and *Thalassodrilides* (Hallett *et al.*, 1999, 2001). Lastly, to our best knowledge there




Fig. 3. Schematic drawing depicting a mature actinospore of the new type of aurantiactinomyxon found infecting the marine oligochaete *Tubificoides pseudogaster* in the lower estuary of the River Minho, as observed in lateral (A) and apical (B) view.

are 5 marine triactinomyxon types described in the literature from naidids of the subfamily Limnodriloidinae Erséus, 1982 (including *Limnodriloides* and *Tectidrilus* spp.) (see Roubal et al., 1997; Hallett et al., 2001; Rangel et al., 2015, 2017). Overall, this suggests that the family Naididae played a preponderant role in the establishment and evolution of myxosporeans in invertebrate communities of estuarine and marine habitats.

Recently, a comprehensive co-phylogenetic study by Holzer et al. (2018) evidenced that oligochaetes and polychaetes are the most ancient hosts of myxosporeans. As such, it can be assumed that the composition, ecology, spatial distribution and susceptibility of these hosts is crucial for myxosporean evolution and diversification. The Naididae is probably the most diverse and cosmopolitan oligochaete family, being present in all biogeographic regions, including sub-Antarctic islands (Timm and Martin, 2015). Its involvement in myxosporean settlement and evolution, therefore, most likely relates to the high availability of its species in different aquatic habitats worldwide. Conjectures to other potential factors possibly influencing the reported range of myxosporean infections in naidids are difficult to be performed based on the limited information currently available for annelid-myxosporean interactions, especially in marine habitats. Future research should, therefore, aim to provide insight into the host-, parasite-, and environmental-related factors that influence myxosporean infection in annelid hosts. The acquisition of further information on this subject will, most certainly, prove to be relevant in improving our knowledge of myxosporean evolution.

Nowadays, the recognition of myxosporean life cycles is mainly based on DNA match between myxosporean and

actinosporean counterparts. This molecular approach for establishing life cycle connections relies on the typically significant interspecific variability and limited intraspecific variability of the SSU rDNA gene of myxosporeans (Køie et al., 2004; Atkinson and Bartholomew, 2009). In some cases, however, it can be difficult to distinguish between different species and different geographic genotypes of one single species. Molecularly inferred life cycles range from 99.4% to 100% similarity between the SSU rDNA sequences of myxosporean and actinosporean counterparts (see Eszterbauer et al., 2015). These values agree with the intraspecific variability generally accepted for myxosporeans, which can range between 0–3.6%, but is usually lower than 1% (Atkinson et al., 2015 and references therein). In this study, comparative molecular analysis of the SSU rDNA gene revealed the Aurantiactinomyxon type infecting *T. pseudogaster* sharing 99.4% of similarity with *P. giardi*, thus inferring its involvement in the life cycle of this myxosporean pathogen. The low genetic difference obtained (0.6%) is most likely representative of intraspecific variability due to the different geographic origin of the biological material, as previously suggested in the life cycle studies of *Myxidium truttae* Léger, 1930, *Myxobolus cultus* Yokoyama et al., 1995 and *Thelohanellus kitauei* Egusa and Nakajima, 1981 (see Holzer et al., 2004; Eszterbauer et al., 2006; Xi et al., 2013; Zhao et al., 2016). In these cases, the emergence of genotypically different subspecies was hypothesized to be the outcome of geographic isolation, with subsequent evolutionary adaptation to distinct environmental factors, and possibly different invertebrate hosts. In order to confirm our assumption of intraspecific variability between geographic isolates of *P. giardi*, a myxozoan survey on *A. anguilla* from the River Minho is being prepared and will aim to acquire molecular information from myxospores. This will allow to verify if genetic variation also occurs within a given geographic location and/or between life cycle counterparts. In this context, acquiring information from molecular markers other than the SSU rDNA gene may prove necessary in order to either confirm or refute the observations of relatedness performed here. Similarly, it would be interesting to sample marine oligochaetes from Icelandic Rivers where infections by *P. giardi* are known to occur.

The information thus far available for myxosporean life cycles demonstrates that there is no obvious correspondence between actinosporean and myxosporean morphotypes. Accordingly, phylogenetic analyses show most collective groups spanning distinct myxosporean suborders, as is the case of aurantiactinomyxon (e.g. Rocha et al., 2019b). Members of aurantiactinomyxon have been implicated in the life cycles of species belonging to the genera *Chloromyxum*, *Henneguya*, *Hoferellus*, *Myxobolus*, *Myxidium* and *Thelohanellus* (Styer et al., 1991; Grossheider and Körting, 1992; Benajiba and Marques, 1993; Lin et al., 1999; Holzer et al., 2004; Eszterbauer et al., 2006; Molnár et al., 2010). Our results further broaden the array of genera associated with the aurantiactinomyxon collective group, by molecularly inferring the involvement of the type described here in the life cycle of *Paramyxidium giardi*. This finding is corroborated by the experimental transmission study performed by Benajiba and Marques (1993), who also identified the actinosporean counterpart of *P. giardi* as an aurantiactinomyxon type, but developing in a freshwater tubificid. Accordingly, Freeman and Kristmundsson (2018) suggested freshwater oligochaetes as the most probable annelid hosts of *Paramyxidium* spp., considering that known fish hosts [European eel and Indo-Pacific tarpon *Megalops cyprinoides* (Broussonet, 1782)], were sampled from freshwater habitats. Nonetheless, both these species are migratory fish that spawn at sea producing leptocephalic larvae that migrate into inland waters to grow. Taking into account that a significant portion of their lives is spent in fresh- and brackish waters, it is possible that

infection is acquired from marine oligochaetes inhabiting estuarine habitats, as our results suggest in the case of *P. giardi*. During our myxozoan survey, molecular evidence was also found for the development of a single type of sphaeractinomyxon in both a freshwater oligochaete species of the genus *Potamotheix* Vejdovský and Mrázek, 1903 and *T. pseudogaster*, suggesting that the myxosporean counterpart of this type uses a fish host that is not entirely restricted to freshwater, most probably a grey mullet (unpublished results). Congruently, the possibility of *P. giardi* having acquired both marine and freshwater oligochaetes as hosts cannot be disregarded, considering that the ecology of the European eel allows it to contact with intrinsically distinct invertebrate communities throughout its life.

The genus *Paramyxidium* Freeman and Kristmundsson, 2018 encompasses species that, whilst being morphologically similar to *Myxidium*, differ by having histozoic development in various tissues of fish from Elopomorpha (see Freeman and Kristmundsson, 2018). The European eel is the sole representative of Elopomorpha in Portuguese waters. Thus far, it has been recognized to host four myxosporean species in this geographic location, including *P. giardi* from infections that occur in multiple organs (Ventura and Paperna, 1984; Azevedo *et al.*, 1989; Saraiva and Chubb, 1989; Saraiva and Eiras, 1996; Hermida *et al.*, 2008). The study by Freeman and Kristmundsson (2018), however, showed that *P. giardi* is not systemic; while myxospore morphology is conserved and, therefore, unreliable for identification at the species-level, the site of infection appears to be an important diagnostic feature for the group. As such, it is expected that the myxozoan survey of *A. anguilla* in Portuguese Rivers will lead to the discovery of other *Paramyxidium* spp.

Author ORCIDs.  S. Rocha, 0000-0001-6230-6142

Acknowledgements. The authors acknowledge Professor Eduardo Rocha and the Laboratory of Histology of the Institute of Biomedical Sciences, University of Porto, for the usage of the Olympus BX50 light microscope, as well as Miguel Pereira for his iconographic assistance.

Financial support. This work was financially supported by the Foundation for Science and Technology (Lisbon, Portugal) within the scope of the Ph.D. fellowship grant attributed to S. Rocha (SFRH/BD/92661/2013) through the programme QREN-POP/FSE; and the Eng^o António de Almeida Foundation (Porto, Portugal).

Conflicts of interest. None.

Ethical standards. Not applicable.

References

- Atkinson SD and Bartholomew JL (2009) Alternate spore stages of *Myxobilatus gasterostei*, a myxosporean parasite of three-spined sticklebacks (*Gasterosteus aculeatus*) and oligochaetes (*Nais communis*). *Parasitology Research* **104**, 1173–1181.
- Atkinson SD, Bartošová-Sojková P, Whipps CM and Bartholomew JL (2015) Approaches for characterising Myxozoan species. In Okamura B, Gruhl A and Bartholomew JL (eds), *Myxozoan Evolution, Ecology and Development*. Switzerland: Springer International Publishing, pp. 111–123.
- Azevedo C, Lom J and Corral L (1989) Ultrastructural aspects of *Myxidium giardi* (Myxozoa, Myxosporidia), parasite of the European eel *Anguilla Anguilla*. *Diseases of Aquatic Organisms* **6**, 55–61.
- Bartholomew JL, Whipple MJ, Stevens DG and Fryer JL (1997) The life cycle of *Ceratomyxa shasta*, a myxosporean parasite of salmonids, requires a freshwater polychaete as an alternate host. *Journal of Parasitology* **83**, 859–868.
- Bartholomew JL, Atkinson SD and Hallett SL (2006) Involvement of *Manayunkia speciosa* (Annelida: Polychaeta: Sabellidae) in the life cycle of *Parvicapsula minibicornis*, a myxozoan parasite of Pacific salmon. *Journal of Parasitology* **92**, 742–748.
- Benajiba MH and Marques A (1993) The alternation of actinomyxidial and myxosporidian sporadic forms in the development of *Myxidium giardi* (parasite of *Anguilla Anguilla*) through oligochaetes. *Bulletin of the European Association of Fish Pathologists* **13**, 100–103.
- Cépède C (1906) *Myxidium giardi* Cépède, et la prétendue immunité des Anguilles à l'égard des infections myxosporidiennes. *Comptes Rendus des Séances de la Société de Biologie et de ses Filiales* **6**, 170–173.
- Copland JW (1981) The occurrence and distribution of *Myxidium giardi* Cépède, 1906 in wild and cultured European eels, *Anguilla Anguilla* L., in England. *Journal of Fish Diseases* **4**, 231–242.
- Dias E, Morais P, Cotter AM, Antunes C and Hoffman JC (2016) Estuarine consumers utilize marine, estuarine and terrestrial organic matter and provide connectivity among these food webs. *Marine Ecology Progress Series* **554**, 21–34.
- Eiras JC (2016) Parasites of marine, freshwater and farmed fishes of Portugal: a review. *Brazilian Journal of Veterinary Parasitology* **25**, 259–278.
- El-Mansy A, Székely C and Molnár K (1998) Studies on the occurrence of actinosporidian stages of fish myxosporidians in a fish farm of Hungary, with the description of triactinomyxon, raabeia, aurantiactinomyxon and neoactinomyxon types. *Acta Veterinaria Hungarica* **46**, 259–284.
- Erséus C, Wetzel MJ and Gustavsson L (2008) ICZN rules – a farewell to Tubificidae (Annelida, Clitellata). *Zootaxa* **1744**, 66–68.
- Eszterbauer E, Székely C, Molnár K and Baska F (2000) Development of *Myxobolus bramae* (Myxosporidia: Myxobolidae) in an oligochaete alternate host, *Tubifex tubifex*. *Journal of Fish Diseases* **23**, 19–25.
- Eszterbauer E, Marton S, Rácz OZ, Letenyei M and Molnár K (2006) Morphological and genetic differences among actinosporidian stages of fish-parasitic myxosporidians (Myxozoa): difficulties of species identification. *Systematic Parasitology* **65**, 97–114.
- Eszterbauer E, Atkinson S, Diamant A, Morris D, El-Matbouli M and Hartikainen H (2015) Myxozoan life cycles: practical approaches and insights. In Okamura B, Gruhl A and Bartholomew JL (eds), *Myxozoan Evolution, Ecology and Development*. Switzerland: Springer International Publishing, pp. 175–198.
- Fiala I (2006) The phylogeny of Myxosporidia (Myxozoa) based on small subunit ribosomal RNA gene analysis. *International Journal for Parasitology* **36**, 1521–1534.
- Freeman MA and Kristmundsson A (2018) Studies of *Myxidium giardi* Cépède, 1906 infections in Icelandic eels identifies a genetically diverse clade of myxosporidians that represents the *Paramyxidium* n. g. (Myxosporidia: Myxidiidae). *Parasites and Vectors* **11**, 551.
- Fujita T (1927) Studies on Myxosporidia of Japan. *Journal of the Faculty of Agriculture of the Hokkaido Imperial University* **16**, 229–247.
- Grossheider G and Körting W (1992) First evidence that *Hoferellus cyprini* (Doflein, 1898) is transmitted by *Nais* sp. *Bulletin of the European Association of Fish Pathologists* **12**, 17–20.
- Hallett SL and Diamant A (2001) Ultrastructure and small-subunit ribosomal DNA sequence of *Henneguya lesteri* n. sp. (Myxosporidia), a parasite of sand whiting *Sillago analis* (Sillaginidae) from the coast of Queensland, Australia. *Diseases of Aquatic Organisms* **46**, 197–212.
- Hallett SL and Lester RJ (1999) Actinosporidians (Myxozoa) with four developing spores within a pansporocyst: *Tetraspora discoidea* n. g. n. sp. and *Tetraspora rotundum* n. sp. *International Journal of Parasitology* **29**, 419–427.
- Hallett SL, Erséus C and Lester RJ (1997) Actinosporidia from Hong Kong marine oligochaeta. In Morton B (ed.), *Proceedings of the Eight International Marine Biological Workshop: The Marine Flora and Fauna of Hong Kong and Southern China*. Hong Kong: Hong Kong University Press, pp. 1–7.
- Hallett SL, O'Donoghue PJ and Lester RJG (1998) Structure and development of a marine actinosporidian, *Sphaeractinomyxon ersei* n. sp. (Myxozoa). *Journal of Eukaryotic Microbiology* **45**, 142–150.
- Hallett SL, Erséus C and Lester RJ (1999) Actinosporidians (Myxozoa) from marine oligochaetes of the Great Barrier Reef. *Systematic Parasitology* **44**, 49–57.
- Hallett SL, Erséus C, O'Donoghue PJ and Lester RJG (2001) Parasite fauna of Australian marine oligochaetes. *Memoirs of the Queensland Museum* **46**, 555–576.
- Hallett SL, Atkinson SD and El-Matbouli M (2002) Molecular characterisation of two aurantiactinomyxon (Myxozoa) phenotypes reveals one genotype. *Journal of Fish Diseases* **57**, 1–14.
- Hallett SL, Atkinson SD, Erséus C and El-Matbouli M (2004) Molecular methods clarify morphometric variation in triactinomyxon spores (Myxozoa) released from different oligochaete hosts. *Systematic Parasitology* **57**, 1–14.

- Hallett SL, Atkinson SD, Erséus C and El-Matbouli M (2006) Myxozoan parasites disseminated via oligochaete worms as live food for aquarium fishes: descriptions of aurantiactinomyxon and raabeia actinospore types. *Diseases of Aquatic Organisms* **69**, 213–225.
- Hermida M, Saraiva A and Cruz C (2008) Metazoan parasite community of a European eel (*Anguilla Anguilla*) population from an estuary in Portugal. *Bulletin of the European Association of Fish Pathologists* **28**, 35–40.
- Hillis DM and Dixon MT (1991) Ribosomal DNA: molecular evolution and phylogenetic inference. *The Quarterly Review of Biology* **66**, 411–453.
- Hine PM (1975) Three new species of *Myxidium* (protozoa: Myxosporidia) parasitic in *Anguilla australis* Richardson, 1848 and *A. dieffenbachii* Gray, 1842 in New Zealand. *Journal of the Royal Society of New Zealand* **5**, 153–161.
- Hine PM (1978) Variations in the spores of *Myxidium zealandicum* Hine, 1975 (Protozoa: Myxosporidia). *New Zealand Journal of Marine and Freshwater Research* **12**, 189–195.
- Hine PM (1980) A review of some species of *Myxidium* bütschli, 1882 (Myxosporidia) from eels (*Anguilla* spp.). *The Journal of Protozoology* **27**, 260–267.
- Holzer AS, Sommerville C and Wootten R (2004) Molecular relationships and phylogeny in a community of myxosporidians and actinosporidians based on their 18S rDNA sequences. *International Journal of Parasitology* **34**, 1099–1111.
- Holzer AS, Sommerville C and Wootten R (2006) Molecular identity, phylogeny and life cycle of *Chloromyxum schurovi* Shul'man & Ieshko 2003. *Parasitology Research* **99**, 90–96.
- Holzer AS, Bartošová-Sojková P, Born-Torrijos A, Lövy A, Hartigan A and Fiala I (2018) The joint evolution of the Myxozoa and their alternate hosts: a cnidarian recipe for success and vast biodiversity. *Molecular Ecology* **27**, 1651–1666.
- Ishii S (1915) Myxosporidiosis of the Japanese eel. *Zoological Magazine of Tokyo* **27**, 372–382.
- Kallert DM, Eszterbauer E, El-Matbouli M, Erséus C and Haas W (2005) The life cycle of *Henneguya nuesslii* Schuberg & Schroder, 1905 (Myxozoa) involves a triactinomyxon-type actinospore. *Journal of Fish Diseases* **28**, 71–79.
- Karlsbakk E and Koie M (2012) The marine myxosporidian *Sigmomyxa sphaerica* (Thélohan, 1895) gen. n., comb. n. (syn. *Myxidium sphaericum*) from garfish (*Belone belone* (L.)) uses the polychaete *Nereis pelagica* L. as invertebrate host. *Parasitology Research* **110**, 211–218.
- Kent ML, Whitaker DJ and Margolis L (1993) Transmission of *Myxobolus arcticus* Pugachev and Khokhlov, 1979, a myxosporidian parasite of Pacific salmon, via a triactinomyxon from the aquatic oligochaete *Stygodrilus heringianus* (Lumbriculidae). *Canadian Journal of Zoology* **71**, 1207–1211.
- Koie M, Whipps CM and Kent ML (2004) *Ellipsomyxa gobi* (Myxozoa: Ceratomyxidae) in the common goby *Pomatoschistus microps* (Teleostei: Gobiidae) uses *Nereis* spp. (Annelida: Polychaeta) as invertebrate hosts. *Folia Parasitologica* **51**, 14–18.
- Koie M, Karlsbakk E and Nylund A (2007) A new genus *Gadimyxa* with three new species (Myxozoa, Parvicapsulidae) parasitic in marine fish (Gadidae) and the two-host life cycle of *Gadimyxa atlantica* n. sp. *Journal of Parasitology* **93**, 1459–1467.
- Koie M, Karlsbakk E and Nylund A (2008) The marine herring myxozoan *Ceratomyxa auerbachii* (Myxozoa: Ceratomyxidae) uses *Chone infundibuliformis* (Annelida: Polychaeta: Sabellidae) as invertebrate host. *Folia Parasitologica* **55**, 100–104.
- Kumar S, Stecher G and Tamura K (2016) MEGA7: molecular Evolutionary Genetics Analysis version 7.0 for bigger datasets. *Molecular Biology and Evolution* **33**, 1870–1874.
- Kvist S, Sarkar IN and Erséus C (2010) Genetic variation and phylogeny of the cosmopolitan marine genus *Tubificoides* (annelida: Clitellata: Naididae: Tubificinae). *Molecular Phylogenetics and Evolution* **57**, 687–702.
- Lin D, Hanson LA and Pote LM (1999) Small subunit ribosomal RNA sequence of *Henneguya exilis* (class Myxosporidia) identifies the actinosporidian stage from an oligochaete host. *Journal of Eukaryotic Microbiology* **46**, 66–68.
- Lom J and Dyková I (2006) Myxozoan genera: definition and notes on taxonomy, life-cycle terminology and pathogenic species. *Folia Parasitologica* **53**, 1–36.
- Lom J, McGeorge J, Feist SW, Morris D and Adams A (1997) Guidelines for the uniform characterisation of the actinosporidian stages of parasites of the phylum Myxozoa. *Diseases of Aquatic Organisms* **30**, 1–9.
- Marcucci C, Caffara M and Goretti E (2009) Occurrence of actinosporidian stages (Myxozoa) in the Nera River system (Umbria, central Italy). *Parasitology Research* **105**, 1517–1530.
- Marques A (1984) Contribution à la connaissance des Actinomyxidies: ultra-structure, cycle biologique, systématique (PhD thesis). Université des Sciences et Techniques de Languedoc, Montpellier, France, pp. 218.
- Marton S and Eszterbauer E (2011) The development of *Myxobolus pavlovskii* (Myxozoa: Myxobolidae) includes an echinactinomyxon-type actinospore. *Folia Parasitologica* **58**, 157–163.
- McGeorge J, Sommerville C and Wootten R (1997) Studies of actinosporidian myxozoan stages parasitic in oligochaetes from the sediments of a hatchery where Atlantic salmon harbour *Sphaerospora truttae* infection. *Diseases of Aquatic Organisms* **30**, 107–119.
- Milanin T, Atkinson SD, Silva MR, Alves RG, Maia AA and Adriano EA (2017) Occurrence of two novel actinospore types (Cnidaria: Myxosporidia) in Brazilian fish farms, and the creation of a novel actinospore collective group, Seisactinomyxon. *Acta Parasitologica* **62**, 121–128.
- Molnár K, Marton S, Székely C and Eszterbauer E (2010) Differentiation of *Myxobolus* spp. (Myxozoa: Myxobolidae) infecting roach (*Rutilus rutilus*) in Hungary. *Parasitology Research* **107**, 1137–1150.
- Negredo C and Mulcahy MF (2001) Actinosporidian infections in oligochaetes in a river system in southwest Ireland with descriptions of three new forms. *Diseases of Aquatic Organisms* **46**, 67–77.
- Negredo C, Dillane E and Mulcahy MF (2003) Small subunit ribosomal DNA characterization of an unidentified aurantiactinomyxon form and its oligochaete host *Tubifex ignotus*. *Diseases of Aquatic Organisms* **54**, 229–241.
- Oumouna M, Hallett SL, Hoffmann RW and El-Matbouli M (2003) Seasonal occurrence of actinosporidians (Myxozoa) and oligochaetes (Annelida) at a trout hatchery in Bavaria, Germany. *Parasitology Research* **89**, 170–184.
- Özer A, Wootten R and Shinn AP (2002) Survey of actinosporidian types (Myxozoa) belonging to seven collective groups found in a freshwater salmon farm in Northern Scotland. *Folia Parasitologica* **49**, 189–210.
- Palumbi S, Martin A, Romano S, McMillan WO, Stice L and Grabowski G (2002) *The Simple Fools Guide to PCR*, Version 2.0. Honolulu: University of Hawaii. Available at <http://palumbi.stanford.edu/SimpleFoolsMaster>.
- Rangel LF, Santos MJ, Cech G and Székely C (2009) Morphology, molecular data, and development of *Zschokkella mugilis* (Myxosporidia, Bivalvulida) in a polychaete alternate host, *Nereis diversicolor*. *Journal of Parasitology* **95**, 561–569.
- Rangel LF, Cech G, Székely C and Santos MJ (2011) A new actinospore type Unicapsulactinomyxon (Myxozoa), infecting the marine polychaete, *Diopatra neapolitana* (Polychaeta: Onuphidae) in the Aveiro estuary, Portugal. *Parasitology* **138**, 698–712.
- Rangel LF, Rocha S, Castro R, Severino R, Casal G, Azevedo C, Cavaleiro F and Santos MJ (2015) The life cycle of *Ortholinea auratae* (Myxozoa: Ortholineidae) involves an actinospore of the triactinomyxon morphotype infecting a marine oligochaete. *Parasitology Research* **114**, 2671–2678.
- Rangel LF, Castro R, Rocha S, Cech G, Casal G, Azevedo C, Székely C, Cavaleiro F and Santos MJ (2016a) Description of new types of sphaeractinomyxon actinospores (Myxozoa: Myxosporidia) from marine tubificid oligochaetes, with a discussion on the validity of the tetraspora and the endocapsa as actinospore collective group names. *Parasitology Research* **115**, 2341–2351.
- Rangel LF, Castro R, Rocha S, Severino R, Casal G, Azevedo C, Cavaleiro F and Santos MJ (2016b) Tetractinomyxon stages genetically consistent with *Sphaerospora dicentrarchi* (Myxozoa: Sphaerosporidae) found in *Capitella* sp. (Polychaeta: Capitellidae) suggest potential role of marine polychaetes in parasite's life cycle. *Parasitology* **143**, 1067–1073.
- Rangel LF, Rocha S, Casal G, Castro R, Severino R, Azevedo C, Cavaleiro F and Santos MJ (2017) Life cycle inference and phylogeny of *Ortholinea labracis* n. sp. (Myxosporidia: Ortholineidae), a parasite of the European seabass *Dicentrarchus labrax* (Teleostei: Moronidae), in a Portuguese fish farm. *Journal of Fish Diseases* **40**, 243–262.
- Rocha S, Casal G, Rangel L, Castro R, Severino R, Azevedo C and Santos MJ (2015) Ultrastructure and phylogeny of *Ceratomyxa auratae* n. sp. (Myxosporidia: Ceratomyxidae), a parasite infecting the gilthead seabream *Sparus aurata* (Teleostei: Sparidae). *Parasitology International* **64**, 305–313.
- Rocha S, Alves Á, Fernandes P, Antunes C, Azevedo C and Casal G (2019a) Description of a new actinosporidian prompts union of the raabeia and echinactinomyxon collective groups (Cnidaria, Myxozoa), due to overlap in actinospore morphology. *Diseases of Aquatic Organisms* (In press).
- Rocha S, Rangel LF, Castro R, Severino R, Azevedo C, Santos MJ and Casal G (2019b) The potential role of the sphaeractinomyxon collective

- group (Cnidaria, Myxozoa) in the life cycle of mugiliform-infecting myxobolids, with the morphological and molecular description of three new types from the oligochaete *Tubificoides insularis*. *Journal of Invertebrate Pathology* **160**, 33–42.
- Roubal FR, Hallett SL and Lester RJG** (1997) First record of triactinomyxon actinosporean in marine oligochaete. *Bulletin of the European Association of Fish Pathologists* **17**, 83–85.
- Saraiva A and Chubb JC** (1989) Preliminary observations on the parasites of *Anguilla Anguilla* (L.) from Portugal. *Bulletin of the European Association of Fish Pathologists* **9**, 88–89.
- Saraiva A and Eiras JC** (1996) Parasite community of european eel *Anguilla Anguilla* (L.) in the river Este, northern Portugal. *Research and Reviews in Parasitology* **56**, 179–183.
- Styer EL, Harrison LR and Burtle GJ** (1991) Communications: experimental production of proliferative gill disease in channel catfish exposed to a myxozoan-infected oligochaete, *Dero digitata*. *Journal of Aquatic Animal Health* **3**, 288–291.
- Székely C, Eiras JC and Eszterbauer E** (2005) Description of a new synactinomyxon type from the River Sousa, Portugal. *Diseases of Aquatic Organisms* **66**, 9–14.
- Székely C, Borkhanuddin MH, Cech G, Kelemen O and Molnár K** (2014) Life cycles of three *Myxobolus* spp. from cyprinid fishes of Lake Balaton, Hungary involve triactinomyxon-type actinospores. *Parasitology Research* **113**, 2817–2825.
- Timm T and Martin PJ** (2015) Clitellata: Oligochaeta. In Thorp JH and Rogers DC (eds), *Thorp and Covich's Freshwater Invertebrates*, 4th Edn. USA: Academic Press, pp. 529–549.
- Treasurer JW and Cox D** (1997) The occurrence of *Myxidium giardi* Cépède, in cultured eels, *Anguilla Anguilla* L., in West Scotland. *Bulletin of the European Association of Fish Pathologists* **17**, 171–173.
- Ventura MT and Paperna I** (1984) Histopathology of *Myxidium giardi* Cépède, 1906 infection in European eels, *Anguilla Anguilla* L., in Portugal. *Aquaculture* **43**, 357–368.
- Whipps CM, Adlard RD, Bryant MS, Lester RJ, Findlay V and Kent ML** (2003) First report of three *Kudoa* species from eastern Australia: *Kudoa thyrssites* from mahi mahi (*Coryphaena hippurus*), *Kudoa amamiensis* and *Kudoa minithyrssites* n. sp. from sweeper (*Pempheris ypsilychnus*). *Journal of Eukaryotic Microbiology* **50**, 215–219.
- Wolf K and Markiw ME** (1984) Biology contravenes taxonomy in the Myxozoa: new discoveries show alternation of invertebrate and vertebrate hosts. *Science* **225**, 1449–1452.
- Xi BW, Zhang JY, Xie J, Pan LK, Xu P and Ge XP** (2013) Three actinosporan types (Myxozoa) from the oligochaete *Branchiura sowerbyi* in China. *Parasitology Research* **112**, 1575–1582.
- Xi BW, Zhou ZG, Xie J, Pan LK, Yang YL and Ge XP** (2015) Morphological and molecular characterization of actinosporans infecting oligochaete *Branchiura sowerbyi* from Chinese carp ponds. *Diseases of Aquatic Organisms* **114**, 217–228.
- Xiao CX and Desser SS** (1998) Actinosporan stages of myxozoan parasites of oligochaetes from Lake Sasajewun, Algonquin Park, Ontario: new forms of echinactinomyxon, neoactinomyxon, aurantiactinomyxon, guyenotia, synactinomyxon and antonactinomyxon. *Journal of Parasitology* **84**, 1010–1019.
- Yokoyama H, Ogawa K and Wakabayashi H** (1993) Involvement of *Branchiura sowerbyi* (Oligochaeta, Annelida) in the transmission of *Hoferellus carassii* (Myxosporea, Myxozoa), the causative agent of kidney enlargement disease (KED) of goldfish *Carassius auratus*. *Fish Pathology* **28**, 135–139.
- Zhao D, Borkhanuddin MH, Wang W, Liu Y, Cech G, Zhai Y and Székely C** (2016) The life cycle of *Thelohanellus kitauei* (Myxozoa: Myxosporea) infecting common carp (*Cyprinus carpio*) involves aurantiactinomyxon in *Branchiura sowerbyi*. *Parasitology Research* **115**, 4317–4325.
- Zhao DD, Zhai YH, Liu Y, Wang SJ and Gu ZM** (2017) Involvement of aurantiactinomyxon in the life cycle of *Thelohanellus testudineus* (Cnidaria: Myxosporea) from allogynogenetic gibel carp *Carassius auratus gibelio*, with morphological, ultrastructural, and molecular analysis. *Parasitology Research* **116**, 2449–2456.