

New taxa and revised stratigraphic distribution of the crinoid fauna from Anticosti Island, Québec, Canada (Late Ordovician-early Silurian)

William I. Ausich¹ and Mario E. Cournoyer²

¹School of Earth Sciences, 155 South Oval Mall, The Ohio State University, Columbus, OH 43210 USA (ausich.1@osu.edu)

²Musée de paléontologie et de l'évolution, 541 Congrégation Street, Montréal, Québec Canada H3K 2J1 (paleovision@videotron.ca)

Abstract.—End-Ordovician extinctions had a profound effect on shallow-water benthic communities, including the Crinoidea. Further, recovery after the extinctions resulted in a macroevolutionary turnover in crinoid faunas. Anticosti Island is the most complete Ordovician-Silurian boundary section recording shallow-water habitats. Both new taxa and changes in Anticosti Island stratigraphic nomenclature are addressed herein. New taxa include *Becsciecrinus groulxi* n. sp., *Bucucrinus isotaloi* n. sp., *Jovacrinus clarki* n. sp., *Plicodendrocrinus petryki* n. sp., *Plicodendrocrinus martini* n. sp., *Thalamocrinus daoustae* n. sp., and *Lateranicrinus saintlaurenti* n. gen. n. sp. The status of *Xenocrinus rubus* as a boundary-crossing taxon is confirmed, range extensions of several taxa are documented, and the distribution of crinoids with the revised stratigraphic nomenclature is documented.

UUID: <http://zoobank.org/19613a44-ec69-47d7-88ab-fcf88ba771f0>.

Introduction

The Ordovician through early Silurian was a significant interval in the evolutionary history of the Crinoidea. In general, echinoderms, including crinoids, experienced an adaptive radiation during the Great Ordovician Biodiversification Event (GOBE) (Guensburg and Sprinkle 2000; Sprinkle and Guensburg, 2004; Webby et al., 2004; Peters and Ausich, 2008; Lefebvre et al., 2013). The most significant crinoid diversification began in earnest during the Sandbian and continued through the Katian, after which crinoids experienced a mass extinction event at the onset of the Hirnantian glaciation (Peters and Ausich, 2008). Profound turnover for all echinoderms occurred during the Late Ordovician, and this event also resulted in a macroevolutionary turnover for crinoid faunas (Baumiller, 1993; Ausich et al., 1994; Ausich and Deline, 2012; Deline et al., 2012). Late Ordovician extinctions ended the early Paleozoic Crinoid Evolutionary Fauna (CEF), and preferential radiation of different crinoid clades during the post-Ordovician radiation yielded the middle Paleozoic CEF. Ordovician faunas were typically dominated by diplobathrid camerate, disparid, and hybocrinid crinoids, which were replaced by faunas typically dominated by monobathrid camerate, cladid, and flexible crinoids.

This extinction occurred during the Late Ordovician glacial epoch, with global climate change and habitat destruction resulting from the global ocean regression during this glaciation. Various drivers for this climate change have been proposed, among others as a consequence of plate tectonic history (Kump et al., 1999; Herrmann et al., 2004; Nardin et al., 2011), high cosmic ray flux (Shaviv and Veizer, 2003), extensive volcanic eruptions that resulted in a large igneous province (Buggisch et al., 2010), massive weathering of volcanic rocks

(Lefebvre et al., 2010), and ocean euxinia (Zou et al., 2018). Whatever the causes, the Late Ordovician mass extinction is generally regarded as the second-most significant Phanerozoic mass extinction (Sepkoski, 1996).

Until recently, the general absence of echinoderm fossil occurrences from the latest Ordovician (Hirnantian)–earliest Silurian (Llandovery) interval has prevented a nuanced understanding of this critical interval of pelmatozoan evolution. Recent discovery of Hirnantian and Rhuddanian crinoids (e.g., Eckert, 1984, 1990; Ausich, 1984a, b, 1985, 1986a, b, c, 1987; Donovan, 1993; Fearnhead and Donovan, 2007; Ausich and Copper, 2010; Ausich et al., 2015b; Ausich and Wilson, 2016) has improved our understanding of crinoids through this interval. However, it was not until the description of Anticosti Island crinoids was completed that sufficient data were available to evaluate crinoid evolutionary dynamics more critically through this biodiversity crisis (Peters and Ausich, 2008; Ausich and Deline, 2012). The extreme importance of Anticosti Island crinoid faunas necessitates the present paper, which describes new material and updates the known stratigraphic distribution of Ordovician to Silurian crinoids from Anticosti Island.

Geography and stratigraphic occurrences of Anticosti Island crinoids

Anticosti Island (Fig. 1) is in the Gulf of St. Lawrence north of the Gaspé Peninsula in Québec, Canada. The strata on Anticosti Island include a nearly complete Katian (Ordovician) through Llandovery (Silurian) stratigraphic section (Fig. 2), which was described initially by Richardson (1857). Perhaps because the section is so complete compared to other regions in North America, the position of Ordovician-Silurian boundary on Anticosti Island has not been settled (e.g., Schuchert and Twenhofel,




Figure 1. Geologic Map Anticosti Island, Québec, Canada.

1910; Twenhofel, 1928; Bolton, 1961; Cocks and Copper, 1981; Petryk, 1981; Long and Copper, 1987; Barnes, 1988; Copper, 1989, 2001; Desrochers et al., 2008, 2010; Jin and Copper, 2008; Copper et al., 2013; and others). Most recently, Copper et al. (2013) regarded the top of the Ordovician as the top of the grainstone facies immediately above the Laframboise reefs.

Because of facies changes in the Ordovician-Silurian boundary interval from east to west across the island and the lack of many key, cosmopolitan, biostratigraphically important fossils, stratigraphic uncertainty has existed. A comprehensive synthesis of fossil stratigraphic distribution, facies, and sequence stratigraphy is the basis for the stratigraphic revisions proposed in Copper et al. (2013). For the present study, the significant changes include recognition that some Late Ordovician lithostratigraphic units do not extend across the island from east to west, new lithostratigraphic names have been applied for Hirnantian and Llandovery strata in western Anticosti Island, and the base of the Katian-Hirnantian boundary was changed.

Revised stratigraphic distribution of Anticosti crinoids

As noted above, revisions to understanding of Hirnantian and Llandovery chronostratigraphy (Fig. 2) of Anticosti Island necessitates reconsideration of the stratigraphic distribution of crinoids, which is listed in Supplementary Tables 1–5. Several changes are noteworthy. One new genus and seven new species are described, including the second species of *Becsciecrinus* and *Bucucrinus*. Hirnantian taxa from the west end of the island previously described from the Grindstone and Velleda members of Ellis Bay are reassigned to new members (Copper et al., 2013) (Supplementary Table 3); and with division of the former Jupiter Formation into the Menier and Jupiter (revised) formations,

crinoid distributions are revised accordingly (Supplementary Tables 4 and 5).

Based on the inferred stratigraphic occurrence of *Gaurocrinus fimbriatus* (Billings, 1859) and *Plicodendrocrinus observationensis* Ausich and Copper, 2010, the range of both of these taxa are extended higher in the Katian to either the Schmitt Creek or Grindstone member of the Vauréal Formation. The existence of *Xenocrinus rubus* Ausich and Copper, 2010 in Silurian strata along the Jupiter River confirms that this species crossed the Ordovician-Silurian boundary, which verifies three boundary-crossing species: *Dendrocrinus leptos* Ausich and Copper, 2010, *Protaxocrinus paraios* Ausich and Copper, 2010, and *X. rubus*. Boundary-crossing genera are *Dendrocrinus*, *Eomyelodactylus*, *Plicodendrocrinus*, *Protaxocrinus*, and *Xenocrinus*. With the exception of *Eomyelodactylus*, all of these boundary-crossing taxa belonged to major clades that would dominate the Middle Paleozoic CEF (Baumiller, 1993; Ausich et al., 1994; Ausich and Deline, 2012). Further, despite the fact that disparids in general were not a typically dominant clade after the Ordovician, the myelodactylid clade radiated during the Silurian.

Plicodendrocrinus petryki n. sp. is the first Silurian species of *Plicodendrocrinus* described from North America. Stratigraphic range extensions include *Becsciecrinus*, which is now known from the Fox Point Member of the Becscie Formation (Rhuddanian) to the Ferrum Member of the Jupiter Formation (Telychian); *Dimerocrinites elegans* Springer, 1928, which is now known from the Cybèle through Pavillon members of the Jupiter Formation (Telychian); and *Aetocrinus gracilis* Ausich and Copper, 2010 and *Jovacrinus*, which are now known from the Ferrum and Pavillon members of the Jupiter Formation (Telychian).

System	Series	Stage	Formation	Member	
Silurian	Llandoverly	Telychian	Chicotte		
			Jupiter	Pavillon	
				Ferrum	
				Cybèle	
			Aeronian	Richardson	
				East Point	
		Goéland			
		Gun River		Macgilvray	
				Sandtop	
				Innommée	
		Rhuddanian	Merrimack		
			Becscie	Chabot	
				Fox Point	
Ordovician	Late	Hirnantian	Ellis Bay	Laframboise	
				Lousy Cove	
				Parastro	Prinsta
				Junction Cliff	
				Fraise	
				Katian	Vauréal
		Grindstone			
		Schmitt Creek			
		Mill Bay			
		Battery			
		Homard			
		Tower			
		Easton			
		Lavache			
	west	east			

Figure 2. Ordovician to Silurian stratigraphic column for Anticosti Island (modified from Copper et al., 2013).

Paleoecological associations

Supplementary Tables 1–5 list the crinoids in each stratigraphic unit; but in most cases, crinoids are typically preserved either individually or with multiple specimens of the same species. Multiple species are rarely preserved on the same bedding surface. Striking exceptions are the crinoid slabs deposited as MPEP718.1 and MPEP718.2 from the Pavillon Member of the Jupiter Formation (Telychian) at Brisants Jumpers (Locality 6) (Fig. 3). These slabs have innumerable specimens present with three dominant taxa: *Dimerocrinites elegans*; *Aetocrinus gracilis*; and *Eomyelodactylus springeri* Ausich and Copper, 2010. Poorly preserved specimens of additional taxa may also be present. *Dimerocrinites elegans* is the larger of the three dominant species and is most evident on initial inspection

of the material (the large pluricolumnals also belong to *Dimerocrinites elegans*). However, *A. gracilis* is the most abundant species, with much of the smaller, articulated or partially articulated crinoidal material belonging to *A. gracilis*. *Eomyelodactylus springeri* is rare but present on both MPEP718.1 and MPEP718.2.

The presence of gastropods on the tegmen or on other locations on crinoids is a well-known association in Paleozoic faunas, although the exact ecological interaction between gastropods and crinoids has engendered much discussion (e.g., Bowsher, 1955; Meyer and Ausich, 1983; Baumiller, 1990, 2002; Baumiller and Gahn, 2003; Gahn and Baumiller, 2003, 2006; Baumiller et al., 2004). MPEP1138.6 is a specimen of *Fibrocrinus phragmos* Ausich and Copper (2010) from the East Point Member of the Menier Formation (Aeronian). It is a partially disarticulated crown that is broken longitudinally



Figure 3. Multispecies bedding surface from Locality 6 (Pavillon Member of the Jupiter Formation); MPEP718.1. Three dominant taxa are present on this bedding surface: (1) *Dimerocrinites elegans* Springer, 1928 (see Fig. 5.3); (2) *Aetocrinus gracilis* Ausich and Copper, 2010 (see Fig. 13.5); and (3) *Eomyelodactylus springeri* Ausich and Copper, 2010 (see Fig. 12.2). Scale bar 5 mm.



Figure 4. Paleocological associations with Anticosti Island crinoids. (1) Longitudinal section through a slightly disarticulated crown of *Fibrocrinus phragmos* Ausich and Copper, 2010; immediately above the tegmen and within the arms is the cross section through a gastropod; East Point Member, Menier Formation; MPEP1138.6; (2) crinoid pluricolumnal completely encased by favositid coral; MPEP308.11. All scale bars 5 mm.

parallel to the oral-aboral axis of the specimen. Present on the tegmen and otherwise concealed within the arms is a gastropod

(Fig. 4.1). This is the first such association recorded in crinoids from Anticosti Island.

A second biotic interaction not previously known on Anticosti Island (Locality 2) is a tabulate coral as an epizoan on a crinoid pluricolumnal (Fig. 4.2). Both examples of this interaction have the coral growing around the pluricolumnal. In MPEP308.11, the coral grew over one end of the pluricolumnal, but the other end of the column is exposed. In MPEP308.10, both ends of the column are exposed. Because the coral colony grew all around the column, it is probable that this association only occurred when the crinoid was still erect. It is possible that MPEP308.11 is an indication that the host crinoid had lost its crown, but the column remained erect (see Ausich and Baumiller, 1993; Oji and Amemija, 1998).

Materials and methods

Localities and specimen preparation.—New specimens reported here are from several new, small collections not available to Ausich and Copper (2010). One new collection is housed in the Royal Ontario Museum (ROMIP), Toronto, Canada. Seven additional collections have been donated to the Musée de paléontologie et l'évolution, Montréal, Canada (MPEP), including those made by David Clark, Nathalie Daoust/Mario Cournoyer, Pierre Groulx, Phillip Isotalo, Markus Martin, Daniel Saint-Laurent, and the Geological Survey of Canada (collected by Allen Petryk). Unfortunately, the location and stratigraphic data for these new collections is variable from precise locations to “along the Jupiter River.” Most of the more recent of these collections have well-documented geographic locations, and the stratigraphic positions of collections are precise or can be inferred with reasonable certainty. Unfortunately, this is not the case for all collections.

The key to the Allen Petryk field numbers did not accompany his specimens. However, when acquired, his collections were still in field bags. As far as possible, the stratigraphic intervals of his collections were inferred by the associated fauna (Supplemental Appendix).

In many cases, insufficient locality and stratigraphic data render a specimen useless for scientific study. However, Anticosti Island is a relatively small island with known stratigraphy and a well-documented fossil record. So, in nearly all cases, stratigraphic occurrences can be established or interpreted with reasonable certainty to at least the stage level. The Supplemental Appendix lists known or inferred geographic and stratigraphic information of the collections studied herein.

All measurements are in mm; * indicates an incomplete measurement or a crushed specimen; and measurement abbreviations are as follows: ACH, aboral cup height; ACW, aboral cup width; dACW, distal aboral cup height; pACW, proximal aboral cup height; AH, arm height; ASH, anal sac height; BH, basal plate height; BW, basal plate width; CaH, calyx height; CaW, maximum calyx width; CoH, column height; CrH, crown height; IH, infrabasal plate height; iIH, infrabasal cirlet height in interradial position; rIH, infrabasal cirlet height in radial position; IW, infrabasal plate cirlet width; RH, radial plate height; and RW, radial plate width. Specimens were photographed either after whitening with NH_4Cl , immersed in alcohol, or with no treatment.

Additional discussion of new occurrence data on Anticosti Island crinoids is in the Supplementary Data.



Figure 5. Anticosti Island diplobathrid camerates. (1, 2, 4) *Gaucrocrinus fimbriatus* (Billings, 1859), MPEP 476.9; (1) B ray lateral view of calyx with arms mostly missing; (2) oral view of tegmen, not small, multiplated tegmen and relatively short anal sac at CD-interray side of tegmen; (4) CD interray view of theca and proximal-most brachials. (3) Lateral view of *Dimerocrinites elegans* Springer, 1928; see Figure 3.1; Locality 6 (Pavillon Member of the Jupiter Formation); MPEP718.1. (5, 6) *Bucucrinus isotaloi* n. sp.; holotype, ROMIP 54202; (5) oral view of cross section through proximal portion of free arms that compare to (6); at arrows, note webbing that connects proximal arms; (6) lateral view of calyx and proximal arms; arrow denotes level of webbing in (5). All scale bars 5.0 mm.

Repositories and institutional abbreviations.—New material is deposited in the Royal Ontario Museum (ROMIP), Toronto, Canada; and Musée de paléontologie et de l'évolution, Montréal, Canada (MPEP). Previously published specimens cited are from the Geological Survey of Canada, Ottawa (GSC); the University of Alberta (UA); and the Yale University Peabody Museum (YPM).

Systematic paleontology

The classification used herein follows the phylogeny-based revision of crinoid higher taxa by Wright et al. (2017). Recent phylogenetic analyses find disparids to be sister to the Cladida, with hybocrinids nested within the Cladida and sister to the

porocrinids (Ausich et al., 2015a; Wright, 2017). At higher taxonomic levels, both disparids and hybocrinids belong to the infraclass Inadunata, which is placed within the newly resurrected subclass Pentacrinoidea (Wright et al., 2017). Morphological terminology follows Ubaghs (1978) and Ausich et al. (1999), with modifications from Ausich (1998), Ausich et al. (2015a), and Webster and Maples (2008, fig. 10.2; for brachial plate terminology). Plating of interrays is given by the number of plates in each range from the proximal-most plate to the last range before the tegmen, if known. In the posterior interray, the primanal is indicated by “P,” and the first interradian plate in regular interrays is indicated by “1.” “?” indicates that more distal plating is unknown.

Class Crinoidea Miller, 1821

Subclass Camerata Wachsmuth and Springer, 1885

Infraclass Eucamerata Cole, 2017

Order Diplobathrida Moore and Laudon, 1943

Family Reteocrinidae Wachsmuth and Springer, 1885

Genus *Gaurocrinus* Miller, 1883

Type species.—*Gaurocrinus nealli* Hall, 1866.

Gaurocrinus fimbriatus (Billings, 1859)

Figure 5.1, 5.2, 5.4

Holotype.—GSC 1994.

Occurrence.—*Gaurocrinus fimbriatus* is known with certainty from the Easton Member, Vauréal Formation (Katian); from several localities on Anticosti Island, Québec, Canada (Ausich and Copper, 2010). New specimens are from Locality 16, which are inferred to be from either the Schmitt Creek or Grindstone members of the Vauréal Formation (Katian).

Materials.—MPEP476.4a, MPEP476.4b, MPEP476.7, and MPEP476.9.

Remarks.—Several specimens of *Gaurocrinus fimbriatus* (Billings, 1859) were reported by Ausich and Copper (2010) with all of the verifiable occurrences from the Easton Member of the Vauréal Formation. Four new, well-preserved specimens are reported herein from the Petryk Collection. MPEP476.9 (Fig. 5.1, 5.2, 5.4) is a particularly instructive specimen because, unlike most other specimens, this has not been flattened during preservation. MPEP476.9 has variability in the number of fixed primibrachials and corresponding differences in the number of fixed intrabrachial plates within half-rays. Further, this specimen confirms the morphology of the tegmen previously known only from one specimen (GSC 12661 h, Ausich and Copper, 2010, pl. 1, fig. 3). The relatively low tegmen is composed of innumerable small plates, and the anal tube is a relatively short, conical structure (Fig. 5.2, 5.4).

New specimens of *G. fimbriatus* co-occur with *Plicodendrocrinus observationensis* in the Petryk collection (Locality 16). The only information accompanying these specimens is “Ordovician.” *Gaurocrinus fimbriatus* was known only from

the Easton Member of the Vauréal Formation, and *P. observationensis* was known from the Easton and Tower Members of the Vauréal Formation (Ausich and Copper, 2010). However, these new crinoids co-occur with aulacerid stromatoporoids, which suggest they are also from higher in the Katian in either the Schmitt Creek or Grindstone member of the Vauréal Formation.

Family Rhodocrinitidae Roemer, 1855

Genus *Bucucrinus* Ausich and Copper, 2010

Type species.—*Bucucrinus saccus* Ausich and Copper, 2010.

Diagnosis.—Rhodocrinitid with high bowl-shaped calyx, only distal corners or all of infrabasal plates visible in lateral view, 1-2 plating in proximal part of regular interrays, P-3 plating in proximal part of CD interray, median ray ridges defined by convexity of ray plates, no anitaxial ridge, primanal heptagonal, no fixed pinnules, 30–40 free arms, brachials cuneate (modified from Ausich and Copper, 2010).

Bucucrinus isotaloi new species

Figure 5.5, 5.6

Holotype.—ROMIP 54202.

Diagnosis.—*Bucucrinus* with full height of infrabasal plates visible in side view, basal plates approximately as high as wide, radial plates approximately as high as wide, interradian plates slightly convex, first interradian plate as high as wide and larger than radial plates, at least 12 ranges of interradian plates, 30 free arms, proximal brachials with typical cuneate uniserial appearance but connected by presumably flexible plating connecting adjacent arms, ~13 typically appearing cuneate uniserial brachial actually a free brachial, free brachials 2.0–4.0 times wider than high.

Occurrence.—Llandoverly (Aeronian or Telychian) from strata along the Jupiter River; Anticosti Island, Québec, Canada (Locality 31).

Description.—Calyx, large; high bowl shaped; broad ray ridge defined by convex ray plates (Fig. 5.6); interradian plates smooth, slightly convex.

Infrabasal circlet very low, visible in side view; 4.0% of calyx height; infrabasal concavity. Basal circlet height estimated to be ~11.0% of calyx height; basal plates five, equal in size; approximately as wide as high. Radial circlet estimated to be ~14.0% of calyx height; radial plates five, hexagonal, approximately as high as wide. Radial circlet interrupted in all interrays by sutural contact of basal plates with first interradian plates.

Normal interrays in contact with tegmen, plate sculpturing as described above; first interradian plate heptagonal, as high as wide, larger than radial plates and first primibrachials; in sutural contact with basal plate in all interrays. Second range typically with two plates; plating 1-2-3-2-3-2-2-1-2-2-2-. Above what appears to be the distal edge of the calyx (brachials assume appearance of typical free uniserial brachials) interray plating

forms what may be a flexible plated membrane connecting adjacent proximal arms that extends through at least the 12th of the “apparent free brachials” (Fig. 5.5).

Primanal not preserved.

First primibrachial hexagonal, approximately as high as wide, approximately the same size as radial plates and primaxil; second primibrachial axillary, septagonal. Second secundibrachial axillary, second tertibrachial axillary, and fourth tertibrachial axillary on medial tertibrachitaxes. Fields of intrabrachial plates between secundibrachials, tertibrachials, and quartibrachials medially within brachitaxes. Apparently flexible plating in inter- and intrarays connecting adjacent proximal arms to approximately the fourth tertibrachial medially and the 12th secundibrachial brachial abmedially. Tegmen not preserved.

Free arms 30, atomous as known, pinnulate. Brachials cuneate, uniserial, aborally convex, ~2.0–4.0 times wider than high. Arms free above flexible plating at approximately the 13th cuneate uniserial brachial.

Column not preserved.

Etymology.—The species name recognizes Phillip Isotalo, who donated specimens for this study.

Materials.—The holotype (ROMIP 54202) is the only known specimen.

Measurements.—CrH, 69.5*; CaH, 43.9; CaW, 40*.

Remarks.—The flexible plating between the proximal cuneate brachials is diagnostic for *B. isotaloi* n. sp. This is an unusual feature that is similar to crinoids such as *Scyphocrinites* (Silurian–Devonian, monobathrid Eucamerata), and may be sufficient to define a new genus; however, this is not considered advisable until more is known about the complete morphology of both *B. saccus* and *B. isotaloi* n. sp. Supplemental Table 6 lists the diagnostic features of the two species of *Bucucrinus*.

Data with the specimen indicate that it is from Silurian strata along the Jupiter River, Anticosti Island, which indicates a Llandovery age. *Bucucrinus saccus* was collected from the Jupiter River Rock Pool cliff section down river from six-mile cabin. Based on the matrix, it is plausible that the holotype of *B. isotaloi* n. sp. is also from either the Menier or Jupiter formation, but we cannot speculate on the member or whether it is from Aeronian or Telychian strata

Family Dimerocrinitidae von Zittel, 1879

Genus *Dimerocrinites* Phillips, 1839

Type species.—*Dimerocrinites decadactylus* Roemer, 1855.

Dimerocrinites elegans Springer, 1928

Figures 3.1, 5.3

Holotype.—YPM 20483.

Materials.—As noted in Ausich and Copper (2010), many specimens are known. Numerous additional specimens are

present in the new material on MPEP718.1–MPEP718.4; multiple specimens are on MPEP718.1 and MPEP718.2; MPEP718.3 and MPEP718.4 are single specimens.

Occurrence.—Cybèle, Ferrum, and Pavillon members, Jupiter Formation (Llandovery, Telychian); Anticosti Island, Québec, Canada.

Remarks.—Numerous new specimens of *Dimerocrinites elegans* Springer, 1928 are from Brisants Jumpers (Pavillon Member, Jupiter Formation, MPEP718) (Locality 6). As noted above, specimens MPEP718.1 and MPEP718.2 are remarkable slabs with numerous complete and partial specimens of *D. elegans* (Figs. 3.1, 5.3) as well as *Aetocrinus gracilis* and *Eomyelodactylus springeri*.

Genus *Becsciecrinus* Ausich and Copper, 2010

Type species.—*Becsciecrinus adonis* Ausich and Copper, 2010.

Becsciecrinus groulxi new species

Figures 6, 7.1

Holotype.—ROMIP 54198a.

Diagnosis.—*Becsciecrinus* with a medium cone-shaped calyx; plate sculpturing with radial ridges and irregular pitting; basal concavity absent; basal cirlet ~30% of calyx height; radial cirlet ~25% of calyx height, radial plates approximately as wide as high; first interradial plate higher than wide, approximately same size as radial plates, larger than first primibrachial; primanal higher than wide, larger than radial plates; column pattern N3231323.

Occurrence.—Cybèle Member, Jupiter Formation (Llandovery, Telychian) at Jupiter-la-Mer, Anticosti Island, Québec, Canada (Locality 7); and from an unknown stratigraphic horizon along the Jupiter River (Locality 28).

Description.—Calyx, small, medium cone-shaped (Fig. 7.1); wide, convex ridge (depressed sutures) from radials to basals and to median ray ridges defined by very convex fixed brachials (Fig. 6.2), basals with large proximal node, otherwise fine pitted plate sculpturing.

Infrabasal cirlet either not visible in interradial position (covered by basal node) or visible radially, where visible ~8.0% of calyx height; without basal concavity. Basal cirlet ~30% of calyx height; basal plates five, hexagonal, more than twice as high as wide, CD basal plate larger than other basal plates. Radial cirlet ~25% of calyx height, interrupted in posterior; radial plates five, heptagonal, approximately as wide as high.

Normal interrays in contact with tegmen, depressed, first interradial hexagonal, higher than wide, approximately same size as radial plates, larger than first primibrachial. Second range with two plates; plating incompletely known.

Primanal hexagonal, higher than wide, anitaxial ridge from primanal onto tegmen (Fig. 6.2, 6.5), larger than radial plates,




Figure 6. *Becciecrinus groulxi* n. sp. (1–6) (1) Holotype and paratype on a single bedding surface, ROMIP 54198; (2, 5, 6) CD interray, lateral view of calyx and proximal column; holotype, ROMIP 54198a; (2) whitened; (5) no whitening; (6) photographed under alcohol; (3, 4) crown with long length of column, paratype, MPEP719.1; (4) enlargement, specimen silicified and coarsely preserved. Scale bars 5.0 mm, unless otherwise labeled.




Figure 7. Camera lucida plate diagrams. (1) *Becciecrinus groulxi* n. sp. (ROMIP 54198a) (scale bar 1.0 mm); (2) *Lateranicrinus saintlaurenti* n. gen. n. sp. Black shading, radial plates; stippled shading, interradial plates; P, primanal; and dotted lines, ray ridges along rays and antiaxial ridge in CD interray.

interrupts the radial circlet; plating in CD interray incompletely known, three plates above primanal; CD interray in contact with tegmen.

First primibrachial hexagonal, approximately same size as radial plates and primaxil; second primibrachial axillary. Small intrabrachial plates may be present between proximal secundibrachials. Fixed brachial plates ~37% of calyx height.

Tegmen unknown.

Free arms 10, high, branch once, pinnulate. Brachials rectangular uniserial, higher than wide; first secundibrachials in lateral contact medially with adjacent first secundibrachial; second secundibrachial with abmedial pinnule, third secundibrachial nonaxillary, fourth secundibrachial axillary. Distal tertibrachials with pinnules on all brachials (probably true for all

tertibrachials) (Fig. 6.4). Secundibrachials and tertibrachials higher than wide; tertibrachials ~ 1.5 times higher than wide. Pinnules long (Fig. 6.3).

Column circular, heteromorphic (Fig. 6.5); column pattern N3231323. Proximally all columnals with convex latus with nodal the widest and highest and progressively decreasing in height and width in successive cycles.

Etymology.—The species name recognizes Pierre Groulx, who donated specimens for this study.

Materials.—Paratypes: ROMIP 54198b, ROMIP 54198c and MPEP719.1.

Measurements.—ROMIP 54198a: CrH, 15*, CaH, 7.0; CaW, 7.3, CoH, 9.6*; ROMIP 54198b: CaH, 10.5*; CaW, 9.0, CoH, 32.0*; ROMIP 54198c: AH, 25.0*; and MPEP719.1: CrH, 46.0, CaH, 10.2; CaW, 8.1, CoH, 90.4*.

Remarks.—The only previously recognized species of this genus was the type species, *Becsciecrinus adonis*, from the Fox Point and Chabot members of the Becscie Formation (Llandoverly, Rhuddanian). The label on ROMIP 54198 indicates Silurian, Jupiter River (Locality 28), but MPEP719.1 from the Markus Martin collection is from the coast at the mouth of the Jupiter River, thus from the Cybèle Member of the Jupiter Formation (Locality 7). This Silurian occurrence extends the known range of *Becsciecrinus* to the Rhuddanian through Telychian. *Becsciecrinus groulxi* n. sp. is the second species of this genus and is compared to *B. adonis* in Supplemental Table 7.

MPEP719.1 is a pyritized specimen with very poorly preserved definition of plate boundaries (Fig. 6.4). Thus, ROMIP 54198a is designated as the holotype despite the imprecise locality data for this specimen. ROMIP 54198a is a partial crown with column attached that exposes the CD interray and adjacent rays (Fig. 6.2, 6.5, 6.6). ROMIP 54198b is a partial calyx with column attached (Fig. 6.1), and ROMIP 54198c is a broken specimen with a considerable amount of the arms preserved. The plating on the holotype is poorly defined, and a plate diagram is given in Figure 7.1.

Genus *Cybelecrinus* Ausich and Copper, 2010

Type species.—*Cybelecrinus ladas* Ausich and Copper, 2010.

Cybelecrinus ladas Ausich and Copper, 2010
Figure 8

Holotype.—GSC 126819.

Occurrence.—*Cybelecrinus ladas* is known from the Goéland Member, Menier Formation and Cybèle Member, Jupiter Formation (Llandoverly, Aeronian to Telychian) of Anticosti Island, Québec, Canada. New material is from Localities 11 and 12.

Materials.—Paratypes: GSC 126820–GSC 126823, UA13623a, UA13623b, UA13624a, and UA13624b. Several new specimens are on MPEP1148.1 (five specimens) (Fig. 8) and MPEP1149.1 (two specimens).




Figure 8. *Cybelecrinus ladas* Ausich and Copper, 2010; lateral view of two partial crowns and associated pluricolumnals; MPEP1148.1; scale bar, 5.0 mm.

Remarks.—Two small slabs (MPEP1148.1 and MPEP1148.2) contain excellently preserved specimens of *C. ladas* and preserve characters of the column (Fig. 8).

Order Monobathrida Moore and Laudon, 1943
Family Xenocrinidae Miller, 1890
Genus *Xenocrinus* Miller, 1881

Type species.—*Xenocrinus penicillus* Miller, 1881.

Xenocrinus rubus Ausich and Copper, 2010
Figure 9.1–9.3, 9.5

Holotype.—GSC 126697.

Occurrence.—Crowns of *X. rubus* were originally known from the Juncliff (in today's lithostratigraphy), Lousy Cove, and Laframboise members of the Ellis Bay Formation, and only columnals were known from the Silurian (lower Chabot

Member, Ellis Bay Formation) (Ausich and Copper, 2010). New specimens represented by crowns reported here expand the range of *X. rubus* to the Katian through at least the Rhuddanian (and perhaps the Aeronian), Anticosti Island, Québec, Canada.

Materials.—Paratypes: GSC 126688a–d, GSC 126689a–f, GSC 126691, GSC 126692b, GSC 126693a–e, GSC 126696a–c, GSC 126698, and GSC 126699. New material includes MPEP504.1 (three specimens), MPEP510.4 (two specimens), and ROMIP 54197.

Remarks.—Six new, well-preserved specimens of *X. rubus* Ausich and Copper, 2010 are in the collections reported on herein. MPEP504.1 is a slab composed primarily of aligned pluricolumnals, many exceeding 50 mm in length (Locality 19). Two of the *X. rubus* specimens are oriented parallel to the orientation of the pluricolumnals, and a third specimen is nearly perpendicular to this orientation (Fig. 9.5).

Ausich and Copper (2010) reported *X. rubus* crowns only from the Hirnantian Ellis Bay Formation (Velleda [now Juncliff], Lousy Cove, and Laframboise members). They also reported the distinctive tetragonal *Xenocrinus* columnals from the Homard Member of the Vauréal Formation (Katian) to the lower Chabot Member of the Becscie Formation (Rhuddanian). Individual columnals were the only basis by which Ausich and Copper (2010) extended the range of *Xenocrinus* across the Ordovician–Silurian boundary, thus making it one of the few species-level taxa that survived Late Ordovician extinctions. Three *Xenocrinus* specimens from the Petryk collection (MPEP504.1) are interpreted to be from the Prinista Member of the Ellis Bay Formation (Hirnantian) (Locality 19). MPEP510.4 is from talus in the Becscie Formation along the Jupiter River, and ROMIP 54197 (Locality 27) is labeled as Silurian from along the Jupiter River, and supports the Silurian occurrence of *X. rubus* (Fig. 9.1–9.3).

Family Carpoecrinidae de Koninck and LeHon, 1854
Genus *Fibrocrinus* Ausich and Copper, 2010

Type species.—*Fibrocrinus phragmos* Ausich and Copper, 2010.

Fibrocrinus phragmos Ausich and Copper, 2010
Figures 4.1, 10

Holotype.—GSC 126768.

Occurrence.—East Point Member, Menier Formation (Aeronian), along the Trans-Anticosti Road, 1.9 km west of the Box-Bell Road junction (Locality 13) (Ausich and Copper, 2010).

Materials.—Paratypes: GSC 126769–GSC 126780; other specimens reported in Ausich and Copper (2010); and new material: MPEP1138.1–MPEP1138.8 and MPEP1139.1.

Remarks.—Numerous new, variously preserved specimens of *F. phragmos* are available from Locality 13, from which this taxon was originally described. Particularly instructive is

MPEP1138.3, which is a broken specimen exposing the interior surface of the tegmen (Fig. 10.2). The solid tegmen of *F. phragmos* is constructed of fixed primary peristomial cover plates, fixed ambulacral cover plates, and fixed interambulacral plates (Fig. 10.1). The interambulacral plates are very small; and based on preservation, it is possible that some of the abaxial-most cover plates are either loosely fixed or not fixed. On the underside of the tegmen (Fig. 10.2), the mouth and ambulacra (oral surface) are immediately beneath the cemented tegmen. A large, central, subelliptical opening is present for the mouth. A biseries of floor plates underlies the ambulacra. Fixed plates bifurcate two times in each ray, and fixed intra-ambulacral plates are present. A small circular pore is present between adjacent ambulacral floor plates. At the adaxial end of the fixed floor plates, two large pores are present in each ray between the oral frame plates and the adaxial-most fixed floor plates. In normal interrays, relatively few, large interradiial plates unite with fixed floor plates to make a solid plated surface. Although not fully exposed, a large opening is present in the CD interray, which housed the hindgut that led to the anal opening on the tegmen surface.

Family Eucalyptocrinitidae Roemer, 1855
Genus *Eucalyptocrinites* Goldfuss, 1831

Type species.—*Eucalyptocrinites rosaceus* Goldfuss, 1831.

Eucalyptocrinites archaios Ausich and Copper, 2010
Figure 11.3

Holotype.—UA 13633.

Occurrence.—Ferrum Member, Jupiter Formation (Llandovery, Telychian), Fire Tower Road, Anticosti Island, Québec, Canada (Ausich and Copper, 2010).

Materials.—A single new specimen MPEP495.3 is present in the new collections.

Remarks.—The holotype of *Eucalyptocrinites archaios* Ausich and Copper, 2010 is an excellently preserved specimen with only one or two columnals preserved (Ausich and Copper, 2010, pl. 6, figs. 2 and 3). The paratypes are all incomplete crowns. Therefore, the new specimen from the Petryk collection that has a portion of the column preserved significantly expands our knowledge of this species. The column of *E. archaios* is heteromorphic and holomeric. The nodal-internodal pattern is N121, with each cycle narrower and lower (Fig. 11.3). The nodals are 3.0 times wider than high, the priminternodals are 4.0 times wider than high, and the secundinternodals are 3.6 times wider than high. The latus on all columnals is strongly convex.

The new specimen is from the Petryk collection and is inferred to be from the Ferrum Member of the Jupiter Formation (Telychian) (Locality 18), which is the stratigraphic horizon from which this species was originally described (Ausich and Copper, 2010).

Family Patelliocrinidae Angelin, 1878
Genus *Jovacrinus* Ausich and Copper, 2010




Figure 9. Anticosti Island monobathrid camerates. (1–3, 5) *Xenocrinus rubus* Ausich and Copper, 2010; (1) enlarged lateral view of specimen A in MPEP510.4; (2) enlarged lateral view of specimen B in MPEP510.4; (3) the two previous specimens on single bedding surface; MPEP510.4; (5) crinoid pluricolumnal “logjam” with three specimens of *X. rubus*; MPEP504.1; (4) *Jovacrinus clarki* n. sp., lateral view of crown, note distinctive pinnules MPEP1146.1. Scale bars, 5.0 mm.




Figure 10. The tegmen of *Fibrocrinus phragmos* Ausich and Copper, 2010. (1) Exterior of well-preserved tegmen, note fixed ambulacral plates; holotype, GSC 126768 (from Ausich and Copper, 2010, pl. 5, fig. 8); the five primary peristomial cover plates are in the center of the tegmen (all slightly broken); the large central plate is the CD primary peristomial cover plate, and the other four primary peristomial cover plates are immediately above; (2) interior of tegmen, note opening of ambulacra that lead into mouth, central mouth, and fixed plates in ambulacra; MPEP1138.3. Scale bar 5.0 mm.

Type species.—*Jovacrinus spinosus*; by original designation.

Occurrence.—*Jovacrinus* is only known from the Jupiter Formation on Anticosti, but with recognition of *J. clarki* n. sp., this genus now ranges from the Ferrum to the Pavillon members of the Jupiter Formation (Telychian). As listed below, several features distinguish species within *Jovacrinus*.

Jovacrinus clarki new species

Figure 9.4

Holotype.—MPEP1146.1.

Diagnosis.—*Jovacrinus* with sculpturing on aboral cup plates scattered nodes, first primibrachials tetragonal, one fixed secundibrachial, at least three ranges of fixed interradial plates, and uniserial to biserial arms.

Occurrence.—Pavillon Member of the Jupiter Formation (Telychian) at the mouth of the Martin River, Anticosti Island, Québec, Canada (Locality 14).

Description.—Crown with conical shape due to arm posture (Fig. 9.4). Calyx small; probably low bowl shaped; thin plated; aboral plate sculpturing scattered nodes, radials and fixed brachials without median ray ridge; calyx plate spines absent or not preserved (Fig. 9.4).

Basal cirlet poorly preserved, probably visible in side view. Radial plates heptagonal, ~1.5 times wider than high, sculpturing as noted above.

Normal interrays in contact with tegmen (Fig. 9.4); first interradial hexagonal, approximately as high as wide, smaller than radials and larger than first primibrachial. Second range typically with two plates; plating 1-2-?.

Primanal probably heptagonal, 2.1 times wider than high, significantly smaller than radial plates, does not interrupt the radial cirlet; plating in CD interray P-3-6(?) (additional plating unknown); posterior interray slightly wider than regular interrays; CD interray in contact with tegmen.

First primibrachial tetragonal, ~1.1 times wider than high, much smaller than radial plates, but larger than primaxil; second primibrachial axillary, pentagonal. First secundibrachial distal-most fixed brachial, second secundibrachial with an abaxial pin-nule somewhat larger than other pinnules. Intrabrachial plates absent.

Tegmen not known.

Free arms 10, atomous. Brachials cuneate uniserial proximally, but biserial in at least distal half of arm. Brachials aborally convex without spines or nodes. Pinnules long, composed of at least six pinnulars.

Column circular; lumen unknown. Priminternodals and secundinternodals each separated by three tertinternodals. Long rhizoids along column (as known) borne on nodals, rhizoids up to 14 mm long. Details of column facets not known; facets of rhizoid segments as wide as nodals are high.

Etymology.—The name recognizes David Clark, who collected this specimen and made it available for study.

Measurements.—MPEP1146.1: CrH, 21.0*; CaH, 4.0*; CaW, 5.0*; RH, 1.8; RW, 2.4.

Remarks.—The relatively small calyx, 10 arms with distinctive stout, long pinnules, and a column with long rhizoids through the column aligns this species with *Jovacrinus*. There is only one specimen of *J. clarki* n. sp., but it is distinctly different from the two other species in this genus, which are compared in Supplemental Table 8. The *J. clarki* n. sp. holotype is slightly crushed or broken because of the thin-plated calyx, so only approximate measurements are given.

Family Marsupiocrinidae Bronn in Bronn and Roemer, 1855

Genus *Lateranicrinus* new genus

Type species.—*Lateranicrinus saintlaurenti* n. gen. n. sp.; by monotypy.




Figure 11. Anticosti Island monobathrid camerates. (1, 2) *Lateranicrinus saintlaurenti* n. gen., n. sp.: (1) bedding surface with several specimens, including holotype and paratypes; ROMIP 54203; (2) enlarged lateral view of holotype, in (11.1) ROMIP 54203a. (3) Lateral view of partially crushed calyx with proximal column of *Eucalyptocrinites archaios* Ausich and Copper, 2010; MPEP495.3. (4) Camerate Indet. B, well-preserved arms and column, but details of calyx not preserved, specimen uncoated, ROMIP 54196. (5) Partial calyx and proximal arms of Camerata indet. A, MPEP311.86. Scale bars 5.0 mm, unless otherwise labeled.

Diagnosis.—Marsupiocrinid with a very low bowl-shaped calyx, convex base with a basal concavity, basal plates confined to the basal concavity, ridge around basal concavity absent; radial plates project distally, in regular interrays three plates with one each in three ranges fixed in calyx, one plate above first interradial plate; two primibrachials, primaxil full width of brachitaxis, 10–15 free arms.

Occurrence.—Silurian (Llandovery), Québec, Canada.

Description.—See species description below.

Etymology.—Lateranus (L., m.) is the god of the hearth, which is a reference to the provenance of this crinoid from along the Brick River.




Figure 12. Anticosti Island myelodactylids and flexibles. (1) Partial crown and proxistele of *Eomyelodactylus springeri*, details obscured by heavy preparation MPEP718.6; (2) partially disarticulated crown with proxistele and mesistele of *Eomyelodactylus springeri* Ausich and Copper, 2010, see on Fig. 3.3; MPEP718.2; (3) disarticulated crown with proxistele and mesistele of *Eomyelodactylus springeri* Ausich and Copper, 2010, MPEP304.42. (4) Partially disarticulated crown with proxistele and short portion of the mesistele of *Eomyelodactylus* sp., MPEP1147.1. (5, 6) *Hormocrinus quebecensis*? Ausich and Copper, 2010, note infrabasal circling, MPEP324.1: (5) CD interray view of crown; (6) A ray view of crown. Scale bars 5.0 mm.

Remarks.—This new, relatively small marsupiocrinid is different based primarily on basal plates confined to the basal concavity, ridge around basal concavity absent; in regular interrays three plates with one each in three ranges fixed in calyx, one plate above first interradial plate; two primibrachials, primaxil full width of brachitaxis. These and other diagnostic characters for the Marsupiocrinidae are listed in Supplemental Table 9.

Lateranicrinus saintlaurenti n. gen. n. sp. is the second Llandovery-aged marsupiocrinid, with the other being *Manticrinus* from the Brassfield Limestone (Aeronian) of Ohio (U.S.A.). We suggest that *Lateranicrinus* n. gen. with more interradial plates in regular interrays and a primaxil that extends the full height of the primibrachitaxis is a more stemward marsupiocrinid than other known genera. However, determination of evolutionary relationships awaits thorough phylogenetic analyses.

Lateranicrinus saintlaurenti new species

Figures 7.2, 11.1, 11.2

Holotype.—ROMIP 54203a.

Diagnosis.—See generic diagnosis above.

Occurrence.—As noted in the Supplemental Appendix, data accompanying this specimen only indicates Silurian Brick River (Locality 32). It is interpreted most likely to be from the Jupiter Formation (Llandovery, Telychian) Anticosti Island, Québec, Canada.

Description.—Crown small, ovate. Calyx small, very low bowl-shaped (Fig. 11.2); arms not grouped; calyx plates convex (plate sculpturing not known).

Basal circling small, beneath proximal columnal and confined to basal concavity. Radial circling wraps around from edge of basal concavity to being visible in lateral view, <10% of calyx height, not interrupted in CD interray; radial plates five, heptagonal, ~1.2 times wider than high.

Regular interrays in narrow contact with tegmen; first interradial plate on upper shoulders of two radial plates, heptagonal, slightly wider than high, approximately same size as radials and much larger than first primibrachials; plating 1-1-1 (Fig. 7.2).

Primanal nonagonal, approximately equal in height and width, approximately same size as radial plates, does not interrupt radial circling; plating in CD interray P-3-?.

First primibrachial tetragonal with arcuate suture with radial plate, ~1.6 times wider than high, much smaller than radial plates and approximately same size as primaxil; all fixed brachials full width of ray, half-ray, or quarter-ray; second primibrachial axillary, pentagonal, ~1.6 times wider than high. Either secundibrachitaxis unbranched or second secundibrachial axillary; fixed third secundibrachial or first tertibrachial distal-most fixed brachial. Intrabrachial plates absent. Tegmen not known.

Either two or three free arms per ray (two most common), atomous. In undivided half-rays, first free brachial cuneate uniserial, approximately as high as wide; in half-ray divided, first free brachial cuneate uniserial much wider than high; remainder

of free brachials chisel biserial, aborally flattened through most of arm height, but in distal portion aborally convex and incurved toward oral-aboral axis. Arms pinnulate.

Column circular, holomeric (proximally), heteromorphic with pattern of N3231323, nodals and priminternodals much pronounced rounded epifacets. Other internodals with less pronounced convex epifacets.

Etymology.—This species name recognizes Daniel Saint-Laurent, who donated specimens for this study.

Materials.—Paratypes: ROMIP 54203b–ROMIP 54203i.

Measurements.—Holotype: ROMIP 54203a***: CrH, 25.0*; CaH, 6.3; CaW, 11.0; CoH, 15.0*; Paratypes: ROMIP 54203b: CrH, 28.1; CaH, 6.4; ROMIP 54203c: CrH, 27.3; CaH, 6.6; CaW, 10.5; CoH, 4.0*.

Remarks.—*Lateranicrinus saintlaurenti* n. gen. n. sp. is represented by one small slab with nine specimens (five of which have lateral views of at least part of the calyx) (Fig. 11.1). The slab was over-prepared so that sculpturing on plate surfaces cannot be determined. No single specimen is preserved so that the entire calyx plating is revealed. Thus, Figure 7.2 is a composite diagram illustrating the calyx plating of *Lateranicrinus saintlaurenti* n. gen. n. sp.

Parvclass Cladida Moore and Laudon, 1943
 Magnaorder Euclidida Wright, 2017
 Clade Articuliformes Wright, 2017
 Family Thalamocrinidae Miller and Gurley, 1895
 Genus *Thalamocrinus* Miller and Gurley, 1895

Type species.—*Thalamocrinus ovatus* Miller and Gurley, 1895; by original designation.

Occurrence.—Silurian (Llandovery to Pridoli) to Devonian (Lochkovian), Canada, United States, and South Africa.

Remarks.—McIntosh and Brett (1988) reviewed *Thalamocrinus*, concluding that seven valid species were present, including: *T. cylindricus* (Hall, 1852) (non Miller and Gurley, 1895); *T. elongatus* Springer, 1926; *T. globosus* Springer, 1926; *T. ovalus* (Rowley, 1904); *T. ovatus* Miller and Gurley, 1895; *T. robustus* McIntosh and Brett, 1988; and *T. strimplei* McIntosh and Brett, 1988. Note that *T. cylindricus* Miller and Gurley, 1895 (non Hall, 1852) was regarded a junior objective homonym, and McIntosh and Brett (1988) designated *T. strimplei* as the replacement name. The senior objective homonym, *T. cylindricus* (Hall, 1852) (non Miller, and Gurley, 1895), was considered valid, but this is not recognized, as it should be, in Webster and Webster (2013).

Thalamocrinus species known in 1988 ranged in age from Silurian (Wenlock) to Devonian (Lochkovian) and were confined to the Laurentia paleocontinent. Jell and Theron (1999) described *T. arenaceous* from the Devonian (Emsian) of Australia; however, this generic assignment should be questioned because *T. arenaceous* has a pentagonal radianal plate rather than a tetragonal radianal plate, which is a diagnostic

characteristic of *Thalamocrinus* (Lane and Moore, 1978; McIntosh and Brett, 1988). *Thalamocrinus daoustae* n. sp. is the oldest known member of *Thalamocrinus*, extending its range to the Llandovery. Species-diagnostic characters for *Thalamocrinus* are relative size, aboral cup shape, nature of aboral cup plate sutures, height to width ratio of infrabasal plates, relative convexity of infrabasal plates, relative convexity of basal plates, relative convexity of radial plates, and proximal width of the infrabasal circlet compared to the width of the proximal columnals. Although not preserved in most species, the nodals and infranodals of *T. daoustae* n. sp. have a flat latus, whereas those of *T. cylindricus* have a very convex latus; and *T. daoustae* n. sp. has a robust anal sac with strongly sculpted plates and *T. robustus* has a much more gracile anal sac.

Thalamocrinus daoustae new species
 Figure 13.1, 13.2

Holotype.—ROMIP 54199.

Diagnosis.—*Thalamocrinus* with relatively small high cone-shaped aboral cup, aboral cup sutures flush, infrabasal plate height:width ratio 1.2, gently convex infrabasal plates, basal plate height:width ratio 1.8, gently convex basal plates, radial plate height:width ratio 1.2, and the width of the proximal aboral cup the same as the width of the proximal columnals.

Occurrence.—The only data accompanying this specimen are Silurian, along the Jupiter River, Anticosti Island, Québec, Canada, which verifies a Llandovery age (either Aeronian or Telychian) (Locality 29).

Description.—Crown small size, conical. Aboral cup high cone shape, height to maximum width ratio ~1.2, plates gently convex, smooth.

Infrabasal plates five, equal in size, infrabasal plate height:width ratio 1.2; infrabasal circlet ~30% of aboral cup height; evenly tapered proximal aboral cup, width equals width of proximal columnals; infrabasal concavity absent. Basal plates five, hexagonal, except for CD which is heptagonal, height:width ratio 1.8; basal plates approximately same size as radial plates, ~37% of aboral cup height; basal plate height:width ratio ~1.8. Radials five, heptagonal in all rays, height:width ratio 1.2; radial circlet ~33% of aboral cup height. Radial facets angustary or peneplenary, occupy ~70% of distal radial plate width; radial facet declivate, rounded abaxially, details of facet topography not known.

Total number of anal plates in cup not preserved, two visible, radianal extends below left part of C radial plate, tetragonal, as high as wide. Presumed anal X extends to distal height of aboral cup. Other aspects of CD interray unknown. Anal sac straight sided, ~3.0 times higher than aboral cup height, shorter than arm height; plates arranged in vertical columns, each plate with a central node and radiating ridges connecting to like ridges on adjoining plates.

Arms branch at least two times isotomously. First primibrachial slightly wider than high; fifth primibrachial axillary, fifth




Figure 13. Eucladid crinoids from Anticosti Island: (1, 2) *Thalamocrinus daoustae* n. sp. lateral view of crown; holotype, ROMIP 54199; (1) partial crown and column through mesistele; note change in morphology on the column from the proxistele through the mesistele; (2) note high infrabasal plates and sculpturing on anal sac plates. (3, 4) *Plicodendrocrinus observationensis* Ausich and Copper, 2010; MPEP476.5: (3) note pentalobate column and a trilobite hyposstome covering part of aboral cup; (4) enlargement of (3). (5) *Aetocrinus gracilis* crown; see Fig. 3.2; MPEP718.1. (6) *Plicodendrocrinus petryki* n. sp.; holotype, ROMIP 54201. (7, 8) *Plicodendrocrinus martini* n. sp., holotype, MPEP1142.1: (7) enlargement of (8); note very low infrabasal circlet height in interradial positions; (8) note long anal sac and plate sculpturing of anal sac plates. All scale bars 5.0 mm.

secundibrachial axillary. Brachials rectangular uniserial, narrow, aborally rounded, brachials distal to first primibrachial higher than wide.

Column circular, holomeric, latus flat. Proxistele heteromorphic, pattern N1, nodals 3.0 times wider than high, internodals 5.0 times wider than high; taper slightly distally to a

homeomorphic mesistele with mesistele columnals slightly higher than wide; outer surface of epifacet rounded; lumen unknown. Columnal facets unknown.

Etymology.—The species name recognizes Nathalie Daoust, who donated specimens for this study.

Measurements.—Holotype: ROMIP 54199: CH, 18.2*; ACH, 5.0; pACW, 1.6; dACW, 4.1; IH, 2.0; BH, 2.4; BW, 1.6; RH, 2.0; RW, 1.8; CoH, 43.0*.

Remarks.—The high, cone-shaped aboral cup of *T. daoustae* n. sp. is unique for *Thalamocrinus*. The depressed aboral cup plate sutures, strongly convex infrabasal and basal plates, and the proximal aboral cup wider than the proximal columnals are unique characters for *T. robustus*. *Thalamocrinus daoustae* n. sp. is most similar to *T. strimplei* McIntosh and Brett, 1988 from the Wenlock and Ludlow of the United States and Canada. These and other diagnostic characters of *Thalamocrinus* species are listed in Supplemental Table 10.

Family Plicodendrocrinidae Jell, 1999
Genus *Plicodendrocrinus* Brower, 1995

Type species.—*Dendrocrinus casei* Meek, 1871.

Occurrence.—Ordovician (Katian) to Silurian (Llandovery), United States, Canada; Ordovician (Katian) to Silurian (Wenlock or Ludlow), United Kingdom; Devonian (Lochkovian), Australia.

Remarks.—Ausich and Copper (2010) described two new Ordovician species of *Plicodendrocrinus*, *P. epinettensis* and *P. observationensis*. Both of these species had broadly plicate/convex plates, and both were from the Vauréal Formation (Katian). Two additional new species of *Plicodendrocrinus* are present in the new material studied here, *P. petryki* n. sp. and *P. martini* n. sp. Both of these taxa have prominent stellate ridges on aboral cup plates, which among other characters, determine these as distinct Anticosti Island taxa. *Plicodendrocrinus petryki* n. sp. is also from the Vauréal Formation. When originally described, *Plicodendrocrinus* was known exclusively from Ordovician taxa. Since, Donovan et al. (2010) described *P. brevis* from the Silurian of the United Kingdom, and Jell (1999) described *P. australis* from the Early Devonian of Australia. A third post-Ordovician species is described herein from Anticosti Island. This specimen lacks locality information beyond Silurian from the Jupiter River. Thus, it can only be considered Llandovery, but this is the youngest known species in North America. Species-diagnostic characters for *Plicodendrocrinus* are listed in Supplemental Table 3.

Donovan et al. (2010) described a species previously known only from an unpublished manuscript from 1963 by W.H.C. Ramsbottom. However, the description is clearly noted as having been written by S.K. Donovan. Thus, this taxon should be regarded as *Plicodendrocrinus brevis* Donovan in Donovan et al., 2010. Donovan et al. (2010) correctly designated *Plicodendrocrinus brevis* Ramsbottom, 1963 as an unpublished nomen nudum (see International Commission of Zoological Nomenclature, 1999, Article, 50.1.1, p. 52). Authorship of this species is not Ramsbottom, as indicated in Webster and Webster (2013).

Plicodendrocrinus observationensis Ausich and Copper, 2010
Figure 13.3, 13.4

Holotype.—GSC 126675.

Occurrence.—Easton, Tower, and Schmitt Creek or Grindstone members, Vauréal Formation (Katian), Anticosti Island, Québec, Canada.

Materials.—Paratype: GSC 126675, new specimen: MPEP476.5.

Remarks.—*Plicodendrocrinus observationensis* was originally described from the Easton and Tower members of the Vauréal Formation. As noted in the Supplemental Appendix and the discussion of *Gaurocrinus fimbriatus*, the occurrence of *P. observationensis* at Locality 16 is inferred to be from either the Schmitt Creek or Grindstone Member of the Vauréal Formation.

Ausich and Copper (2010) described *Plicodendrocrinus epinettensis* and *P. observationensis*, both from the Vauréal Formation (Katian) of Anticosti Island. The primary differences between these taxa are that *P. observationensis* has a more equidimensional aboral cup (H:W ~1.0), wider radial facets, and higher brachial plates (Ausich and Copper, 2010, table 24). MPEP476.5 is another specimen of *P. observationensis*, and it adds to our understanding of this species. The arms on this specimen are nearly complete (Fig. 13.3) and reveal that free arms have two isotomous divisions rather than a single division, which was the extent of the preservation on the holotype (Fig. 13.4). Also, MPEP476.5 has more brachials in the primibrachitaxis than the holotype, and the brachials are much higher than wide rather than having the height and width approximately equal. This species is now understood to have 5–8 primibrachials and 8–10 secundibrachials. Although only partially exposed, the plating of the anal sac is much better preserved on MPEP476.5 than on specimens reported in Ausich and Copper (2010). The anal sac extends through most of the height of the arms. The plicate anal sac plates are typical for *Plicodendrocrinus*, and six columns of plates are present across the preserved anal sac. This new specimen is quite small, and it is possible that the very high brachial plates may represent a juvenile characteristic rather than an attribute of the adult members of the species.

Plicodendrocrinus petryki new species
Figure 13.6

Holotype.—ROMIP 54201.

Diagnosis.—*Plicodendrocrinus* with aboral cup plates with well-defined, single stellate ridges and plate triple junctions depressed; other plate sculpturing absent; width:height ratio of aboral cup 1.5; infrabasal plate radial height/infrabasal plate interradial height ~1.5; anal sac straight sided, approximately five plate columns per side, radial facet ~50% of radial plate width; brachials as wide as high; broadly convex aborally; column pentalobate to strongly pentalobate.

Occurrence.—The single specimen is known only from the Silurian along the Jupiter River (Supplemental Appendix, Locality 30), Anticosti Island, Québec, Canada.



Figure 14. *Fragucrinus bothros* Ausich and Copper, 2010. Note variable number of primibrachials and characteristics of the mesistele; MPEP518.2, photographed under alcohol. Scale bar 2.5 mm.

Description.—Crown medium-sized, conical. Aboral cup high cone-shaped; height:width ratio ~ 1.5 ; plates smooth, with single stellate ridges connecting to those on adjoining plates; broadly convex to plicate; aboral cup plate triple junctions depressed (Fig. 13.6).

Infrabasals five, equal in size; infrabasal circlet $\sim 31\%$ of aboral cup height. Infrabasal plate radial height/infrabasal plate interrarial height ~ 1.5 . Basals probably five, hexagonal, higher than wide, larger than radials; basal circlet $\sim 41\%$ of aboral cup height. Radials probably five, pentagonal in all rays, ~ 1.5 times wider than high; radial circlet $\sim 28\%$ of aboral cup height. Radial facets angustary, horseshoe shaped, declivate, occupy $\sim 50\%$ of distal radial plate width; radial facet topography not known.

Posterior interray and oral surface not known. Anal sac approximately three times higher than aboral cup, straight sided; composed of medium-sized, hexagonal, gently convex, wider than high, plicate plates arranged in vertical columns; at least five columns per side; summit of anal sac unknown.

Arms branch at least two times in an isotomous pattern. Fourth primibrachial axillary; nonaxillary primibrachials ~ 1.5 times wider than high; brachials as wide as high; broadly convex aborally; fourth or fifth secundibrachial axillary; brachials rectangular uniserial.

Proximal column pentalobate to strongly pentalobate, holomeric, homeomorphic outer surface of epifacet rounded; lumen wide, pentalobate, $\sim 45\%$ of column diameter. Columnal facets unknown.

Etymology.—The species name recognizes Allen Petryk, whose collection is included in this study.

Measurements.—Holotype: ROMIP 54201: CrH, 26.9; ACH, 8.1; pACW, 3.2; dACW, 10.4; rIH, 3.1; iIH, 2.0; ICW, 2.7; BH, 4.3; BW, 3.4; RH, 3.0; RW, 4.1; ASH, 19.5*.

Remarks.—Supplemental Table 11 is a listing of species-diagnostic characters for *Plicodendrocrinus*.

Plicodendrocrinus martini new species
Figure 13.7, 13.8

Holotype.—MPEP1142.1.

Diagnosis.—*Plicodendrocrinus* with aboral cup plates with well-defined, single stellate ridges and plate triple junctions depressed; other plate sculpturing absent; width:height ratio of aboral cup 1.25; infrabasal plate radial height/infrabasal plate interrarial height ~ 6.0 ; anal sac straight sided, approximately eight plate columns per side, radial facet $\sim 40\%$ of radial plate width; primibrachials as wide as high and broadly convex aborally with a keel, secundibrachials higher than wide and keeled aborally; column pentalobate to strongly pentalobate.

Occurrence.—Top of falls along Ruisseau Blanc, from the Vauréal Formation and interpreted to be from the Easton Member (Katian) (Locality 10), Anticosti Island, Québec, Canada.

Description.—Crown medium-sized. Aboral cup high cone-shaped; height:width ratio ~ 1.5 ; plates smooth, with single dominant stellate ridge connecting to like ridges on adjoining plates; aboral cup plate triple junctions depressed.

Infrabasals five, equal in size; infrabasal circlet ~37% of aboral cup height. Infrabasal plate radial height/interradial plate height ~6.0 (Fig. 13.7). Infrabasal plates concave. Basals probably five, hexagonal, higher than wide, approximately same size as radial plates, extend proximally nearly to proximal columnal; basal circlet ~34% of aboral cup height. Radials probably five, pentagonal or hexagonal; radial circlet ~28% of aboral cup height. Radial facets angustary, horseshoe shaped, declivate, occupy ~40% of distal radial plate width; radial facet topography not known.

Parts of five anal plates in aboral cup; radianal same size as C radial plate and directly beneath C radial plate, equidimensional, in lateral sutural contact with anal X. Anal X between D radial plate and radianal/C radial plate. Two anal plates above anal X; both partially in aboral cup and sutured to D radial or C radial plates. The fifth anal plate is the second plate above and to the right of the anal X and is sutured to the uppermost shoulder of the C radial plate. Anal sac ~10 times higher than aboral cup height, straight sided; composed of small-sized, hexagonal, gently convex, wider than high, plicate plates arranged in vertical columns (Fig. 13.8); at least eight columns per side; summit of anal sac poorly preserved.

Arms branch at least two times in an isotomous pattern. Fourth or fifth primibrachial axillary; nonaxillary primibrachials as high as wide and broadly convex. Sixth or seventh secundibrachial axillary; nonaxillary secundibrachials higher than wide and aborally keeled.

Proximal column strongly pentalobate, holomeric; other aspects of column not preserved.

Etymology.—The species name recognizes Markus Martin, who collected and prepared this specimen.

Measurements.—MPEP1142.1: ACH, 3.5; pACW, 2.0; dACW, 5.3*; rIH, 1.2; iIH, 0.4; ICW, 1.4; BH, 1.9; BW, 1.9; RH, 1.9; RW, 2.7; ASH, 31.0.

Remarks.—See discussion above and Supplemental Table 11 for comparison to other species of *Plicodendrocrinus*.

Family Botryocrinidae Wachsmuth and Springer, 1886
Genus *Fragucrinus* Ausich and Copper, 2010

Type species.—*Fragucrinus bothros* Ausich and Copper, 2010.

Fragucrinus bothros Ausich and Copper, 2010
Figure 14

Holotype.—GSC 126881.

Occurrence.—Richardson Member of the Jupiter Formation; along Sandtop Main Road (Ausich and Copper, 2010); Anticosti Island, Québec, Canada. The specimen is only noted as from the Silurian (Locality 23).

Materials.—The paratype is GSC 126675, and the new specimens are MPEP518.2a and MPEP518.2b.

Remarks.—One complete crown with column attached and one partial crown with a longer column attached constitute the new

specimens of *F. bothros* (Fig. 14). Previously only known from the holotype, these new specimens demonstrate that the number of primibrachials is variable (three to five) and the mesistele has a N121 pattern of nodals and internodals.

Acknowledgments

This paper is a continuation of fieldwork supported by the National Geographic Society (grant 6789-00) and the National Science Foundation (EAR-0205968). P. Copper, J. Jin, and A. Desrochers were stratigraphic and geographic guides during most Anticosti field seasons. We thank J. Iellamo for preparing specimens from the Petryk and Cournoyer/Daoust collections. We also thank the work and generosity of D. Clark, N. Daoust, the late P. Groulx, P. Isotalo, M. Martin, the late A. Petryk, and D. Saint-Laurent, who collected these specimens and made them available for scientific study. J. Dougherty and M. Coyne (Geological Survey of Canada) donated the Petryk collection to the MPEP. The authors also thank T.E. Guensburg, D.L. Meyer, and R. Mooi, for thoughtful reviews.

Accessibility of supplemental data

Additional information on new occurrences and further discussion of diagnostic characters are given in the Online Supplement on the Dryad Digital Repository: <https://doi.org/10.5061/dryad.2j0b93b>.

Taxa discussed in the Online Supplement but not in the manuscript above are *Camerata* indet. A (Fig. 11.5), *Camerata* indet. B (Fig. 11.4), *Eomyelodactylus foerstei* Ausich and Copper, 2010, *E. springeri* Ausich and Copper, 2010 (Figs. 3.3, 12.1–12.3), *Eomyelodactylus* sp. (Fig. 12.4), *Ladacrinus?* sp., *Hormocrinus quebecensis?* Ausich and Copper, 2010 (Fig. 12.5, 12.6), and *Aetocrinus gracilis* Ausich and Copper, 2010 (Figs. 3.2, 13.5).

References

- Angelin, N.P., 1878, Iconographia Crinoideorum in stratis Sueciae Siluricis fossilium: Holmiae, Samson and Wallin, 62 p.
- Ausich, W.I., 1984a, Calceocrinids from the Early Silurian (Llandoveryan) Brassfield Formation of southwestern Ohio: *Journal of Paleontology*, v. 58, p. 1167–1185.
- Ausich, W.I., 1984b, The genus *Chidochirus* from the Early Silurian of Ohio (Crinoidea, Llandoveryan): *Journal of Paleontology*, v. 58, p. 1341–1346.
- Ausich, W.I., 1985, New crinoids and revision of the superfamily Glyptocrinoidea (Early Silurian, Ohio): *Journal of Paleontology*, v. 59, p. 793–808.
- Ausich, W.I., 1986a, Early Silurian rhodocrinitean crinoids (Brassfield Formation, Ohio): *Journal of Paleontology*, v. 60, p. 84–106.
- Ausich, W.I., 1986b, Early Silurian inadunate crinoids (Brassfield Formation, Ohio): *Journal of Paleontology*, v. 60, p. 719–735.
- Ausich, W.I., 1986c, New camerate crinoids of the Suborder Glyptocrinina from the Lower Silurian Brassfield Formation (southwestern Ohio): *Journal of Paleontology*, v. 60, p. 887–897.
- Ausich, W.I., 1987, Brassfield Compsocrinina (Lower Silurian crinoids) from Ohio: *Journal of Paleontology*, v. 61, p. 552–562.
- Ausich, W.I., 1998, Phylogeny of Arenig to Caradoc crinoids (Phylum Echinodermata) and suprageneric classification of the Crinoidea: *The University of Kansas Paleontological Contributions*, n.s., v. 9, 36 p.
- Ausich, W.I., and Baumiller, T.K., 1993, Column regeneration in an Ordovician crinoid (Echinodermata): paleobiologic implications: *Journal of Paleontology*, v. 67, p. 1068–1070.
- Ausich, W.I., and Copper, P., 2010, The Crinoidea of Anticosti Island, Québec (Late Ordovician to Early Silurian): *Palaeontographica Canadiana*, v. 29, 157 p.

- Ausich, W.I., and Deline, B., 2012, Macroevolutionary transitions in crinoids following the Late Ordovician extinction event (Ordovician to Early Silurian): *Palaeogeography, Palaeoclimatology, Palaeoecology*, v. 361–362, p. 38–48.
- Ausich, W.I., and Wilson, M.A., 2016, Llandovery (early Silurian) crinoids from Hiiumaa Island, western Estonia: *Journal of Paleontology*, v. 90, p. 1139–1147.
- Ausich, W.I., Kammer, T.W., and Baumiller, T.K., 1994, Demise of the middle Paleozoic crinoid fauna: *Paleobiology*, v. 20, p. 345–361.
- Ausich, W.I., Brett, C.E., Hess, H., and Simms, M.J., 1999, Crinoid form and function, in Hess, H., Ausich, W.I., Brett, C.E., and Simms, M.J., *Fossil Crinoids*: Cambridge, UK, Cambridge University Press, p. 3–30.
- Ausich, W.I., Kammer, T.W., Rhenberg, E.C., and Wright, D.F., 2015a, Frontiers in Paleontology: early phylogeny of crinoids within the Pelmatozoan clade: *Palaeontology*, v. 58, p. 937–952.
- Ausich, W.I., Peter, M.E., and Etensohn, F.R., 2015b, Echinoderms from the lower Silurian Brassfield Formation of east-central Kentucky: *Journal of Paleontology*, v. 89, p. 245–256.
- Barnes, C.R., 1988, Stratigraphy and paleontology of the Ordovician-Silurian boundary interval, Anticosti Island, Québec, Canada: *British Museum Natural History Bulletin (Geology)*, v. 43, p. 195–219.
- Baumiller, T.K., 1990, Non-predatory drilling in Mississippian crinoids by platyceratid gastropods: *Palaeontology*, v. 33, p. 743–748.
- Baumiller, T.K., 1993, Crinoid stalks as cantilever beams and the nature of stalk ligament: *Neues Jahrbuch für Geologie und Paläontologie, Abhandlungen*, v. 190, p. 115–133.
- Baumiller, T.K., 2002, Multi-snail infestation of Devonian crinoids and the nature of platyceratid-crinoid interactions: *Acta Palaeontologica Polonica*, v. 47, p. 133–139.
- Baumiller, T.K., and Gahn, F.J., 2003, Predation on crinoids, in Kelley, P.H., Kowalewski, M., and Hanson, T.A., eds., *Predator-Prey Interactions in the Fossil Record*: New York, Kluwer/Plenum Publishers, p. 263–278.
- Baumiller, T.K., Gahn, F.J., and Savill, J., 2004, New data and interpretations of the crinoid-platyceratid relationship, in Heinzeller, T., and Nebelsick, J.H., eds., *Echinoderms*: München, London, Talyor and Francis, p. 393–398.
- Billings, E., 1859, On the Crinoidea of the Lower Silurian rocks of Canada: *Canadian Organic Remains, Decade 4*, Geological Survey of Canada, 72 p.
- Bolton, T.E., 1961, Ordovician and Silurian formations of Anticosti Island, Quebec: *Geological Survey of Canada Paper*, no. 71-19, 45 p.
- Bowsher, A.L., 1955, Origin and adaptation of platyceratid gastropods: *University of Kansas Paleontological Contributions, Mollusca Article 5*, 11 p.
- Bronn, H.G., and Roemer, F. (C.F.), 1851–56, *Lethaea Geognostica oder Abbildung und Beschreibung der für die Gebirgs-Formationen bezeichnendsten Versteinerungen. Dritte stark vermehrte Auflage bearbeitet von H.G. Bronn und F. Roemer, Erster Band. 1. Übersichten, I. Theil. Systematische Übersicht der Fossilien; Schlüssel-Tabellen; Register, von H.G. Bronn. 2. Palaeo-Lethaea. II. Theil. Kohlen-Periode (Silur-, Devon-, Kohlen-, und Zechstein-Formation), von F. Roemer, 788 p., Zweiter Band (1851–52). 3. Meso-Lethaea. III. Theil. Trias-Periode; IV. Theil. Oolithen-Periode; V. Theil. Kreide-Periode*: Stuttgart, E Schweizerbart, 412 p.
- Brower, J.C., 1995, Dendrocrinid crinoids from the Ordovician of northern Iowa and southern Minnesota: *Journal of Paleontology*, v. 69, p. 939–960.
- Buggisch, W., Joachimski, M.M., Lehnert, O., Bergström, S.M., Repetski, J.E., and Webers, G., 2010, Did intense volcanism trigger the first Late Ordovician icehouse?: *Geology*, v. 38, p. 327–330.
- Cocks, L.R.M., and Copper, P., 1981, The Ordovician-Silurian boundary at the eastern end of Anticosti Island: *Canadian Journal of Earth Sciences*, v. 18, p. 1029–1034.
- Cole, S.R., 2017, Phylogeny and morphologic evolution of the Ordovician Camerata (Class Crinoidea, Phylum Echinodermata): *Journal of Paleontology*, 91, p. 815–828.
- Copper, P., 1989, Upper Ordovician and Lower Silurian reefs of Anticosti Island, Quebec: *Canadian Society of Petroleum Geologists Memoir*, v. 13, p. 271–276.
- Copper, P., 2001, Reefs during the multiple crises towards the Ordovician-Silurian boundary: Anticosti Island, eastern Canada, and worldwide: *Canadian Journal of Earth Sciences*, v. 38, p. 153–171.
- Copper, P., Jin, J., and Desrochers, A., 2013, The Ordovician-Silurian boundary (late Katian–Hirnantian) of western Anticosti Island: revised stratigraphy and benthic megafauna correlations: *Stratigraphy*, v. 19, p. 213–227.
- Deline, B., Ausich, W.I., and Brett, C.E., 2012, Comparing taxonomic and geographic scales in the morphologic disparity of Ordovician through Early Silurian Laurentian crinoids: *Paleobiology*, v. 38, p. 538–553.
- Desrochers, A., Farley, C., Achab, A., and Asselin, E., 2008, A high-resolution stratigraphic model to resolve the longstanding issues relative to the correlation and interpretation of the O/S boundary on Anticosti Island, Canada, in Kröger, B., and Servais, T., eds., *Palaeozoic Climates—International Congress: Closing Meeting of the International Geoscience Programme 503 “Ordovician Palaeogeography and Palaeoclimate,” Abstracts*: Lille, Université de Lille, p. 32.
- Desrochers, A., Farley, C., Achab, A., Asselin, E., and Riva, J.F., 2010, A far-field record of the end Ordovician glaciations: the Ellis Bay Formation, Anticosti Island, Eastern Canada: *Palaeogeography, Palaeoclimatology, Palaeoecology*, v. 296, p. 248–263.
- Donovan, S.K., 1993, A Rhuddanian (Silurian, Llandovery) pelmatozoan fauna from south-west Wales: *Geological Journal*, v. 28, p. 1–19.
- Donovan, S.K., Widdison, R.E., Lewis, D.N., and Fearnhead, F.E., 2010, The British Silurian Crinoidea, part 2: addendum to part 1 and Cladida: *Palaeontographical Society*, v. 164, p. 47–133.
- Eckert, J.D., 1984, Early Llandovery crinoids and stelleroids from the Cataract Group (Lower Silurian), southern Ontario, Canada: *Royal Ontario Museum Life Sciences Contributions*, v. 137, 83 p.
- Eckert, J.D., 1990, The Early Silurian myelodactylid crinoid *Eomyelodactylus Foerste*: *Journal of Paleontology*, v. 64, p. 135–141.
- Fearnhead, F.E., and Donovan, S.K., 2007, A cladid crinoid (Echinodermata) from the Llandovery (Lower Silurian) of the Girvan district, SW Scotland: *Scottish Journal of Geology*, v. 43, p. 74–82.
- Gahn, F.J., and Baumiller, T.K., 2003, Infestation of Middle Devonian (Givetian) camerate crinoids by platyceratid gastropods and its implications for the nature of their biotic interaction: *Lethaia*, v. 36, p. 71–82.
- Gahn, F.J., and Baumiller, T.K., 2006, Using platyceratid gastropod behavior to test functional morphology: *Historical Geology*, v. 18, p. 397–404.
- Goldfuss, G.A., 1826–44, *Petrefacta Germaniae, tam ea, Quae in Museo Universitatis Regiae Borussicae Fridericiae Wilhelmae Rhenanea, serventur, quam alia quaecunque in Museis Hoeninghusiano Muensteriano alisque, extant, iconibus et descriptionibus illustrata.—Abbildungen und Beschreibungen der Petrefacten Deutschlands und der Angränzende Länder, unter Mitwirkung des Herrn Grafen Georg zu Münster, herausgegeben von August Goldfuss: v. 1 (1826–1833), Divisio prima. Zoophytorum reliquiae, p. 1–114; Divisio secunda. Radiariorum reliquiae, p. 115–221 [Echinodermata]; Divisio tertia. Annulatorium reliquiae, p. 222–242; v. 2 (1834–40), Divisio quarta. Molluscorum acephalorum reliquiae. I. Bivalvia, p. 65–286; II. Brachiopoda, p. 287–303; III. (1841–44), Divisio quinta. Molluscorum gasteropodum reliquiae, p. 1–121; atlas of plates, 1–199, Düsseldorf, Armz & Co.*
- Guensburg, T.E., and Sprinkle, J., 2000, Ecologic radiation of Cambro-Ordovician echinoderms, in Zhuravlev, A.Y., and Riding, R., eds., *Ecology of the Cambrian Radiation*: New York, Columbia University Press, p. 428–444.
- Hall, J., 1852, *Palaeontology of New York*, v. 2, containing descriptions of the organic remains of the lower middle division of the New-York system: *Natural History of New York*, New York, D. Appleton & Co. and Wiley & Putnam; Boston, Gould, Kendall, & Lincoln, v. 6, 362 p.
- Hall, J., 1866, Descriptions of new species of Crinoidea and other fossils from the Lower Silurian strata of the age of the Hudson-River Group and Trenton Limestone: Albany, 17 p. [preprint]
- Herrmann, A.D., Patkowsky, M.E., and Pollard, D., 2004, The impact of paleogeography, $p\text{CO}_2$, poleward ocean heat transfer, and sea level change on global cooling during the Late Ordovician: *Palaeogeography, Palaeoecology, and Palaeoclimatology*, v. 206, p. 59–74.
- International Commission of Zoological Nomenclature, 1999, *International Code of Zoological Nomenclature, fourth edition*: London, The International Trust for Zoological Nomenclature 1999, 306 p.
- Jell, P.A., 1999, Silurian and Devonian crinoids from central Victoria: *Memoirs of the Queensland Museum*, v. 43, p. 1–114 p.
- Jell, P.A., and Theron, J.N., 1999, Early Devonian echinoderms from South Africa: *Memoirs of the Queensland Museum*, v. 43, p. 115–200.
- Jin, J., and Copper, P., 2008, Response of brachiopod communities to environmental change during the Late Ordovician mass extinction interval, Anticosti Island, Eastern Canada: *Fossils and Strata*, v. 54, p. 41–51.
- Koninck, L.G. de, and Le Hon, H., 1854, *Recherches sur les crinoïdes du terrain carbonifère de la Belgique*: Académie Royal de Belgique, *Memoir*, v. 28, no. 3, 215 p.
- Kump, L.R., Arthur, M.A., Patkowsky, M.E., Gibbs, M.T., Pinkus, D.S., and Sheehan, P.M., 1999, A weathering hypothesis for glaciation at high atmospheric $p\text{CO}_2$ during the Late Ordovician: *Palaeogeography, Palaeoclimatology, Palaeoecology*, v. 152, p. 173–187.
- Lane, N.G., and Moore, R.C., 1978, Suborder Cyathocrinina, in Moore, R.C., and Teichert, K. eds., *Treatise on Invertebrate Paleontology, Echinodermata, Pt. T(2)*: Geological Society of America and University of Kansas Press, Boulder, CO and Lawrence, KS, p. T578–T607.
- Lefebvre, B., Sumrall, C.D., Shroat-Lewis, R.A., Reich, M., Webster, G.D., Hunter, A.W., Nardin, E., Rozhnov, S.V., Guensburg, T.E., and Touzeau, A., 2013, *Palaeobiogeography of Ordovician echinoderms*: Geological Society, London, *Memoirs*, v. 38, p. 173–198.

- Lefebvre, V., Servais, T., François, L., and Averbuch, O., 2010, Did a Katian large igneous province trigger the Late Ordovician glaciation?: Palaeogeography, Palaeoclimatology, Palaeoecology, v. 296, p. 309–310.
- Long, D.G.F., and Copper, P., 1987, Stratigraphy of the Upper Ordovician upper Vauréal and Ellis Bay formations, eastern Anticosti Island, Québec: Canadian Journal of Earth Sciences, v. 24, p. 1807–1820.
- McIntosh, G.C., and Brett, C.E., 1988, Occurrence of the cladid inadunate crinoid *Thalamocrinus* in the Silurian (Wenlockian) of New York and Ontario: Royal Ontario Museum, Life Sciences Contributions, v. 149, 17 p.
- Meek, F.B., 1871, On some new Silurian (Ordovician) crinoids and shells: American Journal of Science, ser. 3, v.1, p. 295–299.
- Meyer, D.L., and Ausich, W.I., 1983, Biotic interactions among Recent and among fossil crinoids, Chapter 9, in Tevesz, M.J.S., and McCall, P.L., eds., Biotic Interactions in Recent and Fossil Benthic Communities: New York, Plenum Publishing Corporation, p. 337–427.
- Miller, J.S., 1821, A Natural History of the Crinoidea, or Lily-shaped Animals; with observations on the genera, *Asteria*, *Euryale*, *Comatula* and *Marsupites*: Bristol, England, Bryan & Co., 150 p.
- Miller, S.A., 1881, Description of some new and remarkable crinoids and some other fossils of the Hudson River Group and notice of *Strotocrinus bloomfieldensis*: Journal of the Cincinnati Society of Natural History, v. 4, p. 69–77.
- Miller, S.A., 1883, The American Palaeozoic Fossils: a catalogue of the genera and species, with names of authors, dates, places of publication, groups of books in which found, and the etymology and signification of the words, and an introduction devoted to the stratigraphical geology of the Palaeozoic rocks, 2nd ed.: Cincinnati, Ohio, published by the author, p. 247–334.
- Miller, S.A., 1890, The structure, classification, and arrangement of American Palaeozoic crinoids into families: American Geologist, v. 6, p. 275–286, 340–357.
- Miller, S.A., and Gurley, W.F.E., 1895, New and interesting species of Palaeozoic fossils: Illinois State Museum Bulletin, v. 7, 89 p.
- Moore, R.C., and Laudon, L.R., 1943, Evolution and classification of Paleozoic crinoids: Geological Society of America Special Paper, v. 46, 151 p.
- Nardin, E., Goddérís, Y., Donnadieu, Y., Le Hir, G., Blakely, R.C., Pucéa, E., and Arze, M., 2011, Modeling the early Paleozoic long-term climate trend: Geological Society of America Bulletin, v. 123, p. 1181–1192.
- Oji, T., and Amemiya, S., 1998, Survival of crinoid stalk fragments and its taphonomic implications: Paleontological Research, v. 2, p. 67–70.
- Peters, S.E., and Ausich, W.I., 2008, A sampling-adjusted macroevolutionary history for Ordovician–Early Silurian crinoids: Paleobiology, v. 3, p. 104–116.
- Petryk, A.A., 1981, Stratigraphy, sedimentology, and paleogeography of the Upper Ordovician–Lower Silurian of Anticosti Island, Québec, in Lespérance, P.J., ed., Subcommission on Silurian Stratigraphy, Ordovician–Silurian Boundary Working Group: Field meeting Anticosti–Gaspé, Québec, Université de Montréal, v. 2, p. 10–39.
- Phillips, J., 1839, Chapter 48. Encrinites and zoophytes of the Silurian System, in Murchison, R.T., The Silurian System: London, John Brown, p. 670–675.
- Richardson, J., 1857, Report of the year 1856: Canadian Survey of Canada Report of Progress for the years 1853–54–55–56, p. 191–245.
- Roemer, C.F., 1855, in Bronn, H.G., and Roemer, C.F. Lethaea Geognostica, oder Abbildung und Beschreibung der für die Gebirgs-Formationen bezeichnendsten Versteinerungen 1851–1856, 3rd ed.: Stuttgart, E. Schweizerbart, v. 2, 788 p.
- Rowley, R.R., 1904, The Echinodermata of the Missouri Silurian and a new brachiopod: American Geologist, v. 34, p. 269–282.
- Schuchert, C., and Twenhofel, W.H., 1910, Ordovician–Silurian section of the Mingan and Anticosti islands, Gulf of St. Lawrence: Geological Society of America Bulletin, v. 21, p. 677–716.
- Sepkoski, J.J., Jr., 1996, Patterns of Phanerozoic extinction: a perspective from global databases, in Walliser, O.H., ed., Global Events and Event Stratigraphy in the Phanerozoic: Berlin, Springer-Verlag, p. 31–51.
- Shaviv, N.J., and Veizer, J., 2003, Celestial drive of Phanerozoic climate?: GSA Today, v. 13/7, p. 4–10.
- Springer, F., 1926, American Silurian Crinoids: Smithsonian Institution Publication, v. 2872, 239 p.
- Springer, F., 1928, Echinodermata, in Twenhofel, W.A., Geology of Anticosti Island: Canada Geological Survey Memoir, v. 154 (1927), 481 p.
- Sprinkle, J., and Guensburg, T.E., 2004, Crinozoan, blastozoan, echinozoan, asterozoan, and homalozoan echinoderms, in Webby, D.B., Paris, F., Droser, M.L., and Percival, I.G., eds., The Great Ordovician Biodiversification Event: New York, Columbia University Press, p. 266–280.
- Twenhofel, W.H., 1928, Geology of Anticosti Island: Geological Survey of Canada Memoirs, v. 154, 481 p.
- Ubahgs, G., 1978, Skeletal morphology of fossil crinoids, in Moore, R.C., and Teichert, K. eds., Treatise on Invertebrate Paleontology, Echinodermata, Pt. T(2): Geological Society of America and University of Kansas Press, Boulder, CO and Lawrence, KS, p. T58–T216.
- Wachsmuth, C., and Springer, F., 1880–1886, Revision of the Palaeocrinoidea: Proceedings of the Academy of Natural Sciences of Philadelphia Pt. I. The families Ichthyocrinidae and Cyathocrinidae (1880), p. 226–378 (separate repaged p. 1–153). Pt. II. Family Sphaeroidocrinidae, with the sub-families Platycrinidae, Rhodocrinidae, and Actinocrinidae (1881), p. 177–411 (separate repaged, p. 1–237). Pt. III, Sec. 1. Discussion of the classification and relations of the brachiate crinoids, and conclusion of the generic descriptions (1885), p. 225–364 (separate repaged, 1–138). Pt. III, Sec. 2. Discussion of the classification and relations of the brachiate crinoids, and conclusion of the generic descriptions (1886), p. 64–226 (separate repaged to continue with section 1, 139–302).
- Webby, G.D., Paris, F., Droser, M.L., and Percival, I.G., 2004, The Great Ordovician Biodiversification Event: New York, Columbia University Press, 484 p.
- Webster, G.D., and Maples, C.G., 2008, Cladid crinoid radial facets, brachials, and arm appendages: a terminology solution for studies of lineage, classification, and paleoenvironment, in Ausich, W.I., and Webster, G.D., eds., Echinoderm Paleobiology: Bloomington, IN, Indiana University Press, p. 196–226.
- Webster, G.D., and Webster, D.W., 2013, Bibliography and index of Paleozoic crinoids, coronoids, and hemistrectocrinids, 1758–2012. <http://crinoids.azurewebsites.net/> (Oct. 2018).
- Wright, D.F., 2017, Bayesian estimation of fossil phylogenies and the evolution of early to middle Paleozoic crinoids (Echinodermata): Journal of Paleontology, v. 91, p. 799–814. doi: 10.1017/jpa.2-16.141
- Wright, D.F., Ausich, W.I., Cole, S.R., Peter, M.E., and Rhenberg, E.C., 2017, Phylogenetic taxonomy and classification of the Crinoidea (Echinodermata): Journal of Paleontology, v. 91, p. 829–846. doi 10.1917/jpa.2016.142
- Zittel, K.A. von., 1876–80, Handbuch der Palaeontologie, v. 1, Palaeozoologie: München, Leipzig, R. Oldenbourg, (1879), 765 p. [Echinoderms, p. 308–560]
- Zou, C., Qui, Z., Poulton, S.W., Dong, D., Wang, H., Chen, D., Lu, B., Shi, Z., and Tao, H., 2018, Ocean euxinia and climate change “double whammy” drove Late Ordovician mass extinction: Geology, v. 46, p. 533–538.

Accepted: 24 April 2019