

Athletes' Age, Sex, and Years of Education Moderate the Acute Neuropsychological Impact of Sports-Related Concussion: A Meta-analysis

Brooke K. Dougan, Mark S. Horswill, AND Gina M. Geffen

School of Psychology, The University of Queensland, Brisbane, Queensland, Australia

(RECEIVED September 8, 2011; FINAL REVISION October 23, 2012; ACCEPTED October 25, 2012; FIRST PUBLISHED ONLINE February 4, 2013)

Abstract

The objective of this study is to determine which pre-existing athlete characteristics, if any, are associated with greater deficits in functioning following sports-related concussion, after controlling for factors previously shown to moderate this effect (e.g., time since injury). Ninety-one independent samples of concussion were included in a fixed+systematic effects meta-analysis ($n = 3,801$ concussed athletes; 5,631 controls). Moderating variables were assessed using analogue-to-ANOVA and meta-regression analyses. Post-injury assessments first conducted 1–10 days following sports-related concussion revealed significant neuropsychological dysfunction, postural instability and post-concussion symptom reporting ($d = -0.54$, -1.10 , and -1.14 , respectively). During this interval, females ($d = -0.87$), adolescent athletes competing in high school competitions ($d = -0.60$), and those with 10 years of education ($d = -1.32$) demonstrated larger post-concussion neuropsychological deficits than males ($d = -0.42$), adults ($d = -0.25$), athletes competing at other levels of competition ($d = -0.43$ to -0.41), or those with 16 years of education ($d = -0.15$), respectively. However, these sub-groups' differential impairment/recovery beyond 10 days could not be reliably quantified from available literature. Pre-existing athlete characteristics, particularly age, sex and education, were demonstrated to be significant modifiers of neuropsychological outcomes within 10 days of a sports-related concussion. Implications for return-to-play decision-making and future research directions are discussed. (*JINS*, 2014, 20, 64–80).

Keywords: Traumatic brain injury, Cognition, Symptoms, Postural balance, Football, Systematic review

INTRODUCTION

A high incidence of sports-related concussion is well documented (e.g., Centers for Disease Control and Prevention, 2007; Guskiewicz, Weaver, Padua, & Garrett, 2000; Tate, McDonald, & Lulham, 1998). Evidence-based evaluation of the impact of concussion on an athlete's functioning, and of the optimal timing for return to play, is recognized as necessary to safe-guard athletes' well-being: unnecessarily delayed return and a loss of competitive advantage must be balanced against the risk of further injury if athletes are returned prematurely (i.e., before recovery). In this regard, the National Academy of Neurology (Moser et al., 2007) and the third International Conference on Concussion in Sport (McCrory et al., 2009) recommend: (1) an individualized approach to the assessment and management of sports-related

concussion, guided by the results of neuropsychological, self-report symptom, and postural stability assessments; and (2) that return-to-play decision-making should take into account possible modifiers of injury outcomes such as pre-existing athlete characteristics (e.g., age, sex, sport and position played, level of competition, and premorbid neurological functioning). However, research remains equivocal regarding specifically which of these factors contribute to the outcomes associated with sports-related concussion, and to what degree.

Variations in incidence, severity and duration to recovery between younger and older athletes, males and females, and those competing in different sports or at different levels of competition have been reported in studies of sports-related concussion (Baillargeon, Lassonde, Leclerc, & Ellemberg, 2012; Daniel, Rowson, & Duma, 2012; Dick, 2009; Guskiewicz et al., 2000; Putukian, Aubry, & McCrory, 2009). Evidence of disparity in outcome according to individual characteristics has also been reported in the mild, moderate and severe traumatic brain injury literatures (Bruce et al., 1981; Farace & Alves, 2000; Hoofien, Vakil, Gilboa, Donovan, & Barak, 2002;

Correspondence and reprint requests to: Brooke Dougan, School of Psychology, University of Queensland, St Lucia, Brisbane, QLD 4072 Australia. E-mail: brooke.dougan@uqconnect.edu.au

Ratcliff et al., 2007). However, contradictory evidence also exists (e.g., Tsushima, Lum, & Geling, 2009). While experts in the field (e.g., McCrory et al., 2009) have critically reviewed this literature, firm conclusions regarding the modifying role of pre-existing athlete characteristics on sports-related concussion outcomes have been constrained by (1) a paucity of well-controlled prospective studies, (2) infrequent recruitment of samples other than adult male athletes competing at the college or professional levels of competition, (3) inconsistent reporting of detailed sample demographic variables, and (4) infrequent stratification of results according to sample characteristics. Likewise, previous meta-analyses of sports-related concussion and of mixed-mechanism concussion in the general population (mild traumatic brain injury, mTBI) have consistently identified several factors that moderate recovery (e.g., time since injury), but have not investigated the contribution of premorbid athlete characteristics to variation in concussion outcomes.

Published meta-analyses typically report significant “small to moderate”¹ neuropsychological deficits ($d = -0.54$, Belanger, Curtiss, Demery, Lebowitz, & Vanderploeg, 2005; -0.49 , Belanger & Vanderploeg, 2005; -0.28 , Rohling et al., 2011; -0.24 , Schretlen & Shapiro, 2003), and minimally increased symptom reports (-0.05 to -0.15 , Panayiotou, Jackson, & Crowe, 2010), when aggregated over all assessments following a concussive injury. However, aggregation of outcomes over broad follow-up epochs potentially obscures variation according to the different assessment and sample characteristics of the included studies (Iverson, 2010). For example, the time elapsed between injury and assessment is recognized as a key moderator of outcome associated with both mTBI and sports-related concussion, such that an inverse association between neuropsychological impairment and days post-injury is reliably observed when effects are aggregated over briefer follow-up intervals: “large” adverse effects observed within 24 hr of injury (e.g., -0.97 , Belanger & Vanderploeg, 2005), typically reduce to “moderate” or “small” effects within days to weeks (e.g., -0.43 to -0.22 , Belanger & Vanderploeg, 2005; -0.39 to -0.32 , Rohling et al., 2011; -0.41 to -0.29 , Schretlen & Shapiro, 2003). Importantly, resolution of post-concussion neuropsychological deficits (defined as a non-significant effect size) usually occurs within 7 to 10 days of sports-related concussion (Belanger & Vanderploeg, 2005), or within 1 to 3 months following mTBI (Rohling et al., 2011; Schretlen & Shapiro, 2003), although deficits may persist for a minority of individuals (Iverson, 2010). Previous meta-analyses also demonstrate that multiple follow-up assessments (*vs.* single) and pre-injury baseline comparisons (*vs.* control group comparisons) are associated with smaller aggregated effect sizes, attributed to the confounding effect of practice arising from repeat assessment (Belanger & Vanderploeg, 2005; Broglio & Puetz, 2008). For example, at first assessments conducted within 14 days of injury, Broglio and Puetz (2008) identified

“large” adverse effects on athletes’ neuropsychological function (-0.81 ; $p < .001$), symptom reports (-3.31 ; $p < .05$), and postural stability (-2.56 , *ns*); yet these effect sizes were substantially reduced upon repeat assessment within the same 14 day period (-0.26 ; $p = .001$; -1.09 ; $p < .05$; and -1.16 ; *ns*, respectively).

In each published meta-analysis, substantial effect size heterogeneity remained unexplained by the moderator variables evaluated. Moreover, the moderating effect of pre-existing athlete characteristics on concussion outcomes, and the extent to which other variables such as the method and timing of post-injury assessment explain this effect, has not been explored. The aim of the current study was, therefore, to redress the shortcomings of previous research by applying meta-analytic techniques to a contemporary sample of the sports-related concussion literature. This was to quantify the effect of sports-related concussion on neuropsychological, symptomatic, and postural functioning, and to identify the key athlete characteristics that moderate the magnitude of this effect, after controlling for the influence of variables known to moderate concussion outcomes (i.e., time since injury, repeat assessment and comparison group).

METHODS

Literature Search and Inclusion Criteria

Online databases (PsychINFO, PUBMED, MEDLINE) were searched for relevant papers published between January 1970 and August 2011.² Studies were located that quantified post-injury outcome from sports-related concussion in adolescent or adult athletes on at least one neuropsychological or cognitive test, measure of postural stability, or self-report symptom checklist relative to a pre-injury baseline and/or an independent control group (see Table 1 for additional criteria). Data included in this study were obtained in compliance with regulations of the University of Queensland.

Data Extraction and Effect Size Calculation

Statistical information required for effect size calculation (group means, standard deviations, and sample sizes) or effect size estimation (descriptive statistics extrapolated from graphs, or inferential statistics such as *F*-test, *t*-test, or *p*-values), as well as assessment and athlete variables required for moderator analyses, were coded in accordance with a detailed protocol. All effects were coded such that a post-injury decline in concussed athletes’ neuropsychological function or postural stability, or an increase in self-reported symptoms, would produce a negative effect size. Those measures for which results were described as “not statistically significant” by study authors, without accompanying descriptive or inferential statistics, were

¹ By convention, effect size (*d*) magnitudes $\geq .80$ are considered large, $.50$ moderate, and $\leq .20$ small (Cohen, 1988).

² For a detailed description of methodology, see online supplementary materials: http://www2.psy.uq.edu.au/~horswill/DouganHorswillGeffen_SupplementaryMaterials.pdf

Table 1. Criteria for inclusion of studies in the meta-analysis

Criterion	Description
Publication type	Published (peer-reviewed journal) Empirical research Methodology and results described in detail (sufficient to support planned effect size and moderator analyses) ^a English language
Participant characteristics	Adolescent and/or adult athletes Engaged in sport or athletic activities at the high-school, college, professional/elite, or amateur/non-professional club level of competition Age or level of competition reported ^b
Injury characteristics	Diagnosed with concussion or mild traumatic brain injury Diagnosed by a medical officer experienced in concussion assessment (e.g., certified athletic trainer or team physician); or identified using a specific clinical definition of concussion or system of grading injury severity (e.g., Quality Standards Subcommittee of the American Academy of Neurology, 1997) Sport-related mechanism of injury (practice or competition)
Outcome measures	Participants assessed with at least one psychometric test of neuropsychological function, experimental cognitive task, measure of dynamic or static postural stability with or without disrupted sensory input, or a self-rated scale or checklist of post-concussion symptom frequency and/or severity ^c Time elapsed between injury and first assessment reported First post-injury assessment conducted within 12 months of injury
Research design	Post-injury performance on outcome measure(s) compared to control group without acute concussion and/or injured athletes' pre-injury baseline performance Evaluation of outcome from injury (cf. intervention or management of injury)
Statistical information	Concussion data presented separately from injuries of greater severity (e.g., moderate to severe traumatic brain injuries), other aetiologies (e.g., falls or motor vehicle accidents), or other clinical conditions (e.g., sub-concussive blows, whiplash, malingering). Sufficient statistical information to calculate or estimate at least one effect size ^{d,e}

Note. ^aCase studies or concussion samples of less than four participants were excluded, as samples of this size would prohibit the calculation of estimates of variability required to conduct a meta-analysis (see Rohling, Beverly, Faust, & Demakis, 2009).

^bIf age was not reported by primary study authors, competition level was taken as a proxy for age such that high school athletes were presumed to be adolescents and professional, college, or amateur athletes were presumed to be adults for the purpose of moderator analyses.

^cReported as scale total scores vs. individual symptom frequencies.

^dIf effect size data were only reported for a subset of variables within a particular study, all effects were coded and only those that could not be estimated were excluded from the analysis.

^eIf an effect was reported only as "not statistically significant" an effect size of zero was entered into the analysis.

entered into the analysis conservatively as an effect size of zero (as per Frencham, Fox, & Maybery, 2005).

Effect sizes were calculated/estimated in Microsoft Office Excel 2003 according to research design-specific formulae for continuous variables described by Lipsey and Wilson (2001). Effect sizes were calculated by dividing the difference between the concussed group mean and the uninjured group (pre-injury baseline or independent control group) mean, by the pooled standard deviation of the concussed and uninjured group means (d_{pooled}).³ Weighted mean effect sizes were then computed by (1) aggregating multiple effects within a given sample (i.e., from two or more outcome measures, cognitive domains or post-injury assessments) by arithmetic mean to create an independent set of effects for each analysis, (2) applying Hedge's small-sample bias correction, (3) weighting by the inverse of the sampling error variance, and (4) aggregating across samples.

³ For results calculated using the standard deviation of the uninjured group mean only ($d_{control}$), see online supplementary materials.

Before aggregation, d_{pooled} effect size estimates were checked for extreme scores using standard techniques; none were identified. After aggregation, one sample-level effect size (from $k = 91$ d_{pooled} effects) was identified as an outlier in the negative direction (see Figure 1); but was retained unaltered for analysis as it was considered a genuine reflection of the data and its inclusion did not substantively influence the weighted mean effect size. The results of the meta-analysis were also found to be robust to the potential effects of both publication bias and selective reporting bias.⁴

Meta-analytic Model and Statistical Analyses

Analyses were conducted using Comprehensive Meta-Analysis Version 2 (Borenstein, Hedges, Higgins, & Rothstein, 2005). A fixed effects model was used to estimate the overall effect of concussion, such that aggregated (sample-level) effect sizes were assumed to estimate a single population effect with

⁴ For further details of these analyses, see online supplementary materials.




Fig. 1. Funnel plot of 91 independent aggregated effect sizes by the standard error of each effect size (weighted mean effect size, $d_{pooled} = -0.54$). The single outlying effect size is indicated by an unfilled data point (see Table 2, note f for sample details). By convention, effect size magnitudes $\geq .80$ are considered large, $.50$ moderate and $\leq .20$ small (Cohen, 1988).

variation arising only from random subject-level sampling error (Lipsey & Wilson, 2001). Weighted mean effect sizes were compared to the null hypothesis using a one-tailed z -test and the precision of the estimate was indicated by 95% confidence intervals. To evaluate the adequacy of these assumptions, homogeneity of effect size variance was tested using the Q statistic with $k-1$ degrees of freedom (Hedges & Olkin, 1985). Where heterogeneity was indicated by a statistically significant Q , the excess variability beyond subject-level sampling error was hypothesized to be attributable to systematic variation in measurable sample characteristics, rather than random variation, best represented by a “fixed effects model with systematic between-study variance” (Lipsey & Wilson, 2001, p. 118)⁵; moderator analyses were planned to examine this variation further.

First, moderator analyses were conducted to replicate the influence of variables previously shown to be important moderators of concussion: namely, outcome (neuropsychological function, self-report symptoms, postural stability), comparison group (pre-injury baseline, independent control, or both⁶), time elapsed between injury and assessment (<24 hr, 1–10 days, 10–30 days, >30 days⁷), and repeat assessment (first, second, third, or fourth post-injury assessment). Follow-up moderator analyses were also planned to investigate the hypothesized contribution of athlete characteristics to residual effect size variation, using either the analogue-to-ANOVA procedure for categorical variables (concussed athletes’ age

⁵ For results using a “mixed effects” model, see online supplementary materials.

⁶ The latter design represents the most rigorous research design, as it controls for both premorbid functioning (pre-injury baseline) and the effect of repeat assessment (control group).

⁷ Intervals were selected for consistency with documented neurometabolic and neurophysiologic recovery periods (Giza & Hovda, 2001, 2004), and with the assessment intervals commonly used within the empirical literature sampled.

group, sex, level of competition, and sport played by 50% or more of the sample; see Table 3), or fixed effect linear regression analyses for continuous variables (concussed athletes’ average age in years and average years of education). Samples that did not report a given moderator variable, or samples that could not be allocated to a single level of a moderator, were excluded from the relevant analysis.

RESULTS

Characteristics of Included Samples

Seventy-eight papers, describing 92 independent samples of sports-related concussion (see Table 2), were identified as eligible for inclusion in the meta-analysis against *a priori* criteria, although only 91 samples comprised of 3,801 concussed athletes and 5,631 controls were available for d_{pooled} analyses (see Table 2, note d). As shown in Table 2, the majority of included samples were recruited in the USA either exclusively from college competitions or from more than one level of competition, and were largely comprised of adult males playing American football or recruited from a variety of sports. The position played by concussed athletes and the mechanism of injury were rarely reported. Thirty-one samples ($n = 1,516$) reported concussed athletes’ average years of education. Other measures of premorbid functioning and neuropsychiatric history were reported too infrequently, too inconsistently or with insufficient variation for analysis.⁸

Overall Effect of Sports-Related Concussion

Aggregated across all outcome measures and post-injury assessments, the overall weighted mean effect size (d_{pooled}) represented a statistically significant “moderate”⁹ decrement in general functioning following sports-related concussion (-0.54 ; 95% confidence interval [CI]: $-0.57, -0.50$); based on 91 independent effects (range: -2.68 to 0.25), 87% of which represented a decline in post-injury functioning and 46% of which were “moderate” or “large” in magnitude. The overall effect was significantly heterogeneous ($Q(90) = 668$; $p < .001$); investigation of moderator variables was, therefore, considered appropriate.

The overall effect was comprised of a “small to moderate” decrement in neuropsychological functioning (-0.40 ; 95% CI: $-0.44, -0.36$; $Q(69) = 405$; $p < .001$), a “moderate to large” increase in self-reported symptoms (-0.66 ; 95% CI: $-0.70, -0.62$; $Q(49) = 603$; $p < .001$), and a “small” but significant decrement in postural stability (-0.11 ; 95% CI: $-0.18, -0.04$; $Q(21) = 55$; $p < .001$), when collapsed across all follow-up assessments. Neuropsychological outcomes varied marginally by comparison group: a “small to moderate” effect was derived from samples using a pre-injury baseline (-0.38 ; 95% CI: $-0.42, -0.34$; $Q(32) = 296$; $p < .001$),

⁸ For further details, see online supplementary materials.

⁹ By convention, effect size magnitudes $\geq .80$ are considered large, $.50$ moderate, and $\leq .20$ small (Cohen, 1988).

Table 2. Characteristics of the 92 sports-related concussion samples identified as eligible for inclusion in the meta-analysis, arranged by comparison group and aggregated effect size (d_{pooled})

Sample ^a	Concussed; controls			Country	Sport	Level	Concussed athletes' history of previous concussions ^b	TSI at each post-injury assessment ^c				Outcome measures					
	Sample size	% Male	Age (in years)					1 st	2 nd	3 rd	4 th	NP	SRS	PS	No. effects	d_{pooled}	
Pre-injury baseline and independent control group comparison																	
Barr & McCrea (2001)	50; 68	100; 100	17.2; 18.1	USA	F	HS/C	—	0					S			1	-1.74
Peterson et al. (2003)	24; 18	75; 75	20.2; 19.3	USA	M (F)	C	Yes	1	2	3	10		PnP	PnP	PC	20	-1.22
Field et al. (2003) – 1	19; 20	88; 88	15.2; 16.6	USA	M (F)	HS	Yes	<1	3	5	7		PnP	PnP		20	-0.87
Sosnoff et al. (2007)	22; 22	91;—	19.8;—	USA	M (-)	C	Yes	2					PC			7	-0.86
Iverson et al. (2003)	41; 56	90; 52	16.8; 17.6	USA	M (F)	A	—	1.3					PC	PC		5	-0.82
Piland et al. (2003)	17; 16	88; 88	19.8; 19.5	USA	M (F)	C	—	1	2	3	10			PnP		8	-0.75
McCrea (2001)	63; 55	100; 100	18.2; 18.2	USA	F	HS/C	—	0	<2				S			10	-0.71
Hinton-Bayre et al. (1997)	10; 10	100; 100	22.1; 19.9	AUS	R	P	Yes	1.5					PnP			5	-0.58
Guskiewicz et al. (1996)	10; 10	100; 100	17.4; 18.6	USA	F	HS/C	—	1	3	5	10				PC	20	-0.55
Lovell et al. (2003)	64; 24	94; 67	—;—	USA	M (F)	HS	Yes	1.5	4.2	7.6			PC	PC		6	-0.40
Cavanaugh et al. (2005)	27; 30	78; 50	19.5; 21.7	USA	M (F)	C	Yes	<2							PC	15	-0.39
Field et al. (2003) – 2	35; 18	94; 94	19.9; 20.1	USA	M (F)	C	Yes	<1	3	5	7		PnP	PnP		12	-0.37
Macciocchi et al. (1996)	183; 48	100; 100	19.0; 19.0	USA	F	C	Yes	1	5	10	84		PnP	PnP		22	-0.35
Guskiewicz et al. (2001)	36; 36	69; 69	19.5; 20.0	USA	M	C	—	1	3	5			PnP		C; PC	36	-0.33
McCrea et al. (2003)	94; 56	100; 100	20.0; 19.2	USA	F	C	Yes	0	<3hrs	1	2		S;	PnP	C	45	-0.25
								2	7	90			PnP				
Sim et al. (2008)	14; 14	79; 77	15.5; 15.7	USA	M (-)	HS	Yes	2.5	6.3	9.9	45		PC			24	-0.25
Echemendia et al. (2001)	29; 20	92; 92	—;—	USA	M	C	—	2hrs	2	7	30		PnP	PnP		81	-0.23
Lovell & Collins (1998)	4; 40	100; 100	—; 19.6	USA	F	M	Yes	<1	180				PnP			20	-0.23
Hinton-Bayre et al. (1999)	20; 13	100; 100	21.1; 19.6	AUS	R	P	Yes	2	10.5	28			PnP			9	-0.13
Collie et al. (2006)	61; 84	100; 100	22.9; 23.4	AUS	ARf	A	Yes	3					PnP; PC			9	-0.06
Makdissi (2001)	6; 7	100; 100	20.5; 20.3	AUS	ARf	C	—	2					PnP; PC			3	0.04
Johnson et al. (2002)	9; 9	44; 60	—;—	USA	M (R)	C	—	1	3	5	10			PnP	PC	8	0.25
Maddocks & Saling (1996)	10; 10	100; 100	—;—	AUS	ARf	P	Yes	5					PnP			4	— ^d
Independent control group comparison only																	
Bruce & Echemendia (2004) – 2	30; 147	100; 100	20.2; 19.0	USA	M	C	Yes	2hrs						PnP		1	-2.01
Bruce & Echemendia (2004) – 1	27; 286	100; 100	20.2; 18.8	USA	M	C	No	2hrs						PnP		1	-1.78
Lovell et al. (2006) – 2 ^c	39; 1,039	—; 77	—;—	USA	M (-)	C	—	2						PC		1	-1.68
Lovell et al. (2006) – 1 ^e	221; 707	—; 83	16.5;—	USA	M (-)	HS	—	2						PC		1	-1.53
Fazio et al. (2007)	122; 70	81; 47	16.7; 17.3	USA	M (F)	HS/C	Yes	1.9					PC			4	-1.16
Schatz et al. (2006)	72; 66	79; 44	16.5; 17.3	USA	M (F)	HS	Yes	2					PC	PC		6	-0.91
Maddocks et al. (1995)	28; 28	100; 100	—;—	AUS	ARf	P	—	10 mins					S			14	-0.83
Pellman et al. (2006) – 1	37; 125	100; 100	15.8; 15.6	USA	F	HS	Yes	1.5	5				PC	PC		10	-0.64
Gosselin et al. (2006)	20; 10	95; 90	25.9; 22.0	CAN	M (IH)	C/P	Yes	71.4					PnP; Exp	PnP		12	-0.61
Elleberg et al. (2007)	10; 12	0; 0	22.7; 22.3	CAN	S	C	No	246					PnP; PC			19	-0.61
Thompson et al. (2005)	12; 12	100; 100	21.0; 21.0	USA	M (-)	C	Yes	89							PC	2	-0.53
Parker et al. (2005)	10; 10	40; 40	20.2; 19.9	USA	M (-)	C	—	1.6							G	20	-0.53
Lovell et al. (2006) – 3	52; 1,746	90; 80	—;—	USA	M (-)	HS/C	—	—	5.6	11.7			PC			2	-0.43

(Continued)

Table 2. Continued

Sample ^a	Concussed; controls			Country	Sport	Level	Concussed athletes' history of previous concussions ^b	TSI at each post-injury assessment ^c				Outcome measures				
	Sample size	% Male	Age (in years)					1 st	2 nd	3 rd	4 th	NP	SRS	PS	No. effects	<i>d</i> _{pooled}
Pellman et al. (2006) – 2	48; 68	100; 100	26.3; 24.3	USA	F	P	Yes	1.2	2.9			PC	PC		10	-0.39
Riemann & Guskiewicz (2000)	16; 16	94; 94	19.2; 22.5	USA	—	C	—	1	3	5	10			C; PC	36	-0.38
Bruce & Echemendia (2003)	19; 19	100; 100	20.1; 19.9	USA	M (F)	C	Yes	2hrs	2	7	30	PnP			24	-0.37
Gosselin et al. (2009)	10; 11	70; 64	24.3; 22.6	CAN	M	C/P	Yes	132				PnP; PC	PnP		26	-0.32
Chen et al. (2008a)	16; 16	100; 100	26.0; 20.0	CAN	M (-)	C/P	Yes	219				Exp	PnP		6	-0.32
Parker et al., (2006)	15; 15	60; 60	20.6; 20.6	USA	M (-)	C	—	1.6	5	14	28			G	80	-0.29
Guskiewicz et al. (1997)	11; 11	73; 73	18.6; 20.2	USA	—	C	—	1	3	5	10	PnP		PC	37	-0.29
Chen et al. (2007)	18; 10	100; 100	28.9; 21.9	CAN	M (IH)	C/P	Yes	156				PC; Exp	PnP		25	-0.28
McCrea et al. (2002)	91; 45	100; 100	17.3; 17.5	USA	F	HS/C	—	—	15 mins	2	90	S			20	-0.24
Moser & Schatz (2002)	14; 21	79; 81	16.4; 16.8	USA	M (-)	HS	Yes	4				PnP			18	-0.12
Cremona-Meteyard & Geffen (1994) – Experiment 1	9; 12	100; 100	23.0; 22.1	AUS	ARf	P	Yes	<14	365			Exp			12	-0.09
Moser et al. (2005)	40; 183	—; —	15.8; 15.7	USA	M (-)	HS	Yes	3.5				PnP	PnP		9	-0.06
Chen et al. (2008b)	9; 6	100; 100	31.5; 20.0	CAN	M (IH)	C	Yes	90	547			Exp			4	-0.01
Dupuis et al. (2000)	20; 10	100; 100	21.5; 21.5	CAN	M (F)	C	Yes	171.8				PnP; Exp			11	0.00
Killam et al. (2005)	5; 9	60; 67	22.6; 22.0	USA	M (-)	C	Yes	73				PnP	PnP		10	0.07
Parker et al. (2008)	14; 14	—; —	20.7; 20.6	USA	M (-)	C	—	1.4	5	14	28			G	32	0.11
Ferguson et al. (1999)	50; 159	100; 100	20.2; 19.6	USA	M (IH)	HS/P	No	180					PnP		1	0.21
Pre-injury baseline comparison only																
McCrary et al. (2000)	23	100	—	AUS	ARf	P	—	15 mins				PnP			1	-2.68 ^f
Collins et al. (2003) – 1	34 (poor post-injury)	85	17.4	USA	M (F)	HS/C	Yes	1.4				PC	PC		2	-2.03
Slobounov, Tutwiler, et al. (2006)	8	100	21.0	USA	M (-)	C	No	3	10	30				PC	15	-1.52
Daniel et al. (2002)	21	100	—	USA	F	C	Yes	0				S			1	-1.30
Broshpek et al. (2005) – 1	37	0	17.5	USA	M	HS/C	Yes	3.8				PC			3	-1.10
Erlanger et al. (2003)	47	57	17.6	USA	M	HS/C	Yes	2.1				PC			3	-1.05
Collins et al. (2006)	136	100	16.1	USA	F	HS	Yes	2.2				PC	PC		5	-0.88
Cavanaugh et al. (2006) – 1	13 (unsteady post-injury)	—	—	USA	M	C	Yes	<2	3				PnP	PC	46	-0.84
Sosnoff et al. (2008)	36	81	21.2	USA	M (-)	C	—	<1				PC	PnP	PC	16	-0.77
Van Kampen et al. (2006)	122	82	16.6	USA	M (F)	HS/C	Yes	2				PC	PC		5	-0.76
Mihalik et al. (2007) – 1	155	84	15.6	USA	M (-)	HS	Yes	3.3				PC	PC		5	-0.75
Erlanger et al. (2001)	26	65	18.6	USA	M (-)	HS/C	—	1.8				PC			3	-0.73
Covassin et al. (2008) – 1	21	67	21.1	USA	M	C	Yes	1.2	5.1			PC	PC		10	-0.69
Slobounov et al. (2008)	12	53	21.2	USA	R	C	No	—	—	30				PC	11	-0.56
Broglio et al. (2007a) – 1	4 (complex concussion)	76	19.8	USA	M (F)	C	Yes	1.2	18.8			PC	PnP		12	-0.55
Mihalik et al. (2007) – 2	26	89	22.1	USA	M (-)	C	Yes	3.3				PC	PC		5	-0.55
McClincy et al. (2006)	104	88	16.1	USA	M (F)	HS/C	Yes	2.4	7.6	14.4		PC	PC		15	-0.54

(Continued)

Table 2. Continued

Sample ^a	Concussed; controls			Country	Sport	Level	Concussed athletes' history of previous concussions ^b	TSI at each post-injury assessment ^c				Outcome measures				
	Sample size	% Male	Age (in years)					1 st	2 nd	3 rd	4 th	NP	SRS	PS	No. effects	d_{pooled}
Covassin et al. (2008) – 2	36	47	20.6	USA	M	C	No	1.2	5.1			PC	PC		10	-0.49
Covassin et al. (2007) – 1	41	100	—	USA	M	C	Yes	1.9	8.1			PC	PC		10	-0.48
Iverson et al. (2006)	30	93	16.1	USA	M (F)	A	Yes	1.5	5.2	10.3		PC	PC		15	-0.48
Broshek et al. (2005) – 2	94	100	19.2	USA	M (F)	HS/C	Yes	2.8				PC			3	-0.48
Broglio et al. (2007b)	75	83	—	USA	M (F)	C	—	<1				PnP; PC	PnP	PC	27	-0.44
Covassin et al. (2007) – 2	39	0	—	USA	M	C	Yes	1.9	8.1			PC	PC		10	-0.40
Iverson et al. (2004) – 2	19	90	17.8	USA	M (F)	A	Yes	1.6				PC	PC		4	-0.40
Warden et al. (2001)	14	100	19.0	USA	B	C	—	—	4			PC			7	-0.34
Cavanaugh et al. (2006) – 2	16 (steady post-injury)	—	—	USA	M	C	Yes	<2	3				PnP	PC	46	-0.32
McCrea et al. (1998)	33	100	—	USA	F	HS/C	—	0	2			S			10	-0.31
Iverson et al. (2004) – 1	19	95	17.9	USA	M (F)	A	No	1.8				PC	PC		4	-0.31
Register-Mihalik et al. (2007) – 1	258 (no headache)	—	16.7	USA	M (-)	HS/C	Yes	1	3	7		S	PnP	C	12	-0.28
Lavoie et al. (2004)	10	100	21.5	CAN	M	C	Yes	51				PnP	PnP		14	-0.24
Slobounov et al. (2007)	38	55	21.2	USA	R	C	No	10	17	30		PnP		PC	11	-0.14
Register-Mihalik et al. (2007) – 2	106 (preseason headache)	—	16.7	USA	M (-)	HS/C	Yes	1	3	7		S	PnP	C	12	-0.07
Broglio et al. (2007a) – 2	17 (simple concussion)	76	19.8	USA	M (F)	C	Yes	1.2	5.6			PC	PnP		12	-0.02
Jantzen et al. (2004)	4	100	20.0	USA	F	C	—	3.5				Exp			4	0.00
Macciocchi et al. (2001)	12	100	19.1	USA	F	C	Yes	—	—	10	84		PnP		2	0.00
Slobounov, Slobounov, & Newell (2006)	10	100	19.5	USA	M (-)	C	No	—	—	30				PC	3	0.00
Lovell et al. (2004)	43	81	15.6	USA	M (F)	HS	Yes	1.4	6.3			PC	PC		8	0.01
Pellman et al. (2004)	95	100	25.4	USA	F	C/P	—	2.2				PnP			10	0.10
Collins et al. (2003) – 2	44 (good post-injury)	91	15.5	USA	M (F)	HS/C	Yes	2				PC	PC		2	0.16

Note. d_{pooled} = weighted mean effect size calculated using the pooled standard deviations of the concussed group and the uninjured comparison group as the denominator – aggregated across all post-injury assessments and all outcome measures; TSI = time since injury. Country: AUS = Australia; CAN = Canada; USA = United States of America. Sport: Arf = Australian Rules football; B = Boxing; F = American football; IH = ice hockey; M = multiple sports at risk of concussive injury (specify if >50% sample from single sport); R = Rugby; S = soccer. Level of competition: A = amateur/non-professional club; HS = high school; C = college; P = professional/elite; M = mixed levels. Outcome Measures: C = Clinical assessment of postural stability; Exp = Experimental/cognitive tasks; G = computerised assessment of gait stability under single and dual-task conditions; NP = neuropsychological tests; PC = computerised assessment; PnP = traditional pen-and-paper assessment; PS = postural stability assessment; S = sideline assessment of mental status; SRS = self-report symptoms.

^aFor full reference see asterisked (*) citations in References section.

^bFor further detail see on-line supplementary materials.

^cThe four columns (1st, 2nd, 3rd, 4th) represent the first four post-injury assessment occasions potentially conducted by a given study, while the number presented within each column represents the time elapsed between injury and that specific assessment occasion (TSI), reported as the average TSI, or mid-point of a reported TSI range; in days unless otherwise indicated.

^d d_{pooled} could not be calculated from the available data (post-injury standard deviations for concussed and control groups were not reported), leaving $k = 91$ for d_{pooled} analyses ($n = 3,801$ concussed athletes and 5,631 controls); cf. $k = 92$ for $d_{control}$ analyses ($n = 3,811$ concussed athletes and 5,641 controls) presented in the online supplementary materials.

^eData also presented separately for 217 males (1,391 controls), $d_{pooled} = -1.85$ and 43 females (355 controls), $d_{pooled} = -1.52$.

^fIdentified as an outlier relative to the overall mean effect size and the mean effect size for studies using a pre-injury baseline comparison only.

while a “moderate” effect was derived from samples using an independent control group (-0.46 ; 95% CI: $-0.58, -0.33$; $Q(18) = 45$; $p < .001$), and from samples using both comparisons (-0.51 ; 95% CI: $-0.62, -0.39$; $Q(17) = 60$; $p < .001$). Postural stability outcomes also varied by comparison group: a non-significant effect was derived from samples using a pre-injury baseline (-0.05 ; 95% CI: $-0.13, 0.02$; $Q(9) = 31$; $p < .001$), while a “small to moderate” effect was derived from samples using an independent control group (-0.34 ; 95% CI: $-0.66, -0.02$; $Q(5) = 2$, *ns*), and from samples using both comparisons (-0.44 ; 95% CI: $-0.66, -0.23$; $Q(5) = 8$, *ns*). In contrast, a “moderate” effect was derived from samples using a pre-injury baseline (-0.56 ; 95% CI: $-0.61, -0.51$; $Q(24) = 286$; $p < .001$), and from samples using both comparisons to assess self-report symptoms (-0.49 ; 95% CI: $-0.65, -0.34$; $Q(9) = 29$; $p < .001$), while a “large” effect was derived from samples using an independent control group only (-1.16 ; 95% CI: $-1.26, -1.07$; $Q(14) = 159$; $p < .001$).

Effect size diminished rapidly with increasing time since injury: a “moderate to large” effect was derived from all assessments conducted within 24 hr of injury (-0.76 ; 95% CI: $-0.82, -0.70$; $Q(29) = 119$; $p < .001$; $M = 12$ hr post-injury), a “small to moderate” effect was observed between 1 and 10 days post-injury (-0.44 ; 95% CI: $-0.47, -0.40$; $Q(66) = 671$; $p < .001$; $M = 3.7$ days), while a “small” homogenous effect was observed between 10 and 30 days post-injury (-0.13 ; 95% CI: $-0.23, -0.03$; $Q(11) = 7$, *ns*; $M = 23.3$ days), which was not significantly different from zero beyond 30 days (-0.06 ; 95% CI: $-0.18, 0.07$; $Q(18) = 18$, *ns*; $M = 143.5$ days). Effect size also diminished with repeat assessment: from “moderate to large” effects at first post-injury assessment (-0.71 ; 95% CI: $-0.75, -0.68$; $Q(85) = 539$; $p < .001$; $M = 18.2$ days post-injury), to “small to moderate” effects at second assessment (-0.24 ; 95% CI: $-0.29, -0.19$; $Q(42) = 145$; $p < .001$; $M = 35.3$ days), non-significant effects at third assessment (-0.01 ; 95% CI: $-0.06, 0.04$; $Q(28) = 85$; $p < .001$; $M = 13.2$ days), and “small” homogenous effects at fourth assessment (-0.15 ; 95% CI: $-0.28, -0.03$; $Q(16) = 40$, *ns*; $M = 29.1$ days).

Consequently, to control for the confound of recovery over time with the effect of repeat assessment, time since injury was re-analyzed including only first post-injury assessments: revealing a “moderate to large” effect within 24 hr of injury (-0.79 ; 95% CI: $-0.85, -0.72$; $Q(29) = 132$; $p < .001$; $M = 12$ hr post-injury) which remained “moderate to large” 1 to 10 days post-injury (-0.71 ; 95% CI: $-0.76, -0.66$; $Q(45) = 377$; $p < .001$; $M = 2.4$ days), but was non-significant and homogenous beyond 30 days (-0.09 ; 95% CI: $-0.30, 0.11$; $Q(10) = 9$, *ns*; $M = 126.5$ days).¹⁰ A regression analysis of first assessments conducted within 10 days post-injury confirmed a significant reduction in effect size magnitude with an increasing number of days since injury ($\beta = 0.06$; 95% CI: $0.03, 0.08$; $p < .001$; $\alpha = -0.84$; $k = 75$). Extrapolating from the model, concussed athletes

first assessed 24 hr following injury produced a “moderate to large” effect ($d_{pooled} = -0.78$), while athletes first assessed 10 days following injury produced a “small to moderate” effect ($d_{pooled} = -0.28$). The relationship between time and concussion effect was stronger than would be expected by chance ($Q_M(1) = 14$; $p < .001$), yet significant between-study variability remained unexplained by this model ($Q_R(73) = 479$; $p < .001$).

When first post-injury assessments were further analyzed by outcome, a “small to moderate” decrement in neuropsychological functioning (-0.38 ; 95% CI: $-0.45, -0.31$; $Q(19) = 141$; $p < .001$), a “large” increase in self-reported symptoms (-0.96 ; 95% CI: $-1.02, -0.89$; $Q(14) = 144$; $p < .001$), and a “moderate” decrement in postural stability (-0.45 ; 95% CI: $-0.52, -0.37$; $Q(13) = 27$; $p < .05$) were observed within 24 hr of injury, while a “moderate” decrement in neuropsychological functioning (-0.54 ; 95% CI: $-0.59, -0.50$; $Q(39) = 237$; $p < .001$), a “large” increase in self-reported symptoms (-1.14 ; 95% CI: $-1.20, -1.08$; $Q(24) = 203$; $p < .001$), and a “large” decrement in postural stability (-1.10 ; 95% CI: $-1.45, -0.75$; $Q(4) = 31$; $p < .001$) were observed 1 to 10 days post-injury (see Table 3). Furthermore, when only samples using both a baseline and control group comparison were included in the analysis (i.e., the most rigorous research design), a “large” decrement in neuropsychological functioning (-0.90 ; 95% CI: $-1.05, -0.76$; $Q(9) = 30$; $p < .001$), a “large” increase in self-reported symptoms (-1.49 ; 95% CI: $-1.72, -1.26$; $Q(6) = 26$; $p < .001$), and a “moderate to large” decrement in postural stability (-0.76 ; 95% CI: $-0.98, -0.54$; $Q(5) = 5$, *ns*) were observed within 24 hr of injury, while a “small to moderate” decrement in neuropsychological functioning (-0.41 ; 95% CI: $-0.57, -0.24$; $Q(8) = 14$; $p < .05$) and a “large” increase in self-reported symptoms (-0.91 ; 95% CI: $-1.23, -0.59$; $Q(1) = 0.1$, *ns*) remained at 1 to 10 days post-injury.¹¹

With the exception of postural stability within 24 hr, self-report symptoms within 1–10 days, and outcomes assessed beyond 10 days from injury, significant heterogeneity remained unexplained by these moderator analyses; additional analyses were, therefore, required. Insufficient samples were available for further analysis of outcomes within 24 hr of injury (e.g., $k = 1$ adolescent or high school athletes, $k = 0$ female athletes). Consequently, subsequent moderator analyses include outcomes first assessed during the 1- to 10-day follow-up interval only.

Athlete Characteristics

Age group

At first assessments conducted 1–10 days following injury, adolescent athletes demonstrated larger post-concussion neuropsychological deficits, on average, than adult athletes

¹⁰ Nil first assessments were conducted between 10 and 30 days from injury in the current meta-analytic sample.

¹¹ Nil postural stability assessments first assessed 1–10 days post-injury were compared to both a baseline and control group in the current meta-analytic sample.

Table 3. Effect size presented as a function of athlete characteristics and type of outcome measure: administered at first post-injury assessments conducted 1–10 days following a sports-related concussion

Athlete characteristics	Sample size		Neuropsychological tests				Self-report symptom scales				Postural stability assessment			
			d_{pooled}		Q		d_{pooled}		Q		d_{pooled}		Q	
	Concussed	Controls	k	Q	k	Q	k	Q	k	Q	k	Q	k	Q
Aggregated effect at 1–10 days	2,212	2,616	40	236.59***	25	202.90***	5	30.99***	25	202.90***	5	30.99***	5	30.99***
Adolescent (≤ 18 years)	641	1,140	9	33.41***	8	56.15***	—	—	8	56.15***	—	—	—	—
Adult (≥ 19 years)	677	1,350	18	48.78***	9	25.70***	—	—	9	25.70***	—	—	5	30.99***
100% Female	119	355	2	2.96	2	10.38***	—	—	2	10.38***	—	—	—	—
100% Male	880	1,766	13	59.29***	4	58.50***	—	—	4	58.50***	—	—	—	—
High school	796	1,140	9	33.41***	8	56.15***	—	—	8	56.15***	—	—	—	—
Professional/elite	87	103	4	0.69	1	—	—	—	1	—	—	—	—	—
College	408	1,156	11	26.88**	8	25.69***	—	—	8	25.69***	—	—	5	30.99***
Amateur	170	140	5	18.16***	4	4.64	—	—	4	4.64	—	—	—	—
American football	1,243	465	21	169.29***	16	155.97***	—	—	16	155.97***	—	—	—	—
Australian rules	76	103	3	0.04	—	—	—	—	—	—	—	—	—	—
Rugby union	68	23	3	3.01	—	—	—	—	—	—	—	—	—	—

Note. d_{pooled} = weighted mean effect size calculated using the pooled standard deviations of the concussed group and the uninjured comparison group as the denominator. By convention, effect size magnitudes $\geq .80$ are considered large, $.50$ moderate, and $\leq .20$ small (Cohen, 1988); k = number of independent sample effect sizes; Q = test of homogeneity of effect size variance. * $p < .05$, ** $p < .01$, *** $p < .001$.

(Table 3: $d_{pooled} = -0.60$ and -0.25 , respectively), and reported marginally more symptoms (Table 3: $d_{pooled} = -1.29$ and -1.07 , respectively), but were not assessed for postural stability. The significant difference in neuropsychological outcomes was not better accounted for by differences between adolescents and adults in average time since injury (Table 4: $M = 2.4$ and 2.7 days, respectively), type of comparison group (Table 4), or sample sex (Table 5). When only samples using both a baseline and control group comparison were included in analysis, adolescents demonstrated greater neuropsychological impairment than adults ($d_{pooled} = -0.69$ and -0.25 , respectively). Adolescent males also demonstrated substantially larger neuropsychological deficits than adult males ($d_{pooled} = -0.75$ and -0.15 , respectively). Adolescent females were not available for comparison to adult females in the current sample.

Age in years

Regression analyses confirmed that each additional year of concussed athletes' average age (range: 15.2 to 31.5 years) corresponded to a significant reduction in the magnitude of the overall effect of concussion ($\beta = 0.04$; 95% CI: 0.03, 0.06; $p < .001$; $\alpha = -1.28$; $k = 76$), and the effect size magnitude when only neuropsychological outcomes, first post-injury assessments conducted 1–10 days from injury, and studies using both baseline and control group comparisons were included in analyses ($\beta = 0.11$; 95% CI: 0.03, 0.18; $p < .01$; $\alpha = -2.50$; $k = 8$). Holding these variables constant, the relationship between age and concussion effect was stronger than would be expected by chance ($Q_M(1) = 8$; $p < .01$), while residual between-study variability was not significant ($Q_R(6) = 5$, *ns*). Extrapolating from the model, athletes at 15 years of age could be expected to demonstrate a “large” decrement in neuropsychological functioning upon first assessment within 1–10 days post-concussion ($d_{pooled} = -0.92$), while adult athletes over the age of 24 years could be expected to demonstrate a minimal effect within the same interval ($d_{pooled} = 0.02$).

Years of education

Regression analyses also indicated that each additional year of concussed athletes' education (range: 9.6 to 16.6 years) corresponded to a significant reduction in the magnitude of the overall effect of concussion ($\beta = 0.16$; 95% CI: 0.12, 0.19; $p < .001$; $\alpha = -2.49$; $k = 31$), and the effect size magnitude when only neuropsychological outcomes, first post-injury assessments conducted 1–10 days from injury, and studies using both baseline and control group comparisons were included in analyses ($\beta = 0.20$; 95% CI: 0.02, 0.37; $p < .05$; $\alpha = -3.28$; $k = 5$). Holding these variables constant, the relationship between years of education and concussion effect was stronger than would be expected by chance ($Q_M(1) = 5$; $p < .05$), while residual between-study variability was not significant ($Q_R(3) = 6$, *ns*). Extrapolating from the model, athletes with 10 years of education could be

Table 4. Effect size presented as a function of athlete characteristics and comparison group: neuropsychological outcome measures administered at first post-injury assessments conducted 1–10 days following sports-related concussion

Athlete characteristics	Aggregated over comparison groups						Comparison group ^a at first assessment 1–10 days post-injury (neuropsychological outcomes only)					
	Both baseline and control group			Independent control group only			Pre-injury baseline only			Pre-injury baseline only		
	d_{pooled}	k	Q	d_{pooled}	k	Q	d_{pooled}	k	Q	d_{pooled}	k	Q
Aggregated effect at 1–10 days	-0.54***	40	236.59***	-0.41***	9	13.94*	-0.59***	7	31.89***	-0.55***	24	187.62***
Adolescent (≤ 18 years)	-0.60***	9	33.41***	-0.69***	2	0.09	-0.44***	4	15.38**	-0.63***	3	14.75***
Adult (≥ 19 years)	-0.25***	18	48.78***	-0.25*	6	6.53	-0.39*	2	0.46	-0.23***	10	41.06***
100% Female	-0.87***	2	2.96	—	—	—	—	—	—	-0.87***	2	2.96
100% Male	-0.42***	13	59.29***	-0.18	5	2.39	-0.48***	3	1.08	-0.45***	5	50.23***
High school	-0.60***	9	33.41***	-0.69***	2	0.09	-0.44***	4	15.38**	-0.63***	3	14.75***
Professional/Elite	-0.43**	4	0.69	-0.53	2	0.02	-0.39*	2	0.46	—	—	—
College	-0.41***	11	26.88**	-0.34*	2	3.48	—	—	—	-0.43***	9	23.13**
Amateur	-0.42***	5	18.16***	-0.34*	2	7.17**	—	—	—	-0.48***	3	10.32**

Note. d_{pooled} = weighted mean effect size calculated using the pooled standard deviations of the concussed group and the uninjured comparison group as the denominator. By convention, effect size magnitudes $\geq .80$ are considered large, .50 moderate and $\leq .20$ small (Cohen, 1988); k = number of independent sample effect sizes; Q = test of homogeneity of effect size variance; TSI = average time elapsed since injury (in days).
^aFor a full break-down of results by athlete characteristics, time since injury, and comparison group see online supplementary materials, Appendix B, Table B2.
 * $p < .05$, ** $p < .01$, *** $p < .001$.

expected to demonstrate a “large” decrement in neuropsychological functioning upon first assessment within 1–10 days post-concussion ($d_{pooled} = -1.32$), while those with 16 years of education could be expected to demonstrate a “small” effect within the same interval ($d_{pooled} = -0.15$). Insufficient samples were available for analysis of the interaction between age, sex, and years of education.

Sex

At first assessments conducted 1–10 days following injury, female athletes demonstrated larger post-concussion neuropsychological deficits, on average, than male athletes (Table 3: $d_{pooled} = -0.87$ and -0.42 , respectively), although males reported more symptoms than females (Table 3: $d_{pooled} = -1.58$ and -1.12 , respectively); females were not assessed for postural stability. The significant difference in neuropsychological outcomes was not better accounted for by differences between females and males in average time since injury (Table 4: $M = 2.9$ and 2.5 days, respectively), type of comparison group (Table 4), or age group (Table 5). When only samples using a baseline comparison were included in analysis, females demonstrated greater neuropsychological impairment than males ($d_{pooled} = -0.87$ and -0.42 , respectively); females were not assessed using both a baseline and control group comparison or control group only. Female adults also demonstrated substantially larger neuropsychological deficits than male adults ($d_{pooled} = -0.62$ and -0.15 , respectively). Female adolescents were not available for comparison to male adolescents in the current meta-analytic sample.

Level of competition

At first assessments conducted 1–10 days following injury, athletes injured during high school competition demonstrated larger post-concussion neuropsychological deficits, on average, than athletes concussed at other levels of competition (Table 3: $d_{pooled} = -0.60$ and range: -0.43 to -0.41 , respectively), and reported marginally more symptoms (Table 3: $d_{pooled} = -1.29$ and range: -1.07 to -1.02 , respectively), but were not assessed for postural stability. The significant difference in neuropsychological outcomes was not better accounted for by differences between levels of competition in average time since injury (Table 4: $M = 2.4$ and range: 1.8 to 2.8 days, respectively) or type of comparison group (Table 4). When only samples using both a baseline and control group comparison were included in analysis, high school athletes demonstrated greater neuropsychological impairment than other levels of competition ($d_{pooled} = -0.69$ and range: -0.34 to ns , respectively). Insufficient samples were available for analysis of the moderating effect of level of competition by sample age, sex, or years of education.

Sport played

At first assessments conducted 1–10 days following injury, samples predominantly recruiting American footballer players

Table 5. Effect size presented as a function of athlete age and sex: neuropsychological outcome measures administered at first post-injury assessments conducted 1–10 days following sports-related concussion

Sex of sample	Age group at first assessment 1–10 days post-injury (neuropsychological outcomes only)							
	Adolescent (≤ 18 years)				Adult (≥ 19 years)			
	d_{pooled}	k	Q	TSI	d_{pooled}	k	Q	TSI
100% Female	—	—	—	—	−0.62**	1	—	1.9
100% Male	−0.75***	2	0.78	1.9	−0.13* ^a	10	15.71	2.6 ^a

Note. d_{pooled} = weighted mean effect size calculated using the pooled standard deviations of the concussed group and the uninjured comparison group as the denominator. By convention, effect size magnitudes $\geq .80$ are considered large, $.50$ moderate and $\leq .20$ small (Cohen, 1988); k = number of independent sample effect sizes; Q = test of homogeneity of effect size variance; TSI = average time elapsed since injury (in days).

^aIf adult male samples are matched to adolescent male and adult female samples on TSI (≥ 3 days excluded): adult male $d_{pooled} = -0.15^*$, $k = 7$, $Q = 15.37^*$, TSI $M = 1.8$ days.

* $p < .05$, ** $p < .01$, *** $p < .001$.

demonstrated “moderate” neuropsychological deficits and a “large” increase in self-report symptoms (Table 3: $d_{pooled} = -0.53$ and -1.08 , respectively), comparable in magnitude to the overall meta-analytic sample. A single sample of Rugby union players demonstrated “large” postural stability deficits within 1–10 days following concussion ($d_{pooled} = -2.25$). However, samples of Australian Rules and Rugby union football players did not demonstrate a statistically significant change in neuropsychological function within the same period of assessment. As the majority of samples included in this meta-analysis recruited athletes from a variety of sports, other sports such as ice hockey, soccer, and boxing were not sufficiently represented within the sample to support individual analysis.

DISCUSSION

This review used meta-analytic techniques to quantify the impact of sports-related concussion on athletes’ neuropsychological, symptomatic, and postural functioning. It used a contemporary sample to replicate and extend findings from both quasi-experimental empirical studies and previous meta-analyses of the concussion literature, included almost double the number of studies previously reviewed, and is the first meta-analysis to investigate the role of pre-existing athlete characteristics in moderating outcomes associated with sports-related concussion.

Consistent with previous meta-analyses, the aggregated effect of concussion was heterogeneous and significantly moderated by outcome, time since injury, repeat assessment, and comparison group¹². Athletes consistently demonstrated a significant post-concussion increase in subjective symptom reports and significant impairment on objective measures of neuropsychological function and postural stability. This pattern of results was observed when outcomes were collapsed across all post-injury assessments, and also (although to a greater magnitude) when only first post-injury assessments conducted within 10 days of injury, and only those studies using both a baseline and independent control group comparison,

were included in analyses to control for the attenuating effects of repeat assessment. The finding of significant post-concussion postural instability is consistent with previous reports of the sensitivity of postural assessment to concussion sequelae (see Davis, Iverson, Guskiewicz, Ptito, & Johnston, 2009), and suggests that this may be a promising, although presently under-used, measure of concussion-related impairment in athletes’ psychomotor function—a domain with particular relevance for athletes’ competitive performance and risk of re-injury if returned to play within the acute post-injury period. While this finding differs from the non-significant postural stability deficits previously reported by Broglio and Puetz (2008), the increased sample of the current meta-analysis and the unmasking of significant deficits by controlling for the confounding effect of repeat assessment across smaller recovery intervals (<24 hr and 1–10 days vs. <14 days) may explain this difference.

At first assessments conducted 1–10 days following sports-related concussion, more severe deficits in neuropsychological functioning were demonstrated by concussed samples comprised of younger athletes (particularly those in their adolescence competing at the high school level of competition), female athletes, and those with fewer years of education, than samples comprised of older athletes, male athletes, or those with more years of education, respectively. This finding was not better accounted for by differences between groups in the number or timing of post-injury assessments or the control comparison made. The moderating effect of age group also remained when sample sex was held constant, and vice versa. Hence, converging evidence from multiple measures (age group, age in years, level of competition, and years of education) indicates that young age may be a reliable indicator for the potential severity of post-concussion neuropsychological deficits within the first days or weeks following injury. Conversely, these results suggest that older age or higher education may represent important protective factors during this early post-injury period (associated with increased brain/cognitive reserve, see Kaplan et al., 2009; Satz, 1993; Stern, 2009).

However, it must be emphasized that our finding of a significant moderating effect of athlete age, sex and education can be reliably applied to neuropsychological outcomes

¹² For further discussion, see online supplementary materials.

within the acute (1–10 days) recovery interval only. With insufficient studies reporting effects outside of this interval, conclusions regarding the immediate (<24 hr) or longer-term (>10 days) impact of athlete variables on sports-related concussion outcomes could not be reliably generated from the extant literature. The lack of follow-up of concussed athletes beyond 10 days and the limited assessment of symptom reports or postural stability also thwarted our investigation of differential rates of recovery, despite indications in the empirical literature that post-concussion recovery may be slower for high school athletes than college athletes (Baillargeon et al., 2012; McClincy, Lovell, Pardini, Collins, & Spore, 2006), or professional athletes (Moser et al., 2007), and that recovery may proceed at differential rates across neuropsychological functioning, subjective symptoms and postural control (Echemendia, Putukian, Mackin, Julian, & Shoss, 2001; Makdissi, 2009; McCrea et al., 2003).

The finding of greater post-concussion neuropsychological deficits in female and young athletes within the first 10 days of injury is nonetheless concerning given increasing rates of participation in contact sports, and hence an increasing exposure to sports-related concussion, in these populations (Dick, 2009; Guskiewicz et al., 2000). Sex- and age-related differences in the severity of early sports-related concussion outcomes may be attributable to a range of physiological, metabolic, hormonal, neurodevelopmental, neuroanatomical, or muscular (especially neck) characteristics that differ between males and females and between adolescents and adults (Anderson & Moore, 1995; Dick, 2009; Lovell & Fazio, 2008; Reddy, Collins, & Gioia, 2008; Viano, Casson, & Pellman, 2007). For example, female brains generally demonstrate greater metabolic requirements than male brains (see Broshek et al., 2005), which, in the presence of acute concussive stimuli, may produce an amplified cellular response to concussion-induced metabolic demands and changes in regional cerebral blood flow (Hovda et al., 1999). Additionally, evidence of altered intracranial blood pressure, prolonged and diffuse cerebral swelling, and excitotoxic sensitivities to concussion-activated neurotransmitters (e.g., glutamate) has been recorded in developmental animal models and following moderate to severe traumatic brain injury in children and adolescents; suggesting a potential vulnerability of the developing adolescent brain to the early effects of concussion (Bruce et al., 1981; McDonald & Johnston, 1990; Prins, Lee, Cheng, Becker, & Hovda, 1996). However, research regarding the underlying cause of sex- and age-related differences in initial severity and recovery from brain injury has generally been limited to animal models or non-athletic populations with more severe injuries, and includes reports of both protective and detrimental characteristics associated with developmental age or female gonadal hormones (Dick, 2009; Reddy et al., 2008). Further research is, therefore, needed to clarify the specific underlying contributors to the moderating effect of athlete characteristics on acute injury outcomes, and to determine whether these factors also contribute to a differential rate of recovery from concussion.

Alternatively, these findings may be explained by behavioral factors that vary systematically with pre-existing athlete characteristics to predict concussion outcome. For example, male and female athletes, athletes of different age groups, or athletes participating at different levels of competition, may adopt a style of play that is more or less aggressive, daring, or reckless—leading to differences in behavioral risk factors and the biomechanics of concussive injuries subsequently sustained. Indeed, a systematic review of studies reporting injury mechanisms within football, ice hockey and basketball, demonstrated that males are more likely to be concussed by player-to-player contact than females, while female athletes are more likely to make contact with a non-human object (Dick, 2009). Differences in the heading behavior of male and female soccer players have also been documented (Kontos, Dolese, Elbin, Covassin, & Warren, 2011). However, the current findings were unable to distinguish whether or not variations in the mechanism of injury were associated with acute concussion effects and/or a post-acute divergence in recovery.

The literature also suggests that demographic differences in measured concussion outcomes may be attributable to differences in athletes' psychosocial response to injury and style of symptom reporting. For example, in the wider population, women are more likely to report an illness, seek medical assistance, or report subjective symptoms than males (see Farace & Alves, 2000). Female gender is also a significant predictor of post-concussion symptom complaints 1 month following mTBI treated in hospital emergency departments (Bazarian et al., 1999). In contrast, female athletes reported fewer symptoms than males in the current meta-analysis, yet were more impaired than males on objective neuropsychological outcome measures at 1–10 days post-injury (Broshek et al., 2005; Covassin, Schatz, & Swanik, 2007). Consistent with this result, a meta-analysis of outcomes from mixed-severity traumatic brain injury in hospital attendees found that females were worse off than males on 85% of outcome variables, including both objective and subjective measures (Farace & Alves, 2000). Further epidemiological research is yet needed to confirm these potential differences in the mechanisms of sports-related concussion, rates of injury notification, and characteristics of subjective symptom reporting across different athletic sub-groups.

It must also be considered that observed differences in effect size magnitude across athletic sub-groups may (1) be an artifact of the specific meta-analytic methodology adopted, (2) be better explained by other potential moderators of concussion, or (3) be a spurious result arising from small sample sizes in certain cells of analysis. The findings of this meta-analysis will, therefore, require replication as the literature base expands and a cautious interpretation is recommended in the interim.

We argue, however, that the current findings are unlikely to be an artifact of variation in meta-analysis methodology, as we deliberately selected methods that were consistent with published meta-analyses in the field and/or could be expected to produce the most reliable results (Lipsey & Wilson, 2001).

Moreover, our findings were not dependent upon the specific effect size formulae and/or statistical model used.¹³ Additionally, our analyses yielded homogenous effects from assessments conducted beyond 10 days from injury, and from analyses of age group/in years, age by sex, years of education, and level of competition when only first neuropsychological assessments conducted 1–10 days post-injury using both baseline and control comparisons were included. However, some heterogeneity remained unexplained in the current study despite the addition of new moderator and confound analyses. Sex, age, years of education, and level of competition are likely to be inter-correlated, yet there were insufficient samples in the current study to analyze the independent effect of each of these variables. Homogenous effects may also have been revealed had there been a sufficient number of samples to further stratify results according to the cognitive domain assessed, computerized *versus* pen-and-paper assessment, or finer follow-up intervals. Other infrequently reported variables may also be required to account for this unexplained variance, for example, sample prevalence of neuropsychiatric factors including history of previous head injuries, learning disorders or attention-deficit/hyperactivity disorder (Solomon & Haase, 2008), indicators of injury severity including immediate post-concussion signs and symptoms (Alla, Sullivan, Hale, & McCrory, 2009), or objective biomarkers of injury such as measures of postural stability, or as-yet-experimental electrophysiological, genetic, or blood markers of injury (Barr, Prichep, Chabot, Powell, & McCrea, 2012; Davis et al., 2009).

Moreover, samples in this field of research were disproportionately comprised of American male athletes in their early adulthood, competing at mixed levels of competition and across mixed sports for which a breakdown of results by sample characteristics were infrequently reported. Female athletes, adolescent athletes competing in high school competitions, and those playing sports other than American football were underrepresented throughout the empirical literature, limiting the evidence-base available to inform injury management for these athletes. Similarly, preseason screening of athletes' premorbid neuropsychological characteristics (e.g., years of education, academic achievement and relevant developmental, medical or neurological history), use of objective postural stability measures, comparison to both a baseline and independent control group, and follow-up assessment beyond 10 days were infrequently reported. Consequently, our conclusions regarding the moderating effect of athlete characteristics on concussion outcomes are based upon only a small subset of the literature and may represent a difference of only marginal *clinical* significance. We, therefore, echo repeated calls for future research to (1) target these underrepresented athletic sub-groups, (2) use objective outcome measures across the full period of follow-up (0–90 days), (3) compare post-concussion performance to both a pre-injury baseline and independent control group,

and (4) stratify results according to athlete demographic variables (age, sex, years of education, level of competition, and sport played) as well as other pre-existing or injury-related factors to support future replication and extension of the current findings.

CONCLUSION

The findings of this meta-analytic review (1) verify the significant moderating effect of athlete characteristics on acute concussion outcomes identified in the empirical literature, (2) provide evidence for the importance of an individualized and conservative approach to the assessment and management of concussive injuries, particularly during the first 10 days of injury when important return-to-play decisions are often made, and (3) encourage the targeted development of empirically-derived assessment protocols and management guidelines tailored specifically for sub-groups of potentially vulnerable athletes.

Clinical decision-making should take into consideration key risk factors for greater deficits in neuropsychological function within the first 10 days following sports-related concussion, namely an athlete's young age, female sex, fewer years of education, or high school level of competition to mitigate the risk of premature return to play and repeat injury during this critical recovery interval. However, uncertainty remains with regard to the association between athlete characteristics and differential rates of symptom reporting, postural instability, or neuropsychological deficits beyond 10 days given insufficient empirical research targeting these outcomes. Future research should adopt rigorous research designs (large-scale prospective studies using both baseline and matched controls), use objective measures of post-concussion outcome (neuropsychological and postural stability assessments) in addition to subjective symptom reports, control for the impact of repeat assessment throughout the acute and post-acute phases of injury, and carefully document variations in response to injury according to pre-existing athlete characteristics to support further examination of the rate of recovery from sports-related concussion specific to each athletic sub-group.

ACKNOWLEDGMENTS

This research forms part of the doctoral thesis of the first author, submitted to the University of Queensland, and was supported by an Australian Postgraduate Award. This research received no specific grant from any other funding agency, commercial or not-for-profit sectors. We confirm that the information in this manuscript and the manuscript itself has never been published either electronically or in print and that none of the authors has any financial or other conflict of interest affecting this manuscript. We acknowledge the help of John Guimelli in checking the data for accuracy.

Supplementary material

To view supplementary material for this article, please visit <http://dx.doi.org/10.1017/S1355617712001464>

¹³ For further details, see online supplementary materials.

REFERENCES

References marked with an asterisk (*) indicate papers included in the meta-analysis.

- Alla, S., Sullivan, S.J., Hale, L., & McCrory, P. (2009). Self-report scales/checklists for the measurement of concussion symptoms: A systematic review. *British Journal of Sports Medicine*, 43(Suppl. 1), i3–i12.
- Anderson, V., & Moore, C. (1995). Age at injury as a predictor of outcome following pediatric head injury: A longitudinal perspective. *Child Neuropsychology*, 1, 187–202.
- Baillargeon, A., Lassonde, M., Leclerc, S., & Ellemberg, D. (2012). Neuropsychological and neurophysiological assessment of sport concussion in children, adolescents and adults. *Brain Injury*, 26(3), 211–220.
- *Barr, W.B., & McCrea, M. (2001). Sensitivity and specificity of standardized neurocognitive testing immediately following sports concussion. *Journal of the International Neuropsychological Society*, 7(6), 693–702.
- Barr, W.B., Prichep, L.S., Chabot, R., Powell, M.R., & McCrea, M. (2012). Measuring brain electrical activity to track recovery from sport-related concussion. *Brain Injury*, 26(1), 58–66.
- Bazarian, J.J., Wong, T., Harris, M., Leahy, N., Mookerjee, S., & Dombrov, M. (1999). Epidemiology and predictors of post-concussion syndrome after minor head injury in an emergency population. *Brain Injury*, 13(3), 173–189.
- Belanger, H.G., Curtiss, G., Demery, J.A., Lebowitz, B.K., & Vanderploeg, R.D. (2005). Factors moderating neuropsychological outcomes following mild traumatic brain injury: A meta-analysis. *Journal of the International Neuropsychological Society*, 11(3), 215–227.
- Belanger, H.G., & Vanderploeg, R.D. (2005). The neuropsychological impact of sports-related concussion: A meta-analysis. *Journal of the International Neuropsychological Society*, 11(4), 345–357.
- Borenstein, M., Hedges, L., Higgins, J., & Rothstein, H. (2005). *Comprehensive Meta-Analysis (Version 2) [Computer software]*. Englewood, NJ: Biostat.
- *Broglia, S.P., Macciocchi, S.N., & Ferrara, M.S. (2007a). Neurocognitive performance of concussed athletes when symptom free. *Journal of Athletic Training*, 42(4), 504–508.
- *Broglia, S.P., Macciocchi, S.N., & Ferrara, M.S. (2007b). Sensitivity of the concussion assessment battery. *Neurosurgery*, 60(6), 1050–1058.
- Broglia, S.P., & Puetz, T.W. (2008). The effect of sport concussion on neurocognitive function, self-report symptoms and postural control: A meta-analysis. *Sports Medicine*, 38(1), 53–67.
- *Broshek, D.K., Kaushik, T., Freeman, J., Erlanger, D., Webbe, F., & Barth, J.T. (2005). Sex differences in outcome following sports-related concussion. *Journal of Neurosurgery*, 102(5), 856–863.
- Bruce, D.A., Alavi, A., Bilaniuk, L., Dolinskas, C.A., Obrist, W.D., & Uzzell, B. (1981). Diffuse cerebral swelling following head injuries in children: The syndrome of “malignant brain edema”. *Journal of Neurosurgery*, 54(2), 170–178.
- *Bruce, J.M., & Echemendia, R.J. (2003). Delayed-onset deficits in verbal encoding strategies among patients with mild traumatic brain injury. *Neuropsychology*, 17(4), 622–629.
- *Bruce, J.M., & Echemendia, R.J. (2004). Concussion history predicts self-reported symptoms before and following a concussive event. *Neurology*, 63(8), 1516–1518.
- *Cavanaugh, J.T., Guskiewicz, K.M., Giuliani, C., Marshall, S., Mercer, V., & Stergiou, N. (2005). Detecting altered postural control after cerebral concussion in athletes with normal postural stability. *British Journal of Sports Medicine*, 39(11), 805–811.
- *Cavanaugh, J.T., Guskiewicz, K.M., Giuliani, C., Marshall, S., Mercer, V.S., & Stergiou, N. (2006). Recovery of postural control after cerebral concussion: New insights using approximate entropy. *Journal of Athletic Training*, 41(3), 305–313.
- Centers for Disease Control and Prevention. (2007). Nonfatal traumatic brain injuries from sports and recreation activities – United States, 2001–2005. *MMWR Morbidity and Mortality Weekly Report*, 56, 733–737.
- *Chen, J.K., Johnston, K.M., Collie, A., McCrory, P.R., & Ptito, A. (2007). A validation of the post concussion symptom scale in the assessment of complex concussion using cognitive testing and functional MRI. *Journal of Neurology, Neurosurgery, & Psychiatry*, 78(11), 1231–1238.
- *Chen, J.K., Johnston, K.M., Petrides, M., & Ptito, A. (2008a). Neural substrates of symptoms of depression following concussion in male athletes with persisting post-concussion symptoms. *Archives of General Psychiatry*, 65(1), 81–89.
- *Chen, J.K., Johnston, K.M., Petrides, M., & Ptito, A. (2008b). Recovery from mild head injury in sports: Evidence from serial functional magnetic resonance imaging studies in male athletes. *Clinical Journal of Sport Medicine*, 18(3), 241–247.
- Cohen, J. (1988). *Statistical power analysis for the behavioral sciences* (2nd ed.). Hillsdale, NJ: Erlbaum.
- *Collie, A., Makdissi, M., Maruff, P., Bennell, K., & McCrory, P.R. (2006). Cognition in the days following concussion: Comparison of symptomatic versus asymptomatic athletes. *Journal of Neurology, Neurosurgery, and Psychiatry*, 77(2), 241–245.
- *Collins, M.W., Iverson, G.L., Lovell, M.R., McKeag, D.B., Norwig, J., & Maroon, J. (2003). On-field predictors of neuropsychological and symptom deficit following sports-related concussion. *Clinical Journal of Sport Medicine*, 13(4), 222–229.
- *Collins, M.W., Lovell, M.R., Iverson, G.L., Ide, T., & Maroon, J. (2006). Examining concussion rates and return to play in high school football players wearing newer helmet technology: A three-year prospective cohort study. *Neurosurgery*, 58(2), 275–286.
- *Covassin, T., Schatz, P., & Swanik, C.B. (2007). Sex differences in neuropsychological function and post-concussion symptoms of concussed collegiate athletes. *Neurosurgery*, 61(2), 345–351.
- *Covassin, T., Stearne, D., & Elbin, R. (2008). Concussion history and postconcussion neurocognitive performance and symptoms in collegiate athletes. *Journal of Athletic Training*, 43(2), 119–124.
- *Cremona-Meteyard, S.L., & Geffen, G.M. (1994). Persistent visuospatial attention deficits following mild head injury in Australian rules football players. *Neuropsychologia*, 32(6), 649–662.
- *Daniel, C., Nassiri, J.D., Wilckens, J., & Land, B.C. (2002). The implementation and use of the Standardized Assessment of Concussion at the U.S. Naval Academy. *Military Medicine*, 167(10), 873–876.
- Daniel, R.W., Rowson, S., & Duma, S.M. (2012). Head impact exposure in youth football. *Annals of Biomedical Engineering*, 40, 976–981. doi:10.1007/s10439-012-0530-7
- Davis, G.A., Iverson, G.L., Guskiewicz, K.M., Ptito, A., & Johnston, K. (2009). Contributions of neuroimaging, balance testing, electrophysiology and blood markers to the assessment of sport-related concussion. *British Journal of Sports Medicine*, 43(Suppl. 1), i36–i45.

- Dick, R.W. (2009). Is there a gender difference in concussion incidence and outcomes? *British Journal of Sports Medicine*, 43(Suppl. 1), i46–i50.
- *Dupuis, F., Johnston, K.M., Lavoie, M., Lepore, F., & Lassonde, M. (2000). Concussions in athletes produce brain dysfunction as revealed by event-related potentials. *Neuroreport: For Rapid Communication of Neuroscience Research*, 11(18), 4087–4092.
- *Echemendia, R.J., Putukian, M., Mackin, R.S., Julian, L., & Shoss, N. (2001). Neuropsychological test performance prior to and following sports-related mild traumatic brain injury. *Clinical Journal of Sport Medicine*, 11(1), 23–31.
- *Elleberg, D., Leclerc, S., Couture, S., & Daigle, C. (2007). Prolonged neuropsychological impairments following a first concussion in female university soccer athletes. *Clinical Journal of Sport Medicine*, 17(5), 369–374.
- *Erlanger, D., Kaushik, T., Cantu, R.C., Barth, J.T., Broshek, D.K., Freeman, J.R., & Webbe, F.M. (2003). Symptom-based assessment of the severity of a concussion. *Journal of Neurosurgery*, 98(3), 477–484.
- *Erlanger, D., Saliba, E., Barth, J., Almquist, J., Webright, W., & Freeman, J. (2001). Monitoring resolution of postconcussion symptoms in athletes: Preliminary results of a web-based neuropsychological test protocol. *Journal of Athletic Training*, 36(3), 280–287.
- Farace, E., & Alves, W.M. (2000). Do women fare worse: A metaanalysis of gender differences in traumatic brain injury outcome. *Journal of Neurosurgery*, 93(4), 539–545.
- *Fazio, V.C., Lovell, M.R., Pardini, J.E., & Collins, M.W. (2007). The relation between post concussion symptoms and neurocognitive performance in concussed athletes. *NeuroRehabilitation. Special Issue: Sports and Concussion*, 22(3), 207–216.
- *Ferguson, R.J., Mittenberg, W., Barone, D.F., & Schneider, B. (1999). Postconcussion syndrome following sports-related head injury: Expectation as etiology. *Neuropsychology*, 13(4), 582–589.
- *Field, M., Collins, M.W., Lovell, M.R., & Maroon, J. (2003). Does age play a role in recovery from sports-related concussion? A comparison of high school and collegiate athletes. *The Journal of Pediatrics*, 142(5), 546–553.
- Frencham, K.A.R., Fox, A.M., & Maybery, M.T. (2005). Neuropsychological studies of mild traumatic brain injury: A meta-analytic review of research since 1995. *Journal of Clinical and Experimental Neuropsychology*, 27(3), 334–351.
- Giza, C.C., & Hovda, D.A. (2001). The neurometabolic cascade of concussion. *Journal of Athletic Training*, 36(3), 228–235.
- Giza, C.C., & Hovda, D.A. (2004). Chapter 4: The pathophysiology of traumatic brain injury. In M. R. Lovell, R. J. Echemendia, J. T. Barth & M. W. Collins (Eds.), *Traumatic brain injury in sports: An international neuropsychological perspective* (pp. 45–70). Lisse: Swets & Zeitlinger.
- *Gosselin, N., Lassonde, M., Petit, D., Leclerc, S., Mongrain, V., Collie, A., & Montplaisir, J. (2009). Sleep following sport-related concussions. *Sleep Medicine*, 10(1), 35–46.
- *Gosselin, N., Theriault, M., Leclerc, S., Montplaisir, J., & Lassonde, M. (2006). Neurophysiological anomalies in symptomatic and asymptomatic concussed athletes. *Neurosurgery*, 58(6), 1151–1161.
- *Guskiewicz, K.M., Perrin, D.H., & Gansneder, B.M. (1996). Effect of mild head injury on postural stability in athletes. *Journal of Athletic Training*, 31(4), 300–306.
- *Guskiewicz, K.M., Riemann, B.L., Perrin, D.H., & Nashner, L.M. (1997). Alternative approaches to the assessment of mild head injury in athletes. *Medicine & Science in Sports & Exercise*, 29(Suppl. 7), S213–S221.
- *Guskiewicz, K.M., Ross, S.E., & Marshall, S.W. (2001). Postural stability and neuropsychological deficits following concussion in collegiate athletes. *Journal of Athletic Training*, 36(3), 263–273.
- Guskiewicz, K.M., Weaver, N., Padua, D.A., & Garrett, W.E., Jr. (2000). Epidemiology of concussion in collegiate and high school football players. *American Journal of Sports Medicine*, 28(5), 643–650.
- Hedges, L.V., & Olkin, I. (1985). *Statistical methods for meta-analysis*. Sydney: Academic Press.
- *Hinton-Bayre, A.D., Geffen, G.M., Geffen, L.B., McFarland, K.A., & Frijs, P. (1999). Concussion in contact sports: Reliable change indices of impairment and recovery. *Journal of Clinical and Experimental Neuropsychology*, 21(1), 70–86.
- *Hinton-Bayre, A.D., Geffen, G.M., & McFarland, K.A. (1997). Mild head injury and speed of information processing: A prospective study of professional rugby league players. *Journal of Clinical and Experimental Neuropsychology*, 19(2), 275–289.
- Hoofien, D., Vakil, E., Gilboa, A., Donovan, P.J., & Barak, O. (2002). Comparison of the predictive power of socio-economic variables, severity of injury and age on long-term outcome of traumatic brain injury: Sample-specific variables versus factors as predictors. *Brain Injury*, 16(1), 9–27.
- Hovda, D.A., Prins, M., Becker, D.P., Lee, S., Bergsneider, M., & Martin, N. (1999). Neurobiology of concussion. In J. E. Bailes, M. Lovell, J. C. Maroon (Eds.), *Sports-related concussion* (pp. 12–51). St Louis, MO: Quality Medical Publishing.
- Iverson, G.L. (2010). Mild traumatic brain injury meta-analyses can obscure individual differences. *Brain Injury*, 24(10), 1246–1255.
- *Iverson, G.L., Brooks, B.L., Collins, M.W., & Lovell, M.R. (2006). Tracking neuropsychological recovery following concussion in sport. *Brain Injury*, 20(3), 245–252.
- *Iverson, G.L., Gaetz, M., Lovell, M.R., & Collins, M.W. (2004). Cumulative effects of concussion in amateur athletics. *Brain Injury*, 18(5), 433–443.
- *Iverson, G.L., Lovell, M.R., & Collins, M.W. (2003). Interpreting change on ImPACT following sport concussion. *The Clinical Neuropsychologist*, 17(4), 460–467.
- *Jantzen, K.J., Anderson, B., Steinberg, F.L., & Kelso, J.A.S. (2004). A prospective functional MR imaging study of mild traumatic brain injury in college football players. *AJNR American Journal of Neuroradiology*, 25(5), 738–745.
- *Johnson, P.D., Hertel, J., Olmsted, L.C., Denegar, C.R., & Putukian, M. (2002). Effect of mild brain injury on an instrumented agility task. *Clinical Journal of Sport Medicine*, 12(1), 12–17.
- Kaplan, R.F., Cohen, R.A., Moscufo, N., Guttman, C., Chasman, J., Buttaro, M., ... Wolfson, L. (2009). Demographic and biological influences on cognitive reserve. *Journal of Clinical and Experimental Neuropsychology*, 31(7), 868–876.
- *Killam, C., Cautin, R.L., & Santucci, A.C. (2005). Assessing the enduring residual neuropsychological effects of head trauma in college athletes who participate in contact sports. *Archives of Clinical Neuropsychology*, 20(5), 599–611.
- Kontos, A.P., Dolese, A., Elbin, R.J., Covassin, T., & Warren, B.L. (2011). Relationship of soccer heading to computerized neurocognitive performance and symptoms among female and male youth soccer players. *Brain Injury*, 25(12), 1234–1241.
- *Lavoie, M.E., Dupuis, F., Johnston, K.M., Leclerc, S., & Lassonde, M. (2004). Visual P300 effects beyond symptoms in concussed college athletes. *Journal of Clinical and Experimental Neuropsychology*, 26(1), 55–73.

- Lipsey, M.W., & Wilson, D.B. (2001). *Practical meta-analysis*. London: Sage.
- *Lovell, M.R., & Collins, M.W. (1998). Neuropsychological assessment of the college football player. *Journal of Head Trauma Rehabilitation, 13*(2), 9–26.
- *Lovell, M.R., Collins, M.W., Iverson, G.L., Field, M., Maroon, J.C., Cantu, R.C., ... Fu, F.H. (2003). Recovery from mild concussion in high school athletes. *Journal of Neurosurgery, 98*(2), 296–301.
- *Lovell, M.R., Collins, M.W., Iverson, G.L., Johnston, K.M., & Bradley, J.P. (2004). Grade 1 or 'ding' concussions in high school athletes. *The American Journal of Sports Medicine, 32*(1), 47–54.
- Lovell, M.R., & Fazio, V. (2008). Concussion management in the child and adolescent athlete. *Current Sports Medicine Reports, 7*(1), 12–15.
- *Lovell, M.R., Iverson, G.L., Collins, M.W., Podell, K., Johnston, K.M., Pardini, D., ... Maroon, J.C. (2006). Measurement of symptoms following sports-related concussion: Reliability and normative data for the post-concussion scale. *Applied Neuropsychology, 13*(3), 166–174.
- *Macciocchi, S.N., Barth, J.T., Alves, W., Rimel, R.W., & Jane, J.A. (1996). Neuropsychological functioning and recovery after mild head injury in collegiate athletes. *Neurosurgery, 39*(3), 510–514.
- *Macciocchi, S.N., Barth, J., Littlefield, L., & Cantu, R.C. (2001). Multiple concussions and neuropsychological functioning in collegiate football players. *Journal of Athletic Training, 36*(3), 303–306.
- *Maddocks, D.L., Dicker, G.D., & Saling, M.M. (1995). The assessment of orientation following concussion in athletes. *Clinical Journal of Sport Medicine, 5*(1), 32–35.
- *Maddocks, D.L., & Saling, M.M. (1996). Neuropsychological deficits following concussion. *Brain Injury, 10*(2), 99–103.
- Makdissi, M. (2009). Is the simple versus complex classification of concussion a valid and useful differentiation? *British Journal of Sports Medicine, 43*(Suppl. 1), i23–i27.
- *Makdissi, M., Collie, A., Maruff, P., Darby, D.G., Bush, A., McCrory, P.R., & Bennell, K. (2001). Computerised cognitive assessment of concussed Australian Rules footballers. *British Journal of Sports Medicine, 35*(5), 354–360.
- *McClincy, M.P., Lovell, M.R., Pardini, J.E., Collins, M.W., & Spore, M.K. (2006). Recovery from sports concussion in high school and collegiate athletes. *Brain Injury, 20*(1), 33–39.
- *McCrea, M. (2001). Standardized mental status testing on the sideline after sport-related concussion. *Journal of Athletic Training, 36*(3), 274–279.
- *McCrea, M., Guskiewicz, K.M., Marshall, S.W., Barr, W., Randolph, C., Cantu, R.C., ... Kelly, J.P. (2003). Acute effects and recovery time following concussion in collegiate football players: The NCAA Concussion Study. *Journal of the American Medical Association, 290*(19), 2556–2563.
- *McCrea, M., Kelly, J.P., Randolph, C., Cisler, R., & Berger, L. (2002). Immediate neurocognitive effects of concussion. *Neurosurgery, 50*(5), 1032–1040.
- *McCrea, M., Kelly, J.P., Randolph, C., Kluge, J., Bartolic, E., Finn, G., & Baxter, B. (1998). Standardized Assessment of Concussion (SAC): On-site mental status evaluation of the athlete. *Journal of Head Trauma Rehabilitation, 13*(2), 27–35.
- *McCrory, P.R., Ariens, M., & Berkovic, S.F. (2000). The nature and duration of acute concussive symptoms in Australian football. *Clinical Journal of Sport Medicine, 10*(4), 235–238.
- McCrory, P.R., Meeuwisse, W., Johnston, K., Dvorak, J., Aubry, M., Molloy, M., & Cantu, R. (2009). Consensus statement on concussion in sport: The 3rd International Conference on Concussion in Sport held in Zurich, November 2008. *British Journal of Sports Medicine, 43*(Suppl. 1), i76–i84.
- McDonald, J.W., & Johnston, M.V. (1990). Physiological and pathophysiological roles of excitatory amino acids during central nervous system development. *Brain Research Reviews, 15*, 41–70.
- *Mihalik, J.P., McCaffrey, M.A., Rivera, E.M., Pardini, J.E., Guskiewicz, K.M., Collins, M.W., & Lovell, M.R. (2007). Effectiveness of mouthguards in reducing neurocognitive deficits following sports-related cerebral concussion. *Dental Traumatology, 23*(1), 14–20.
- Moser, R.S., Iverson, G.L., Echemendia, R.J., Lovell, M.R., Schatz, P., Webbe, F.M., ... NAN Policy Planning Committee. (2007). NAN Position Paper: Neuropsychological evaluation in the diagnosis and management of sports-related concussion. *Archives of Clinical Neuropsychology, 22*(8), 909–916.
- *Moser, R.S., & Schatz, P. (2002). Enduring effects of concussion in youth athletes. *Archives of Clinical Neuropsychology, 17*(1), 91–100.
- *Moser, R.S., Schatz, P., & Jordan, B.D. (2005). Prolonged effects of concussion in high school athletes. *Neurosurgery, 57*(2), 300–306.
- Panayiotou, A., Jackson, M., & Crowe, S.F. (2010). A meta-analytic review of the emotional symptoms associated with mild traumatic brain injury. *Journal of Clinical and Experimental Neuropsychology, 32*(5), 463–473.
- *Parker, T.M., Osternig, L.R., Lee, H.J., Donkelaar, P., & Chou, L.S. (2005). The effect of divided attention on gait stability following concussion. *Clinical Biomechanics, 20*(4), 389–395.
- *Parker, T.M., Osternig, L.R., Van Donkelaar, P., & Chou, L.-S. (2006). Gait stability following concussion. *Medicine and Science in Sport and Exercise, 38*(6), 1032–1040.
- *Parker, T.M., Osternig, L.R., van Donkelaar, P., & Chou, L.S. (2008). Balance control during gait in athletes and non-athletes following concussion. *Medical Engineering & Physics, 30*(8), 959–967.
- *Pellman, E.J., Lovell, M.R., Viano, D.C., & Casson, I.R. (2006). Concussion in professional football: Recovery of NFL and high school athletes assessed by computerized neuropsychological testing – Part 12. *Neurosurgery, 58*(2), 263–274.
- *Pellman, E.J., Lovell, M.R., Viano, D.C., Casson, I.R., & Tucker, A.M. (2004). Concussion in professional football: Neuropsychological testing – Part 6. *Neurosurgery, 55*(6), 1290–1303.
- *Peterson, C.L., Ferrara, M.S., Mrazik, M., Piland, S., & Elliott, R. (2003). Evaluation of neuropsychological domain scores and postural stability following cerebral concussion in sports. *Clinical Journal of Sport Medicine, 13*(4), 230–237.
- *Piland, S.G., Motl, R.W., Ferrara, M.S., & Peterson, C.L. (2003). Evidence for the factorial and construct validity of a self-report concussion symptoms scale. *Journal of Athletic Training, 38*(2), 104–112.
- Prins, M.L., Lee, S.M., Cheng, C.L.Y., Becker, D.P., & Hovda, D.A. (1996). Fluid percussion brain injury in the developing and adult rat: A comparative study of mortality, morphology, intracranial pressure and mean arterial blood pressure. *Developmental Brain Research, 95*(2), 272–282.
- Putukian, M., Aubry, M., & McCrory, P.R. (2009). Return to play after sports concussion in elite and non-elite athletes? *British Journal of Sports Medicine, 43*(Suppl. 1), i28–i31.
- Quality Standards Subcommittee of the American Academy of Neurology. (1997). Practice Parameter. The management of concussion in sports (summary statement). Report of the Quality Standards Subcommittee of the American Academy of Neurology. *Neurology, 48*(3), 581–585.

- Ratcliff, J.J., Greenspan, A.I., Goldstein, F.C., Stringer, A.Y., Bushnik, T., Hammond, F.M., ... Wright, D.W. (2007). Gender and traumatic brain injury: Do the sexes fare differently? *Brain Injury*, 21(10), 1023–1030.
- Reddy, C.C., Collins, M.W., & Gioia, G.A. (2008). Adolescent sports concussion. *Physical Medicine and Rehabilitation Clinics of North America*, 19(2), 247–269.
- *Register-Mihalik, J., Guskiewicz, K.M., Mann, J.D., & Shields, E.W. (2007). The effects of headache on clinical measures of neurocognitive function. *Clinical Journal of Sport Medicine*, 17(4), 282–288.
- *Riemann, B.L., & Guskiewicz, K.M. (2000). Effects of mild head injury on postural stability as measured through clinical balance testing. *Journal of Athletic Training*, 35(1), 19–25.
- Rohling, M.L., Beverly, B., Faust, M.E., & Demakis, G.J. (2009). Effectiveness of cognitive rehabilitation following acquired brain injury: A meta-analytic re-examination of Cicerone et al.'s (2000, 2005) systematic reviews. *Neuropsychology*, 23(1), 20–39.
- Rohling, M.L., Binder, L.M., Demakis, G.J., Larrabee, G.J., Ploetz, D.M., & Langhinrichsen-Rohling, J. (2011). A meta-analysis of neuropsychological outcome after mild traumatic brain injury: Re-analysis and reconsiderations of Binder et al. (1997), Frencham et al. (2005), and Pertab et al. (2009). *The Clinical Neuropsychologist*, 25(4), 608–623.
- Satz, P. (1993). Brain reserve capacity on symptom onset after brain injury: A formulation and review of evidence for threshold theory. *Neuropsychology*, 7(3), 273–295.
- *Schatz, P., Pardini, J.E., Lovell, M.R., Collins, M.W., & Podell, K. (2006). Sensitivity and specificity of the ImPACT Test Battery for concussion in athletes. *Archives of Clinical Neuropsychology*, 21(1), 91–99.
- Schretlen, D.J., & Shapiro, A.M. (2003). A quantitative review of the effects of traumatic brain injury on cognitive functioning. *International Review of Psychiatry*, 15(4), 341–349.
- *Sim, A., Terryberry-Spohr, L., & Wilson, K.R. (2008). Prolonged recovery of memory functioning after mild traumatic brain injury in adolescent athletes. *Journal of Neurosurgery*, 108(3), 511–516.
- *Slobounov, S., Cao, C., Sebastianelli, W., Slobounov, E., & Newell, K. (2008). Residual deficits from concussion as revealed by virtual time-to-contact measures of postural stability. *Clinical Neurophysiology*, 119(2), 281–289.
- *Slobounov, S., Slobounov, E., & Newell, K. (2006). Application of virtual reality graphics in assessment of concussion. *CyberPsychology & Behavior*, 9(2), 188–191.
- *Slobounov, S., Slobounov, E., Sebastianelli, W., Cao, C., & Newell, K. (2007). Differential rate of recovery in athletes after first and second concussion episodes. *Neurosurgery*, 61(2), 338–344.
- *Slobounov, S., Tutwiler, R., Sebastianelli, W., & Slobounov, E. (2006). Alteration of postural responses to visual field motion in mild traumatic brain injury. *Neurosurgery*, 59(1), 134–139.
- Solomon, G.S., & Haase, R.F. (2008). Biopsychosocial characteristics and neurocognitive test performance in National Football League players: An initial assessment. *Archives of Clinical Neuropsychology*, 23(5), 563–577.
- *Sosnoff, J.J., Broglio, S.P., & Ferrara, M.S. (2008). Cognitive and motor function are associated following mild traumatic brain injury. *Experimental Brain Research*, 187(4), 563–571.
- *Sosnoff, J.J., Broglio, S.P., Hillman, C.H., & Ferrara, M.S. (2007). Concussion does not impact intraindividual response time variability. *Neuropsychology*, 21(6), 796–802.
- Stern, Y. (2009). Cognitive reserve. *Neuropsychologia*, 47, 2015–2028.
- Tate, R., McDonald, S., & Lulham, J.M. (1998). Incidence of hospital-treated traumatic brain injury in an Australian community. *Australian and New Zealand Journal of Public Health*, 22(4), 419–425.
- *Thompson, J., Sebastianelli, W., & Slobounov, S. (2005). EEG and postural correlates of mild traumatic brain injury in athletes. *Neuroscience Letters*, 377(3), 158–163.
- Tsushima, W.T., Lum, M., & Geling, O. (2009). Sex differences in the long-term neuropsychological outcome of mild traumatic brain injury. *Brain Injury*, 23(10), 809–814.
- *Van Kampen, D.A., Lovell, M.R., Pardini, J.E., Collins, M.W., & Fu, F.H. (2006). The 'value added' of neurocognitive testing after sports-related concussion. *The American Journal of Sports Medicine*, 34(10), 1630–1635.
- Viano, D.C., Casson, I.R., & Pellman, E.J. (2007). Concussion in professional football: Biomechanics of the struck player – Part 14. *Neurosurgery*, 61, 313–327.
- *Warden, D.L., Bleiberg, J., Cameron, K.L., Ecklund, J., Walter, J., Sparling, M.B., ... Arciero, R. (2001). Persistent prolongation of simple reaction time in sports concussion. *Neurology*, 57(3), 524–526.