

# Cardiovascular disease and meaning in life: A systematic literature review and conceptual model

## Review Article

**Cite this article:** Vos J (2021). Cardiovascular disease and meaning in life: A systematic literature review and conceptual model. *Palliative and Supportive Care* **19**, 367–376. <https://doi.org/10.1017/S1478951520001261>

Received: 14 September 2020

Revised: 19 October 2020


Accepted: 8 November 2020

### Key words:

Counseling; Existential; Health psychology; Heart attack; Meaning in life; Psychotherapy

### Author for correspondence:

Joel Vos, Metanoia Institute, Ealing, London, UK. E-mail: [Joel.Vos@metanoia.ac.uk](mailto:Joel.Vos@metanoia.ac.uk)

Joel Vos, PH.D., M.SC., M.A., CPSYCHOL., F.H.E.A. 

Metanoia Institute, Ealing, London, UK

### Abstract

**Objectives.** Previous studies have shown that psychological stress and mental health problems increase the risk for cardiovascular disease (CVD) events, such as heart attack or stroke. Furthermore, after CVD events, the majority of patients report large stress. However, psychological treatments have only modest effects in CVD patients. Therefore, it has been argued that new conceptual models are needed to understand the aetiology of stress and mental health problems in CVD patients. Therefore, this study included a systematic literature review and a conceptual model on the role of meaning in life for psychological stress, mental health, and CVD risks.

**Methods.** A systematic literature review was conducted on relationships between CVD and meaning in life. PRISMA/MOOSE review guidelines were followed. These findings were used to build a conceptual model.

**Results.** The literature review included 113 studies on meaning and CVD. The included studies described meaning as a predictor of cardiovascular risks and health, meaning-centered needs of patients in conversations with medical staff, meaning-centered changes after CVD events, meaning-centered coping with CVD, meaning as motivator of CVD-related lifestyle changes, and meaning as an element in psychological treatments of CVD patients. In sum, the literature showed that a central clinical concern for patients is their question how to live a meaningful life despite CVD. Meaning-centered concerns seem to lead to lower motivation to make lifestyle changes, more psychological stress, lower quality-of-life, worse physical well-being, and increased CVD risk. The ability to live a meaningful life after CVD events is related with lower stress, better mental health, and several biomarkers.

**Significance of results.** An evidence-based conceptual framework was developed for the relationship between meaning and CVD. It may be hypothesized CVD patients may benefit from psychological therapies focused on meaning.

## Introduction

Cardiovascular disease (CVD) is one of the most prevalent life-threatening health conditions in western countries, associated with relatively high mortality rates. Cardiovascular events such as heart attack or stroke often lead to heightened levels of psychological stress, depression, and anxiety (Konstam et al., 2005). It is also widely recognized that reducing stress and mental health problems is crucial to the prevention and recovery of cardiovascular events (Whalley et al., 2011). Therefore, national health guidelines, for instance by the British National Health Service, recommend offering psychological counseling to CVD patients (Gillham and Clark, 2011; Smith and Kneebone, 2016).

Despite the importance of mental health care for CVD, only 24 psychological clinical trials have been published (Whalley et al., 2011; Tan and Morgan, 2015). These studies show that these treatments have only small effects on anxiety and depression and no consistent positive effects on outcomes such as quality-of-life and physical well-being. The statistical heterogeneity between studies is large, which makes it difficult to generalize findings. Interventions mainly include behavioral and cognitive exercises, such as relaxation, self-awareness, self-monitoring, risk education, emotional support, homework, behavior change guidance, and cognitive techniques (Whalley et al., 2011). In most studies, the therapeutic mechanisms leading to therapeutic change are not directly measured, thus it remains unknown how these interventions precisely lead to improvement. Thus, most psychological interventions in CVD have insufficient validity and efficacy. Therefore, it has been argued that new conceptual models are needed to understand the aetiology of stress and mental health problems in CVD patients. Such a new model may help to develop new — potentially more effective — psychological therapies (Carroll and Nuro, 2002; Rounsaville et al., 2006; Moore et al., 2015).

The aim of this study is to conduct a systematic literature review on the relationship between meaning and CVD, which will be used to develop a conceptual model, — which may be used as foundations for the development of new psychological treatments focusing

on meaning. This focus on meaning follows from the growing literature on the relationship between CVD and positive psychological attributes, and particularly meaning-making (Park, 2010; Boehm and Kubzansky, 2012; Dubois et al., 2012). This meaning-centered paradigm could explain the relative lack of effects of the usual care for CVD. For example, cognitive-behavioral therapy does not systematically address the central question that most patients with a chronic or life-threatening physical disease such as CVD ask: “How can I live a meaningful and satisfying life despite my disease?” (Vos, 2016a). Their primary concerns seem more related to their quest for meaning than to cognitive or behavioral issues. Research indicates that physically ill patients need to make sense of the disease and fit this into their overall perspective on life; they could subsequently experience psychological stress due to the tensions between these situational and global meanings (Park and Folkman, 1997). Table 1 summarizes the empirical evidence from previous reviews for the relevance of meaning in coping with a chronic or life-threatening physical

disease other than CVD (Vos, 2016a, 2016b). For example, this shows how meaning is an evidence-based psychological construct, which can be differentiated from other phenomena (Park and Folkman, 1997; Park, 2010). A large body of the literature shows that in the general population, a lack of meaning leads to lower quality-of-life, higher levels of stress, and worse physical health such as higher blood pressure and suppressed immune system (Ryff et al., 2004; Vos, 2016b).

## Method

Several authors have published systematic literature reviews on general psychological causes and consequences of CVD (Rozanski et al., 1999; Dimsdale, 2008). A systematic meta-analysis on psychological treatments was published by Whalley et al. (2014). Other reviews also describe the literature on general positive psychology in CVD (Boehm and Kubzansky, 2012). Some reviews describe meaning amongst other factors (Rozanski et al.,

**Table 1.** Overview of evidence-based characteristics of meaning in life (based on the review in Vos, 2016a, 2017)

Characteristic	Description
Definition	The meaning construct consists of seven highly correlated experiences: <ol style="list-style-type: none"> <li>1. Motivation (purpose, goals, direction),</li> <li>2. Values,</li> <li>3. Understanding of life events and surrounding world (sense of coherence),</li> <li>4. Self-worth,</li> <li>5. Commitment to action,</li> <li>6. Self-regulation (control, setting and adjusting goals), and</li> <li>7. Accepting existential limitations.</li> </ol>
Types of meaning	<i>A world-wide review showed individuals experience 5 types and 27 sub-types of meaning:</i> <ul style="list-style-type: none"> <li>• Material-hedonic type (material conditions, professional-educational success, hedonic-experiential activities, and health)</li> <li>• Self-oriented type (resilience, self-efficacy, self-acceptance, autonomy, creative self-expression, self-care, and authenticity)</li> <li>• Social type (connections, belonging, altruism, and children)</li> <li>• Larger type (higher purpose, personal growth, temporal coherence, justice and ethics, and spirituality and religion)</li> <li>• Existential-philosophical type (being-alive, uniqueness, embeddedness, freedom, gratitude, and responsibility)</li> </ul>
Large correlations in general population	<i>A strong sense of meaning is highly correlated with:</i> <ul style="list-style-type: none"> <li>• Low level of psychological stress (e.g., anxiety and depression),</li> <li>• High level of acceptance of life's limitations and low level of existential stress (e.g., fear of death),</li> <li>• High level of quality-of-life (e.g., life satisfaction),</li> <li>• Better self-reported physical well-being (e.g., pain perception) and biomarkers (e.g., immune system, blood pressure, and survival rate).</li> </ul>
Meaning-centered changes after diagnosis of a chronic or life-threatening disease	<i>Changes in specific meanings</i> <ul style="list-style-type: none"> <li>• All specific pre-disease meanings stay the same</li> <li>• A specific pre-disease meaning becomes more valuable</li> <li>• A specific pre-disease meaning feels less valuable</li> <li>• A specific pre-disease goal is adjusted</li> <li>• A specific pre-disease meaning becomes unattainable</li> <li>• A specific new meaning develops</li> </ul> <i>Changes in general perspectives</i> <ul style="list-style-type: none"> <li>• General priorities in life change</li> <li>• Experiencing life more intensively</li> <li>• Focusing more on authentic meanings</li> <li>• Such a narrow focus on life's limitations, that other meanings are forgotten or denied</li> <li>• Hyper-reflection or hyper-intention (i.e., wanting a specific meaning very strong and/or strongly reflecting on it)</li> <li>• Experiencing meaninglessness</li> <li>• More realistic about life's limitations</li> <li>• Existential stress, e.g., fear of death</li> </ul>
Effects of meaning-oriented therapy	<i>Meaning-oriented therapy has large to very large significant effects in 60 trials in physically ill and/or mentally ill populations, compared with active interventions:</i> <ul style="list-style-type: none"> <li>• Larger quality-of-life;</li> <li>• Larger acceptance and lower existential stress;</li> <li>• Lower psychological stress;</li> <li>• Better self-reported physical well-being and biomarkers (24 trials).</li> </ul>

2005; Leegaard and Fagermoen, 2008; Cohen et al., 2016; Sin, 2016). However, these studies do not review all meaning-centered aspects of meaning in CVD. Therefore, a systematic literature review was conducted on meaning and CVD. Meaning was defined as described in Table 1.

A systematic scoping literature review was conducted in consecutive rounds. A review protocol was developed in line with PRISMA and MOOSE-guidelines (Stroup et al., 2000; Liberati et al., 2009), which can be requested from the authors. The aim of this review was to search for any studies with or without interventions, with or without comparisons on the relationship between meaning in life and CVD. Meaning in life was defined as in Table 1. CVD was defined as any type of cardiovascular event or cardiovascular surgery (e.g., acute coronary syndrome, acute myocardial infarction, coronary artery bypass grafting, recovery of coronary artery or heart disease, and cerebrovascular accident).

First, databases were searched and included Pubmed, Web-of-Knowledge, PsycInfo, PsycTest, Medline, Embase, scholar.google.com, and Scopus. Two sets of search terms were combined. The set of terms regarding meaning was derived from previous reviews (Vos and Vitali, 2018) and included: “meaning in life” or “meaning of life” or “life meaning” or “meaningful life” or “living meaningful\*” or “noetic” or “purpose in life” or “purpose of life” or “life purpose” or “purposes in life” or “purposes of life” or “life’s purpose” or “goal in life” or “goal of life”

or “goals of life” or “goals in life” or “life’s goal\*” or “value\* in life” or “life’s value\*” or “significance of life” or “life destiny” or “destiny in life” or “destiny of life” or “life\* essence” or “essence of life” or “sense of life” or “aims in life” or “aims of life” or “life\* aims” or “meaning-making” or “existential” or “positive psychol\*.” The set of terms regarding CVD included: “cardiovascular” or “heart” or “stroke” or “cardiac” or “cardiolog\*” or “coronary” or myocardial or cerebrovascular\*. Given the large number of findings, we added PubMed Mesh terms ([counseling] OR [psychotherapy] OR [psychology]) and capped scholar.google.com results at 10,000 hits. Additional studies were identified via the reference lists. Second, meaning-centered experts excluded articles through thorough reading of the titles and abstracts. Third, studies were excluded on basis of full-text manuscripts. Fourth, all studies are summarized and presented in the text. Fifth, risk of bias was assessed via the Cochrane risk of bias tool. Studies with a large risk of bias are described separately in the Findings section (Table 2).

Sixth, articles were synthesized via thematic analyses (Clarke et al., 2015). We did not limit the research findings to any specific variables, as we wanted to get a full overview of the relationships between CVD and meaning in life. We did not conduct meta-analyses due to the heterogeneity of instruments and study designs; therefore, we did also not present the detailed characteristics of each study. However, in the reporting of the studies, we summarize the average effects reported in the studies based on

**Table 2.** Overview of studies on meaning in CVD

Study topic	Number of studies	Reference
<i>Meaning as predictor of cardiovascular health and disease risk in general population</i>	11	Edmondson et al., 2005; Skrabski et al., 2005; Ishida and Okada, 2006; Sone et al., 2008; Shirai et al., 2009; Tanno et al., 2009; Mezick et al., 2010; Kim et al., 2013; Rohleder, 2014; Yu et al., 2015; Cohen et al., 2016
<i>Positive well-being (associated with meaning) as predictor of cardiovascular health and disease risk in general population</i>	19	Rozanski and Kubzansky, 2005; Chida and Steptoe, 2008; Chida and Hamer, 2008; Kožul et al., 2009; Davidson et al., 2010; Low and Lam., 2013; Boehm et al., 2011, 2015; Hamer and Chida, 2011; Boehm and Kubzansky, 2012; DuBois et al., 2012; Steptoe et al., 2012; Falk et al., 2013; Haukkala et al., 2013; Hoen et al., 2013; Schwerdtfeger and Gerteis, 2014; Lambiase et al., 2015; Marteinsdottir et al., 2016; Sin, 2016.
<i>Meaning-centered needs in conversation about CVD with doctors and nurses</i>	4	Schaufel et al., 2009; Kohlmann et al., 2013; McClung, 2013; Strang et al., 2013; Barello et al., 2015
<i>Meaning-centered changes and needs after CVD event</i>	30	Mårtensson et al., 1998; Popogrebsky, 1998; Clarke et al., 1999; Nilsson et al., 1999; Ecohard et al., 2001; Davidson et al., 2003; Secrest and Zeller, 2003; Faircloth et al., 2004; Lobeck et al., 2005; Secrest and Zeller, 2006; Cortis and Williams, 2007; Leegaard and Fagermoen, 2008; Schou and Egerod, 2008; Wann-Hansson et al., 2008; Bremer et al., 2009; Hefferon et al., 2009; Rassin et al., 2009; Eilertsen et al., 2010; Kutner, 2010; Silva et al., 2011; Lawrence and Kinn, 2012; Andersson et al., 2013; Ross and Austin, 2013; Junehag et al., 2014; Leeming et al., 2014; Buck et al., 2015; DuBois et al., 2015; Hansen et al., 2016; Satink et al., 2016; Littooj et al., 2016.
<i>Meaning-centered coping with congenital heart disease</i>	4	Mårtensson et al., 1997; Amianto et al., 2013; Berghammer et al., 2015; Apers et al., 2016.
<i>Meaning and meaning-centered coping as predictor of long-term physical and psychological well-being in CVD patients</i>	18	Muirhead et al., 1992; Evangelista et al., 2003; Rozanski et al., 2005; Park et al., 2007; Balon et al., 2008; Ai et al., 2009, 2010, 2012; Owolabi, 2009; Vollman et al., 2009; Sirri et al., 2010; Bekke-Hansen et al., 2013; Kim et al., 2013; Shao et al., 2013; Grohn et al., 2014; Sacco et al., 2014; Simonj et al., 2015; Huffman et al., 2016b; Sarfo et al., 2017.
<i>Meaning and meaning-associated positive psychological factors as predictor of CVD-related lifestyle changes</i>	15	Laerum et al., 1991; Tinker et al., 2007; Holahan et al., 2008; Cuffee, 2010; Ogedegbe et al., 2012; Peterson et al., 2012; Proyer et al., 2013; Versteeg et al., 2013; Charlson et al., 2014; Ronaldson et al., 2015; Røysland and Friberg, 2015; Simonj et al., 2015; Sin et al., 2015; Huffman et al., 2016a, 2016b; Van Montfort et al., 2016.
<i>Meaning-centered and meaning-associated positive psychological treatments of CVD patients</i>	12	Krucoff et al., 2001, 2005; Kennedy et al., 2007; Lantz and Gregoire, 2003; Seskevich et al., 2004; Burton et al., 2010; Roncella et al., 2013; Huffman et al., 2016a; Lee et al., 2016; Nikrahan et al., 2016a, 2016b; Sanjuán et al., 2016.

their unweighted average. For example, we describe  $R^2$  of 0.75 as large, 0.50 moderately large, and 0.25 small. However, this presentation of effect size should be read as indicative only.

On the basis of this literature review, a conceptual model was developed, by combining this with a pre-existing review of meaning in other chronic and life-threatening diseases (Vos, 2016a) and a review of the mechanisms of meaning-centered treatments (Vos, 2016b, 2017). Several authors have argued that a conceptual model needs to integrate all relevant literature while being concise, and this describes the central clinical phenomenon, aetiology, mechanisms of change, and impact on patients (Wampold, et al., 1997; Kazdin, 2009; Moore et al., 2015; Vos, 2017). The logical model is visualized.

## Findings

### Findings

17,084 publications were found, of which 113 were selected, reflecting 21,509 patients (see Figure 1).

Eleven studies described meaning as a moderately strong predictor of cardiovascular risks. Nineteen studies described meaning as part of positive well-being and showed how positive well-being is a moderately strong predictor of cardiovascular health and lower CVD risk in the general population. Four studies described that CVD patients spontaneously ask questions about meaning with doctors and nurses. Thirty studies described changes in meaning in life and questions about meaning in life after a CVD event. Four studies described the positive effects of a meaning-oriented coping-style on adjusting to living with a congenital heart disease and reducing psychological stress, such as focusing on what is meaningful in life instead of on the

limitations in life. Eighteen studies described meaning-centered coping as a predictor of long-term physical and psychological well-being in CVD patients. Fifteen studies described that experiencing life as meaningful and/or using meaning-oriented coping-styles predict CVD-related lifestyle changes in CVD patients, such as dieting and exercising. Twelve studies described psychological intervention studies in CVD patients which had one or more meaning-oriented aspects (most of which were positive psychology interventions), and which indicated moderate to large improvements in psychological well-being.

Analyses of risk of bias for the correlational studies indicated good or acceptable risk for all studies. Analyses of risk of bias revealed a significant risk for 10 out of 12 clinical trials, due to lack of control groups, unclear blinding and in 3 studies incomplete data reporting.

### Meaning-centered conceptual model

The literature from the previous step was combined with other studies on the psychological aspects of coping with CVD, and on meaning in other chronic and life-threatening physical diseases. Figure 2 summarizes the meaning-centered model of CVD. This is followed by a description of the treatment manual that is based on this meaning-centered model.

### Predictors of CVD event

Empirical studies have suggested many factors which seem to increase the risk of CVD events. This includes, amongst others, biomedical risks such as genetics and body weight; lifestyle factors such as diet and exercise; level of psychological stress; stress reactivity and anxiety (Roest et al., 2010); personality such as type A




Fig. 1. Flowchart of included and excluded studies.






Fig. 2. Logical meaning-centered model of CVD.

and hostility (Chida and Steptoe, 2008); and quality-of-life such as optimism and meaning (Dubois et al., 2012). The effect of socio-demographic and life history risk factors on stress and CVD risks are mediated by meaning-centered coping (Chen et al., 2015; Su et al., 2015). For example, meaning-centered coping-styles reduce the level of psychological stress (Kopp et al., 2007; Bennett et al., 2014b). These factors are likely intertwined, such as high stress levels leading to chronic inflammation which could contribute to CVD risks (Rohleder, 2014; type A personality is associated with high levels of stress and low quality-of-life; physical exercise may also reduce stress; Scully et al., 1998).

**Post-CVD event**

After patients have experienced a CVD event, they may be required to make lifestyle changes, possibly because their disease makes it impossible to do the activities they did in the past, or they need to lower their CVD risk, for instance by exercising or stopping smoking. However, research shows that many patients find it difficult to make these changes, particularly due to a lack of motivation; therefore, intervention studies suggest that their motivation could be improved by focusing on the positive meaning of these activities for their life in general (Tinker et al., 2007). Many patients experience CVD events also as psychologically traumatic, particularly because it confronts them with the limitations and finitude of life, which may subsequently increase their level of psychological stress and decrease their quality-of-life and increase their risk of another CVD event (Nilsson et al., 1999; Stoll et al., 2000; Coughlin, 2011). Like other potentially life-threatening diseases, the CVD event has suddenly confronted them with their limitations and finitude, they need to change parts of their life, and they may develop new general perspectives

on life (LeMay and Wilson, 2008). Qualitative studies show that many patients feel stuck in a vicious cycle of limitation and resignation, and they need to “use self-care” and “see the possibilities that exist in everyday life” (Mårtensson et al., 1997). People start to search for meaning in response to the existential limitations that CVD has confronted them with (Koizumi et al., 2008). Thus, the core psychological question of coping with a CVD event and its aftermath seems to be their question: “how can I live a meaningful and satisfying life, despite the disease and the changes in my life?”

**Meaning-oriented coping**

Patients may have a broad range of answers to this question, as reviewed elsewhere (Vos, 2016a, 2017). This model focuses on the core component of meaning, which is mentioned in many of the dominant medical and health psychology paradigms, such as stress-coping and appraisal models (Park and Folkman, 1997; Lee et al., 2016), terror management theory (Solomon et al., 2015), motivation psychology and motivational interviewing (Lundahl et al., 2013), post-traumatic growth (Tedeschi et al., 1998), and the recovery model (Bennett et al., 2014a). Similar to other diseases, several studies have shown that positive well-being, including meaning, is associated with lower cardiovascular risks and ill-being, albeit with modest effect sizes possibly because the effects are mediated by other factors such as lifestyle changes, stress level and biophysical changes (Boehm et al., 2011).

**Long-term impact**

When individuals can make sense of their disease and its treatment, they are more motivated for the required lifestyle changes (ibidem). Meaning-centered coping skills follow from the concept

of meaning, which is an empirically validated experience, associated with motivation, values, self-worth, understanding and self-insight, self-worth, self-regulation, and goal-setting (Vos, 2016a, 2017). A review of 45,000 individuals world-wide show that individuals find meaning via material-hedonic, self-oriented, social, larger, or existential-philosophical types of meaning (ibidem). Research indicates that patients directly benefit from exploring a wide number of meanings (ibidem). Thus, meaning-centered coping-styles include a well-established range of skills, of which the most important are: accepting the disease and its limitations; experiencing meaning despite change (i.e., continued meanings, new meanings, transcending the situation, and new perspectives on life); using meaning as resilience and stress-buffer; beneficial coping-styles; hope and optimism; self-efficacy; and motivation to lifestyle change (ibidem).

## Discussion

This systematic literature review has shown how meaning is a moderately strong predictor of the occurrence of a CVD event and of the short-term and long-term psychological and physical impact of CVD. To make sense of this literature, other studies were integrated in the development of a logical meaning-centered model of CVD. This model indicated the central role that meaning plays for patients before and after CVD events.

Systematic studies need to further confirm the meaning-centered model, and particularly examine how individuals create and adjust meaning after CVD events. There is a risk of confirmation bias due to reviewing only literature on meaning, and not searching for other literature on other CVD-related mechanisms.

Following the systematic literature review, CVD patients may be recommended to improve their sense of meaning in life, to reduce their CVD risk, and to improve their mental and physical well-being on short-term and long-term. As an improved sense of meaning in life seems to have moderately strong positive effects on these outcomes. Our review suggests that the effect size of meaning on CVD is similar or larger than the effect sizes for usual recommendations to improve the outcomes after a CVD event, such as stopping with smoking, reducing body weight, or using statins (Kannell and Higgins, 1990; Law et al., 2003; Lee et al., 2008; Rizo et al., 2012). Although more systematic and longitudinal studies are warranted to confirm the strengths of these relationships, these findings indicate the importance of meaning in life in the recovery after CVD events. One of the ways to improve this could be to offer meaning-oriented therapy (Vos, 2016a, 2017).

**Funding.** The author does not claim any funding for this study.

**Conflict of interest.** The author does not claim any competing interests.

## References

- Ai AL, Seymour EM, Tice TN, et al. (2009) Spiritual struggle related to plasma interleukin-6 prior to cardiac surgery. *Psychology of Religion and Spirituality* 1(2), 112–128. doi:10.1037/a0015775
- Ai AL, Pargament K, Kronfol Z, et al. (2010) Pathways to postoperative hostility in cardiac patients. *Journal of Health Psychology* 15(2), 186–195. doi:10.1177/1359105309345556
- Ai AL, Hopp F, Tice TN, et al. (2012) Existential relatedness in light of eudemonic well-being and religious coping among middle-aged and older cardiac patients. *Journal of Health Psychology* 18(3), 368–382. doi:10.1177/1359105311434754
- Amianto F, Bellicanta A, Bergui GC, et al. (2013) Growing up with congenital heart disease: Neurocognitive outcome, psychopathology and quality of life. *Rivista di psichiatria* 48(6), 415.
- Andersson EK, Borglin G and Willman A (2013) The experience of younger adults following myocardial infarction. *Qualitative Health Research* 23(6), 762–772. doi:10.1177/1049732313482049
- Apers S, Rassart J, Luyckx K, et al. (2016) Bringing Antonovsky's salutogenic theory to life: A qualitative inquiry into the experiences of young people with congenital heart disease. *International Journal of Qualitative Studies on Health and Well-being* 11, 29346–29346. doi:10.3402/qhw.v11.29346
- Balon YE, Then KL, Rankin JA, et al. (2008) Looking beyond the biophysical realm to optimize health: Results of a survey of psychological well-being in adults with congenital cardiac disease. *Cardiology in the Young* 18(5). doi:10.1017/s1047951108002606
- Barello S, Graffigna G, Vegni E, et al. (2015) 'Engage me in taking care of my heart': A grounded theory study on patient-cardiologist relationship in the hospital management of heart failure. *BMJ Open* 5(3), e005582. doi:10.1136/bmjopen-2014-005582
- Bekke-Hansen S, Pedersen CG, Thygesen K, et al. (2013) The role of religious faith, spirituality and existential considerations among heart patients in a secular society: Relation to depressive symptoms 6 months post acute coronary syndrome. *Journal of Health Psychology* 19(6), 740–753. doi:10.1177/1359105313479625
- Bennett B, Breeze J and Neilson T (2014a) Applying the recovery model to physical rehabilitation. *Nursing Standard* 28(23), 37–43. doi:10.7748/ns2014.02.28.23.37.e8292
- Bennett DA, Arnold SE, Valenzuela MJ, et al. (2014b) Cognitive and social lifestyle: Links with neuropathology and cognition in late life. *Acta Neuropathologica* 127(1), 137–150. doi:10.1007/s00401-013-1226-2
- Berghammer MC, Brink E, Rydberg AM, et al. (2015) Committed to life: Adolescents' and young adults' experiences of living with Fontan circulation. *Congenital Heart Disease* 10(5), 403–412. doi:10.1111/chd.12244
- Boehm JK and Kubzansky LD (2012) The heart's content: The association between positive psychological well-being and cardiovascular health. *Psychological Bulletin* 138(4), 655–691. doi:10.1037/a0027448
- Boehm JK, Peterson C, Kivimaki M, et al. (2011) A prospective study of positive psychological well-being and coronary heart disease. *Health Psychology* 30(3), 259–267. doi:10.1037/a0023124
- Boehm JK, Winning A, Segerstrom S, et al. (2015) Variability modifies life satisfaction's association with mortality risk in older adults. *Psychological Science* 26(7), 1063–1070. doi:10.1177/0956797615581491
- Bremer A, Dahlberg K and Sandman L (2009) Experiencing out-of-hospital cardiac arrest: Significant others' lifeworld perspective. *Qualitative Health Research* 19(10), 1407–1420. doi:10.1177/1049732309348369
- Buck HG, McAndrew L, Dionne-Odom JN, et al. (2015) What were they thinking? *Journal of Hospice & Palliative Nursing* 17(3), 249–256. doi:10.1097/njh.0000000000000156
- Burton NW, Pakenham KI and Brown WJ (2010) Are psychologists willing and able to promote physical activity as part of psychological treatment? *International Journal of Behavioral Medicine* 17(4), 287–297. doi:10.1007/s12529-010-9087-8
- Carroll KM and Nuro KF (2002) One size cannot fit all: A stage model for psychotherapy manual development. *Clinical Psychology: Science and Practice* 9(4), 396–406.
- Charlson ME, Wells MT, Peterson JC, et al. (2014) Mediators and moderators of behavior change in patients with chronic cardiopulmonary disease: The impact of positive affect and self-affirmation. *Translational Behavioral Medicine* 4(1), 7–17. doi:10.1007/s13142-013-0241-0
- Chen E, McLean KC and Miller GE (2015) Shift-and-persist strategies. *Psychosomatic Medicine* 77(4), 371–382. doi:10.1097/psy.0000000000000157
- Chida Y and Hamer M (2008) Chronic psychosocial factors and acute psychological responses to laboratory-induced stress in healthy populations: A quantitative review of 30 years of investigations. *Psychological Bulletin* 134(6), 829–885. doi:10.1037/a0013342
- Chida Y and Steptoe A (2008) Positive psychological well-being and mortality: A quantitative review of prospective observational studies. *Psychosomatic Medicine* 70(7), 741–756. doi:10.1097/psy.0b013e31818105ba

- Clarke PJ, Black SE, Badley EM, *et al.* (1999) Handicap in stroke survivors. *Disability and Rehabilitation* 21(3), 116–123. doi:10.1080/096382899297855
- Clarke V, Braun V and Hayfield N (2015) *Thematic Analysis. Qualitative Psychology: A Practical Guide to Research Methods*, 222–248.
- Cohen R, Bavishi C and Rozanski A (2016) Purpose in life and its relationship to all-cause mortality and cardiovascular events. *Psychosomatic Medicine* 78(2), 122–133. doi:10.1097/psy.0000000000000274
- Cortis JD and Williams A (2007) Palliative and supportive needs of older adults with heart failure. *International Nursing Review* 54(3), 263–270. doi:10.1111/j.1466-7657.2007.00558.x
- Coughlin SS (2011) Post-traumatic stress disorder and cardiovascular disease. *The Open Cardiovascular Medicine Journal* 5, 164–170. doi:10.2174/1874192401105010164
- Cuffee Y, Ogedegbe C, Williams NJ, *et al.* (2014) Psychosocial risk factors for hypertension: An update of the literature. *Current Hypertension Reports* 16(10), 483.
- Davidson PM, Cockburn J, Webster JK, *et al.* (2003) Patients with chronic heart failure rate psychosocial and existential needs higher than physical needs when recently discharged from hospital. *Journal of Cardiac Failure* 9(5), S105. doi:10.1016/s1071-9164(03)00202-1
- Davidson KW, Mostofsky E and Whang W (2010) Don't worry, be happy: Positive affect and reduced 10-year incident coronary heart disease: The Canadian Nova Scotia Health Survey. *European Heart Journal* 31(9), 1065–1070. doi:10.1093/eurheartj/ehp603
- Dimsdale JE (2008) Psychological stress and cardiovascular disease. *Journal of the American College of Cardiology* 51(13), 1237–1246. doi:10.1016/j.jacc.2007.12.024
- DuBois CM, Beach SR, Kashdan TB, *et al.* (2012) Positive psychological attributes and cardiac outcomes: Associations, mechanisms, and interventions. *Psychosomatics* 53(4), 303–318. doi:10.1016/j.psym.2012.04.004
- DuBois CM, Lopez OV, Beale EE, *et al.* (2015) Relationships between positive psychological constructs and health outcomes in patients with cardiovascular disease: A systematic review. *International Journal of Cardiology* 195, 265–280. doi:10.1016/j.ijcard.2015.05.121
- Ecohard R, Colin C, Rabilloud M, *et al.* (2001) Indicators of myocardial dysfunction and quality of life, one year after acute infarction. *European Journal of Heart Failure* 3(5), 561–568. doi:10.1016/s1388-9842(01)00171-4
- Edmondson KA, Lawler KA, Jobe RL, *et al.* (2005) Spirituality predicts health and cardiovascular responses to stress in young adult women. *Journal of Religion and Health* 44(2), 161–171. doi:10.1007/s10943-005-2774-0
- Eilertsen G, Kirkevold M and Bjørk IT (2010) Recovering from a stroke: A longitudinal, qualitative study of older Norwegian women. *Journal of Clinical Nursing* 19(13–14), 2004–2013. doi:10.1111/j.1365-2702.2009.03138.x
- Evangelista LS, Doering L and Dracup K (2003) Meaning and life purpose: The perspectives of post-transplant women. *Heart & Lung* 32(4), 250–257. doi:10.1016/s0147-9563(03)00042-6
- Faircloth CA, Boylstein C, Rittman M, *et al.* (2004) Sudden illness and biographical flow in narratives of stroke recovery. *Sociology of Health and Illness* 26(2), 242–261. doi:10.1111/j.1467-9566.2004.00388.x
- Falk H, Ekman I, Anderson R, *et al.* (2013) Older patients' experiences of heart failure — An integrative literature review. *Journal of Nursing Scholarship*. doi:10.1111/jnu.12025
- Gillham S and Clark L (2011) Psychological care after stroke: Improving stroke services for people with cognitive and mood disorders. *NHS Improvement*. [https://www.nice.org.uk/media/default/sharedlearning/531\\_strokepsychologicalsupportfinal.pdf](https://www.nice.org.uk/media/default/sharedlearning/531_strokepsychologicalsupportfinal.pdf)
- Grohn B, Worrall L, Simmons-Mackie N, *et al.* (2014) Living successfully with aphasia during the first year post-stroke: A longitudinal qualitative study. *Aphasiology* 28(12), 1405–1425. doi:10.1080/02687038.2014.935118
- Hamer M and Chida Y (2011) Life satisfaction and inflammatory biomarkers: The 2008 Scottish Health Survey. *Japanese Psychosomatic Research* 53(2), 133–139. doi:10.1111/j.1468-5884.2011.00460.x
- Hansen TB, Zwisler AD, Berg SK, *et al.* (2016) Cardiac rehabilitation patients' perspectives on the recovery following heart valve surgery: A narrative analysis. *Journal of Advanced Nursing* 72(5), 1097–1108. doi:10.1111/jan.12904
- Haukka A, Kontinen H, Lehto E, *et al.* (2013) Sense of coherence, depressive symptoms, cardiovascular diseases, and all-cause mortality. *Psychosomatic Medicine* 75(4), 429–435. doi:10.1097/psy.0b013e31828c3fa4
- Hefferon K, Grealy M and Mutrie N (2009) Post-traumatic growth and life threatening physical illness: A systematic review of the qualitative literature. *British Journal of Health Psychology* 14(2), 343–378. doi:10.1348/135910708x332936
- Hoehn PW, Denollet J, de Jonge P, *et al.* (2013) Positive affect and survival in patients with stable coronary heart disease. *The Journal of Clinical Psychiatry* 74(07), 716–722. doi:10.4088/jcp.12m08022
- Holahan CK, Holahan CJ and Suzuki R (2008) Purposiveness, physical activity, and perceived health in cardiac patients. *Disability and Rehabilitation* 30(23), 1772–1778. doi:10.1080/10428190701661508
- Huffman JC, Beale EE, Celano CM, *et al.* (2016a) Effects of optimism and gratitude on physical activity, biomarkers, and readmissions after an acute coronary syndrome: The gratitude research in acute coronary events study. *Circulation Cardiovascular Quality and Outcomes* 9(1), 55–63. doi:10.1161/CIRCOUTCOMES.115.002184
- Huffman JC, DuBois CM, Mastromauro CA, *et al.* (2016b) Positive psychological states and health behaviors in acute coronary syndrome patients: A qualitative study. *Journal of Health Psychology* 21(6), 1026–1036. doi:10.1177/1359105314544135
- Ishida R and Okada M (2006) Effects of a firm purpose in life on anxiety and sympathetic nervous activity caused by emotional stress: Assessment by psycho-physiological method. *Stress and Health* 22(4), 275–281. doi:10.1002/smi.1095
- Junehag L, Asplund K and Svedlund M (2014) A qualitative study: Perceptions of the psychosocial consequences and access to support after an acute myocardial infarction. *Intensive and Critical Care Nursing* 30(1), 22–30. doi:10.1016/j.iccn.2013.07.002
- Kannel WB and Higgins M (1990) Smoking and hypertension as predictors of cardiovascular risk in population studies. *Journal of hypertension. Supplement: official journal of the International Society of Hypertension* 8(5), S3–8.
- Kazdin AE (2009) Understanding how and why psychotherapy leads to change. *Psychotherapy Research* 19(4–5), 418–428. doi:10.1080/10503300802448899
- Kennedy S (2007) Nurses' perception of their learning needs to deliver CVD prevention programmes. *British Journal of Cardiac Nursing* 2(12), 571–579.
- Kim ES, Sun JK, Park N, *et al.* (2013) Purpose in life and reduced incidence of stroke in older adults: 'The Health and Retirement Study'. *Journal of Psychosomatic Research* 74(5), 427–432. doi:10.1016/j.jpsychores.2013.01.013
- Kohlmann S, Kilbert MS, Ziegler K, *et al.* (2013) Supportive care needs in patients with cardiovascular disorders. *Patient Education and Counseling* 91(3), 378–384. doi:10.1016/j.pec.2013.01.002
- Koizumi M, Ito H, Kaneko Y, *et al.* (2008) Effect of having a sense of purpose in life on the risk of death from cardiovascular diseases. *Journal of Epidemiology* 18(5), 191–196. doi:10.2188/jea.je2007388
- Konstam V, Moser DK and De Jong MJ (2005) Depression and anxiety in heart failure. *Journal of Cardiac Failure* 11(6), 455–463. doi:10.1016/j.cardfail.2005.03.006
- Kopp MS, Skrabski A, Szekely A, *et al.* (2007) Chronic stress and social changes: Socioeconomic determination of chronic stress. *Annals of the New York Academy of Sciences* 1113(1), 325–338. doi:10.1196/annals.1391.006
- Kožul K, Vidović K, Heinzelman-Kožul H, *et al.* (2009) The heart frequency and its variability in hypertensive patients considering A/B type of behaviour and eight basic emotions and levels of anger expression. *Collegium Antropologicum* 33(2), 409–416.
- Krucoff MW, Crater SW, Green CL, *et al.* (2001) Integrative noetic therapies as adjuncts to percutaneous intervention during unstable coronary syndromes: Monitoring and Actualization of Noetic Training (MANTRA) feasibility pilot. *American Heart Journal* 142(5), 760–769. doi:10.1067/mhj.2001.119138
- Krucoff MW, Crater SW, Gallup D, *et al.* (2005) Music, imagery, touch, and prayer as adjuncts to interventional cardiac care: The Monitoring and Actualisation of Noetic Trainings (MANTRA) II randomised study. *The Lancet* 366(9481), 211–217. doi:10.1016/s0140-6736(05)66910-3
- Kutner JS (2010) An 86-year-old woman with cardiac cachexia contemplating the end of her life. *JAMA* 303(4), 349. doi:10.1001/jama.2009.2015
- Laerum E, Johnsen N, Smith PAL, *et al.* (1991) Positive psychological and life-style changes after myocardial infarction: A follow-up study after 2–4 years. *Family Practice* 8(3), 229–233. doi:10.1093/fampra/8.3.229



- Lambiase MJ, Kubzansky LD and Thurston RC** (2015) Positive psychological health and stroke risk: The benefits of emotional vitality. *Health Psychology* **34**(10), 1043–1046. doi:10.1037/hea0000228
- Lantz J and Gregoire T** (2003) *Journal of Contemporary Psychotherapy* **33**(1), 19–33. doi:10.1023/a:1021451610123
- Law MR, Wald NJ and Rudnicka AR** (2003) Quantifying effect of statins on low density lipoprotein cholesterol, ischaemic heart disease, and stroke: Systematic review and meta-analysis. *BMJ* **326**(7404), 1423–1423. doi:10.1136/bmj.326.7404.1423
- Lawrence M and Kinn S** (2012) Needs, priorities, and desired rehabilitation outcomes of family members of young adults who have had a stroke: Findings from a phenomenological study. *Disability and Rehabilitation* **35**(7), 586–595. doi:10.3109/09638288.2012.711895
- Lee CMY, Huxley RR, Wildman RP, et al.** (2008) Indices of abdominal obesity are better discriminators of cardiovascular risk factors than BMI: A meta-analysis. *Journal of Clinical Epidemiology* **61**(7), 646–653. doi:10.1016/j.jclinepi.2007.08.012
- Lee WWM, Choi KC, Yum RWY, et al.** (2016) Effectiveness of motivational interviewing on lifestyle modification and health outcomes of clients at risk or diagnosed with cardiovascular diseases: A systematic review. *International Journal of Nursing Studies* **53**, 331–341. doi:10.1016/j.ijnurstu.2015.09.010
- Leegaard M and Fagermoen MS** (2008) Patients' key experiences after coronary artery bypass grafting: A synthesis of qualitative studies. *Scandinavian Journal of Caring Sciences* **22**(4), 616–628. doi:10.1111/j.1471-6712.2007.00556
- Leeming A, Murray SA and Kendall M** (2014) The impact of advanced heart failure on social, psychological and existential aspects and personhood. *European Journal of Cardiovascular Nursing* **13**(2), 162–167. doi:10.1177/1474515114520771
- Lemay K and Wilson K** (2008) Treatment of existential distress in life threatening illness: A review of manualized interventions. *Clinical Psychology Review* **28**(3), 472–493. doi:10.1016/j.cpr.2007.07.013
- Liberati A, Altman DG, Tetzlaff J, et al.** (2009) The PRISMA statement for reporting systematic reviews and meta-analyses of studies that evaluate health care interventions: Explanation and elaboration. *PLoS Medicine* **6**(7), e1000100. doi:10.1371/journal.pmed.1000100
- Littooij E, Dekker J, Vloothuis J, et al.** (2016) Global meaning in people with stroke: Content and changes. *Health Psychology Open* **3**(2). 10.1177/2055102916681759
- Lobeck M, Thompson AR and Shankland MC** (2005) The experience of stroke for men in retirement transition. *Qualitative Health Research* **15**(8), 1022–1036. doi:10.1177/1049732305280772
- Low TT and Lam CS** (2013) Women and heart failure: An emerging Venus-Mars concept. *Current Cardiovascular Risk Reports* **7**(3), 212–216.
- Lundahl B, Moleni T, Burke BL, et al.** (2013) Motivational interviewing in medical care settings: A systematic review and meta-analysis of randomized controlled trials. *Patient Education and Counseling* **93**(2), 157–168. doi:10.1016/j.pec.2013.07.012
- Marteinsdottir I, Ernerudh J, Jonasson L, et al.** (2016) Psychological resources are independently associated with markers of inflammation in a middle-aged community sample. *International Journal of Behavioral Medicine* **23**(5), 611–620. doi:10.1007/s12529-016-9553-z
- Mårtensson J, Karlsson J-E and Fridlund B** (1997) Male patients with congestive heart failure and their conception of the life situation. *Journal of Advanced Nursing* **25**(3), 579–586. doi:10.1046/j.1365-2648.1997.1997025579.x
- Mårtensson J, Karlsson JE and Fridlund B** (1998) Female patients with congestive heart failure: How they conceive their life situation. *Journal of Advanced Nursing* **28**(6), 1216–1224. doi:10.1046/j.1365-2648.1998.00827.x
- McClung JA** (2013) End-of-life care in the treatment of advanced heart failure in the elderly. *Cardiology in Review* **21**(1), 9–15. doi:10.1097/crd.0b013e31826d23ea
- Mezick EJ, Matthews KA, Hall M, et al.** (2010) Low life purpose and high hostility are related to an attenuated decline in nocturnal blood pressure. *Health Psychology* **29**(2), 196–204. doi:10.1037/a0017790
- Moore GF, Audrey S, Barker M, et al.** (2015) Process evaluation of complex interventions: Medical Research Council guidance. *BMJ* **350**, h1258. doi:10.1136/bmj.h1258
- Muirhead J, Meyerowitz BE, Leedham B, et al.** (1992) Quality of life and coping in patients awaiting heart transplantation. *The Journal of heart and lung transplantation: the official publication of the International Society for Heart Transplantation* **11**(2 Pt 1), 265–271.
- Nikrahan GR, Laferton JAC, Asgari K, et al.** (2016a) Effects of positive psychology interventions on risk biomarkers in coronary patients: A randomized, wait-list controlled pilot trial. *Psychosomatics* **57**(4), 359–368. doi:10.1016/j.psych.2016.02.007
- Nikrahan GR, Suarez L, Asgari K, et al.** (2016b) Positive psychology interventions for patients with heart disease: A preliminary randomized trial. *Psychosomatics* **57**(4), 348–358. doi:10.1016/j.psych.2016.03.003
- Nilsson I, Jansson I and Norberg A** (1999) Crisis phenomena after stroke reflected in an existential perspective. *The International Journal of Aging and Human Development* **49**(4), 259–277. doi:10.2190/uuae-tq52-luq3-jvrl
- Ogedegbe GO, Boutin-Foster C, Wells MT, et al.** (2012) A randomized controlled trial of positive-affect intervention and medication adherence in hypertensive African Americans. *Archives of Internal Medicine* **172**(4), 322–326. doi:10.1001/archinternmed.2011.1307
- Owolabi MO** (2009) What are the consistent predictors of generic and specific post-stroke health-related quality of life? *Cerebrovascular Diseases* **29**(2), 105–110. doi:10.1159/000262305
- Park CL** (2010) Making sense of the meaning literature: An integrative review of meaning making and its effects on adjustment to stressful life events. *Psychological Bulletin* **136**(2), 257–301. doi:10.1037/a0018301
- Park CL and Folkman S** (1997) Meaning in the context of stress and coping. *Review of General Psychology* **1**(2), 115–144. doi:10.1037/1089-2680.1.2.115
- Park CL, Malone MR, Suresh DP, et al.** (2007) Coping, meaning in life, and quality of life in congestive heart failure patients. *Quality of Life Research* **17**(1), 21–26. doi:10.1007/s11136-007-9279-0
- Peterson JC, Charlson ME, Hoffman Z, et al.** (2012) A randomized controlled trial of positive-affect induction to promote physical activity after percutaneous coronary intervention. *Archives of Internal Medicine* **172**(4), 329–336. doi:10.1001/archinternmed.2011.1311
- Popogrebsky AP** (1998) Influence of myocardial infarction on the persons' meaning of life. *Psikhologicheskii Zhurnal* **19**(5), 113–120.
- Proyer RT, Gander F, Wellenzohn S, et al.** (2013) What good are character strengths beyond subjective well-being? The contribution of the good character on self-reported health-oriented behavior, physical fitness, and the subjective health status. *The Journal of Positive Psychology* **8**(3), 222–232. doi:10.1080/17439760.2013.777767
- Rassin M, Zilcha L and Gross D** (2009) 'A pacemaker in my heart' — Classification of questions asked by pacemaker patients as a basis for intervention. *Journal of Clinical Nursing* **18**(1), 56–62. doi:10.1111/j.1365-2702.2008.02432.x
- Rizos EC, Ntzani EE, Bika E, et al.** (2012) Association between omega-3 fatty acid supplementation and risk of major cardiovascular disease events. *JAMA* **308**(10), 1024. doi:10.1001/2012.jama.11374
- Roest AM, Martens EJ, de Jonge P, et al.** (2010) Anxiety and risk of incident coronary heart disease. *Journal of the American College of Cardiology* **56**(1), 38–46. doi:10.1016/j.jacc.2010.03.034
- Rohleder N** (2014) Stimulation of systemic low-grade inflammation by psychosocial stress. *Psychosomatic Medicine* **76**(3), 181–189. doi:10.1097/psyc.0000000000000049
- Ronaldson A, Molloy GJ, Wikman A, et al.** (2015) Optimism and recovery after acute coronary syndrome: A clinical cohort study. *Psychosomatic Medicine* **77**(3), 311–318. doi:10.1097/PSY.0000000000000155
- Roncella A, Pristipino C, Cianfrocca C, et al.** (2013) One-year results of the randomized, controlled, short-term psychotherapy in acute myocardial infarction (STEP-IN-AMI) trial. *International Journal of Cardiology* **170**(2), 132–139. doi:10.1016/j.ijcard.2013.08.094
- Ross L and Austin J** (2013) Spiritual needs and spiritual support preferences of people with end-stage heart failure and their carers: Implications for nurse managers. *Journal of Nursing Management* **23**(1), 87–95. doi:10.1111/jonm.12087
- Rounsaville BJ, Carroll KM and Onken LS** (2006) A stage model of behavioral therapies research: Getting started and moving on from stage I. *Clinical Psychology: Science and Practice* **8**(2), 133–142. doi:10.1093/clipsy.8.2.133
- Roysland IØ and Friberg F** (2015) Unexplained chest pain and physical activity. *Qualitative Health Research* **26**(2), 215–226. doi:10.1177/1049732315570129



- Rozanski A and Kubzansky LD (2005) Psychologic functioning and physical health: A paradigm of flexibility. *Psychosomatic Medicine* 67, S47–S53.
- Rozanski A, Blumenthal JA and Kaplan J (1999) Impact of psychological factors on the pathogenesis of cardiovascular disease and implications for therapy. *Circulation* 99(16), 2192–2217. doi:10.1161/01.cir.99.16.2192
- Rozanski A, Blumenthal JA, Davidson KW, et al. (2005) The epidemiology, pathophysiology, and management of psychosocial risk factors in cardiac practice. *Journal of the American College of Cardiology* 45(5), 637–651. doi:10.1016/j.jacc.2004.12.005
- Ryff CD, Singer BH and Diener G (2004) Positive health: Connecting well-being with biology. *Philosophical Transactions of the Royal Society of London. Series B: Biological Sciences* 359(1449), 1383–1394.
- Sacco SJ, Park CL, Suresh DP, et al. (2014) Living with heart failure: Psychosocial resources, meaning, gratitude and well-being. *Heart & Lung* 43(3), 213–218. doi:10.1016/j.hrtlng.2014.01.012
- Sanjuán P, Montalbetti T, Pérez-García AM, et al. (2016) A randomised trial of a positive intervention to promote well-being in cardiac patients. *Applied Psychology: Health and Well-Being* 8(1), 64–84. doi:10.1111/aphw.12062
- Sarfo FS, Jenkins C, Singh A, et al. (2017) Post-stroke depression in Ghana: Characteristics and correlates. *Journal of the Neurological Sciences* 379, 261–265. doi:10.1016/j.jns.2017.06.032
- Satink T, Josephsson S, Zajec J, et al. (2016) Self-management develops through doing of everyday activities—a longitudinal qualitative study of stroke survivors during two years post-stroke. *BMC Neurology* 16(1), 1–13.
- Schaufel MA, Nordrehaug JE and Malterud K (2009) “So you think I’ll survive?": A qualitative study about doctor-patient dialogues preceding high-risk cardiac surgery or intervention. *Heart* 95(15), 1245–1249. doi:10.1136/hrt.2008.164657
- Schou L and Egerod I (2008) A qualitative study into the lived experience of post-CABG patients during mechanical ventilator weaning. *Intensive and Critical Care Nursing* 24(3), 171–179. doi:10.1016/j.iccn.2007.12.004
- Schwerdtfeger AR and Gerteis AKS (2014) The manifold effects of positive affect on heart rate variability in everyday life: Distinguishing within-person and between-person associations. *Health Psychology* 33(9), 1065–1073. doi:10.1037/hea0000079
- Scully D, Kremer J, Meade MM, et al. (1998) Physical exercise and psychological well-being: A critical review. *British Journal of Sports Medicine* 32(2), 111–120. doi:10.1136/bjsm.32.2.111
- Secrest JS and Zeller R (2003) Measuring continuity and discontinuity following stroke. *Journal of Nursing Scholarship* 35(3), 243–247. doi:10.1111/j.1547-5069.2003.00243.x
- Secrest J and Zeller R (2006) Replication and extension of the continuity and discontinuity of self scale (CDSS). *Journal of Nursing Scholarship* 38(2), 154–158. doi:10.1111/j.1547-5069.2006.00093.x
- Seskevich JE, Crater SW, Lane JD, et al. (2004) Beneficial effects of noetic therapies on mood before percutaneous intervention for unstable coronary syndromes. *Nursing Research* 53(2), 116–121. doi:10.1097/00006199-200403000-00007
- Shao J, Zhang Q, Lin T, et al. (2013) Well-being of elderly stroke survivors in Chinese communities: Mediating effects of meaning in life. *Aging & Mental Health* 18(4), 435–443. doi:10.1080/13607863.2013.848836
- Shirai K, Iso H, Ohira T, et al. (2009) Perceived level of life enjoyment and risks of cardiovascular disease incidence and mortality. *Circulation* 120(11), 956–963. doi:10.1161/circulationaha.108.834176
- Silva J, Ownsworth T, Shields C, et al. (2011) Enhanced appreciation of life following acquired brain injury: Posttraumatic growth at 6 months post-discharge. *Brain Impairment* 12(2), 93–104. doi:10.1375/brim.12.2.93
- Simonj CP, Pedersen BD, Dreyer P, et al. (2015) Dealing with existential anxiety in exercise-based cardiac rehabilitation: A phenomenological-hermeneutic study of patients’ lived experiences. *Journal of Clinical Nursing* 24(17–18), 2581–2590. doi:10.1111/jocn.12867
- Sin NL (2016) The protective role of positive well-being in cardiovascular disease: Review of current evidence, mechanisms, and clinical implications. *Current Cardiology Reports* 18(11), 106–106. doi:10.1007/s11886-016-0792-z
- Sin NL, Moskowitz JT and Whooley MA (2015) Positive affect and health behaviors across 5 years in patients with coronary heart disease: The heart and soul study. *Psychosomatic Medicine* 77(9), 1058–1066. doi:10.1097/PSY.0000000000000238
- Sirri L, Potena L, Masetti M, et al. (2010) Psychological predictors of mortality in heart transplanted patients: A prospective, 6-year follow-up study. *Transplantation* 89(7), 879–886. doi:10.1097/tp.0b013e3181ca9078
- Skrabski A, Kopp M, Rózsa S, et al. (2005) Life meaning: An important correlate of health in the Hungarian population. *International Journal of Behavioral Medicine* 12(2), 78–85. doi:10.1207/s15327558ijbm1202\_5
- Smith E and Kneebone I (2016) Assessment and psychological interventions for depression comorbid with cardiovascular disease. In *Cardiovascular Diseases and Depression*. Cham: Springer International Publishing, pp. 351–364.
- Solomon S, Greenberg J and Pyszczynski T (2015) *The Worm at the Core: On the Role of Death in Life*. New York: Random House.
- Sone T, Nakaya N, Ohmori K, et al. (2008) Sense of life worth living (Ikigai) and mortality in Japan: Ohsaki Study. *Psychosomatic Medicine* 70(6), 709–715. doi:10.1097/psy.0b013e31817e7e64
- Stephoe A, Demakos P, de Oliveira C, et al. (2012) Distinctive biological correlates of positive psychological well-being in older men and women. *Psychosomatic Medicine* 74(5), 501–508. doi:10.1097/psy.0b013e31824f82c8
- Stoll C, Schelling G, Goetz AE, et al. (2000) Health-related quality of life and post-traumatic stress disorder in patients after cardiac surgery and intensive care treatment. *The Journal of Thoracic and Cardiovascular Surgery* 120(3), 505–512.
- Strang S, Hensch I, Danielson E, et al. (2013) Communication about existential issues with patients close to death—nurses’ reflections on content, process and meaning. *Psycho-Oncology* 23(5), 562–568. doi:10.1002/pon.3456
- Stroup DF, Berlin JA, Morton SC, et al. (2000) Meta-analysis of observational studies in epidemiology: A proposal for reporting. *Jama* 283(15), 2008–2012.
- Su S, Jimenez MP, Roberts CTF, et al. (2015) The role of adverse childhood experiences in cardiovascular disease risk: A review with emphasis on plausible mechanisms. *Current Cardiology Reports* 17(10), 88. doi:10.1007/s11886-015-0645-1
- Tan MP and Morgan K (2015) Psychological interventions in cardiovascular disease. *Current Opinion in Psychiatry* 28(5), 371–377. doi:10.1097/ycp.0000000000000181
- Tanno K, Sakata K, Ohsawa M, et al. (2009) Associations of ikigai as a positive psychological factor with all-cause mortality and cause-specific mortality among middle-aged and elderly Japanese people: Findings from the Japan Collaborative Cohort Study. *Journal of Psychosomatic Research* 67(1), 67–75. doi:10.1016/j.jpsychores.2008.10.018
- Tedeschi RG, Park CL and Calhoun LG (Eds.) (1998) *Posttraumatic Growth: Positive Changes in the Aftermath of Crisis*. New York: Routledge.
- Tinker LF, Rosal MC, Young AF, et al. (2007) Predictors of dietary change and maintenance in the women’s health initiative dietary modification trial. *Journal of the American Dietetic Association* 107(7), 1155–1165. doi:10.1016/j.jada.2007.04.010
- van Montfort E, Denollet J, Widdershoven J, et al. (2016) Interrelation and independence of positive and negative psychological constructs in predicting general treatment adherence in coronary artery patients — Results from the THORESCI study. *Journal of Psychosomatic Research* 88, 1–7. doi:10.1016/j.jpsychores.2016.06.009
- Versteeg H, Hoogwegt MT, Hansen TB, et al. (2013) Depression, not anxiety, is independently associated with 5-year hospitalizations and mortality in patients with ischemic heart disease. *Journal of Psychosomatic Research* 75(6), 518–525. doi:10.1016/j.jpsychores.2013.10.005
- Vollman MW, LaMontagne LL and Wallston KA (2009) Existential well-being predicts perceived control in adults with heart failure. *Applied Nursing Research* 22(3), 198–203. doi:10.1016/j.apnr.2008.02.001
- Vos J (2016a) Working with meaning in life in chronic or life-threatening disease: A review of its relevance and the effectiveness of meaning-centred therapies. In *Clinical Perspectives on Meaning*. Springer International Publishing, pp. 171–200.
- Vos J (2016b) Working with meaning in life in mental health care: A systematic literature review of the practices and effectiveness of meaning-centred therapies. In Russo-Netzer P, Schulenberg SE and Batthyany A (eds.), *Clinical Perspectives on Meaning: Positive and Existential Psychotherapy*. New York: Springer International Publishing, pp. 59–87.

- Vos J** (2017) *Meaning in Life: An Evidence-Based Handbook for Practitioners*. London: Macmillan International Higher Education.
- Vos J and Vitali D** (2018) The effects of psychological meaning-centered therapies on quality of life and psychological stress: A metaanalysis. *Palliative and Supportive Care* **16**(5), 608–632. doi:10.1017/s1478951517000931
- Wampold BE, Mondin GW, Moody M, et al.** (1997) A meta-analysis of outcome studies comparing bona fide psychotherapies: Empirically, “all must have prizes”. *Psychological Bulletin* **122**(3), 203–215. doi:10.1037/0033-2909.122.3.203
- Wann-Hansson C, Rahm Hallberg I, Klevsgård R, et al.** (2008) The long-term experience of living with peripheral arterial disease and the recovery following revascularisation: A qualitative study. *International Journal of Nursing Studies* **45**(4), 552–561. doi:10.1016/j.ijnurstu.2006.11.006
- Whalley B, Rees K, Davies P, et al.** (2011) Psychological interventions for coronary heart disease. In *Cochrane Database of Systematic Reviews*. John Wiley & Sons, Ltd.
- Whalley B, Thompson DR and Taylor RS** (2014) Psychological interventions for coronary heart disease: Cochrane systematic review and meta-analysis. *International Journal of Behavioral Medicine* **21**(1), 109–121.
- Yu L, Boyle PA, Wilson RS, et al.** (2015) Purpose in life and cerebral infarcts in community-dwelling older people. *Stroke* **46**(4), 1071–1076. doi:10.1161/STROKEAHA.114.008010