


## DIRECT DATING OF THE HUMAN NAVICULAR FROM THE CUEVA DE LOS TORREJONES (GUADALAJARA, SPAIN)

Adrián Pablos<sup>1,2\*</sup>  • Nohemi Sala<sup>1,2</sup> • Alfonso Arribas<sup>3</sup>

<sup>1</sup>Centro Nacional de Investigación sobre la Evolución Humana-CENIEH, Burgos, Spain

<sup>2</sup>Centro Mixto UCM-ISCIH de Investigación sobre Evolución y Comportamiento Humanos, Madrid, Spain

<sup>3</sup>Estación paleontológica Valle del río Fardes, Instituto Geológico y Minero de España (IGME), Tres Cantos-Madrid, Spain

**ABSTRACT.** Pleistocene human remains are rare inland on the Iberian Peninsula. Most are considered Neandertals, but anthropological analyses and direct dating are rare. Recently, we published a study of a navicular from this region found in the Torrejones Cave. The results showed it differed from that of Neandertals and it was re-identified as *Homo sapiens*. Following the previous stratigraphic and biochronologic descriptions, we suggested that it could correspond to an Upper Paleolithic human, since the navicular was apparently recovered in the Late Pleistocene from an *in situ* unit. Direct radiocarbon dating from this fossil (4855–5036 cal BP), believed to be the only Paleolithic *Homo sapiens* from inland Iberia, as well as other hominin and faunal remains from the site, show that the human bones actually date to the Chalcolithic. The unexpectedly recent chronology for the navicular implies that there is no evidence of human fossils from the Upper Paleolithic in Torrejones Cave. Thus, any date from the Middle/Upper Paleolithic human record should be taken with caution until in-depth paleoanthropological, stratigraphical and/or direct dating studies are conducted. Extraordinary caution is recommended when human remains are recovered from apparently Paleolithic units in contexts bearing Holocene sepulchral units on the uppermost levels and/or some evidence of bioturbation.

**KEYWORDS:** feet, Holocene/Pleistocene, Iberian Peninsula, navicular, radiocarbon dating.

### INTRODUCTION

Human remains corresponding to Paleolithic *Homo sapiens* are not abundant in the European archaeological record in general, and especially not in certain regions of the Iberian Peninsula, as is the case of inland Iberia. A paleoanthropological study was recently published regarding a human navicular (foot bone) from the Cueva de los Torrejones (Tamajón, Guadalajara, Spain), which was previously considered *Homo* sp. (Arribas et al. 1995), showing that it actually belongs to *Homo sapiens* (Pablos et al. 2018). This human bone (T93-S3-27) was recovered from the Lithostratigraphic Unit-Sumidero 3 (LU-S3) during the 1993 field excavation associated with faunal remains from the Late Pleistocene (Arribas et al. 1995, 1997; Díez Fernández-Lomana et al. 1998; Carrión et al. 2007).

An in-depth and detailed anthropological study of this foot bone ruled out the possibility of it belonging to Neandertals or their ancestors (Pablos et al. 2017), suggesting that it could correspond to a *H. sapiens* fossil (Pablos et al. 2018). In fact, if the stratigraphical and biochronological assignment were confirmed, this foot fossil would become the only Upper Paleolithic human fossil found in the interior of the Iberian Peninsula. Therefore, the implications of this new taxonomic reassignment are particularly relevant due to the absence of Paleolithic *Homo sapiens* remains in inland Iberia (Martínez et al. 2017; Pablos et al. 2018). In addition, Upper Paleolithic occupation sites in this region are scarce. In fact, evidence of late Neandertal or Mousterian survival south of the Ebro River on the Iberian Peninsula, which has been reliably dated using the Ultrafiltration method, ceases more than 40–42 ka BP (Kehl et al. 2013; Wood et al. 2013; Higham et al. 2014; Alcaraz-Castaño et al. 2015, 2017a; Wolf et al. 2018, but see Zilhão et al. 2017 for a discussion). The oldest evidence of Upper Paleolithic sites (Gravettian) within the Iberian Peninsula is dated to around 25–26 ka BP at

\*Corresponding author. Email: [adrizaino@yahoo.es](mailto:adrizaino@yahoo.es)

the Peña Capón (Guadalajara, Spain) site (Alcaraz-Castaño et al. 2013, 2017b, 2019a), which could indicate a long period with no evidence of human presence inland on the Iberian Peninsula. Nevertheless, La Boja rockshelter site displays early or basal Aurignacian chronologies of around 36.5 ka cal BP, though it is located on the coast (Zilhão et al. 2017).

For these reasons, the implications of these findings required a more accurate chronological framework for this human fossil. In this study, we present the direct dating of the human navicular found in the Lithostratigraphic Unit S3 (LU-S3), as well as other hominin and faunal bones and teeth from different sectors and levels from the Cueva de los Torrejones using the AMS radiocarbon ( $^{14}\text{C}$ ) method with ultrafiltration pretreatment.

### Historical Background

The Cueva de los Torrejones (Tamajón, Guadalajara, Spain) was excavated from 1993 until 1995 (Arribas et al. 1995, 1997, 2005). Excavation at this site was divided into four different sectors in the cavity: Entrada (entrance), Sumidero (sinkhole), Pasillo (corridor), and Tejones (badgers). These excavations documented two chronological episodes: the upper units corresponding to the Holocene and the lower units assigned to the Upper Pleistocene (Arribas et al. 1995, 1997, 2005). The fauna recovered from the lower levels of the cave included hyena (*Crocuta crocuta*), bear (*Ursus* sp.), steppe rhinoceros (*Stephanorhinus* cf. *hemitoechus*), and leopard (*Panthera pardus*) remains, among other species, which suggested an age of between 60 and 80 thousand years (ka) for the Pleistocene deposit (Arribas et al. 1995, 1997; Arribas 1997, 2005). A few lithic tools preliminarily assigned to the late Middle Paleolithic were recovered from the lower part of the sequence (Arribas et al. 1995; Carrión et al. 2007). A palynological analysis indicated very recent dates for three badger coprolites from the surface levels, which were directly dated to 150–250 BP (Carrión et al. 2005). This suggests a continuous presence of badgers in the cave from the Pleistocene until recent times.

Human remains were found, along with the rest of the recovered fauna, most of which came from the disturbed upper levels and their chronological ascription corresponded to the Chalcolithic. In some exceptional cases, some human remains found in the superficial and reworked units were assigned to *H. sapiens* cf. *neanderthalensis* due to their plesiomorphic characters (Arribas et al. 1995; Arribas and Jordá 1999). A hominin foot remain (a right navicular bone found in the LU-S3 from the Sumidero sector) was particularly emphasized. Although this element was assigned as *Homo* sp. due to the absence of diagnostic characteristics, its stratigraphic ascription in an *in situ* unit (associated with hyenids and ursids) suggested an older chronology for the rest of the human remains, assigning it to a Pleistocene, late Middle Pleistocene or early Late Pleistocene chronology (Arribas et al. 1995). In 2017, a new excavation and multidisciplinary research project was initiated. The objectives of that project, which remains ongoing, were to gain new insight into the geology, stratigraphy, taphonomy, taxonomy, anthropology, archaeology, and chronology of the site. During three short field seasons from 2017 until 2019, a small surface area of the upper levels of the Sumidero sector was excavated, reaching the top of LU-S3 where the human navicular supposedly came from. The association of the human navicular with Pleistocene fauna suggested a potential Upper Palaeolithic chronology. Given the scarcity of Palaeolithic hominin remains throughout Europe and their absence in central Iberia, the specific objective of this article is to verify the human navicular chronology through direct dating. Moreover, other human and faunal remains were directly dated to provide context for the human navicular.

## MATERIAL AND METHODS

A huge effort has been made at this site to provide the most accurate chronological contextualization possible, focusing on the use of radiocarbon techniques, specifically the ultrafiltration method with the ABA pretreatment, on the fossil remains to effectively remove contaminants from the bone collagen (Bronk Ramsey et al. 2004b; Higham et al. 2006; Higham 2011), with special emphasis on the most relevant taxa from the Sumidero sector. Seven samples (bones and teeth) were sent for AMS-ultrafiltration dating purposes to different labs: Oxford Radiocarbon Accelerator Unit (ORAU)-OxA in England (Brock et al. 2010), Beta Analytic-BETA in the United States ([www.radiocarbon.com](http://www.radiocarbon.com)), and Curt-Engelhorn-Zentrum Archaeometrie-MAMS in Germany (Kromer et al. 2013). However, only four of them yielded an age. Moreover, two additional elements (ursid and hyenid bones/teeth from LU-S3) were selected for possible direct dating, but they did not weigh enough to be sampled. We directly sampled several *Homo* and equid bones from the upper units (LU-S1 and LU-S2, Figure 1) in the Sumidero sector, which were recovered throughout the course of this current project. Furthermore, we sampled an equid tooth and the human navicular, both recovered from level LU-S3 during the excavation carried out in the 90s. Finally, a *C. crocuta* premolar recovered in 2017 from a surface level (level LU-T1) of the “Tejones” sector was also sampled.

All  $^{14}\text{C}$  dates presented in this paper were obtained using ultrafiltration pretreatments to reduce environmental contamination and remove short-chained proteins (see Brock et al. 2007, 2010; Higham et al. 2006 for details of the method of ultrafiltration pretreatment) and have been calibrated using the OxCal v.4.3 software (Bronk Ramsey 2009) against the IntCal13curve (Reimer et al. 2013). The  $^{14}\text{C}$  ages offered in this study are given in BP (before present), meaning years before 1950, and are calibrated (cal BP) to provide absolute calendar ages. The results of the calibration are shown as cal BP (2  $\sigma$ ; 95.4% probability). To verify the state of preservation of the collagen in the different bones to be dated, the C:N ratio, %C and %N values must be evaluated. The C:N ratio should be between 2.9 and 3.6 (Ambrose 1990; van Klinken 1999; Higham et al. 2014). Once the samples met the collagen quality criteria, they were graphitized and dated according to the methods of each lab using ultrafiltration pretreatment in each case (Bronk Ramsey et al. 2004a; Bronk Ramsey 2009).

The methodology carried out by the three labs for AMS radiocarbon dating with which we deal followed the Longin protocol (Longin 1971), modified by Brown et al. (1988). All the three labs applied the ABA (acid-base-acid) pretreatment after the scraping of the surface to eliminate soil contaminants (Table 1). The ABA method is carried out to effectively remove sedimentary and other contaminant carbonates (Bronk Ramsey et al. 2004b; Brock et al. 2010). After the pretreatment, a gelatinization process according to Longin (1971) is done, and the obtained gelatin is filtered with Eeze-like filters to remove small (<80  $\mu\text{m}$ ) solid particles (Brock et al. 2010). The gelatin obtained in the former steps is ultrafiltered with ultrafilters Vivaspin-like of 30 KDa previous to combustion and graphitization in all of the three labs (Brock et al. 2010; Wacker et al. 2010a, 2010b). In every lab, a rigorous precleaning procedure of the ultrafilters is done to effectively remove the carbon-containing humectants present in the filters prior to the ultrafiltration of the bone gelatin (Brock et al. 2007). The filters are precleaned in the three labs by ultrasonication in ultrapure water (Table 1), and then passed through fresh ultrapure water prior to use (Brock et al. 2010). The Beta lab furthermore applies a previous bath of NaOH to the ultrafilters.




Figure 1 A) geographic location and digital terrain model showing the position of the Cueva de los Torrejones site (source LiDAR-IGN). B) Stratigraphic column of the Sumidero Sector from Cueva de los Torrejones showing the different lithostratigraphic units (LU). The colors used in the stratigraphic column correspond to those of the Munsell® color system for wet sediment. Scale in cm. The  $^{14}\text{C}$  samples (TRJ-1 and TRJ-2; Table 2) that are currently being excavated are indicated in their approximate location by an asterisk (\*). The samples recovered from the prior excavation are not shown due to the ambiguity of the original position. C) Topographic scheme of the site plan showing the different sectors. Modified from Arribas et al. (2005). D) Distal (left) and proximal view (right) of the human navicular bone (T93-S3-27), modified from Pablos et al. (2018). Scale in cm.

Table 1 Comparison of the ABA pretreatment and the precleaning of the filters carried out at all the three labs considered in this study.

Lab*	Material	Physical cleaning	ABA pretreatment			Longin protocol	Filter (Eeze-filter)	UF 30 kDa (Vivaspin)	Reference
			Acid (HCL)	Base (NaOH)	Acid (HCL)				
ORAU	Bone/tooth	Scraping	0.5 M	0.1 M	0.5 M	Yes	Yes	Yes	Bronk Ramsey et al. 2004b; Brock et al. 2007, 2010
Beta	Bone/tooth	Scraping	0.2 N	1–2%	1.22 N	Yes	Yes	Yes	<a href="http://www.radiocarbon.com">www.radiocarbon.com</a> & pers. comm. <sup>1</sup>
MAMS	Bone/tooth	Scraping	4%	0.4%	4%	Yes	Yes	Yes	Wacker et al. 2010a, 2010b; Kromer et al. 2013 & pers. comm. <sup>2</sup>

Precleaning of ultrafilters					
	NAOH	H2O	Ultrasonic	H2O centrifugation	Reference
ORAU	No	Yes	Yes	Yes	Brock et al. 2007
Beta	Yes (2%)	Yes	Yes	Yes	Personal communication <sup>1</sup>
MAMS	No	Yes	Yes	Yes	Personal communication <sup>2</sup>

<sup>1</sup>Personal communication, Chris Patrick (Vice President of Laboratory Operations-Beta).

<sup>2</sup>Personal communication, Ronny Friedrich (Laborleiter-Curt-Engelhorn-Zentrum Archäometrie gmbH-MAMS).

\*Beta = Beta analytic Inc. ORAU = Oxford Radiocarbon Accelerator Unit. MAMS = Curt-Engelhorn-Centre for Archaeometry. UF = Ultrafiltration.

Table 2 AMS radiocarbon dates using ultrafiltration pretreatment from different sectors and levels from Cueva de los Torrejones.\*

Site inventory	Sample	Lab /reference	Level	Material	Age (BP)	cal BP (2 $\sigma$ )	C:N	Wt %C	Wt %N
MUPA-TMJ-0006	TRJ-1	Beta-474503	S1	<i>Homo</i> (rib)	4150 $\pm$ 30	4773–4577 (76.6%) 4825–4778 (18.8 %)	+3.1	+38.2	+14.2
MUPA-TMJ-0049	TRJ-3	Beta	S1	<i>Homo</i> (cranium)	Failed	—	—	—	—
MUPA-TMJ-115	TRJ-2	Beta-474504	S2	<i>Homo</i> (rib)	4420 $\pm$ 30	5064–4870 (82.7%) 5215–5185 (7.1%) 5270–5221 (5%) 5120–5112 (0.6%)	+3.2	+41.2	+15.2
MUPA-TMJ-0079	TRJ-4	Beta	S2	<i>Equus</i> (radius)	Failed	—	—	—	—
T93-S3.27	TRJ-6	MAMS-34581	S3	<i>Homo</i> (navicular)	4366 $\pm$ 30	4979–4855 (87.7%) 5036–5009 (7.7%)	+2.9	+38.4	—
T93-S3.24	TRJ-5	Beta-481425	S3	<i>Equus</i> (lower molar)	26,080 $\pm$ 100	30,750–29,950 (95.4 %)	+3.2	+41.8	+15.1
MUPA-TMJ-169	TRJ-7	ORAU	T1	<i>Crocota</i> (premolar)	Failed	—	—	—	—

\*All ages are calibrated with the OxCal v4.3 software (Bronk Ramsey 2009) against the IntCal13 curve providing a probability range of 95.4% (Reimer et al. 2013). For the samples, we included the isotopic values and C:N ratios. Beta = Beta analytic Inc. ORAU = Oxford Radiocarbon Accelerator Unit. MAMS = Curt-Engelhorn-Centre for Archaeometry. BP = before present (years before 1950). cal BP = calibrated years before present (all probability ranges are expressed at 95.4%: 2  $\sigma$ ).

## RESULTS

In order to provide an accurate framework for determining the chronology of the Cueva de los Torrejones human fossil, a revised stratigraphy has been performed on the archaeological pit where it was recovered. In the Sumidero sector, all the profiles exposed in the test pit during the excavations in the 90s were studied. Three main levels have been defined, which coincide with the previously proposed description (Arribas et al. 1995, 1997, 2005); they have been subdivided, however, into additional sub-units (Figure 1).

$^{14}\text{C}$  dates for the different lithostratigraphic units from the Sumidero sector are presented in Table 2. Two dates on human remains from around 4500–5300 cal BP confirm Chalcolithic ages for the uppermost levels (LU-S1 and LU-S2). It was not possible to date two of the samples (one from each of these levels), as not enough collagen was recovered. These dates are compatible with the lithic industry, the micromammal association, and the presence of pottery in these two upper levels.

One Pleistocene date of 30,750–29,950 cal BP was obtained for an equid lower molar found in level LU-S3 (Beta-481425). However, the human navicular recovered from the LU-S3 provided a Holocene age of 4855–5036 cal BP (MAMS-34581, mass of extracted collagen on the raw bone mass (%Coll) = 6.3 referred to the >30 kDa fraction). A hyenid premolar recovered from surface LU-T1 was sampled to be dated and did not provide positive results because not enough collagen was preserved.

## CONCLUDING DISCUSSION

The unexpectedly recent chronology of the navicular T93-S3-27 implies that there is no evidence of human fossils from the Upper Paleolithic in the Cueva de los Torrejones. Rather, this human bone is coetaneous with the human remains recovered from the overlying levels (LU-S1 and LU-S2). However, the direct dating of one horse fossil remain from the same unit (found together with the navicular bone) provided an age of ca. 30,000 cal BP, thus confirming the Pleistocene chronology for the LU-S3.

This chronological inconsistency suggests a mixture of different units, perhaps due to bioturbation processes as was previously observed, or due to inclusions of the LU-S2 in the LU-S3 as described in previous studies (Arribas et al. 1995, 2005). In any case, direct dating of the human bone indicates an obvious mixture of chronologies. The causes of this mixture will be investigated in future studies through geological, taphonomic, biostratigraphical, and micromorphological analyses.

Holocene human remains in Late Pleistocene contexts are not rare in the karstic environments from the European archaeological record (Conard et al. 2004; Gómez-Olivencia 2013; Sala et al. 2013 among others). The closest known case is found in the Cueva de la Zarzamora site in Segovia (Spain), where remains of Chalcolithic individuals have been found incorporated into a hyena den deposit corresponding to the Marine Isotopic Stage (MIS) 3 (Sala et al. 2011, 2013). As a cautionary note, the human remains belonging to *H. sapiens* apparently recovered from Paleolithic cave contexts, which were subsequently used as sepulcher during recent Prehistory (Neolithic/ Chalcolithic/ Bronze), must be studied from a Paleoanthropological point of view and/or directly dated.

As a final remark, evidence of occupation of inland Iberia during the Middle-Upper Paleolithic period is increasingly growing in the areas surrounding the Cueva de los

Torrejones (Alcaraz-Castaño et al. 2013, 2017b, 2019a, 2019b; Alcolea González et al. 1995, 1997; Sala et al. 2020) and, therefore, it is necessary to continue working in the region to expand upon the information currently available.

## ACKNOWLEDGMENTS

This work could not have been carried out without the collaboration of C. Posth and J. Krause, who contributed with the direct dating of the navicular T93-S3-27. We are deeply grateful to the Tamajón-Torrejones research and excavation team. We really would like to acknowledge the Instituto Geológico y Minero de España (IGME) and its personnel, especially S. Menéndez, for allowing access to the fossils under their care. The effort put forth and work carried out by the excavation team during the 1993–1995 seasons in the Cueva de los Torrejones is very much appreciated. We are grateful for the constructive and fruitful discussion provided by M. Alcaraz-Castaño. Especial thanks to Chris Patrick (Beta) and Ronny Friedrich (MAMS) for methodological information provided. Logistic and administrative support from the Ayuntamiento de Tamajón is deeply appreciated.

Part of this research was funded by the Spanish projects of the “Ministerio de Ciencia e Innovación (MCI)” PGC2018-093925-B-C33 and PGC2018-093925-B-C31 (MCI/AEI/FEDER, UE) and “Junta de Comunidades de Castilla la Mancha” (SBPLY/17/180801/000103, SBPLY/18/180801/000034 and SBPLY/19/180801/000015). N-S was supported by a Juan de la Cierva Incorporación postdoctoral contract (IJCI-2017-32804). Lauren Ames kindly reviewed a previous English version of the manuscript.

Finally, we appreciate the helpful comments and suggestions provided by Associate Editor Christine Hatté and two anonymous reviewers, which significantly improved the article.

## REFERENCES

- Alcaraz-Castaño M, Alcolea-González J, Kehl M, Albert R-M, Baena-Preysler J, de Balbín-Behrmann R, Cuartero F, Cuenca-Bescós G, Jiménez-Barredo F, López-Sáez J-A, Piqué R, Rodríguez-Antón D, Yravedra J, Weniger G-C. 2017a. A context for the last Neandertals of interior Iberia: Los Casares cave revisited. *PLoS ONE* 12(7):e0180823.
- Alcaraz-Castaño M, Alcolea-González J, Pablos A, Sala N. 2019a. A Solutrean territory in Central Iberia: new field evidence in the context of recent climate modelling. Workshop: The Last Glacial Maximum in Europe state of knowledge in geosciences and archaeology; Erlangen (Germany). Friedrich-Alexander-Universität Erlangen-Nürnberg. p. 7–8.
- Alcaraz-Castaño M, Alcolea-González J, Balbín-Behrman R, Kehl M, Weniger G-C. 2019b. Recurrent human occupations in Central Iberia around the Last Glacial Maximum. The Solutrean sequence of Peña Capón updated, in: Schmidt I, Cascalleira J, Bicho N, Weniger G-C (Eds.), *Human adaptations to the Last Glacial Maximum: The Solutrean and its neighbors*. 3rd international conference on the Solutrean. Cambridge Scholars Publishing, Faro (Portugal). p. 148–170.
- Alcaraz-Castaño M, Alcolea González J, Balbín Rd, García Valero MÁ, Yravedra J, Baena J. 2013. Los orígenes del Solutrense y la ocupación pleniglaciario del interior de la Península Ibérica: implicaciones del nivel 3 de Peña Capón (valle del Sorbe, Guadalajara). *Trabajos de Prehistoria* 70(1):28-53.
- Alcaraz-Castaño M, López-Recio M, Tapias F, Cuartero F, Baena J, Ruiz-Zapata B, Morín J, Pérez-González A, Santonja M. 2017b. The human settlement of Central Iberia during MIS 2: New technological, chronological and environmental data from the Solutrean workshop of Las Delicias (Manzanares River valley, Spain). *Quaternary International* 431(Part B):104–124.
- Alcaraz-Castaño M, Weniger G-C, Alcolea J, de Andrés-Herrero M, Baena J, de Balbín R, Bolin V, Cuartero F, Kehl M, López A, López-Sáez JA, Martínez Mendizabal I, Pablos A, Rodríguez-Antón D, Torres C, Vizcaíno J, Yravedra J. 2015. Regreso a la



- Cueva de los Casares (Guadalajara). Un nuevo proyecto de investigación para el yacimiento del seno A. *Arqueología y Prehistoria del Interior Peninsular* 2:68–89.
- Alcolea González JJ, Balbín Rd, García MA, Jiménez PJ. 1997. Nouvelles découvertes d'Art Pariétal Paléolithique á la Meseta: La grotte del Reno (Valdesotos, Guadalajara). *L'Anthropologie* 101: 144–163.
- Alcolea González JJ, de Balbín R, García Valero MÁ, Cruz Naimi LA. 1995. La Cueva del Turismo (Tamajón, Guadalajara): Un nuevo yacimiento rupestre Paleolítico en la meseta Castellana. In: De Balbín R, Valiente J, Mussat T, editors. *Arqueología en Guadalajara*. Guadalajara (Spain): Servicio de Publicaciones. Junta de Comunidades de Castilla la Mancha. p. 127–136.
- Ambrose SH. 1990. Preparation and characterization of bone and tooth collagen for isotopic analysis. *Journal of Archaeological Science* 17:431–451.
- Arribas A. 1997. Un leopardo, *Panthera pardus* (Linnaeus, 1758), en el Pleistoceno de la Cueva de los Torrejones (Tamajón, Guadalajara, España). *Geogaceta* 22:19–22.
- Arribas A, Díez Fernández-Lomana C, Jordá Pardo JF. 1997. Primeras ocupaciones en los depósitos Pleistocenos de la Cueva de los Torrejones (Sistema Central español, Tamajón, Guadalajara): Litografía y actividad biológica. *Cuaternario y Geomorfología* 11(1-2):55–66.
- Arribas A, Díez JC, Jordá FJ. 1995. El yacimiento cuaternario de la Cueva de Los Torrejones (Tamajón, Guadalajara, Castilla-La Mancha): resultados preliminares. In: Balbín R, Valiente J, Musat MT, editors. *Arqueología en Guadalajara*. Guadalajara (Spain): Patrimonio Histórico-Arqueología Castilla-La Mancha. p. 97–110.
- Arribas A, Garrido G, García Solano JA. 2005. Depósitos re trabajados y fósiles Pleistocenos reelaborados en un karst de la provincia de Guadalajara (Cueva de los Torrejones). Evidencias de ocupaciones biológicas Pleistocenas y Holocenas, de reactivación kárstica post-calcolítica y de una ocupación biológica histórica. In: Martinell J, Doménech R, Gibert JMeditors. 2nd International Meeting Taphos; Barcelona (Spain). Universitat de Barcelona.
- Arribas A, Jordá JF. 1999. Los mamíferos del Cuaternario kárstico de Guadalajara (Castilla-La Mancha, España). In: Aguirre E, Rábano I, editors. *La Huella del Pasado Fósiles de Castilla-La Mancha*. Patrimonio Histórico. *Arqueología Castilla-La Mancha*. p. 327–353.
- Brock F, Bronk Ramsey C, Higham T. 2007. Quality assurance of ultrafiltered bone dating. *Radiocarbon* 49:187–192.
- Brock F, Higham T, Ditchfield P, Bronk Ramsey C. 2010. Current pretreatment methods for AMS radiocarbon dating at the Oxford Radiocarbon Accelerator Unit (ORAU). *Radiocarbon* 52: 103–112.
- Bronk Ramsey C, Higham T, Leach P. 2004a. Towards high-precision AMS: Progress and limitations. *Radiocarbon* 46:17–24.
- Bronk Ramsey C, Higham T, Bowles A, Hedges R. 2004b. Improvements to the pretreatment of bone at Oxford. *Radiocarbon* 46:155–163.
- Bronk Ramsey C. 2009. Bayesian analysis of radiocarbon dates. *Radiocarbon*. 51(1):337–360.
- Brown TA, Nelson DE, Vogel JS, Southon JR. 1988. Improved collagen extraction by modified Longin method. *Radiocarbon* 30:171–177.
- Carrión JS, Gil G, Rodríguez E, Fuentes N, García-Antón M, Arribas A. 2005. Palynology of badger coprolites from central Spain. *Palaeogeography, Palaeoclimatology, Palaeoecology* 226(3–4): 259–271.
- Carrión JS, Scott L, Arribas A, Fuentes N, Gil-Romera G, Montoya E. 2007. Pleistocene landscapes in central Iberia inferred from pollen analysis of hyena coprolites. *Journal of Quaternary Science* 22(2):191–202.
- Conard NJ, Grootes PM, Smith FH. 2004. Unexpectedly recent dates for human remains from Vogelherd. *Nature* 430(6996):198–201.
- Díez Fernández-Lomana C, Jordá Pardo JF, Arribas Herrera A. 1998. Torrejones (Tamajón, Guadalajara, Spain). A hyaena den on human occupation. *Economie préhistorique: les comportements de subsistance au Paléolithique*. Editions APDCA, Sophia Antipolis. p. 63–72.
- Gómez-Olivencia A. 2013. Estudio de los restos humanos del yacimiento de Arlanpe (Lemoa, Bizkaia). In: Rios-Garaizar J, Garate Maidagan D, Gómez-Olivencia A, editors. *La cueva de Arlanpe (Lemoa): Ocupaciones humanas desde el Paleolítico Medio Antiguo hasta la Prehistoria Reciente*. Diputación Foral de Bizkaia, Bilbao: Kobie Serie BAI 3. p. 161–175.
- Higham TFG, Jacobi RM, Bronk Ramsey CB. 2006. AMS radiocarbon dating of ancient bone using ultrafiltration. *Radiocarbon* 48:179–195.
- Higham T. 2011. European Middle and Upper Palaeolithic radiocarbon dates are often older than they look: problems with previous dates and some remedies. *Antiquity* 85(327):235–249.
- Higham T, Douka K, Wood R, Bronk Ramsey C, Brock F, Basell L, Camps M, Arrizabalaga A, Baena J, Barroso-Ruiz C, Bergman C, Boitard C, Boscato P, Caparros M, Conard NJ, Draily C, Froment A, Galvan B, Gambassini P, Garcia-Moreno A, Grimaldi S, Haesaerts P, Holt B, Iriarte-Chiapusso M-J, Jelinek A, Jorda Pardo JF, Maillo-Fernandez J-M, Marom A, Maroto J, Menendez M, Metz L, Morin E, Moroni A, Negrino F, Panagopoulou E, Peresani M, Pirson S, de la Rasilla M, Riel-Salvatore J, Ronchitelli A, Santamaria D, Semal P, Slimak L, Soler J, Soler N, Villaluenga A, Pinhasi R, Jacobi R. 2014. The timing and spatiotemporal patterning of Neanderthal disappearance. *Nature* 512(7514):306–309.

- Kehl M, Burow C, Hilgers A, Navazo M, Pastoors A, Weniger G-C, Wood R, Jordá Pardo JF. 2013. Late Neanderthals at Jarama VI (central Iberia)? *Quaternary Research* 80(2):218–234.
- Kromer B, Lindauer S, Synal H-A, Wacker L. 2013. MAMS – A new AMS facility at the Curt-Engelhorn-Centre for Achaemetry, Mannheim, Germany. *Nuclear Instruments and Methods in Physics Research Section B: Beam Interactions with Materials and Atoms* 294:11–13.
- Longin, R. 1971. New method of collagen extraction for radiocarbon dating. *Nature* 230:241–242.
- Martínez I, Pablos A, Sala N, Pantoja-Pérez A, Conde-Valverde M. 2017. Hitos y lugares claves para la evolución humana en Iberia. *Enseñanzas de las Ciencias de la Tierra* 25(1):59–64.
- Pablos A, Pantoja-Pérez A, Martínez I, Lorenzo C, Arsuaga JL. 2017. Metric and morphological analysis of the foot in the Middle Pleistocene population of Sima de los Huesos (Sierra de Atapuerca, Burgos, Spain). *Quaternary International* 433(Part A):103–113.
- Pablos A, Sala N, Arribas A. 2018. Taxonomic reassignment of the Paleolithic human navicular from Cueva de los Torrejones (Guadalajara, Spain). *Archaeological and Anthropological Sciences* 10(8):1867–1880.
- Reimer PJ, Bard E, Bayliss A, Beck JW, Blackwell PG, Bronk Ramsey C, Buck CE, Cheng H, Edwards RL, Friedrich M, Grootes PM, Guilderson TP, Hafliðason H, Hajdas I, Hatté C, Heaton TJ, Hoffmann DL, Hogg AG, Hughen KA, Kaiser KF, Kromer B, Manning SW, Niu M, Reimer RW, Richards DA, Scott EM, Southon JR, Staff RA, Turney CSM, van der Plicht J. 2013. IntCal13 and Marine13 radiocarbon age calibration curves 0–50,000 years cal BP. *Radiocarbon* 55(4):1869–1887.
- Sala N, Algaba M, Gómez-Olivencia A, Pablos A, Bonmatí A, Rodríguez L, García R, Arsuaga JL. 2013. Nuevos restos humanos procedentes de la Cueva de la Zarzamora (Segovia, España). *Munibe Antropología-Arkeología* 64:105–116.
- Sala N, Arsuaga JL, Laplana C, Zapata BR, Gil García MJ, García N, Aranburu A, Algaba M. 2011. Un paisaje de la meseta durante el Pleistoceno Superior. Aspectos paleontológicos de la Cueva de la Zarzamora (Segovia, España). *Boletín de la Real Sociedad Española de Historia Natural* 105:67–85.
- Sala N, Pablos A, Gómez-Olivencia A, Sanz A, Villalba M, Pantoja-Pérez A, Laplana C, Arsuaga JL, Algaba M. 2020. Central Iberia in the Middle MIS 3. Paleocological inferences during the period 34–40 cal kyr BP. *Quaternary Science Reviews* 228:106027.
- van Klinken GJ. 1999. Bone collagen quality indicators for palaeodietary and radiocarbon measurements. *Journal of Archaeological Science* 26:687–695.
- Wacker L, Němec M, Bourquin J. 2010a. A revolutionary graphitisation system: Fully automated, compact and simple. *Nuclear Instruments and Methods in Physics Research Section B: Beam Interactions with Materials and Atoms* 268:931–934.
- Wacker L, Bonani G, Friedrich M, Hajdas I, Kromer B, Němec M, Ruff M, Suter M, Synal HA, Vockenhuber C. 2010b. MICADAS: Routine and high-precision radiocarbon dating. *Radiocarbon* 52:252–262.
- Wolf D, Kolb T, Alcaraz-Castaño M, Heinrich S, Baumgart P, Calvo R, Sánchez J, Ryborz K, Schäfer I, Bliedner M, Zech R, Zöller L, Faust D. 2018. Climate deteriorations and Neanderthal demise in interior Iberia. *Scientific Reports* 8(1):7048.
- Wood RE, Barroso-Ruiz C, Caparrós M, Jordá Pardo JF, Galván Santos B, Higham TFG. 2013. Radiocarbon dating casts doubt on the late chronology of the Middle to Upper Palaeolithic transition in southern Iberia. *Proceedings of the National Academy of Sciences* 110(8):2781–2786.
- Zilhão J, Anesin D, Aubry T, Badal E, Cabanes D, Kehl M, Klasen N, Lucena A, Martín-Lerma I, Martínez S, Matias H, Susini D, Steier P, Wild EM, Angelucci DE, Villaverde V, Zapata J. 2017. Precise dating of the Middle-to-Upper Paleolithic transition in Murcia (Spain) supports late Neanderthal persistence in Iberia. *Heliyon* 3:e00435.