

THE DIRICHLET PROBLEM FOR THE MINIMAL SURFACES EQUATION AND THE PLATEAU PROBLEM AT INFINITY

LAURENT MAZET

*Laboratoire Emile Picard, Université Paul Sabatier, 118, route de Narbonne,
31062 Toulouse, France (mazet@picard.ups-tlse.fr)*

(Received 24 February 2003; accepted 1 September 2003)

Abstract In this paper, we shall study the Dirichlet problem for the minimal surfaces equation. We prove some results about the boundary behaviour of a solution of this problem. We describe the behaviour of a non-converging sequence of solutions in term of lines of divergence in the domain. Using this second result, we build some solutions of the Dirichlet problem on unbounded domain. We then give a new proof of the result of Cosín and Ros concerning the Plateau problem at infinity for horizontal ends.

Keywords: minimal surface; Dirichlet problem; boundary behaviour

AMS 2000 *Mathematics subject classification:* Primary 53A10

0. Introduction

One classical way to construct minimal surfaces in \mathbb{R}^3 is to see them as the graph of a function u over a domain $\Omega \subset \mathbb{R}^2$ (see, for example, the paper of Karcher [8]). The graph of a function u is a minimal surface if u satisfies the elliptic partial differential equation called the minimal surfaces equation,

$$\operatorname{div} \left(\frac{\nabla u}{\sqrt{1 + |\nabla u|^2}} \right) = 0. \quad (\text{MSE})$$

The problem which is associated to this point of view is the Dirichlet problem for the equation (MSE): for a domain Ω and a function f on $\partial\Omega$, this problem consists in finding a continuous function u on $\bar{\Omega}$ which is a solution of the minimal surfaces equation in Ω and such that $u = f$ on the boundary of Ω . One of the most general answers to the Dirichlet problem for bounded domain has been given by Jenkins and Serrin in [7]. They give a nice condition on the domain to solve for any function f ; moreover, their result allows us to give infinite value for the boundary data f . For unbounded domain, the Dirichlet problem is still an open problem. We know that, in the general case, we lose the uniqueness of solution. In this paper, using a new approach, we develop some tools for the study of this problem.

Another interesting and still open problem concerning minimal surfaces is the Plateau problem at infinity, which is the following: finding a minimal surface for a given asymptotic behaviour. More precisely, we know that, if a complete minimal surface has finite total curvature and embedded ends, each end of this minimal surface is asymptotic to a plane or to a half-catenoid; besides, we can associate to each end a vector in \mathbb{R}^3 , this vector is called the flux vector of the end. These vectors satisfy the following condition: the sum of the flux vectors over all ends is zero. So the problem is given a finite number of vectors such that their sum is zero, can we find a minimal surface which has these vectors as flux vectors? Our answer comes from the following idea: seeing a solution of the Plateau problem at infinity as the conjugate surface of a solution of the Dirichlet problem on an unbounded domain.

In [2], Cosín and Ros give a description of the space of solutions of the Plateau problem at infinity with an asymptotic behaviour which is symmetric with respect to an horizontal plane (i.e. all the flux vectors are horizontal). They also restrict themselves to the case of Alexandrov embedded minimal surfaces; this condition implies that no flux vector is zero and that there is a natural order on the ends of the surface. Since the flux vectors are horizontal and their sum is zero, these vectors draw a polygon in \mathbb{R}^2 . Cosín and Ros give a necessary and sufficient condition on this polygon to have a solution (see §4 for more explanations about their work).

In this paper, we give a more constructive proof of the result of Cosín and Ros. Our method is based on the Dirichlet problem on an unbounded ‘domain’ Ω . When the polygon given by the flux vectors is convex, Ω can be defined as the polygonal domain bounded by the flux polygon to which we glue a half-strip on each edge. We denote by L_i^+ and L_{i+1}^- the two sides of each half-strip S_i , alternating the sign $+$ and $-$ such that each vertex of the polygon is common to some L_i^- and L_i^+ . When the flux polygon is non-convex and satisfies the condition of Cosín and Ros, we need to use the concept of multi-domain for defining Ω (see Definition 1.1 for this concept).

Our main result for the Dirichlet problem for this kind of domain Ω is then (see Theorem 5.1) as follows.

There exists a solution u of the minimal surfaces equation on Ω such that u tends to $+\infty$ on L_i^+ and $-\infty$ on L_i^- . Besides, the solution is unique up to an additive constant.

This statement is a new result for the Dirichlet problem on unbounded domains that are similar to the ones studied by Rosenberg and Sa-Earp in [12].

The function u in this result is built as the limit of solutions of the Dirichlet problem on bounded domain. We describe the possible divergences that can occur for a sequence of solutions of (MSE). In fact, we prove that if the sequence diverges at a point, it must diverge along a line passing by this point. This result is a generalization of the results that Jenkins and Serrin use in [7]. Our result allows us to do the same discussion that Jenkins and Serrin made in the particular case of monotone sequences of solutions of (MSE); this is our main tool to prove the existence part of Theorem 5.1.

The solution to the Plateau problem at infinity is then the conjugate surface to the graph of u . In order to know the geometry of the conjugate surface along its boundary, we need to understand the behaviour of the graph in the neighbourhood of the vertices

of Ω , which are the vertices of the polygon. Some results are known for such a problem concerning the Dirichlet problem in the convex case. For example, consider f a datum on the boundary of a domain Ω . We suppose that f has a finite discontinuity at a point P where the boundary is convex (i.e. we suppose that $f(Q)$ has a limit if we tend to P by the right-hand side or by the left-hand side and that the difference of these two limits is finite). Then we know that the graph of a solution u over Ω of the Dirichlet problem with f as boundary value has a vertical segment over P in its boundary (as was proved in [10]). In our case, we can prove that the boundary of the graph is the vertical straight line passing by the vertex; although the domain is locally an angular sector that does not need to be convex and the boundary data takes the values $+\infty$ on one side of the sector and $-\infty$ on the other side.

The paper is organized as follows. In the first section, we define multi-domains and extend the result of Jenkins and Serrin to bounded multi-domains. The multi-domains are necessary to express the condition of Cosín and Ros. This result will be our first tool in the proof of our main theorem.

The second section is devoted to the proof of our result concerning the boundary behaviour of solutions of the Dirichlet problem.

In § 3, we study the sequences of solutions of (MSE) and define the lines of divergence.

In § 4, we explain the result of Cosín and Ros, and recall some elements of their proof. In the last section, we give the proof of our main result. We then use it to give a new proof of the result of Cosín and Ros.

Let us fix some notation. In the following, when u is a function on a domain of \mathbb{R}^2 we shall write $W = \sqrt{1 + |\nabla u|^2}$. We shall also use the classical following notation for partial derivatives:

$$p = \frac{\partial u}{\partial x}, \quad q = \frac{\partial u}{\partial y}, \quad r = \frac{\partial^2 u}{\partial x^2}, \quad s = \frac{\partial^2 u}{\partial x \partial y} \quad \text{and} \quad t = \frac{\partial^2 u}{\partial y^2}.$$

Besides, for the graph of u , we shall always chose the downward pointing normal to give an orientation to the graph.

In this paper, we shall call an extraction a strictly increasing function from \mathbb{N} to \mathbb{N} . If a_n is a sequence and θ is an extraction, the sequence $a_{\theta(n)}$ is a subsequence of a_n and every subsequence of a_n can be written in this way. If θ_1 is an extraction, we shall say that θ_2 is a sub-extraction of θ_1 if there exists an extraction α such that $\theta_2 = \theta_1 \circ \alpha$.

1. The Dirichlet problem on multi-domains

In this section, we shall give a generalization of the results of Jenkins and Serrin [7] for the Dirichlet problem on bounded domain. First we have to generalize the notion of domain of \mathbb{R}^2 . Let us consider a pair (Ω, φ) where Ω is a simply connected two-dimensional complete flat manifold with piecewise smooth boundary and $\varphi : \Omega \rightarrow \mathbb{R}^2$ is a local isometry. The map φ is called the *developing map* and the points where the boundary $\partial\Omega$ are not smooth are called *vertices*.

Definition 1.1. A pair (Ω, φ) , where Ω is a simply connected two-dimensional complete flat manifold with piecewise smooth boundary and $\varphi : \Omega \rightarrow \mathbb{R}^2$ is a local isometry, is a *multi-domain* if each connected component of the smooth part of $\partial\Omega$ is a convex arc.

If (Ω, φ) is as above and a part of $\partial\Omega$ is linear then we add two vertices at the endpoints of this linear part and we call this new part an *edge*.

Let (Ω, φ) be a multi-domain. If u is a smooth function on Ω we shall call the *graph* of u the surface in \mathbb{R}^3 given by $\{(\varphi(x), u(x))\}_{x \in \Omega}$. If u is a solution of the minimal surfaces equation (MSE), the graph of u is a minimal surface of \mathbb{R}^3 . The Dirichlet problem on a multi-domain consists in the determination of a function u satisfying the equation (MSE) on Ω and taking on assigned values on the boundary of Ω .

As in the case of a domain in \mathbb{R}^2 , if u is a solution of (MSE) on Ω , we can define a differential form $d\Psi_u$ on Ω which corresponds to the differential of the third coordinate of the conjugate surface of the graph of u . Using the charts given by the developing map φ , we have $d\Psi_u = (p/W) dy - (q/W) dx$. $d\Psi_u$ is a closed form by (MSE) and, since Ω is simply connected, we can define a function Ψ_u on Ω which is 1-Lipschitz continuous. We call this function the *conjugate function* to u . One important result concerning $d\Psi_u$ is the following lemma.

Lemma 1.2. *Let Ω be a domain bounded in part by a straight segment T , oriented such that the right-hand normal to T is the outer normal to Ω . Let u be a solution of (MSE) in Ω which assumes the boundary value $+\infty$ on T . Then*

$$\int_T d\Psi_u = |T|.$$

This is Lemma 4 of [7]. For other properties of Ψ_u and $d\Psi_u$, we refer the reader to this paper.

When Ω is compact there is a finite number of connected components of the smooth part of $\partial\Omega$; let us call them C_1, \dots, C_n . When the data on the boundary is bounded, we have the following result.

Theorem 1.3. *Let (Ω, φ) be a compact multi-domain with boundary arcs C_1, \dots, C_n and let u_1, \dots, u_n be bounded continuous functions respectively on C_1, \dots, C_n . Then there exists a unique solution u of the minimal surfaces equation on Ω such that $u|_{C_i} = u_i$.*

Proof. The proof of the uniqueness is a particular case of the proof of Theorem 1.5, so we make it later.

The existence of the solution on multi-domain is due to a Perron process. Let us recall some elements of this method. If v is a continuous function on Ω and D is a disc in Ω , we denote by $u_{v,D}$ the solution of (MSE) in D which takes the value v on ∂D . We also denote by $M_D[v]$ the continuous function which coincides with v on $\Omega \setminus D$ and $u_{v,D}$ on D . Let u_1, \dots, u_n be the data on the boundary of Ω . We say that v is a sub-solution of the Dirichlet problem if $v \leq u_i$ on C_i and $v \leq M_D[v]$ for all discs D in Ω . Since the u_i are bounded by a constant M , the class \mathcal{F} of all sub-solutions is non-empty, as it contains

the constant function $-M$; besides, each sub-solution v verifies $v \leq M$. So we can define a function u by

$$u(P) = \sup_{v \in \mathcal{F}} v(P) \quad \forall P \in \Omega. \quad (1.1)$$

By standard argument, we can show that u is a solution of (MSE). Since in our definition of multi-domains we suppose that the boundary is locally convex, there exist barrier functions on the boundary (they are constructed in using the Scherk surface). So we can ensure that u takes the value u_i on C_i . For more details on Perron processes, we refer the reader to the book of Gilbarg and Trudinger [4] or the one of Courant and Hilbert [3], which illustrate this method for the classical Laplacian Dirichlet problem; there is also the book of Nitsche [10], which studies the case of the minimal surfaces equation. \square

The work of Jenkins and Serrin allows infinite data on the boundary. By the Straight Line Lemma [7], we know that infinite data can only be allowed on linear parts of the boundary.

Definition 1.4. Let (Ω, φ) be a multi-domain. A *polygonal domain* \mathcal{P} of Ω is a connected compact subset of Ω such that (\mathcal{P}, φ) is a multi-domain, the boundary of \mathcal{P} is only composed of edges and the vertices of \mathcal{P} are drawn from the vertices of Ω .

We want to solve the Dirichlet problem with infinite data, so let us call A_1, \dots, A_k and B_1, \dots, B_l the edges of Ω such that we assign the value $+\infty$ on A_j and $-\infty$ on B_j . We call C_1, \dots, C_n the remaining arcs on which we assign continuous data.

Let \mathcal{P} be a polygonal domain of Ω . We denote by α and β , respectively, the total length of the edges A_j and the one of the edges B_j which belong to the boundary of \mathcal{P} and we denote by γ the perimeter of \mathcal{P} . We then have the following generalization of the result of Jenkins and Serrin.

Theorem 1.5. Let (Ω, φ) be a compact multi-domain with the families $\{A_j\}$, $\{B_j\}$ and $\{C_j\}$ as above.

If the family $\{C_j\}$ is non-empty, then there exists a solution of the minimal surface equation in Ω which assumes the value $+\infty$ on each A_j , the value $-\infty$ on each B_j and arbitrarily assigned continuous data on each C_j , if and only if

$$2\alpha < \gamma \quad \text{and} \quad 2\beta < \gamma \quad (*)$$

for each polygonal domain \mathcal{P} of Ω . If a solution exists, it is unique.

If the family $\{C_j\}$ is empty, then a solution exists, if and only if

$$\alpha = \beta$$

when \mathcal{P} coincides with Ω and $(*)$ holds for all other polygonal domains of Ω . In this case, if a solution exists, it is unique up to an additive constant.

Proof. To prove the existence of a solution, we can use the same arguments as Jenkins and Serrin, so we refer to [7].

The proof of the uniqueness in [7] also works but we give another proof which we can apply in other situations. Let u_1 and u_2 be different solutions of (MSE) with the same data on the boundary. In the case where the family (C_j) is empty, we suppose that $u_1 - u_2$ is not constant; besides, in considering $u_i - u_i(P)$ (where $P \in \Omega$), we can assume that $\{u_1 < u_2\}$ and $\{u_1 > u_2\}$ are non-empty. In choosing sufficiently small $\varepsilon > 0$, we have $\Omega_\varepsilon = \{u_1 - u_2 > \varepsilon\} \neq \emptyset$, besides the choice of ε is such that $\partial\Omega_\varepsilon$ is regular. We write $d\tilde{\Psi} = d\tilde{\Psi}_{u_1} - d\tilde{\Psi}_{u_2}$. Since $d\tilde{\Psi}$ is closed, we have $\int_{\partial\Omega_\varepsilon} d\tilde{\Psi} = 0$. Because u_1 and u_2 have the same data on the boundary, $\partial\Omega_\varepsilon$ does not intersect $\bigcup_j C_j$, so $\partial\Omega_\varepsilon$ is composed of three parts: one is included in $\bigcup_j A_j \cup \bigcup_j B_j$ on which $d\tilde{\Psi} = 0$ (this is a consequence of Lemma 1.2); one is included in Ω ; and a last part which is composed of some vertices of Ω but its contribution to the integral is zero. On the second part, let us call it $\partial\tilde{\Omega}_\varepsilon$, $\nabla u_1 - \nabla u_2$ points in Ω_ε , this part is then oriented by the non-direct normal to $\nabla u_1 - \nabla u_2$ so, by Lemma 2 of Collin and Krust in [1], $\int_{\partial\tilde{\Omega}_\varepsilon} d\tilde{\Psi} < 0$; this gives us a contradiction. \square

2. A result of regularity at the vertices

The aim of this section is to understand what geometrically happens at a vertex of a multi-domain where two edges A_j and B_j converge.

For $\beta_1 < \beta_2$ and $R > 0$, we consider

$$\Omega_{\beta_1}^{\beta_2}(R) = \{(r, \theta) \mid 0 \leq r \leq R, \beta_1 \leq \theta \leq \beta_2\},$$

with the metric $ds^2 = dr^2 + r^2 d\theta^2$ (we identify all the points $(0, \theta)$ and this point will be called the vertex of $\Omega_{\beta_1}^{\beta_2}(R)$). We also define on $\Omega_{\beta_1}^{\beta_2}(R)$ the map $\varphi : (r, \theta) \mapsto (r \cos \theta, r \sin \theta)$. Then $(\Omega_{\beta_1}^{\beta_2}(R), \varphi)$ is a multi-domain, it is a description of a neighbourhood of a vertex where two edges converge. We call $L(\beta)$ the set of points in $\Omega_{\beta_1}^{\beta_2}(R)$ such that $\theta = \beta$. We are interested in the geometrical ‘configuration’ of the graph of a solution u of (MSE) such that u tends to $-\infty$ on $L(\beta_2)$ and $+\infty$ on $L(\beta_1)$; such a solution u will be called a solution of the problem \mathcal{P} .

The first thing we have to do to understand a solution u of the problem \mathcal{P} is being able to bound the function u on each radius $L(\beta)$. Our arguments are based on the comparison with the Scherk surface.

Let us consider ABC an isosceles triangle ($|AB| = |AC| = R$). We consider the solution w of the Dirichlet problem on ABC such that $w = 0$ on $[A, B]$ and $[A, C]$ and tends to $+\infty$ on $[B, C]$. This function exists by Theorem 1.5. When ABC is right-angled w is the Scherk surface, after dilatation, w is given by

$$w(x, y) = h(x, y) = -\ln \cos x + \ln \cos y. \quad (2.1)$$

In the general case, the solution w will be called a *pseudo-Scherk surface*.

We shall use the Scherk surface to control solutions of the problem \mathcal{P} . We first consider the case where ABC is right-angled. In fact, a neighbourhood of B in ABC can be isometrically parametrized by $\Omega_{-\pi/4}^0(R)$ and h is a solution of (MSE) on $\Omega_{-\pi/4}^0(R)$ such that $h = 0$ on $L(0)$, $+\infty$ on $L(-\frac{1}{4}\pi)$ and some positive function on the third part of the

boundary. Since we have an expression for h , we can see that h is uniformly bounded on $\Omega_\alpha^0(R) \subset \Omega_{-\pi/4}^0(R)$ for every $-\frac{1}{4}\pi < \alpha < 0$.

We do not suppose now that ABC is right-angled, but we suppose that the angle at the vertex A is greater than $\frac{1}{2}\pi$. In this case we can choose a point A' such that $A'BC$ is isosceles and right-angled and $A'BC$ contains ABC . We consider in ABC the pseudo-Scherk surface w and h the Scherk surface on $A'BC$. Since h is positive in $A'BC$, we have $h > w$. As above, a neighbourhood of B in ABC can be isometrically parametrized by $\Omega_\beta^0(R)$ with $\beta < 0$ and w can be seen as the solution of (MSE) on $\Omega_\beta^0(R)$ such that $w = 0$ on $L(0)$, $+\infty$ on $L(\beta)$ and some positive function on the third part of the boundary. Since $w < h$, w is uniformly bounded on $\Omega_\alpha^0(R)$ for every $\beta < \alpha < 0$.

By our expression for h , there exists $m \in \mathbb{R}$ such that $h \leq m$ on $[A, B]$ and $[A, C]$. This proves that $h - m \leq w$ in ABC . Then in our parametrization of a neighbourhood of B , for every $M \in \mathbb{R}$ there exist α such that $w \geq M$ in $\Omega_\beta^\alpha(R)$.

Lemma 2.1. *Let $\beta_1 < \beta_2$ and $R > 0$. We consider a solution u of the problem \mathcal{P} on $\Omega_{\beta_1}^{\beta_2}(R)$. Then for every $\beta_1 < \alpha < \beta_2$, there exist M and M' in \mathbb{R} such that $u \leq M$ in $\Omega_{\beta_1}^{\beta_2}(\frac{1}{4}R)$ and $u \geq M'$ on $\Omega_{\beta_1}^{\beta_2}(\frac{1}{4}R)$. For every $M \in \mathbb{R}$, there exist α and α' in $]\beta_1, \beta_2[$ such that $u \geq M$ in $\Omega_{\beta_1}^\alpha(\frac{1}{4}R)$ and $u \leq M$ in $\Omega_{\alpha'}^{\beta_2}(\frac{1}{4}R)$.*

Proof. Let us consider $\alpha_1 < \alpha_2$ and $R' > 0$. We consider v the solution of the problem \mathcal{P} on $\Omega_{\alpha_1}^{\alpha_2}(R')$ such that $v = 0$ on the third part of the boundary. v exists because the hypotheses of Theorem 1.5 are fulfilled. The isometry of $\Omega_{\alpha_1}^{\alpha_2}(R') \times \mathbb{R}$ defined by $(r, \theta, z) \mapsto (r, \alpha_1 + \alpha_2 - \theta, -z)$ does not change the boundary data so v is invariant by this isometry because of the uniqueness of such a solution. This proves that $v = 0$ on $L(\frac{1}{2}(\alpha_1 + \alpha_2))$. Then, by the maximum principle, we have $v > 0$ between $L(\alpha_1)$ and $L(\frac{1}{2}(\alpha_1 + \alpha_2))$ and $v < 0$ between $L(\frac{1}{2}(\alpha_1 + \alpha_2))$ and $L(\alpha_2)$.

Let us consider $\alpha_1 < \alpha < \alpha_2$. Let us prove that there exists a constant M such that $v \leq M$ in $\Omega_\alpha^{\alpha_2}(\frac{1}{2}R')$. If $\alpha \geq \frac{1}{2}(\alpha_1 + \alpha_2)$, $M = 0$ works. We write $\tilde{\alpha} = \frac{1}{2}(\alpha_1 + \alpha_2)$. We suppose that $\alpha \leq \tilde{\alpha}$ then we take a sufficiently big n such that $(\tilde{\alpha} - \alpha)/n \leq \frac{1}{4}\pi$ and $(\tilde{\alpha} - \alpha)/n \leq \alpha - \alpha_1$. We denote by B the vertex of $\Omega_{\alpha_1}^{\alpha_2}(R')$. For $k \leq 2n + 1$ we write $\alpha(k) = \tilde{\alpha} - k((\tilde{\alpha} - \alpha)/2n)$ and for $k \leq 2n - 1$ we denote by A_k the points of coordinates $(\frac{1}{2}R', \alpha(k))$ and C_k the point of second coordinate $\alpha(k + 2)$ such that A_kBC_k is an isosceles triangle at A_k (see Figure 1). We have $v = 0$ on $[B, A_0]$ and v is bounded on $[A_0, C_0]$. Then, by adding a constant, we can put a pseudo-Scherk surface above v over A_0BC_0 . This proves that v is upper-bounded in $\Omega_{\alpha(1)}^{\alpha_2}(\frac{1}{2}R')$. Since v is upper-bounded on $[B, A_1]$ and $[A_1, C_1]$, we can put a pseudo-Scherk surface above v over A_1BC_1 then v is bounded on $\Omega_{\alpha(2)}^{\alpha_2}(\frac{1}{2}R')$. We can do this for every k . Then we obtain that v is uniformly upper-bounded on $\Omega_\alpha^{\alpha_2}(\frac{1}{2}R')$.

With the same method, we can prove that there exists M' such that $v \geq M'$ on $\Omega_{\alpha_1}^\alpha(\frac{1}{2}R')$.

Let us now consider our original problem. We have u and α and we want to prove the existence of M . We consider $\beta_1 < \alpha' < \alpha$. Since u tends to $-\infty$ along $L(\beta_2)$, there exists m such that $u \leq m$ at all the points of coordinates $(\frac{1}{2}R, \theta)$ with $\alpha' \leq \theta < \beta_2$. We consider on $\Omega_{\alpha'}^{\beta_2}(\frac{1}{2}R)$ the solution v that we have studied above. By the maximum principle, we




Figure 1.

have $u \leq v + m$ on $\Omega_{\alpha'}^{\beta_2}(\frac{1}{2}R)$. We then have the existence of M because of the result on v . We construct M' in the same way.

Let us now consider u a solution of \mathcal{P} and $M \in \mathbb{R}$. We consider $\beta_1 < \beta < \beta_2$ such that $\beta - \beta_1 \leq \frac{1}{4}\pi$, we consider the point A of coordinates $(\frac{1}{2}R, \beta)$ and the point C on $L(\beta_1)$ such that ABC is an isosceles triangle (where B is the vertex of $\Omega_{\beta_1}^{\beta_2}$). By what we have just proved, u is lower-bounded on $[B, A]$ and $[A, C]$. Then we can put a pseudo-Scherk surface under u . The existence of α is due to the last remark that we made about pseudo-Scherk surfaces. □

Using this result, we can prove the following geometrical result.

Theorem 2.2. *Let (Ω, φ) be a multi-domain and P a vertex of Ω such that two edges L_1 and L_2 have P as endpoint (L_1 and L_2 are enumerated with respect to the orientation). Let u be a solution of (MSE) on Ω such that u tends to $-\infty$ on L_1 and $+\infty$ on L_2 . We consider Ψ_u the conjugate function to u normalized such that $\Psi_u(P) = 0$. Then, if Ψ_u is non-negative in a neighbourhood of P , the vertical straight line passing through $\varphi(P)$ is the boundary of the graph of u above a neighbourhood of P .*

First, we remark that, if Q is a point on L_1 or L_2 , then $\Psi_u(Q) = |PQ| \geq 0$ by Lemma 1.2. This proves that, if the angle at P is strictly less than π , the hypothesis on Ψ_u is always verified. So we have the result for a convex corner.

Proof. By a translation and a rotation, we can isometrically parametrize a neighbourhood of P by $\Omega_\beta^0(R)$ for $\beta < 0$ and R small enough. Then u can be seen as a solution of the problem \mathcal{P} . We suppose that $\Psi_u \geq 0$ in $\Omega_\beta^0(R)$.

Part 1. First, we prove that there exists $M_1 \in \mathbb{R}$ such that $\varphi(P) \times]-\infty, M_1[$ is a part of the boundary of the graph. We take $-\frac{1}{2}\pi < \alpha < 0$. We suppose that $\alpha > \frac{1}{2}\beta$. Then $\Omega_\alpha^0(R) \subset \Omega_\beta^0(R)$ can be parametrized by Euclidean parameters (x, y) . In fact, $\Omega_\alpha^0(R)$ is embedded in \mathbb{R}^2 . The idea is to see the part of the graph which is over $\Omega_\alpha^0(R)$ as a graph over the vertical plane given by the equation $y = 0$. Let $R' < \frac{1}{2}R$. Then for all $Q \in \Omega_\alpha^0(R')$ the nearest point from Q on $\partial\Omega$ is on L_1 . If we take R' small enough and α such that $\tan \alpha > -\frac{1}{8}$, then every point of $\Omega_\alpha^0(R')$ verifies the hypothesis of Lemma 1 in [7]. This lemma implies that, at every point of $\Omega_\alpha^0(R')$, $q = \partial u / \partial y < 0$. Using our Euclidean parameters, we write, for $(x, y) \in \Omega_\alpha^0(R')$, $\Theta(x, y) = (x, u(x, y))$. We have

$$d\Theta = \begin{pmatrix} 1 & 0 \\ \partial u / \partial x & \partial u / \partial y \end{pmatrix}.$$

Since $q < 0$, this proves that Θ is a local diffeomorphism. Since u strictly decreases when y increases, Θ is injective. By Lemma 2.1, we know that there exists $K \in \mathbb{R}$ such that $u \geq K$ on $L(\alpha)$. We put $x_1 = R' \cos \alpha$. We then have $]0, x_1[\times]-\infty, K[\subset \Theta(\Omega_\alpha^0(R'))$. We note that $\chi = \Theta^{-1}$ on $]0, x_1[\times]-\infty, K[$. We then have $y = \chi_{(2)}(x, z)$ on the graph of u (we denote by $\chi_{(2)}$ the second coordinate function of χ). Then $\chi_{(2)}$ verifies (MSE). When $x \rightarrow 0$, we have $y = \chi_{(2)}(x, z) \rightarrow 0$, which is due to the shape of $\Omega_\alpha^0(R')$. From Lemma 2.1, there exist $\alpha < \alpha' < 0$ and r such that $\Omega_{\alpha'}^0(r) \subset \text{Im } \chi$. By results of boundary regularity, $\chi_{(2)}$ is regular at the boundary. Actually, we can extend $\chi_{(2)}$ by making a reflection with respect to the axis $\{x = 0, y = 0\}$. We now show that a part of this axis is a part of the boundary of the whole graph. By Lemma 2.1, there exists M' such that $u \geq M'$ in $\Omega_{\beta'}^0(r)$. We note that $M_1 = M' - 1 < K$. Then if a sequence of points of the graph of u over $\Omega_{\beta'}^0(r)$ tends to a point of $\varphi(P) \times]-\infty, M_1[$, we have (x, y) in $\text{Im } \chi$ after a certain rank. Then the graph of u over $\text{Im } \chi$ is a neighbourhood of $\varphi(P) \times]-\infty, M_1[$. As $\chi_{(2)}$ is regular through the boundary, $\varphi(P) \times]-\infty, M_1[$ is a part of the boundary.

With the same arguments, we can show that there exists M_2 such that $\varphi(P) \times]M_2, +\infty[$ is a part of the boundary.

Part 2. The first part proves that outside a compact the graph of u has a good behaviour above the point $\varphi(P)$. Now we prove that we can extend, by reflection, this compact part through the vertical straight line passing by $\varphi(P)$.

From what we have just done, there exist $\beta < \alpha_2 < \alpha_1 < 0$ such that the graph above $\Omega_{\beta}^{\alpha_2}(R)$ and $\Omega_{\alpha_1}^0(R)$ is regular above P . We choose M_1 and M_2 as in the first part such that

$$(\varphi(P), M_1) \in \partial \text{Graph}(u|_{\Omega_{\alpha_1}^0(R)}) \quad \text{and} \quad (\varphi(P), M_2) \in \partial \text{Graph}(u|_{\Omega_{\beta}^{\alpha_2}(R)}).$$

We construct a curve Γ as follows. We start from the point $A_1 = (0, 0, M_1)$ ($(0, 0) = \varphi(P)$), we go down vertically to the point $A_2 = (0, 0, M_1 - 1)$, then we go to some point




Figure 2.

$A_3 = (\varepsilon \cos \theta, \varepsilon \sin \theta, M_1 - 1)$, following the level curve $\{u = M_1 - 1\}$ (we suppose that ε is small and that $\theta > \alpha_1$). We then follow the curve

$$t \mapsto (\varepsilon \cos t, \varepsilon \sin t, u(\varepsilon \cos t, \varepsilon \sin t)),$$

we let t decrease to some $\theta' < \alpha_2$ such that $u(\varepsilon \cos \theta', \varepsilon \sin \theta') = M_2 + 1$ (we denote by A_4 the endpoint). Following the level curve $\{u = M_2 + 1\}$, we go to the point $A_5 = (0, 0, M_2 + 1)$ and finally we go down to the point $A_6 = (0, 0, M_2)$. We can smooth Γ at the points A_2, A_3, A_4 and A_5 such that A_2 and A_5 are always in the smooth Γ and the new Γ is embedded in the graph of u . The vertical projection of Γ on $\Omega_\beta^0(R)$ bounds a domain $\tilde{\Omega}$. We denote by Σ the graph of u above $\tilde{\Omega}$. Because of our choice of Γ , Σ extends in a minimal surface Σ' through Γ . (The only problem is through $[A_1, A_2]$ and $[A_5, A_6]$, but the first part says that we can extend Σ through these two segments by symmetry.) Because Σ is a graph, Σ is simply connected and its boundary is not empty; the same is true for Σ' . This remark says us that we have conformal parametrization $h_1 : D \rightarrow \Sigma'$ and $h_2 : D \rightarrow \Sigma$ (D is the unit disc). We put $\tilde{D} = h_1^{-1}(\Sigma)$ and $\tilde{h} : \tilde{D} \rightarrow D$ defined by $\tilde{h} = h_2^{-1} \circ h_1$; \tilde{h} is a biholomorphic map. As $h_1^{-1}(\Gamma)$ is embedded in D , the property of Schönflies is verified at every point. By Carathéodory's Theorem, \tilde{h} extends to an homeomorphism of $\tilde{D} \cup h_1^{-1}(\Gamma)$ into $D \cup V$ where V is part of the boundary of D (for all of this argument we refer to Appendix A). This proves that we can extend h_2 in an homeomorphism of $D \cup V$ into $\Sigma \cup \Gamma$. Let us consider $f : D \rightarrow D^-$ (where $D^- = \{(x, y) \in D \mid y < 0\}$) a biholomorphic map. Then f extends to the boundary. Let us consider the following points on Γ : $A_{1.5} = (0, 0, M_1 - 0.5)$ and $A_{5.5} = (0, 0, M_2 + 0.5)$. We write $X = h_2 \circ H \circ f^{-1}$, where H is a Möbius transformation of the unit disc. We note that $B_i = X^{-1}(A_i)$ for every i . Then, for a suitable choice of H , we can have the situation described by Figure 2.

Let us show that X extends to the whole disc. We shall denote by x_1, x_2 and x_3 the three coordinates of X . These three functions are harmonic since Σ is minimal. First we observe that x_1 and x_2 tend to 0 when $z \in D^-$ tends to $D^0 = \{z \in D \mid z \in \mathbb{R}\}$; this is due to the shape of $\tilde{\Omega}$. Then, by the Schwarz reflection principle, x_1 and x_2 extend to D in harmonic functions. Let us consider x_3^* the harmonic conjugate to x_3 on D^- ; we normalized x_3^* by $x_3^*(B_2) = 0$. By our choice of normalization, for every $z \in D^-$, we have $\Psi_u(X(z)) = x_3^*(z)$. This proves that x_3^* tends to 0 when z tends to D^0 . We can extend

x_3^* by reflection to D . By taking the conjugate function, we have proved that x_3 extends to D . We then have constructed a minimal immersion X on D ; then Σ extends through $[A_1, A_6]$. This extension is given by the reflection with respect to $[A_1, A_6]$.

Part 3. The last thing we have to show is that the minimal immersion X has no branch point. If it has a branch point then it must be on D^0 , since, on the other part, the surface is a graph and then there is no branch point. Let z_0 be a branch point, then $\nabla x_3^*(z_0) = 0$. Since x_3^* is harmonic, its local behaviour is quite similar to the one of $\operatorname{Re}(z - z_0)^p$ with $p \geq 2$ (in fact, in some holomorphic charts, we have $x_3^*(z) = \operatorname{Re}(z - z_0)^p$). This implies that there exists z in D^- such that $x_3^*(z) < 0$, but this contradicts our hypothesis $\Psi_u \geq 0$.

We then have proved that there is no branch point, so the vertical straight line passing by $\varphi(P)$ is the boundary of the graph. \square

Remark 2.3. We can remark that in the first two parts we do not use the hypothesis on Ψ_u . So in such a situation we can always extend the graph by making a reflection with respect to the vertical axis. But what we obtain is a minimal surface with, maybe, a finite number of branch points on the vertical axis.

Remark 2.4. We can make another remark. If we consider a vertex P , two edges L_1 and L_2 having P as endpoint and u such that u assumes the data $+\infty$ (or $-\infty$) on L_1 and L_2 , the hypothesis on Ψ_u did not make any more sense and the angle at the vertex P must be greater than π . But we can always apply the two first parts of the proof. The only problem is that we need a result similar to Lemma 2.1; this is given by Theorem 10.3 in [11]. So we can affirm that on the boundary of the graph of u we have a half straight line with a finite number of branch points. Obviously, we must have a branch point at the endpoint of the half straight line.

3. Convergence and divergence of sequence of solutions of (MSE)

In this section we shall explain what we can say about the convergence of a sequence (u_n) of solutions of (MSE): can we make converge a subsequence by some compactness result? What are the different ways of divergence? In [7], it is shown that for a monotone sequence, it appears lines which separate domains of convergence and domains of divergence (this works only for subsequence). We shall show that such lines always appear (Theorem 3.3).

First, we have to determine the domain on which we can make converge a sequence. Since each surface is a graph, if we want the limit to be a graph, the normal to the surface needs to stay close by the vertical unit vector and then W_n have to be bounded. We have the following lemma.

Lemma 3.1. *Let Ω be a domain and (u_n) a sequence of solutions of (MSE) on Ω . Let $P \in \Omega$. We suppose that $W_n(P)$ is bounded by a constant M . Then there exists $R > 0$ which depends only of M and the distance of P to $\partial\Omega$ such that W_n is bounded by $2M$ on the disc of centre P and radius R .*

Proof. We fix an index n . We know (see [9]) that there exists a constant c such that if u is a solution of (MSE) on the disc $\{(x, y) \mid x^2 + y^2 < R^2\}$ we have

$$r^2(0) + 2s^2(0) + t^2(0) \leq \frac{c}{R^2}W^4(0). \tag{3.1}$$

Let R be such that $2R = d(P, \partial\Omega)$. Then, for all Q in $D(P, R)$, the above equation gives $r^2(Q) + 2s^2(Q) + t^2(Q) \leq (c/R^2)W^4(Q)$. We have $\nabla W = ((rp + sq)/W, (sp + tq)/W)$, so, in $D(P, R)$, we have $\|\nabla W\| \leq \tilde{C}W^2$ with \tilde{C} which depends only of R . Let z be the function such that $z(0) = M$ and $z' = \tilde{C}z^2$, z is defined on $[0, 1/(M\tilde{C})[$ by

$$\frac{1}{M} - \frac{1}{z} = \tilde{C}r. \tag{3.2}$$

Because of our estimate on $\|\nabla W\|$, we have, in polar coordinates, $W(r, \theta) \leq z(r)$. Then W is bounded by $2M$ on $D(P, \min(R, 1/(2M\tilde{C})))$. \square

Let (u_n) be a sequence of solutions of (MSE) on a domain Ω . We then define $\mathcal{B}(u_n)$ as the set of the point $Q \in \Omega$ such that $(W_n(Q))$ is bounded. Lemma 3.1 says us that $\mathcal{B}(u_n)$ is an open set and that W_n is uniformly bounded on each compact included in $\mathcal{B}(u_n)$. Then if D is a connected component of $\mathcal{B}(u_n)$ and $P \in D$ there exists an extraction θ such that $u_{\theta(n)} - u_{\theta(n)}(P)$ converges uniformly on each compact of D to a solution u of (MSE). Here, we use some classical compactness results (see [9]). This proves that the divergence of the sequence is due to the behaviour of the sequence over $\Omega \setminus \mathcal{B}(u_n)$.

If $P \in \Omega \setminus \mathcal{B}(u_n)$, there exists a subsequence $u_{n'}$ such that $W_{n'}(P) \rightarrow +\infty$. As the normal N_n to the graph at $(P, u_n(P))$ is given by

$$N_n(P) = \left(\frac{p_n}{W_n}(P), \frac{q_n}{W_n}(P), -\frac{1}{W_n}(P) \right), \tag{3.3}$$

we can suppose that $N_{n'}(P)$ converges to an horizontal unit vector. The following theorem describes what happens locally.

Theorem 3.2. *Let r be positive. Let (u_n) be a sequence of solutions of (MSE) on the disc $D(0, r)$. We suppose that $N_n(0)$ converges to $(1, 0, 0)$. Let $\alpha \in]0, 1[$. Then there exists an extraction θ such that $N_{\theta(n)}$ converges to $(1, 0, 0)$ at almost every point of $\{0\} \times [-\alpha r, \alpha r]$.*

Proof. Let $n \in \mathbb{N}$. We know (see [7, 11]) that there exists $\Phi_n : (x, y) \mapsto (\xi, \eta)$ with $\Phi_n(0, 0) = (0, 0)$ and

$$d\xi = \left(1 + \frac{1 + p_n^2}{W_n} \right) dx + \frac{p_n q_n}{W_n} dy, \tag{3.4}$$

$$d\eta = \frac{p_n q_n}{W_n} dx + \left(1 + \frac{1 + q_n^2}{W_m} \right) dy. \tag{3.5}$$

We know that Φ_n increases distance so it is bijective on its image. This image contains the disc of centre $(0, 0)$ and radius r . Besides, we know that (ξ, η) are conformal parameters for the graph of u_n . On the $\xi\eta$ disc $D(0, r)$ we then have the Gauss map $g_n(\xi + i\eta)$

which corresponds to the stereographic projection of N_n ; g_n is holomorphic. We write $z_n = g_n(0)$. By hypothesis, we have $z_n \rightarrow 1$. We write $z = \xi + i\eta$. By our choice of normal $g_n : D(0, r) \rightarrow D(0, 1)$, there exists $h_n : D(0, r) \rightarrow D(0, 1)$ holomorphic with $h_n(0) = 0$ such that

$$g_n(z) = \frac{h_n(z) + z_n}{1 + \bar{z}_n h_n(z)}. \tag{3.6}$$

Since $z_n \rightarrow 1$, the sequence of holomorphic functions $z \mapsto (z + z_n)/(1 + \bar{z}_n z)$ converges simply to 1 on $D(0, 1)$ and uniformly on the disc $D(0, \alpha)$ for all $\alpha < 1$. But, by the Schwarz Lemma, we have, for all $n \in \mathbb{N}$, $h_n(D(0, \alpha r)) \subset D(0, \alpha)$. We then have uniform convergence of g_n to 1 on $D(0, \alpha r)$. Using (3.3), this proves that for every ε , if n is big enough, we can say that $p_n/W_n \geq 1 - \varepsilon$ and $|q_n|/W_n \leq \varepsilon$ in $\Phi_n^{-1}(D(0, \alpha r))$. So, to conclude, we need to understand the shape of $\Phi_n^{-1}(D(0, \alpha r))$. We shall see that these sets are concentrating along the segment $\{0\} \times [-\alpha r, \alpha r]$.

For all $n \in \mathbb{N}$, we consider, in the $\xi\eta$ disc, the path $\gamma_n :]-r, r[\rightarrow D(0, r)$ defined by $\gamma_n(0) = 0$ and $\gamma'_n = \nabla y_n / \|\nabla y_n\|$ where y_n is the second coordinate of Φ_n^{-1} . We have (see [11])

$$\nabla y_n = \left(-\frac{p_n q_n}{J_n W_n}, \frac{W_n + 1 + p_n^2}{J_n W_n} \right), \tag{3.7}$$

where $J_n = \det(d\Phi_n) = W_n + 2 + 1/W_n$. Because (3.3), ∇y_n converges uniformly to $(0, 1)$ on the disc $D(0, \alpha r)$ for all $\alpha < 1$.

Let $\alpha \in]0, 1[$. We denote by A_α and B_α the points in the xy disc $D(0, r)$ of respective coordinates $(0, -\alpha r)$ and $(0, \alpha r)$. In the following, we prove that $\int_{[A_\alpha, B_\alpha]} d\Psi_{u_n} \rightarrow 2\alpha r$.

Let $\tilde{\alpha} > \alpha$. Then for n big enough, we have $\|\nabla y_n\| > \alpha/\tilde{\alpha}$ in $D(0, \tilde{\alpha}r)$. Because, for $t \in [-\tilde{\alpha}r, \tilde{\alpha}r]$, $\gamma_n(t) \in D(0, \tilde{\alpha}r)$, there exists $\tilde{\alpha}r \leq t_0^n < t_1^n \leq \tilde{\alpha}r$ such that $y_n(\gamma_n(t_0^n)) = -\alpha r$ and $y_n(\gamma_n(t_1^n)) = \alpha r$. Along $[t_0^n, t_1^n]$, $y_n \circ \gamma_n$ increases strictly from $-\alpha r$ to αr . Then the path $\Gamma_n = \Phi_n^{-1} \circ \gamma_n$ on $[t_0^n, t_1^n]$ can be parametrized by $y \in [-\alpha r, \alpha r]$. We have a function f_n on $[-\alpha r, \alpha r]$ such that, for $t \in [t_0^n, t_1^n]$, $x_n(\gamma_n(t)) = f_n(y_n(\gamma_n(t)))$. We have

$$\begin{aligned} |x_n \circ \gamma_n(t)| &\leq \left| \int_0^t |(x_n \circ \gamma_n)'(u)| \, du \right| \\ &\leq \left| \int_0^t \|\nabla x_n\|(\gamma_n(u)) \, du \right| \\ &= \left| \int_0^t \left(\frac{1 + q_n^2}{(1 + W_n)^2} \right)^{1/2} (\Gamma_n(u)) \, du \right| \end{aligned} \tag{3.8}$$

(for the last equality, see [11]). Then for n big enough $|x_n \circ \gamma_n|$ on $[t_0^n, t_1^n]$ can be bounded by a constant uniformly small; this is due to the fact that $|q_n|/W_n \leq \varepsilon$ for big n . We then have proved that the path Γ_n is close by the segment $[A_\alpha, B_\alpha]$ for big n .

Let us now calculate $\int_{\Gamma_n} d\Psi_{u_n}$. We have

$$\int_{\Gamma_n} \frac{p_n}{W_n} \, dy - \frac{q_n}{W_n} \, dx = \int_{-\alpha r}^{\alpha r} \left(\frac{p_n}{W_n} (f_n(y), y) - \frac{q_n}{W_n} (f_n(y), y) f'_n(y) \right) \, dy. \tag{3.9}$$

We have $|f'_n(y_n(\gamma_n(t)))| \leq (\|\nabla x_n\|/\|\nabla y_n\|)(\gamma_n(t)) \rightarrow 0$, the convergence is uniform so f'_n tends uniformly to 0 on $[-\alpha r, \alpha r]$. This proves that $\int_{\Gamma_n} d\Psi_{u_n} \rightarrow 2\alpha r$.

Let us consider the path $\tilde{\Gamma}_n$ which consists of the segment $[A_\alpha, B_\alpha]$, then the segment $[B_\alpha, \Gamma(t_1^n)]$, then $-\Gamma_n$, then, finally, the segment $[\Gamma_n(t_0^n), A_\alpha]$. Let $\varepsilon > 0$. For n big enough, we can suppose that $\int_{\Gamma_n} d\Psi_{u_n} \geq 2\alpha r - \varepsilon$ and $((1 + q_n^2)/((1 + W_n)^2))^{1/2} \leq \varepsilon$ on $\Phi_n^{-1}(D(0, \tilde{\alpha}r))$. As $d\Psi_{u_n}$ is closed, $\int_{\tilde{\Gamma}_n} d\Psi_{u_n} = 0$. We then have

$$\begin{aligned} 2\alpha r &\geq \int_{[A_\alpha, B_\alpha]} d\Psi_{u_n} \\ &= - \int_{[B_\alpha, \Gamma(t_1^n)]} d\Psi_{u_n} - \int_{-\Gamma_n} d\Psi_{u_n} - \int_{[\Gamma_n(t_0^n), A_\alpha]} d\Psi_{u_n} \\ &\geq \int_{\Gamma_n} d\Psi_{u_n} - |x_n(\gamma(t_1^n))| - |x_n(\gamma(t_0^n))| \\ &\geq 2\alpha r - \varepsilon - \varepsilon|t_1^n| - \varepsilon|t_0^n| \\ &\geq 2\alpha r - \varepsilon(1 + 2\tilde{\alpha}r). \end{aligned} \tag{3.10}$$

This proves that $\int_{[A_\alpha, B_\alpha]} d\Psi_{u_n} \rightarrow 2\alpha r$. We have

$$\int_{[A_\alpha, B_\alpha]} d\Psi_{u_n} = \int_{[A_\alpha, B_\alpha]} (p_n/W_n) dy.$$

Because $p_n/W_n \leq 1$, the preceding equalities prove that p_n/W_n converges to 1 in $\mathbb{L}^1([A_\alpha, B_\alpha])$. Then there exists an extraction θ such that $p_{\theta(n)}/W_{\theta(n)}$ converges simply to 1 at almost every point in $[A_\alpha, B_\alpha]$. Thus the proposition is proved. \square

This proposition gives us a local result and we have the following global result.

Theorem 3.3. *Let (Ω, φ) be a multi-domain. Let (u_n) be a sequence of solutions of (MSE) on Ω . Let $P \in \Omega$ and N a unit horizontal vector and D the geodesic of Ω passing at P and normal to N . If the sequence $(N_n(P))$ converges to N , then $N_n(Q)$ converges to N at every point of D .*

As Ω is locally isometric to \mathbb{R}^2 , we have allowed us to call N the parallel transport of N along D .

Proof. We first get a parametrization of D by arclength with P as origin point. Then D is parametrized by $]a, b[,]-\infty, b[,]a, +\infty[$ or $] -\infty, +\infty[$; we shall suppose that we are in the case $]a, +\infty[$ (the other cases are similar). We then consider the set \mathcal{F} of $t \in \mathbb{R}_+^*$ such that, if θ_1 is an extraction, there exists a sub-extraction θ_2 such that $N_{\theta_2(n)}(Q)$ converge to N at almost every Q of the part of D parametrized by $]a + |a|/(t + 1), t[$ ($a < 0$). Let us prove that $\mathcal{F} = \mathbb{R}_+^*$. First, we observe that, if $t_1 \in \mathcal{F}$ and $t_2 < t_1$, then $t_2 \in \mathcal{F}$. From Theorem 3.2, there exists $t > 0$ such that $t \in \mathcal{F}$. Let $t_0 = \sup \mathcal{F}$ and suppose that $t_0 < +\infty$. We consider P_1 and P_2 the points on D parametrized by $a + |a|/(1 + t_0)$ and t_0 . We choose $R > 0$ such that $D(P_i, R) \subset \Omega$ for $i = 1, 2$. Let θ_1 be an extraction. Since $t_0 = \sup \mathcal{F}$, there exist $Q_1 \in D(P_1, \frac{1}{3}R) \cap D$, $Q_2 \in D(P_2, \frac{1}{3}R) \cap D$ and a sub-extraction θ_2 such that $N_{\theta_2(n)}$ converges to N at Q_1 and Q_2 . We have $D(Q_i, \frac{2}{3}R) \subset \Omega$. We then

apply Theorem 3.2 to points Q_1 and Q_2 with $\alpha = \frac{3}{4}$. We then have a sub-extraction θ_3 such that N_{θ_3} converges to N at almost every point of $D(Q_i, \frac{1}{2}R) \cap D$ for $i = 1, 2$. This proves that t_0 is not $\sup \mathcal{F}$, because N_{θ_3} converges to N at almost every point of the part of D parametrized by an open interval that contains the segment $[a + |a|/(t_0 + 1), t_0]$.

By a standard diagonal process, we can then construct an extraction θ such that $N_{\theta(n)}$ converges to N at almost every point of D . Let Q be in D and we consider N' a cluster point of the sequence $N_{\theta(n)}(Q)$. If the third coordinate of N' is negative, then there exists a sub-extraction θ' such that $W_{\theta'(n)}(Q)$ is bounded, but this is impossible since, by Lemma 3.1, $W_{\theta'(n)}$ would be bounded in a neighbourhood of Q and $W_{\theta(n)}$ diverges at almost every point of D . Thus the third coordinate of N' is 0. If $N' \neq N$, applying what we have already proved, it appears a second geodesic D' passing by Q normal to N' and an extraction θ' such that $N_{\theta'(n)}$ converges to N' at almost every point of D' . We parametrized D and D' by arclength in using the orientation given by the direct normal to N and N' . We choose Q as the origin point. Let $\varepsilon > 0$. We denote by A the point on D of coordinate $-\varepsilon$ and by B the point on D' of coordinate ε . For ε small enough, the triangle AQB is in Ω and then $\int_{AQB} d\Psi_{u_{\theta'(n)}} = 0$. We let n tend to $+\infty$ and then obtain $|AC| + |BC| \leq |AB|$ which contradicts the triangle inequality. We then have proved that $N_{\theta(n)}$ converges to N at every point of D . We then have proved that for every extraction θ we can construct a sub-extraction θ' such that $N_{\theta'(n)}$ converges to N at every point of D .

To finish the proof, we take a point Q in D and suppose that $N_{\alpha(n)}(Q)$ converge to N' with α an extraction. Since $N_{\alpha(n)}(P) \rightarrow N$, there exists a sub-extraction α' such that $N_{\alpha'(n)}$ converges to N at every point of D , in particular at Q . Then $N = N'$. \square

Remark 3.4. We then understand the structure of the complementary of $\mathcal{B}(u_n)$; it is a set of geodesics of Ω . One of these geodesics will be called a *line of divergence*. Then when we have a sequence of solutions of (MSE), the problem of the convergence of the sequence is linked to the understanding of what lines of divergence are possible? The answer is, in general, given by the behaviour at the boundary. The existence of such lines will permit us to use arguments that are similar to the ones used by Jenkins and Serrin in [7].

The behaviour of the normal along a line of divergence says that the limit of $\int_T d\Psi_{u_n}$, where T is a segment of a line of divergence with the orientation given by the limit normal, is $|T|$. In the following, we shall draw this limit normal on the figures to explain our arguments.

4. The Plateau problem at infinity

In this section, we explain the problem studied by Cosín and Ros in [2] and give the main results of their paper with some elements of proofs.

Let M be a complete minimal surface with finite total curvature in \mathbb{R}^3 . We know that M is isometric to a compact Riemann surface minus a finite number of points (we can refer to [11]). M then has a finite number of annular ends. When these ends are embedded they are asymptotic either to a half-catenoid or to a plane. A properly immersed minimal surface with r embedded ends will be called a r -noid. We can associate to each end a vector which characterizes the direction and the growth of the asymptotic half-catenoid (when the end is asymptotic to a plane this vector is zero). This vector is called the *flux* of the end (for a precise definition of the flux, see [6]). If v_1, \dots, v_r are the fluxes at each end, we have the following balancing condition:

$$v_1 + \dots + v_r = 0. \quad (4.1)$$

This condition tells us that the total flux of the system vanishes. If v_1, \dots, v_r are vectors in \mathbb{R}^3 such that (4.1) is verified and g is a non-negative integer, the Plateau problem at infinity for these data is to find an r -noid of genus g which has v_1, \dots, v_r as fluxes at its ends.

Let $\psi : M \rightarrow \mathbb{R}^3$ be an r -noid. M is conformally equivalent to a compact surface \bar{M} minus a finite number of points p_1, \dots, p_r . We will say that M is *Alexandrov-embedded* if \bar{M} bounds a compact 3-manifold $\bar{\Omega}$ and the immersion ψ extends to a proper local diffeomorphism $f : \bar{\Omega} \setminus \{p_1, \dots, p_r\} \rightarrow \mathbb{R}^3$. An Alexandrov-embedded surface has a canonical orientation; we choose the Gauss map to be the outward pointing normal. An Alexandrov-embedded r -noid cannot have a planar end (see [2]). We call \mathcal{M}_r the space of Alexandrov-embedded r -noids of genus 0 and r horizontal catenoidal ends. We identify two elements in \mathcal{M}_r which differ by a translation. In [2], Cosín and Ros give a nice description of the space \mathcal{M}_r .

Let $\psi : M \rightarrow \mathbb{R}^3$ be a non-flat immersion of a connected orientable surface M and Π be a plane in \mathbb{R}^3 , normalized to be $\{x_3 = 0\}$. We denote by S the Euclidean symmetry with respect to Π and consider the subsets

$$\begin{aligned} M^+ &= \{p \in M \mid x_3(p) > 0\}, \\ M^- &= \{p \in M \mid x_3(p) < 0\}, \\ M^0 &= \{p \in M \mid x_3(p) = 0\}. \end{aligned}$$

With this notation, we have the following definition.

Definition 4.1. We shall say that M is strongly symmetric with respect to Π if

- (i) there exists an isometric involution $s : M \rightarrow M$ such that $\psi \circ s = S \circ \psi$;
- (ii) $\{p \in M \mid s(p) = p\} = M^0$;
- (iii) the third coordinate N_3 of the Gauss map of M takes positive (respectively, negative) values on M^+ (respectively, M^-).

In [2], Cosín and Ros prove the following result.

Proposition 4.2. *Let M be an r -noid with horizontal ends. Then M is strongly symmetric with respect to an horizontal plane if and only if M is Alexandrov-embedded.*

We then use the notion of strong symmetry to study \mathcal{M}_r . In the following, we always suppose that the plane of strong symmetry is the plane $\{x_3 = 0\}$. If $M \in \mathcal{M}_r$, s extends to $\bar{M} = \bar{\mathbb{C}}$, the involution s is $z \mapsto 1/\bar{z}$ and the points p_1, \dots, p_r are in $\{z \in \mathbb{C} \mid |z| = 1\}$. We then have an order on $\{p_1, \dots, p_r\}$. Let us suppose that p_1, \dots, p_r are put in this order. Let v_1, \dots, v_r be vectors in \mathbb{R}^3 such that $2v_i$ is the flux at the end p_i . We have $v_1 + \dots + v_r = 0$, so if we draw the vectors consecutively in the plane, we get a piecewise linear closed curve; a polygon. We denote this polygon by $F(M)$.

We say that a polygon V bounds an immersed polygonal disc if there exists a compact multi-domain (\mathcal{P}, φ) such that $\partial\mathcal{P}$ is only composed of edges and $\varphi(\partial\mathcal{P}) = V$.

Then the most important result in [2] is the following.

Theorem 4.3. *Let v_1, \dots, v_r be horizontal vectors such that $v_1 + \dots + v_r = 0$ and V the associated polygon. Then there exists $M \in \mathcal{M}_r$ such that $F(M) = V$ if and only if V bounds an immersed polygonal disc.*

Besides, we have as much $M \in \mathcal{M}_r$ such that $F(M) = V$ as immersed polygonal discs bounded by V . Let V be a polygon and (\mathcal{P}, φ) a compact multi-domain such that $\varphi(\mathcal{P})$ is an immersed polygonal disc bounded by V . Let P_1, \dots, P_r be the vertices of \mathcal{P} which are identified with the ones of V . We put $P_1 = P_{r+1}$. Let $i \in \{1, \dots, r\}$. We can glue to \mathcal{P} along $[P_i, P_{i+1}]$ a half-strip S_i isometric to $[P_i, P_{i+1}] \times \mathbb{R}_+$. We get a multi-domain which we call $\Omega(\mathcal{P})$. The boundary of $\Omega(\mathcal{P})$ is composed of $2r$ half straight lines, we call L_i^- (respectively, L_i^+) the half line in the boundary which has P_i as endpoint and is in S_{i-1} (respectively, S_i).

Let M be in \mathcal{M}_r . We consider $(M^+)^*$ the conjugate surface to M^+ for the outward pointing normal. In [2], the authors prove that there exists (\mathcal{P}, φ) , a multi-domain bounded by $F(M)$ such that $(M^+)^*$ is a graph over the multi-domain $\Omega(\mathcal{P})$. The normal to the graph is the upward pointing normal by Definition 4.1. If u is the function on $\Omega(\mathcal{P})$ that gives $(M^+)^*$, they prove that u tends to $+\infty$ (respectively, $-\infty$) on L_i^+ (respectively, L_i^-). Cosín and Ros use these arguments to prove that if the Plateau problem at infinity has a solution, the flux polygon $F(M)$ bounds an immersed polygonal disc. For the other implication, they prove that the map $F : M \mapsto F(M)$ is a covering map. To conclude, they use a compactness argument and prove that the space \mathcal{M}_r has a smooth structure.

In the next section, we shall solve on $\Omega(\mathcal{P})$ the Dirichlet problem for the boundary data $+\infty$ on L_i^+ and $-\infty$ on L_i^- . We shall then take the conjugate of the graph of the solution for the downward pointing normal and so build the solution to the Plateau problem at infinity. The change of orientation makes that we get the surface we want.

5. The construction of a solution of the Plateau problem at infinity

The first part of this section will be devoted to the proof of our main result.

Theorem 5.1. *Let V be a polygon which bounds an immersed polygonal disc (\mathcal{P}, φ) . We define $\Omega(\mathcal{P})$ as in the preceding section. Then there exists a solution u of (MSE) on $\Omega(\mathcal{P})$ such that u tends to $+\infty$ on L_i^+ and $-\infty$ on L_i^- . Besides, the solution is unique up to an additive constant.*

Proof. Let us first consider u a solution of (MSE) on the half-strip $[0, a] \times \mathbb{R}_+$ such that u tends to $-\infty$ on $\{a\} \times \mathbb{R}_+^*$ and $+\infty$ on $\{0\} \times \mathbb{R}_+^*$. This situation describes the behaviour in the r half-strips S_i . Then by Lemma 1 in [7] we have

$$\frac{|q|}{W}(x, y) \geq 1 - \frac{a^2}{x^2}, \tag{5.1}$$

$$\frac{|p|}{W}(x, y) \leq \sqrt{2} \frac{a}{x}, \tag{5.2}$$

when $x \geq 4a$. We consider now the general problem.

We begin in proving the uniqueness part of Theorem 5.1. Let u_1 and u_2 be two different solutions of the problem (i.e. $u_1 - u_2$ is non-constant). As in the proof of Theorem 1.5, we can suppose that $\{u_1 > u_2\}$ and $\{u_1 < u_2\}$ are non-empty. Let us call Ω_l the subset of $\Omega(\mathcal{P})$ which is the union of \mathcal{P} and the set of points in each S_i that are at a distance less than l from $[P_i, P_{i+1}]$. We define $\Omega_l^+ = \Omega_l \cap \{u_1 > u_2\}$. Let us consider

$$I = \int_{\partial\Omega_l^+} d\tilde{\Psi}, \tag{5.3}$$

where $d\tilde{\Psi} = d\Psi_{u_1} - d\Psi_{u_2}$. Since $d\tilde{\Psi}$ is closed, we have $I = 0$. $\partial\Omega_l^+$ is composed of a part which is included on $(\bigcup_i L_i^+) \cup (\bigcup_i L_i^-)$ where $d\tilde{\Psi} = 0$, a part included in the interior of $\Omega(\mathcal{P})$, denoted by Γ_l , and a part in $I_{i,l}$ which is the part in S_i parametrized by $[P_i, P_{i+1}] \times \{l\}$. On the part included in $I_{i,l}$ if l is big enough the integral of $d\tilde{\Psi}$ is less than $2\sqrt{2}(|P_i P_{i+1}|^2/l)$ by (5.2). We then have

$$0 = I \leq \int_{\Gamma_l} d\tilde{\Psi} + \sum_{i=1}^r 2\sqrt{2} \frac{|P_i P_{i+1}|^2}{l}. \tag{5.4}$$

By Lemma 2 in [1], $\int_{\Gamma_l} d\tilde{\Psi}$ is negative and decreases as l increases. Because

$$\sum_{i=1}^r \sqrt{2} \frac{|P_i P_{i+1}|^2}{l} \xrightarrow{l \rightarrow +\infty} 0,$$

we get a contradiction. This proves that, if u_1 and u_2 are two solutions of our Dirichlet problem, there exists $c \in \mathbb{R}$ such that $u_1 = u_2 + c$.

We now prove the existence of the solution. We fix a point P_0 in \mathcal{P} . Let us consider in S_i the point Q_i^k which is the middle point of $I_{i,k}$. We then define Ω_k to be the compact subdomain of $\Omega(\mathcal{P})$ bounded by the segments $[P_i, Q_i^k]$ and $[Q_i^k, P_{i+1}]$. Let \mathcal{G}_i^k be the set of the points Q in Ω_k such that $d(Q, Q_i^k) < d(P_i, Q_i^k)$. If k is big enough the sets \mathcal{G}_i^k are disjoint, this proves that the conditions of Theorem 1.5 are fulfilled for big k . Then by Theorem 1.5, we can build a function u_k on Ω_k such that u_k tends




Figure 3.

to $-\infty$ (respectively, $+\infty$) on $[Q_i^k, P_{i+1}]$ (respectively, $[P_i, Q_i^k]$) and $u_k(P_0) = 0$. Following Remark 3.4 in § 3, we shall prove that this sequence (u_k) of solutions of (MSE) has no line of divergence. Then the limit u of (u_k) will be our solution. We shall make discussions that are similar to the ones made by Jenkins and Serrin. We write $d\Psi_{u_k} = d\Psi_k$. We recall that, if T is a segment included in a line of divergence, $|\int_T d\Psi_k|$ converges to the length of T for a subsequence.

Suppose there exists a line of divergence L . We first prove that L can not have an endpoint in the interior of a L_i^+ or a L_i^- . Suppose that L has an endpoint D in L_i^- (the same argument works for L_i^+). Let A be a point in $L \cap S_{i-1}$, we orient L by \overrightarrow{AD} , we suppose that the limit normal along L points on the right-hand side of L . We chose a point B of L_i^- on the right-hand side of D . Because of the triangle inequality, there exists a point C on $[A, D]$ such that $|AC| + |DB| > |CD| + |BA|$. for k big enough we have A and C in Ω_k we then put $D_k = [A, D] \cap [P_i, Q_{i-1}^k]$ and $B_k = [A, B] \cap [P_i, Q_{i-1}^k]$ (see Figure 3).

Let T_k be the triangle $AD_k B_k$ with this orientation. We then have

$$\begin{aligned}
 0 &= \int_{T_k} d\Psi_k \\
 &\geq \int_{[A,C]} d\Psi_k - |CD_k| + |D_k B_k| - |B_k A| \\
 &\geq \int_{[A,C]} d\Psi_k - |CD| + |D_k B_k| - |BA|.
 \end{aligned}
 \tag{5.5}$$

But $|D_k B_k| \rightarrow |DB|$ and $\int_{[A,C]} d\Psi_k \rightarrow |AC|$ for the subsequence that makes L appear; this gives us a contradiction.

We have now only a finite number of possibilities for a line of divergence. If it has an endpoint, it must be one P_i . By construction, we have $\int_{[P_i, P_{i+1}]} d\Psi_k = 0$ (because the integral of $d\Psi_k$ along the triangle $P_i Q_i^k P_{i+1}$ is zero and we know $d\Psi_k$ along $[P_i, Q_i^k]$ and $[Q_i^k, P_{i+1}]$ by Lemma 1.2) so if Γ is a curve joining P_i to P_j we have $\int_\Gamma d\Psi_k = 0$. Then, by passing to the limit, if Γ is a line of divergence, we obtain $|P_i P_j| = 0$ which is not possible. This proves that a line of divergence has at most one endpoint. Suppose that a line of divergence L has no endpoint, we are in the situation of Figure 4. Let




Figure 4.

A and B be point on L as in Figure 4 such that $|AB| > |P_i P_{i+1}|$. We denote by D (respectively, C) the projection of A (respectively, B) on L_{i+1}^- . For k big enough we note that $C_k = [B, C] \cap [P_{i+1}, Q_i^k]$ and $D_k = [A, D] \cap [P_{i+1}, Q_i^k]$. We then have

$$\begin{aligned}
 0 &= \int_{ABC_k D_k} d\Psi_k \\
 &\geq \int_{[A, B]} d\Psi_k - |BC_k| + \int_{[C_k, D_k]} d\Psi_k - |D_k A| \\
 &\geq \int_{[A, B]} d\Psi_k - 2|P_i P_{i+1}| + |C_k D_k|.
 \end{aligned} \tag{5.6}$$

We have $|C_k D_k| \rightarrow |CD| = |AB|$ and $\int_{[A, B]} d\Psi_k \rightarrow |AB|$ for a subsequence, so we get a contradiction and a line of divergence must have one endpoint.

Let L be a line of divergence. We know that we are in the case where L has P_i as endpoint and goes to infinity in one S_j . By what we have done just above, we have only one possibility for the limit normal: we are in the same situation as in the semi-strip S_j in Figure 4. Then, by changing L if necessary, we can suppose that the part of L in S_j is parametrized by $\{A\} \times \mathbb{R}_+$ with $A \in]P_j, P_{j+1}[$ and the domain $\tilde{\Omega}$ parametrized by $[A, P_{j+1}] \times \mathbb{R}_+$ is in $\mathcal{B}(u_k)$. Let θ be an extraction that makes L appear, since $\tilde{\Omega} \subset \mathcal{B}(u_k)$, there exists an extraction θ' such that $u_{\theta'(k)}$ is a subsequence of $u_{\theta(k)}$ and $u_{\theta'(k)} - u_{\theta'(k)}(K)$ (where $K \in \tilde{\Omega}$) converges to v a solution of (MSE) on $\tilde{\Omega}$.

We shall now prove that v tends to $-\infty$ on L_{j+1}^- and $+\infty$ on L . Let $B \in [A, P_{j+1}]$ and C and D be the points which are respectively parametrized by (B, c) and (B, d) ($c < d$). We make use of the following notation.

- (i) E the projection of D on L_{j+1}^- .
- (ii) F the projection of C on L_{j+1}^- .
- (iii) G the projection of D on L .
- (iv) H the projection of C on L .

We also note that, when k is big enough, $E_k = [D, E] \cap [P_{j+1}, Q_j^k]$ and $F_k = [C, F] \cap [P_{j+1}, Q_j^k]$. Because $d\Psi_k$ is closed, we have

$$\left| \int_{[C,D]} d\Psi_{\theta'(k)} - |F_{\theta'(k)}E_{\theta'(k)}| \right| \leq 2|BP_{j+1}|, \tag{5.7}$$

$$\left| \int_{[C,D]} d\Psi_{\theta'(k)} - \int_{[H,G]} d\Psi_{\theta'(k)} \right| \leq 2|BA|. \tag{5.8}$$

Thus, letting k tends to infinity, we obtain

$$\left| \int_{[C,D]} d\Psi_v - |CD| \right| \leq 2|BP_{j+1}|, \tag{5.9}$$

$$\left| \int_{[C,D]} d\Psi_v - |CD| \right| \leq 2|BA|. \tag{5.10}$$

So we can calculate $d\Psi_v$ on L and L_{j+1}^- . We remark that $d\Psi_v$ has the same behaviour as if v assumes the boundary values $+\infty$ on L and $-\infty$ on L_{j+1}^- . We prove that this is, in fact, the case. We consider now two points A_1 and A_2 on $L \cap \tilde{\Omega}$ and two points A_3 and A_4 on L_{j+1}^- . There exists a solution v' of (MSE) on the domain bounded by the polygon $A_1A_2A_3A_4$ such that $v' = v$ on $[A_1, A_4]$ and $[A_2, A_3]$, v' tends to $+\infty$ on $[A_1, A_2]$ and tends to $-\infty$ on $[A_3, A_4]$. Since we know the value of $d\Psi_v$ on $[A_1, A_2]$ and $[A_3, A_4]$, the uniqueness part of the proof of Theorem 1.5 proves that $v = v'$. We have then proved that v tends to $+\infty$ on L and $-\infty$ on L_{j+1}^- .

We shall now get a contradiction to the existence of the line of divergence L . We have

$$\int_{[A,P_{j+1}]} d\Psi_v = \int_{\tilde{\Omega} \cap I_{j,l}} d\Psi_v.$$

Then, by (5.2) and letting l tend to infinity, we get $\int_{[A,P_{j+1}]} d\Psi_v = 0$. If we follow L between P_i and A and the segment $[A, P_{j+1}]$, we get a path joining P_i to P_{j+1} . Then we have

$$0 = \int_{[P_i,A]} d\Psi_{\theta'(k)} + \int_{[A,P_{j+1}]} d\Psi_{\theta'(k)}. \tag{5.11}$$

Let k tend to infinity, we get $0 = |P_iA| + \int_{[A,P_{j+1}]} d\Psi_v = |P_iA|$, but $P_i \notin [P_j, P_{j+1}]$. This is our contradiction.

We then have to prove that $\mathcal{B}(u_k) = \Omega(\mathcal{P})$, as $u_k(P_0) = 0$ for all k there exists a subsequence $u_{k'}$ which converges to a solution u of (MSE). The same arguments that we used just above for v prove that u tends to $+\infty$ (respectively, $-\infty$) on L_i^- (respectively, L_i^+). We have then established Theorem 5.1. \square

We are then able to build the solution to the Plateau problem at infinity. Let V be a polygon and (\mathcal{P}, φ) a polygonal disc bounded by V . We consider the solution u of the Dirichlet problem given by Theorem 5.1. We denote by P_1, \dots, P_r the vertices of V and we consider Ψ_u normalized by $\Psi_u(P_1) = 0$. From the proof above, we have $\Psi_u(P_i) = 0$ for all i . Then on L_i^+ and L_i^- we have $\Psi_u(Q) = |QP_i|$. Since Ψ_u is 1-Lipschitz continuous,

we have $\Psi_u(Q) \geq 0$ for all $Q \in S_i$. Suppose that $\{Q \in \Omega(\mathcal{P}) \mid \Psi_u(Q) \leq 0\}$ is not reduced to $\{P_1, \dots, P_r\}$. Then there exists a point in the interior of \mathcal{P} such that Ψ_u is minimal at this point. But Ψ_u corresponds to x_3^* , the third coordinate on M^* the conjugate surface to the graph of u . Since x_3^* is harmonic on M^* , it can not have a minimum in the interior of M^* . We then have proved that $\Psi_u > 0$ in the interior of $\Omega(\mathcal{P})$.

By Theorem 2.2, the boundary of the graph of u is composed of the r vertical lines over the points $\varphi(P_i)$. Let M be the graph of u with these r vertical lines. We consider M^* the conjugate surface to M . The boundary of M^* is composed of r horizontal planar geodesic curves, since $\Psi_u(P_i) = 0$ for all i the r curves are all in the plane $\{x_3 = 0\}$. Finally, we consider Σ the union of M^* and of its symmetry by $\{x_3 = 0\}$. The surface Σ is a regular minimal surface, it is complete and its flux polygon is V by construction. By construction, we know also that Σ is strongly symmetric with respect to $\{x_3 = 0\}$.

The last thing we have to prove about Σ for being sure that it is the solution of the Plateau problem at infinity is that it has finite total curvature.

We know (see [11]) that there exists a constant c such that if u is a solution of (MSE) on a domain D and $A \in D$, M is the graph of u and d is the distance along M of the point in M over A to the boundary of S then the curvature K of M at the point over A is bounded by $c/(d^2W^2(A))$.

Let us consider a half-strip $S = [0, a] \times \mathbb{R}_+$ and u a solution of (MSE) on S such that u takes the value $+\infty$ (respectively, $-\infty$) on $\{0\} \times \mathbb{R}_+^*$ (respectively, $\{a\} \times \mathbb{R}_+^*$). The boundary of the graph of u is over $[0, a] \times \{0\}$. We then have $K(x, y) \leq c/(x^2W^2(x, y))$. We consider the part $S' \subset S$ such that $x \geq x_0 > 0$. For a domain D we denote by $K(D)$ the total curvature of the graph over D . We then have

$$\begin{aligned} K(S') &= \int_{S'} K(x, y)W(x, y) \, dx \, dy \\ &\leq \int_{S'} \frac{c}{x^2W(x, y)} \, dx \, dy \\ &\leq \int_{x_0}^{+\infty} \frac{ca}{x^2} \, dx \\ &= \frac{ca}{x_0} \\ &< +\infty. \end{aligned}$$

We now use arguments that are similar to the first part of the proof of Theorem 2.2. We consider, for $\alpha \in]0, \frac{1}{2}\pi[$, $S(\alpha) = \{(x, y) \in S \mid a - y \leq x \tan \alpha\}$. Let us take α such that $\tan \alpha < \frac{1}{8}$. Then Lemma 1 of [7] proves that for every $(x, y) \in S(\alpha)$, $q(x, y) < 0$. We denote by $L(\alpha)$ the segment in $S(\alpha)$ such that $a - y = x \tan \alpha$. By Lemma 2.1, u is lower-bounded by m_1 on $L(\alpha)$ and upper-bounded by m_2 on the part of L such that $x < x_0 < a$. We then define $\Theta : (x, y) \mapsto (x, u(x, y))$. Θ is a diffeomorphism of $S(\alpha)$ into its image $\text{Im } \Theta$. We define $\chi = \Theta^{-1}$. Then $\chi_{(2)}$ is a solution of MSE on $\text{Im } \Theta$. We observe that χ extends smoothly to $\Theta(L)$. We have $\chi_{(2)}(x, z)$ tends to a as x tends to 0 so we can extend $\chi_{(2)}$ by symmetry to $\text{Im } \Theta \cup \{(x, z) \in \mathbb{R}^2 \mid (-x, z) \in \text{Im } \Theta\}$. To compute the total curvature of the graph of u over $D(\alpha) \cap \{x < x_0\}$, we use its parametrization as a

graph over $\text{Im } \Theta \cap \{0 \leq x \leq x_0\}$,

$$\begin{aligned} K(D(\alpha) \cap \{x < x_0\}) &= K(\text{Im } \Theta \cap \{0 \leq x \leq x_0\}) \\ &= \int_{\text{Im } \Theta \cap \{0 \leq x \leq x_0\} \cap \{z \leq m_1 - 1\}} K(x, z)W(x, z) \, dx \, dz \\ &\quad + \int_{\text{Im } \Theta \cap \{0 \leq x \leq x_0\} \cap \{z \geq m_1 - 1\}} K(x, z)W(x, z) \, dx \, dz \\ &\leq \int_{\text{Im } \Theta \cap \{0 \leq x \leq x_0\} \cap \{z \leq m_1 - 1\}} \frac{c}{(z - m_1)^2} W(x, z) \, dx \, dz \\ &\quad + \int_{\text{Im } \Theta \cap \{0 \leq x \leq x_0\} \cap \{z \geq m_1 - 1\}} K(x, z)W(x, z) \, dx \, dz \\ &\leq \int_1^{+\infty} \frac{cx_0}{z^2} \, dz + C \\ &< +\infty. \end{aligned}$$

The last line is true because $\overline{\text{Im } \Theta} \cap \{0 \leq x \leq x_0\} \cap \{z \geq m_1 - 1\}$ is compact.

We can do the same work for $\{(x, y) \in S \mid y \leq x \tan \alpha\}$.

We then control the curvature on each semi-strip S_i . There is a last part in $\Omega(\mathcal{P})$. This part is compact and by Lemma 2.1, u is bounded on this part; besides, the graph is regular at the boundary. So the graph above this last part is a compact part of the whole graph then it has finite total curvature. We then have proved that the graph M has finite total curvature. Since M^* is isometric to M , it has finite total curvature and then Σ has finite total curvature because it is twice as many as the one of M^* .

Appendix A. Carathéodory's Theorem

In this section, we give some explanations on an argument of the proof of Theorem 2.2. The problem is when we have a biholomorphic map between two open sets of \mathbb{C} : can we extend it to the boundary?

We consider U an open set included in \mathbb{C} and P a point of ∂U . We say that P has the property of Schönflies if, for all radii R , there exists a radius $r = r(R)$ such that for all two points in $U \cap D(P, r)$ there exists a path in $U \cap D(P, R)$ joining these two points.

We then have the following theorem that we use in our proof.

Theorem A.1 (Carathéodory). *Let U be a simply connected open set in \mathbb{C} and V an open set of the boundary of U . We consider $f : U \rightarrow D = \{z \in \mathbb{C} \mid |z| < 1\}$ a biholomorphic map. We suppose that every point of V has the property of Schönflies, then f extends to an homeomorphism from $U \cup V$ into $D \cup C$ where $C \subset \partial D$.*

A proof of this theorem can be found in [5]. In our proof, we have to verify the property of Schönflies at the points of a part of the boundary. We know that this part of the boundary is embedded in \mathbb{C} so we can build neighbourhoods of every point of the boundary in using ε -tubular neighbourhood of the boundary. These neighbourhoods prove that we have the property of Schönflies.

References

- [1] P. COLLIN AND R. KRUST, Le problème de Dirichlet pour l'équation des surfaces minimales sur des domaines non bornés, *Bull. Soc. Math. France* **119** (1991), 443–462.
- [2] C. COSÍN AND A. ROS, A Plateau problem at infinity for properly immersed surfaces with finite total curvature, *Indiana Univ. Math. J.* **50** (2001), 847–879.
- [3] R. COURANT AND D. HILBERT, *Methods of mathematical physics*, vol. II (Interscience, 1962).
- [4] D. GILBARG AND N. S. TRUDINGER, *Elliptic partial differential equations of second order*, 2nd edn (Springer, 1983).
- [5] M. HERVÉ, *Les fonctions analytiques* (Presses Universitaires de France, 1982).
- [6] D. HOFFMAN AND H. KARCHER, Complete embedded minimal surfaces of finite total curvature, in *Geometry*, vol. V, Encyclopaedia of Mathematical Sciences, vol. 90, pp. 5–93 (Springer, 1997).
- [7] H. JENKINS AND J. SERRIN, Variational problems of minimal surface type, II, *Arch. Ration. Mech. Analysis* **21** (1966), 321–342.
- [8] H. KARCHER, Embedded minimal surfaces derived from Scherk's examples, *Manuscr. Math.* **62** (1988), 83–114.
- [9] J. C. C. NITSCHKE, On new results on the theory of minimal surfaces, *Bull. Am. Math. Soc.* **71** (1965), 195–270.
- [10] J. C. C. NITSCHKE, *Vorlesung über Minimalflächen*, Die Grundlehren der Mathematischen Wissenschaften, no. 199 (Springer, 1975).
- [11] R. OSSERMAN, *A survey on minimal surfaces*, Van Nostrand Mathematical Studies (1969).
- [12] H. ROSENBERG AND R. SA-EARP, The Dirichlet problem for the minimal surface equation on unbounded planar domains, *J. Math. Pures Appl.* **68** (1989), 163–183.