

Current knowledge of the family Cardiliidae (Bivalvia, Mactroidea)

Javier H. Signorelli¹ and J.G.M. (Han) Raven²

¹Laboratorio de Reproducción y Biología Integrativa de Invertebrados marinos, LARBIM – IBIMAR – CENPAT – CONICET. Bvd. Brown 2915, U9120ACD Puerto Madryn, Chubut, Argentina (jsignorelli@cenpa-conicet.gov.ar)

²National Center for Biodiversity Naturalis, Leiden, The Netherlands (schelp56@hotmail.com)

Abstract.—The family Cardiliidae has been scarcely studied. It was historically placed in the superfamily Mactroidea. Members of this family are characterized by a cordiform shell with a typical mactrid hinge, posterior adductor muscle placed into a myophore and three ornamental areas on the external surface of the shell. Six extant and 14 exclusively fossil species have been previously mentioned in the literature as belonging to the genus *Cardilia*. The geographical distribution, stratigraphic range, type material and type locality of each extant and fossil species are provided. In this work, four extant species and 11 exclusively fossil species belonging to the genus *Cardilia* are recognized. Extant species are from the western Pacific Ocean, Indian Ocean and eastern Atlantic Ocean, while fossil taxa are recorded from deposits of middle Eocene to late Pliocene in Europe and Asia. One of them is formally described herein as *Cardilia edwardsi* new species.

Introduction

Cardiliidae bivalves are uncommon and have been rarely mentioned in recent literature. The family name was introduced by Fischer (1887, p. 1120) and historically placed in the superfamily Mactroidea (Dall, 1895; Lamy, 1917; Keen in Moore, 1969; Bieler et al., 2010; Carter et al., 2011, among others). Diagnostic characters are a cordiform shell with a typical mactrid hinge, internal resilium and external ligament, posterior adductor muscle placed into a myophore and the absence of pallial sinus. Currently six extant and fourteen exclusively fossil species have been mentioned in the literature (Lamarck, 1819; Newton, 1891; Deshayes, 1844; Nicklès, 1955; Hörnes, 1859; Fischer, 1861; Oostingh, 1934; Otuka, 1934; Beets, 1944; Van Regteren Altena and Beets, 1945; Tsuda, 1959, among others). Fossil material representing extinct species has been registered from several localities of Europe and Asia whereas fresh (partially live) specimens (representing extant species) were sampled from the Indo-Pacific and the eastern Atlantic Ocean. Only for one extant species a fossil record exists, which is restricted to the Quaternary.

During an ongoing revision of the superfamily Mactroidea it became necessary to give a synopsis of the current taxonomy and distribution of recent and fossil cardiliids.

Materials and methods

This study is based on an exhaustive literature search of described species belonging to the family Cardiliidae. All original descriptions were checked. Geographical distribution, stratigraphic range, type material and type locality of each extant and fossil species are provided. Valid species herein

revised are listed in a stratigraphic chart with temporal and geographic distribution (Table 1).

Repositories and institutional abbreviations.—Type material of all nominal species is deposited in the following institutions: Muséum national d'Histoire naturelle (MNHN), Paris; The Natural History Museum (NHMUK), London; Zoologisk Museum (ZMUC), Copenhagen; Natural History Museum of Wien (NHMW); Geology Museum of Bandung, Indonesia (GMBI); Naturalis Biodiversity Center (RGM). Museum of the Institute of Geology and Mineralogy of Faculty of Science, University of Kyoto (JC); the University Museum, The University of Tokyo (UMUT). Where possible this type material was reviewed by the authors.

Systematic paleontology

Superfamily Mactroidea
Family Cardiliidae Fischer, 1887

Diagnosis.—Shell small, equivalve, higher than long, extremely inflated, slightly inequilateral, ventrally extended; dorso-ventrally sub-ovate; anteriorly ovate to broadly ovate, dorsally strongly sloping downward anteriorly and posteriorly; ventrally ovate. Posteroventral shell margins slightly digitate in some members. Sculpture of only growth lines or closely spaced, rounded radial ribs over the posterior part of the shell. Inner shell margins smooth. Shell not gaping. Umbos strongly prosogyrate, dorsally projecting, subspiral. Hinge plate strongly arched; very thin anteriorly and posteriorly, thickened only on strong, restricted projection below beaks, containing both ligamental resilifer and cardinal dentition. Hinge with strong, inverted V-shaped cardinal in the left valve, triangular and

Table 1. Stratigraphic chart with temporal and geographic distribution of valid species of *Cardilia*.

	Era	System period	Series Epoch	Stage Age	Age Ma (million of years)	Species	Locality	
Phanerozoic	Cenozoic	Quaternary*	Holocene		0	<i>C. semisulcata</i> <i>C. inermis</i> <i>C. martini</i> <i>C. atlantica</i>	Indo–Pacific ocean coast of Sumatra Malacca strait coast of Ghana	
				Pleistocene	Upper “Ionian”	0.0117		
					Calabrian	0.126		
					Gelasian	0.781		
			Neogene	Pliocene	Piacenzian	2,588	<i>C. michellottii</i>	Asti, Italy
					Zanclean	3,600		
				Miocene	Messinian	5,332	<i>C. sundaica</i> <i>C. ludwigi</i> <i>C. krawangensis</i> <i>C. bruneiana</i>	West Java Purwakarta Purwakarta Brunei
					Tortonian	7,246	<i>C. palembangensis</i>	South Sumatra
					Serravallian	11,608	<i>C. deshayesi</i> <i>C. palembangensis</i> <i>C. toyamaensis</i>	Vienna basin Sarawak Toyama, Japan
			Paleogene	Oligocene	Burdigalian	15.97		
					Aquitainian	20.43		
					Chatian	23.03		
				Eocene	Rupelian	28.4 ± 0.1		
					Priabonian	33.9 ± 0.1		
					Bartonian	37.2 ± 0.1	<i>C. michelini</i> <i>C. edwardsi</i> <i>C. laeviuscula</i>	Paris Basin Barton Beds, England Bracklesham Bay, England
					Lutetian	40.4 ± 0.2		
				Paleocene	Ypresian	48.6 ± 0.2		
					Thanetian	55.8 ± 0.2		
					Selandian	58.7 ± 0.2		
				Danian	61.1			

*Chart taken from the International stratigraphic chart (ICS) Definition of the Quaternary and revision of the Pleistocene are under discussion. Base of the Pleistocene is at 1.81 Ma (base of Calabrian), but may be extended to 2.59 Ma (base of Gelasian). The historic “Tertiary” comprises the Paleogene and Neogene, and has no official rank.

weaker lamellar cardinal in the right valve, both positioned on a buttress below the resilifer; one lateral tooth present in the right valve. Lamellar part of the ligament not observed. Fibrous resilium submarginal/internal, triangular to tear-drop shaped, slightly posteroventrally inclined, extending onto ventrally projecting chondrophore in each valve. Anterior adductor muscle scar narrow, elongate, positioned close to the anterior shell margin; posterior adductor muscle scar positioned on strongly projecting myophoric lamina, extending ventrally from the postero-dorsal shell margin. Pallial line entire, without sinus. Shells aragonitic and internally porcelaneous, but details of shell microstructure unknown (Taylor et al., 1973).

Remarks.—The family Cardiliidae was traditionally classified into the superfamily Mactroidea. However, the examined species of this family do not really resemble other mactroideans. The characters shared with other mactroidean species are the internal resilium as well as the inverted V-shaped cardinal tooth in the left valve, being this last character, the main diagnostic that defines the superfamily. However, the presence of three different ornamental areas (OA) on shell external surface, the strongly inflated umbo, the shell outline and posterior adductor muscle placed into a myophore are exclusive features of cardiliids. Until new material become available for anatomical and molecular studies we prefer to be conservative and retain Cardiliidae as a separate family within Mactroidea.

Genus *Cardilia* Deshayes in Lamarck, 1835

Type species.—*Isocardia semisulcata* Lamarck, 1819, by subsequent designation (Deshayes, 1844). Usually this type

designation was ascribed to Herrmannsen, 1846 (e.g., Keen in Moore, 1969). However, Deshayes in Lamarck, 1835 placed both *Isocardia semisulcata* and *Isocardia michelini* in his new genus *Cardilia* without type designation. Later Deshayes (1844) assigned *Isocardia semisulcata* as type species of the genus (Huber, 2010).

Occurrence.—Europe, Indo–Pacific, Japan, South eastern Atlantic; middle Eocene – Recent.

Remarks.—The following genus level names are considered in this work as junior synonym of *Cardilia*: *Hemicyclonosta* Deshayes in Blainville, 1827 (p. 660; nomen nudum); *Hemiclostera* Bronn, 1838 (p. 806; error pro *Hemicyclonosta* Deshayes in Blainville, 1827); *Hemicyclostoma* Gray, 1840 (p. 136; error pro *Hemicyclonosta* Deshayes in Blainville, 1827); *Cardilla* Lycett, 1848 (p. 258; error pro *Cardilia* Deshayes, 1835); *Hemicyclodonta* Deshayes, 1850 (p. 251; error pro *Hemicyclonosta* Deshayes in Blainville, 1827); *Hemicycloster* Paetel, 1875 (error pro *Hemiclostera* Bronn, 1838); *Leptina* Pictet, 1855 (p. 500; based on manuscript by Bonelli) (non. Meigen, 1830, p. 283, obj.). Besides the genus *Cardilia* and its synonyms, the Oligocene *Cardilona* Marwick, 1943 (type *C. bensoni* Marwick, 1943) had been referred to this family (Keen, 1969, p. N608). However, it is currently assigned to Lyonsiellidae as a junior synonym of *Pecchiolia* Savi and Meneghini in Murchison, 1850 (Maxwell, 1978; Beu and Maxwell, 1990).

Cardilia semisulcata (Lamarck, 1819)
Figure 1.1–1.8




Figure 1. (1–8) *Cardilia semisulcata* Lamarck, 1819: (1, 2) holotype MNHN 23202; (3–8) *Cardilia semisulcata*, Jousseume collection, MNHN IM-2014-6485. Abbreviations: l: accessory lamellae; 2b: left cardinal tooth; 3b: right cardinal tooth; LAI: anterior lateral teeth; m: myophore; n: nymph. Scale bar (1, 2) 3 mm; (3–8) 1 cm.

Type material.—One syntype MNHN 23202, length 24 mm. Les mers de Nouvelle-Hollande, à l'île St. Pierre et St. François, Australia (locality erroneously registered according to Huber, 2010).

Remarks.—*Cardilia semisulcata* has been sampled from north Queensland and Western Australia by Lamprell and Healy, 1998. Deshayes (1844) recorded it from the Malacca Strait and Lamy (1917) from the Philippines and Japan. Additional records of this species are from off Port Blair, Andaman Islands (Smith, 1906); Persian Gulf (Melville and Standen, 1907; Al-Khayat, 1997); and offshore Oman (Oliver, 1995) and Northern Territory, New South Wales (both Australia), Fiji, New Caledonia, Papua New Guinea (Atlas of Living Australia). Cotton (1961, p. 20) stated that this species is not present in South Australia from where it was described, and also the Atlas of Living Australia does not provide any records from that area. Huber (2010) concluded that the type locality is erroneous.

Cardilia atlantica Nicklès, 1955
Figure 2.1–2.3

Type material.—One syntype, ZMUC 1512, one right valve collected from muddy bottom; collection date: 15/1/1946; height 14 mm, length 8.6 mm, width 5.5 mm. Station 70 of Danish Atlantide expedition to West Africa, Lat: 4.83 N, Long: 2.82 W, 60–65 m depth.

Remarks.—*Cardilia atlantica* Nicklès, 1955 is a valid living species that comes from the eastern Atlantic Ocean. It was reported by Nicklès (1955) as fossil from Quaternary deposits of Port-Gentil, Gabon and as living specimens from Liberia, Ghana (then called Gold Coast) and Nigeria. Similarities with *C. inermis* Deshayes from the Pacific Ocean and to *C. krawangensis* Oostingh from the Neogene of Java were mentioned, although Oostingh's species has a larger shell with a less prosogyrous umbo.

Cardilia bruneiana Beets, 1944
Figure 2.4–2.6

Type material.—Holotype RGM 783503, articulated specimen; height 20 mm; length 14.7 mm; width 18 mm.; one paratype RGM 783535, height 9.4 mm. Late Miocene deposits (Seria Formation) sample Tutong 13, NW of Tutong, Brunei, NW Borneo. The paratype is from sample Tutong 15B slightly lower in the same formation. The Seria Formation was deposited during the Tortonian as was recently reconfirmed by microfossils from an outcrop of a younger and shallower part of the formation (Bukit Ambug, latest Tortonian; Roslim et al., 2016).

Remarks.—*Cardilia bruneiana* was registered from deposits of similar age than those where *C. krawangensis* and *C. ludwigi* came from. The differences compared to other Indonesian species have been analyzed (Beets, 1944). In his publication, two new names were introduced: *Cardilia bruneiana* and *C. palembangensis*. The first species has been described from Tutong, Brunei and has a less oval shell than *C. krawangensis*, with the radial ribs of the posterior area extending further forward ventrally but less so on the umbo. In addition, two undetermined

species were mentioned by Beets (1944): *Cardilia* sp. 1 from Padang, Sumatra and *Cardilia* sp. 2 from Rembang, Central Java Province.

Cardilia deshayesi Hörnes, 1859
Figure 2.7, 2.8

Type material.—Holotype NHM Wien 1855/XLV/286, a single specimen collected from middle Miocene deposits of the Vienna Basin; height 9 mm, length 6 mm, and width 8 mm. Badenian, middle Miocene deposits exposed at Steinebrunn, Vienna Basin, Austria.

Remarks.—*Cardilia deshayesi* is a valid species widely recorded from Badenian (middle Miocene) exposures in the Vienna Basin (Karrer, 1877; Handmann, 1888; Sieber, 1958; Schultz, 2003, among others). It was additionally registered from different localities in east Europe like Smarzowa, Poland (Friedberg, 1936), Lontov, Slovakia (Tejkal et al., 1967), Mikulov, Czech Republic (Studencka et al., 1998), and Venetia, Italy (Stefanini 1916; Venzo, 1934).

Cardilia gemmulata Gould, 1861

Type material.—Lost, not reported in the original description as deposited in any American institution (Johnson, 1964), height 2.5 mm, length 2 mm, width 2 mm. China seas.

Remarks.—*Cardilia gemmulata* was described from the China Sea, only 2 mm length. It should not be considered a nomen oblitum as concluded by Higo et al. (1999) because it was mentioned after 1899 (art. 23.9.2 of the code) by several authors (Lamy, 1917; Johnson, 1964; Ruhoff, 1980; Huber, 2010). However, the type material is probably lost as it was not mentioned by Johnson (1964) as deposited in any institution. *Cardilia gemmulata* should be considered *nomen dubium*, as suggested by Huber (2010), until new material can be collected.

Cardilia inermis Deshayes, 1844
Figures 3.1–3.3, 4.3, 4.4

Type Material.—Not found. Coast of Sumatra, Indonesia.

Additional material examined.—Four valves, two broken, NHMUK 20160445, H. Cuming collection.

Remarks.—According to Huber (2010), the species *C. inermis* is the smaller and more squarish living species. Originally described from Sumatra, its distribution has been expanded to South China, Thailand and Philippine Islands (Fischer, 1861; Lynge, 1909).

Cardilia krawangensis Oostingh, 1934
Figure 5.1, 5.2

Type material.—Not found, probably deposited at GMBI; height 11.5 mm, width 7.5 mm, thickness 10.5 mm. Besides the holotype, there are two specimens from the same locality. Topographical/Geological Map Sheet 30 (Purwakarta, Java Province, Indonesia), Locality 1998, Cikao, Cidap Formation




Figure 2. (1–3) *Cardilia atlantica* Nicklès, 1955, syntype, ZMUC 1512; (4–6) *Cardilia bruneiiana* Beets, 1944, holotype RGM 783503; (7, 8) *Cardilia deshayesi* Hörnes, 1859, holotype NHM Wien 1855/XLV/286. Scale bar (1, 2) 3 mm; (4–6) 1 cm; (7, 8) 2 mm.




Figure 3. (1–3) *Cardilia inermis* Deshayes, 1844, original illustrations; (4, 5) *Cardilia martini* Deshayes, 1844, original illustrations. Scale bar 5 mm.

(previously known as Tjidadap Beds, late Miocene to early Pliocene, sensu Dr. O. Ludwig).

Remarks.—Although *C. krawangensis* superficially resembles *C. palembangensis*, it differs clearly in shape, having a more rounded posterior side and being considerably smaller. The escutcheon is broader and smooth. The number of ribs is higher and they do not reach the middle of the shell along the ventral side as in *C. palembangensis*.

Cardilia laeviuscula Sowerby in Dixon, 1850
Figure 5.5–5.14

Type material.—Lectotype herein designated: NHMUK PI OR 73007b, the specimen figured in Dixon, 1850 (pl. 2, fig. 6a; three views); paralectotypes: three specimens NHMUK PI OR73007a from Bracklesham Bay, middle Eocene. Locality: Selsey, Bracklesham Bay, West Sussex, England. Lithostratigraphy: Unit S10 (Curry et al., 1978), Selsey Formation, Bracklesham Group; confined to this horizon (J. Todd, personal communication, 2017). Age: late Lutetian, zone NP16, middle Eocene.

Remarks.—In this work, two species from the fossil record of England are considered valid. The first one is *Cardilia laeviuscula* described by Sowerby in Dixon, 1850 (p. 165, pl. 2, fig. 6a) from middle Eocene deposits exposed at Bracklesham Bay, West Sussex, UK. The second one is *Cardilia edwardsi* n. sp.

Cardilia ludwigi Oostingh, 1934
Figure 5.3, 5.4

Type material.—Not found, probably deposited at GMBI; height 9.5 mm, width 7.8 mm thickness 8.7 mm. Two additional specimens registered from the same locality. The hard molds do not allow observation of the hinge. Purwakarta, Java province, Indonesia, Locality 1998, Cikao, Cidapad Formation (previously known as Tjidadap Beds), late Miocene to early Pliocene, sensu Dr. O. Ludwig.

Remarks.—*Cardilia ludwigi* has a circular outline with finer radial ribs and narrower interspaces. Oostingh's species, *C. krawangensis*, and *C. ludwigi*, came from one outcrop and thus have the same age and lived as sympatric species. Their age is very similar to that of *C. bruneiana*. The descriptions of these two species are based on Oostingh's descriptions and drawings because the type material could not be accessed.

Cardilia martini Deshayes, 1844
Figures 3.4, 3.5, 4.5–4.13

Type material.—Not found. Malacca Strait, SE Asia.

Additional material examined.—Three valves, one broken, NHMUK 20160446. Measures of largest specimen: height 16 mm, length 12 mm.

Remarks.—*Cardilia martini*, described from the Malacca Strait has been recorded from the coast of China and the Philippines (Adams and Adams, 1853–1858; Sowerby, 1873; Paetel, 1883). It is the narrowest living species with stronger ribs (Huber, 2010). However, no modern records of this species were found in the literature.

Cardilia michelini (Deshayes in Michelin, 1825).
Figure 5.15, 5.16

Type material.—MNHN A09474, one well-preserved syntype. La Chapelle-en-Serval, Oise, France, Sables de Beauchamp, Bartonian (Auversien), middle Eocene. Syntype measures: height 16 mm, length 11 mm.

Remarks.—*Cardilia michelini* was introduced by Deshayes in Michelin (1825) and illustrated on a lithographic plate with its respective explanation. Sensu Keen (1969), the evidence for valid publication by Michelin is inconclusive, however his contemporaries accepted it. Later, *C. michelini* was recorded as fossil in France by other authors (Deshayes, 1827; Deshayes and Milne Edwards, 1835; Fischer, 1861; Cossmann and Pissarro, 1906; Le Renard and Pacaud, 1995). It was additionally reported from the Eocene of Gerona, Spain (Marcet I Riba, 1955).




Figure 4. (1–4) *Cardilia inermis* Deshayes, 1844, NHMUK 20160445; (5–13) *Cardilia martini* Deshayes, 1844, NHMUK 20160446. Scale bar (1, 2) 5 mm; (3, 4) 2 mm; (5–13) 10 mm.




Figure 5. (1, 2) *Cardilia krawangensis* Oostingh, 1934, original illustrations; (3, 4) *Cardilia ludwigi* Oostingh, 1934, original illustrations; (5–14) *Cardilia laeviuscula* J. de C. Sowerby in Dixon, 1850; (5, 6) Lectotype; (7–14) additional examined material NHMUK 73007b; (15, 16) *Cardilia michelini* Deshayes in Michelin, 1825, MNHN A09474, syntype, image photographed by Jacques Mouchart (Project: E-Recolnat). Scale bar (1–14) 5 mm; (15, 16) 5 mm.

Cardilia michelottii Deshayes, 1844

Type material.—Not found. Type measures reported in the original description: height 23 mm, length 18 mm, width 20 mm. Collected by M. Michelotti from Pliocene deposits of Asti, Piedmont, Italy.

Remarks.—*Cardilia michelottii* is a valid species subsequently registered by several authors from the Piacenzian, late Pliocene of Italy (Bronn, 1848; d'Orbigny, 1852; Manzoni, 1868; Pantanelli, 1892; Sacco, 1901).

Cardilia palembangensis Beets, 1944

Figure 6.1–6.12

Type material.—Originally deposited at Geological Museum, University of Utrecht, now RGM. Holotype RGM 822551, articulated specimen height 16.4 mm, length 12.5 mm, width 14.8 mm; RGM.822549 paratype 1; RGM.1007835 Paratype 2 (single valve); RGM.1007836 a paratype, single valve (part of larger sample of paratypes with number RGM.1007836, containing three articulated specimens and this single valve). Test pit Talang Abab, province South Sumatra, Indonesia; Air Benakat Formation, Lower Palembang Member; middle to late Miocene.

Remarks.—Table 2 compares the five species from SE Asia. Of all the SE Asian fossil species, the largest number of specimens is available for *C. palembangensis*. Besides the four articulated specimens and two valves in the sample from the type locality, 15 articulated specimens were collected from the Pendopo oilfield, 10 miles E of Talang Akar, early to middle Miocene, border between lower and upper part of the Telisa Formation, sensu Huyse, 15 articulated specimens, RGM820.364 and one articulated specimen from the Kampong Tengah C/D outcrops, S. of Miri, Sarawak, Malaysia, Sibuti Formation, Langhian, middle Miocene (sample F41.23 in J.G.M. Raven's collection). Because there are adult and juvenile specimens, the variation in this species is well understood (Table 3 gives measurements for a selection of these). Of the other species from this area, only specimens of *Cardilia sundaica* are available. Additional material will help both in the differentiation among the species and understanding their stratigraphic ranges.

Cardilia reeveana Hidalgo y Rodriguez, 1903

Type material.—Does not exist. The taxon is based on Sowerby's illustration (in Reeve and Sowerby, 1843–1878, pl. 1, fig. 2) of *Cardilia inermis* (non Deshayes). Philippines.

Remarks.—*Cardilia reeveana* Hidalgo y Rodriguez, 1903, based on Sowerby's illustration of *C. inermis*, seems to be unnecessary. Huber (2010) noted that Sowerby's illustration was accepted by Lynge (1909) as belonging to the earlier described *C. inermis*, which expanded the distribution of that species previously given by Fischer (1861) from southern China and east Thailand, to also include the Philippines.

Cardilia sundaica Van Regteren Altena and Beets, 1945

Figure 6.13–6.15

Type material.—Holotype RGM 820549, single valve, height 15.2 mm, length 11 mm. Originally the type was stored at the Geological Institute, University of Amsterdam; it is now at RGM. Left bank Ci Gugur River, north of Koleberes plantation, SW of Bandung province, West Java, Indonesia; late Miocene.

Remarks.—*Cardilia sundaica* Van Regteren Altena and Beets, 1945 was recognized by its authors as an intermediate form between *C. ludwigi* and *C. krawangensis*. It has a narrower anterior surface (AO3) without shell sculpture, delimited posteriorly by radial ribs that are transversely wrinkled. The furrows and ridges are all approximately equal in width, but the ribs are narrower than the furrows. The lunule is significantly thinner than in *C. krawangensis*. The fine radial shell sculpture runs almost parallel to the ribs, but is much narrower.

Cardilia toyamaensis Tsuda, 1959

Figure 6.16

Type material.—Holotype, left valve JC1400026. Type measures: height 11.7 mm, length 9 mm, thickness 5.5 mm; three paratypes JC 1400027, JC140028, one right valve and two contact valves. Kurosedani Formation, late early Miocene. Toyama Prefecture on the new River County Daze, Japan.

Remarks.—The Japanese *Cardilia toyamaensis* appears to be common because it has been recorded from different Japanese Miocene outcrops (Okamoto and Terachi, 1974; Takayasu, 1981; Taguchi, 2002; Nakagawa, 2009, among others). This species differs from the recent *C. semisulcata* in its smaller shell without radial sculpture on the posterior area and in its smaller number of ribs.

Cardilia edwardsi new species

Figure 7.1–7.5

Type material.—Holotype, NHMUK PI TB 14589 (1), length 5.4 mm, height 5.6 mm; paratype, one broken shell, length 7.3 mm, height 7.6 mm, NHMUK PI TB (2), both Dennis Curry collection.

Diagnosis.—Shell small, radial ribs crossed by fine concentric striation, disposed along the postero-dorsal axis from the umbo to the ventral edge, external surface areas poorly defined; on the anterior side of this line the external surface is reticulated over the umbonal area, and with fine and irregular growth lines along the ventral edge; posterior myophore well developed, trigonal with a circular posterior adductor muscle scar.

Occurrence.—Type locality: Barton-on-Sea, Hampshire, England. Lithostratigraphy: 'Chama Bed' (Bed H of Burton, 1933), Becton Formation, Barton Group. Age: late Bartonian, middle Eocene.

Description.—Shell small, fragile, subcircular, height up to 8 mm; external surface with three areas; dorso-posterior, with irregular and small concentric striation from the posterior edge to the first ribs of the adjacent area; external area with radial and narrow ribs disposed along the postero-dorsal





Figure 6. (1–12) *Cardilia palembangensis* Beets, 1944 type series from Talang Alab, Palembang; (1–4) paratype 1 RGM 822549; (5–8) holotype RGM 822551; (9–12) paratype 2 RGM 1007835. (13–15) *Cardilia sundaica* Van Regteren Altena and Beets, 1945, holotype RGM 820549. (16) *Cardilia toyamaensis* Tsuda, 1959, holotype, left valve JC1400026. OA: Ornamental area. Scale bar (1–16) 10 mm.

Table 2. Morphological comparison of the five extinct species from SE Asia.

Species	<i>krawangensis</i>	<i>palembangensis</i>	<i>sundaica</i>	<i>ludwigi</i>	<i>bruneiana</i>
Examined material	Three specimens from one locality	>20 specimens from three localities	A single valve	Three specimens from one locality	Four specimens from two close localities in single stratigraphic interval
Shell	Fully equivalve	Fully equivalve	unknown (single valve)		
Max. size (H, L, W in mm)	11.5, 7.5, 10.5	17.5, 12.9, 15.2	15.4, 10.4, 13.6	9.5, 7.8, 8.7	20, 14.7, 18
Shape	Much higher than long. Anterior edge passes smoothly into dorsal edge; dorsal side of ventral edge slightly concave; short straight dorsal edge.	Much higher than long. Anterior edge passes smoothly into dorsal edge; dorsal side of ventral edge slightly concave; short straight dorsal edge.	Much higher than long. Ellipsoid outline with rounded ventral edge and short straight dorsal edge.	Slightly higher than long. Circular outline with short, straight dorsal edge.	Slightly higher than long. Anterior edge rounded, with marked bend to more ellipsoid shaped posterior edge.
Sculpture (OA1 – Escutcheon, depression posterior to beaks)	Broad, lancet shaped, slightly sunken; smooth; slightly wider on the left valve	Narrow zone with radial lamellae; 2.4–2.9 mm wide in adult specimens	Wide zone (8.6 mm) with radial lamellae	Very narrow smooth area bordered by wider zone with radial lamellae	Narrow zone of a few granulated ribs.
Sculpture (OA2)	21 ribs (Oostingh's [1934] drawings). Posterior ribs consist of rows on nodules. Ribs do not reach the middle of the shell, neither along the ventral edge nor the umbonal area Ribs and grooves equally wide with zigzag lines (as along the edge): inclined to the back and ventral side on the ribs and to the front and ventral side in the grooves.	11–19 ribs. Ribs extend to middle of the shell along the ventral edge (albeit last ribs often are faint) but do not reach the middle in the umbonal area. On posterior side grooves as wide as or wider than ribs, towards anterior side ribs become flatter and wider whereas grooves become narrower. Lying S-shaped stripes on the ribs and to the front and ventral side in the grooves. Ribs protrude slightly over the ventral edge.	11 ribs (single specimen). Ribs extend beyond middle of the shell along the ventral edge and the umbonal area. Flat ribs, especially the anterior ones are much wider than the grooves.	10 ribs (Oostingh's [1934] drawings). Posterior ribs consist of rows of nodules. Ribs extend to middle of the shell along the ventral but do not reach the middle of the umbonal area. Ribs slightly wider than grooves but posterior ribs narrower than the grooves.	17 ribs (last one fine). Ribs extend to middle of the shell along the ventral edge (just beyond the flexure) but do not reach the middle in the umbonal area. Ribs slightly wider than grooves but posterior ribs narrower than the grooves.
Sculpture (OA3)	Fine radial lines parallel to ribs near ventral edge, at slight angle nearer the umbo	Fine diagonal lines abutting at clear angle against the ribs	Only near the ribs; fine diagonal lines abutting at clear angle against the ribs	Weak growth lines	Growth lines only

Table 3. Measurements of *C. palembangensis* showing its intraspecific variation and comparison with *C. sundaica*. Size measurements in mm.

Locality	Height	Length	Width	Number of ribs at edge	Number of ribs 1/3rd from umbo	Width ribbed field (left valve)	Remarks
<i>Cardilia palembangensis</i> Beets, 1944							
Talang Alab (types)	16.5	12.2	14.5	13	11	2.4	Holotype
	12.9	9.7	11.7	17	11	1.6	Paratype 1
	17.2	12.4	7.2 (half)	15	14	2.7 estimate	Paratype 2 (valve)
	14.4	10.7	11.9	13	11	1.9	Paratype
	12.5	9.5	10.4	11	11	1.6	Paratype
	17.5	11.0	8.0 (half)	14	14	2.7	Paratype (valve)
	16.8	12.4	15.1	14	12	2.5	
	17.1	12.2	15.2	13	11	2.9	
	15.8	11.4	13.6	19	16	2.5	
	16.7	12.9	12.2	13	13	2.6	
Pendopo	17.2	12.6	13.7	14 (2 weak)	12	2.4	
	12.1	7.9	10.5	11	11	1.8	
	10.1	7.4	9.0	12	12	?	
<i>Cardilia sundaica</i> Van Regteren Altena and Beets, 1945							
Left bank Ci Gugur river	15.4	10.4	6.8 (half)	11	11	4.3	Holotype (valve)

**Figure 7.** (1–5) *Cardilia edwardsi* n. sp. from the Becton Formation (late Bartonian) of Barton, Hampshire, UK; (1) holotype, NHMUK PI TB 14589, (2) paratype, one broken shell, NHMUK PI TB 14589, both from the Curry collection; (3, 5) NHMUK PI OR 73008; (4) NHMUK PI OR 73008, additional material from the F.E. Edwards collection. Scale bar (1–5) 5 mm.

axis from the umbo to the ventral edge; all ribs crossed by additional shell sculpture of concentric and irregular growth lines; straight line that separates the external surface areas poorly defined; area of the external surface covering a major portion of the shell, composed by fine and irregular growth lines less evident near the ventral edge and reticulated shell

sculpture over the umbonal area. This area extends from the straight line towards the anterior side of the shell; internally mactroid hinge plate with an oblique chondrophore ventrally projected and a posterior trigonal myophore well developed, with a circular posterior adductor muscle scar well visible; pallial sinus absent.

Etymology.—Honoring Frederick E. Edwards, the discoverer of this species.

Remarks.—This species was mentioned previously by a few authors as *Cardilia radiata* from exposures of the Barton and the Bracklesham Beds (Tennant, 1847, p. 29; Beet–Jukes, 1857, p. 530; Newton, 1891, p. 85). Although this species was attributed to either Edwards or Sowerby, it is not listed by Petit (2009) as a taxon named by Sowerby and not found published in the “Quarterly Journal of the Geological Society of London” by Edwards (1854). *Cardilia radiata* appears in a manuscript name of F.E. Edwards that was never validated. In this work, it is formally described as *Cardilia edwardsi* new species from the Becton Formation, Barton Group.

Discussion

Literature search allows an understanding of the geographical and temporal radiation of the family Cardiliidae. Extant species were historically sampled from the western Pacific Ocean in the Philippines, Indonesia, China, and Malaysia and the Indian Ocean/Persian Gulf in Oman and Qatar. From this huge area, only three valid species are recognized: *C. semisulcata*, *C. inermis*, and *C. martini*. *Cardilia atlantica* is the fourth living species registered in the Atlantic Ocean. On the other hand, from 14 names found in the literature for species only known as fossils, 11 are recognized as valid. Five valid species were registered from the fossil record of Europe, one of them formally described herein. They were registered from different localities in England, France, Austria, Italy, and Russia, among others. The stratigraphic range of these records goes from middle Eocene to late Pliocene. They are: *Cardilia deshayesi* from middle Miocene exposures at Steinebrunn, Vienna Basin, Austria; *Cardilia michelottii* from Pliocene deposits at Asti, Italy; *Cardilia michelini* described from Bartonian (middle Eocene) deposits exposed at La Chapelle-en-Serval, Oise, France; *Cardilia laeviuscula* from Lutetian (middle Eocene) strata, Bracklesham Bay, West Sussex, England; and *Cardilia edwardsi* n. sp. from the Becton Formation, late Bartonian, middle Eocene of Hampshire, England.

Finally, the family Cardiliidae was recorded in the fossil record of Asia from where seven species have been described: four from Indonesia, one from Brunei and two from Japan. These are *C. sundaica* Van Regteren Altena and Beets, 1945; *C. bruneiana* and *C. palembangensis*, both described by Beets in 1944; *C. krawangensis* and *C. ludwigi*, both described by Oostingh in 1934; and *Cardilia yudaensis* Otuka, 1934 and *Cardilia toyamaensis* Tsuda, 1959 from Japan. The species *Cardilia yudaensis*, described by Okuta (1934, p. 620, pl. 48, figs. 46–48), was reported from lower Kadonosawa Series, Yuda, Iwate Prefecture, Japan. The author placed this species with doubts within the genus *Cardilia*, although he considered it as the most adequate. However, we conclude that after revision of the original description and illustration, the species *Cardilia yudaensis* must be excluded from the genus *Cardilia* due to the inequivalve form of the type material.

Two nominal subspecies (*Cardilia michelini asiana* and *Cardilia michelini georgiana*) are recorded in the Global Names Index portal (http://www.gni.globalnames.org/name_strings),

as described by Korobkov (1971). However, these names are not found within the publication (S. Popov, personal communication, 2016). In addition, type material of either taxon has not been deposited into the collection of Saint Petersburg Central Geological Museum, where Korobkov’s types are (S. Popov, personal communication, 2016). Therefore these names are considered invalid.

All species belonging to the genus *Cardilia* have three different ornamental areas (OA) on the external surface of the shell. The first one (OA1) is the escutcheon: dorso–posterior, concentric, uniformly separated, lamellar, nodular towards the top, and intersecting the first ribs of the second ornamental area. It extends from the posterior side of the umbo to the mid-posterior end of the shell. The second area (OA2) has narrow ribs disposed along the postero-dorsal axis from the umbo to the ventral edge and is delimited by a straight line that separates this area from the contiguous one. The number of ribs of this area is variable. The third and final area (OA3) covers almost the entire external surface and is composed fine and irregular growth lines that become invisible along the ventral edge. This area extends from the straight line towards the anterior side of the shell. The morphologies of these areas have taxonomic value for the identification of the different species of the genus. In addition, the outline of the shell plays an important role in the identification of *Cardilia* species. Hinge morphology of all examined species is very conservative with the strong inverted V-shaped cardinal tooth obliquely anteriorly directed in the left valve, which is complementary to the prominent, triangular, and bifid cardinal tooth (3a) of the right valve; additionally, with a second cardinal tooth (3b) in the right valve and a small and rudimentary anterior lateral tooth (LAI).

Many works had been revised morphologically and genetically the phylogeny of the related families Mactridae and Mesodesmatidae (Giribet and Wheeler, 2002; Taylor et al., 2007; Combosch et al., 2017, among others). The superfamily Mactroidea was traditionally considered to be a well-supported monophyletic group with a sister group, Cardioidea. Taylor et al. (2007) agreed with the monophyletic concept of Mactroidea, but suggested a potential sister group with Ungulinidae, and the group Veneridae/Corbiculidae/Arcticidae, and Chamiidae, but with no connection with Cardioidea. Within this context, the phylogenetic position of Cardiliidae within Mactroidea is uncertain.

The first records of mactrids date from the Early Cretaceous (Aptian). In a detailed study of hinge development, Saul (1973) suggested that mactrids are derived from the Arcticidae. Cretaceous Mactridae are characterized by a moderate development of resilifer with two cardinals in each valve and long laterals. This kind of hinge was also observed in the Arcticidae, however in Cenozoic Mactridae this similarity is masked due to the enlargement of the resilifer. The systematic position of Cardiliidae within Mactroidea was historically based on similarities of the hinge structure and resilifer. As noted by Keen (in Moore, 1969) and subsequent authors, these similarities include a similar pattern of cardinal and lateral teeth with an external ligament small or wanting, and an internal ligament seated in a resilifer.

There are no new data on anatomy, reproduction, and ecology of Cardiliidae. Thus, the conventional position within

Mactroidea is unconfirmed, but retained. Besides type material of described species, scarce additional specimens are available in malacological collections. The position of *Cardilia* between Mactridae and Myidae, mentioned by Marwick (1943), was never tested. Currently, the superfamily Mactroidea includes four families (Signorelli and Carter, 2016). They are Mactridae, which originated in the Late Cretaceous, Anatinellidae, Mesodesmatidae, and Cardiliidae, all recorded from the Eocene. All families are characterized as shallow or deep burrowers, occurring in the intertidal zone or on the shelf.

This work is part of an ongoing revision of superfamily Mactroidea, including all recent and fossil species. This study includes morphological data both from literature and from new observations of type material and non-type specimens, and states the basis for future studies on recent and fossil species of Cardiliidae.

Conclusions

The Cardiliidae comprise a single genus, *Cardilia*, which is known from middle Eocene to Recent. Most species are rare and material available is very limited. Eleven extinct species (including one that is described in this paper) and four extant species (for which fossil material is remarkably scarce) are recognized. The placement of the family Cardiliidae in the superfamily Mactroidea requires confirmation.

Acknowledgments

Special thanks to V. Herós and P. Maestrati (MNHN); A. Salvador, K. Way, and J. Todd (NHMUK); T. Schiøtte (ZMUC); B. Hoeksema, F. Wesselingh, and R. Pouwer (Naturalis); and M. Harzhauser and O. Mandic (NHMW) for their assistance in the revision of type material and additional examined specimens. Many thanks to G. Rosenberg and S. Popov for their comments. JHS acknowledges CONICET of Argentina. This work was partially supported by the PICT 640 of the Agencia Nacional de Investigaciones Científicas y Técnicas. This is publication N° 89 of the Laboratorio de Reproducción y Biología Integrativa de Invertebrados Marinos (LARBIM).

References

Adams, H., and Adams, A., 1853–1858, The genera of Recent Mollusca; arranged according to their organization: London, John van Voorst, Paternoster Row, v. 2, 661 p.

Al-Khayat, J.A., 1997, The marine Mollusca of the Qatari waters, Arabian Gulf: Journal of Qatar University of Science, v. 17, p. 479–491.

Atlas of Living Australia website at <http://biocache.ala.org.au/occurrences/search?q=lsid:urn:lsid:biodiversity.org.au:afd.taxon:bf4000e1-beee-4362-af45-6a01029c2ad5>. [accessed 30 May 2017]

Beete Jukes, J., 1857, The Student's Manual of Geology: Edinburgh, Adam and Charles Black, 607 p.

Beets, C., 1944 [1943], Notizen über die Gattung *Cardilia* Deshayes: Verhandlungen Geologisch Mijnbouwkundig Genootschap voor Nederland en Koloniën, v. 13, p. 355–362.

Beu, A., and Maxwell, P.A., 1990, Cenozoic Mollusca of New Zealand: New Zealand Geological Survey Paleontological Bulletin, v. 58, p. 1–518.

Bieler, R., Carter, J., and Coan, E., 2010, Classification of bivalve families, in Bouchet, P., and Rocroi, J.-P., eds., Nomenclator of Bivalve Families: Malacologia, v. 52, p. 1–184.

Bronn, H.G., 1834, 1837, 1838, Lethaea Geognostica oder Abbildungen und Beschreibungen der für die Gebirgs-Formationen bezeichnendsten Versteinerungen: Stuttgart, Schweizerbart, v. 1, p. 1–544 [1834, 1837]; v. 2, p. 545–1350 [1837, 1838].

Bronn, H.G., 1848–1849, Index Palaeontologicus oder Übersicht der bis Jetzt Bekannten Fossilen Organismen, bearbeitet unter Mitwirkung der Herren Prof. H. R. Göppert and Herm. v. Meyer, bearbeitet von Dr. H. G. Bronn. A. Nomenclator Palaeontologicus, in Alphabetischer Ordnung [two halves, each in a separate volume]: Stuttgart, Schweizerbart, Erste Hälfte, A–M, p. 1–775 [1848]; Zweite Hälfte, N–Z, p. 776–1381 [1848].

Carter, J.G., et al. 2011, A Synoptical Classification of the Bivalvia (Mollusca): Paleontological Contributions, v. 4, p. 1–47. <https://doi.org/10.17161/PC.1808.8287>.

Combosch, D.J., Collins, T.M., Glover, E., Graf, D.L., Harper, E., Healy, J.M., Kawauchi, G.Y., Lemer, S., McIntyre, E., Strong, E., Taylor, J.D., Zardus, J.D., Mikkelsen, P., Giribet, G., and Bieler, R., 2017, A family-level Tree of Life for bivalves based on a Sanger-sequencing approach: Molecular Phylogenetics and Evolution, v. 107, p. 191–208.

Cossmann, A.E.M., and Pissarro, G., 1906, Iconographie complète des coquilles fossiles de l'Éocène des environs de Paris: Paris, H. Bouillants, v. 1, 45 pl.

Cotton, B.C., 1961, South Australia Mollusca—Pelecypoda. Handbook of the Flora and Fauna of South Australia Series: South Australian Branch of British Science Guild: Adelaide, W. L. Hawes, Government Printer, 363 p.

Dall, W.H., 1895, Synopsis of a review of the genera of Recent and Tertiary Mactridae and Mesodesmatidae: Malacological Society of London, Proceedings, v. 1, p. 203–213.

Deshayes, G.P., 1825, in Hardouin Michelin, M., Collection de M. Hardouin Michelin. Coquilles Fossiles du Bassin Parisien: 1 pl. + expl.

Deshayes, G.P., 1827, in Blainville, H.M.D. de. 1825, 1827, Manuel de Malacologie et de Conchyliologie: Paris, F. G. Levrault, v. 1, 664 p.

Deshayes, G.P., 1827, Figures de quelques coquilles de la collection de M. Hardouin Michelin: Bulletin des sciences naturelles et de Géologie, v. 11, p. 446–447.

Deshayes, G.P., 1844, Magasin de Zoologie, d'Anatomie Comparée et de Paléontologie: série 2, v. 6, n. 8, p. 1–8.

Deshayes, G.P., 1850, Traité Élémentaire de Conchyliologie avec les Applications de cette Science à la Géologie Tome Premier, pt. 2, Conchifères Dimyaires, “1843–1850”: Paris, Victor Masson, v. 2, p. 129–824.

Deshayes, G.P., and Milne Edwards, H., 1835, Histoire des Mollusques, in Lamarck, J. B. P. A. de., Histoire naturelle des animaux sans vertèbres: présentant les caractères généraux et particuliers de ces animaux, leur distribution, leurs classes, leurs familles, leurs genres, et la citation des principales espèces qui s'y rapportent: précédée d'une introduction offrant la détermination des caractères essentiels de l'animal, sa distinction du végétal et des autres corps naturels, enfin, l'exposition des principes fondamentaux de la zoologie., 2nd edition, revised, edited, and augmented: Londres, J.B. Baillière, v. 6, 600 p.

Dixon, F., 1850, The Geology and Fossils of the Tertiary and Cretaceous Formations of Sussex: London, Richard and John Edward Taylor [also Longman, Brown, Green & Longmans], 422 p.

Edwards, F.E., 1854, List of the fossil shells from the Eocene strata of Bracklesham Bay, Sussex: The Quarterly Journal of the Geological Society of London, v. 10, p. 450–454.

Fischer, P., 1861, Liste monographique des especes du genre *Cardilia*: Journal de Conchyliologie, v. 9, p. 335–344.

Fischer, P., 1880–1887, Manuel de Conchyliologie et de Paleontologie Conchyliologique ou Histoire Naturelle des Mollusques Vivants et Fossiles: Paris, Savy, 1369 p.

Friedberg, W., 1934–1936, Mięczaki miocenijskie Ziemi Polskich. Część I – Małże (Mollusca miocenaica Poloniae. Pars II Lamellibranchiata): Kraków, Polskie Towarzystwo Geologiczne, 274 p.

Giribet, G., and Wheeler, W., 2002, On bivalve phylogeny: a high-level analysis of the Bivalvia (Mollusca) based on combined morphology and DNA sequence data: Invertebrate Biology, v. 121, p. 271–324.

Gould, A.A., 1861, Descriptions of shells collected by the North Pacific Exploring Expedition: Proceedings of the Boston Society of Natural History, v. 8, p. 14–40.

Gray, J.E., 1840, Shells of molluscous animals, Eastern Zoological Gallery, in Synopsis of the Contents of the British Museum: London, G. Woodfall & Son, p. 89–156.

Handmann, R., 1888, Die Neogenablagerungen des österreichisch-ungarischen Tertiärbeckens: Münster, Aschendorff, 71 p.

Herrmannsen, A.N., 1846–1852, Indicis Generum Malacozoorum Primordia. Nomina Subgenerum, Generum, Familiarum, Tribuum, Ordinum, Classium; Adjectis Auctoribus, Temporibus, Locis Systematicis, Atque Literariis, Etymis, Synonymis. Praetermittuntur Cirripedia, Tunicata et Rhizopoda: Cassel, Theodori Fischer, v. 1, p. 1–232 [1846], p. 233–637 [1847]; v. 2, p. 1–352 [1847], p. 353–492 [1848], p. 493–717 [1849]; Supplementa et Corrigenda, p. 1–140 [1852].

Hidalgo y Rodriguez, J.G., 1903, Obras malacológicas. (4ª Parte) Estudios preliminares sobre la fauna Malacológica de las islas Filipinas: Memorias de la Real Academia de Ciencias Exactas, Físicas y Naturales de Madrid, v. 21, p. 1–400.

- Higo, S., Callomon, P., and Goto, Y., 1999, Catalogue and Bibliography of the Marine Shell-Bearing Mollusca of Japan. Gastropoda, Bivalvia, Polyplacophora, Scaphopoda: Ōsaka–fu, Yao–shi, Suchiro–chō, Japan, Elle Scientific Publications, 749 p.
- Hörnes, M., 1859, Die Fossilien Mollusken des Tertiär–Beckens von Wien unter der Mitwirkung von Paul Partsch: Abhandlungen der kaiserlich–königlichen Geologischen Reichsanstalt, v. 4, p. 1–477.
- Huber, M., 2010, Compendium of Bivalves. A Full-color Guide to 3,300 of the World's Marine Bivalves. A Status on Bivalvia after 250 Years of Research: Hackenheim, Germany ConchBooks, 901 p.
- Johnson, R., 1964, The Recent mollusca of August Addison Gould: Bulletin of the United States National Museum, v. 239, p. 1–172.
- Karrer, F., 1877, Geologie der Kaiser Franz Josefs Hochquellen–Wasserleitung. Eine Studie in den Tertiär–Bildungen am Westrande des alpinen Theiles der Niederung von Wien: Abhandlungen der Kaiserlich–Königlichen Geologischen Reichsanstalt, v. 9, p. 1–420.
- Keen, A.M., 1969, Superfamily Mactracea Lamarck, 1809, in Moore, R.C., ed., Treatise on Invertebrate Paleontology, Part N. Mollusca 6: Boulder, Colorado and Lawrence, Kansas, Geological Society of America and University of Kansas Press, p. N595–N610.
- Korobkov, I., 1971, Paleontological Description: Leningrad, Nedra, 200 p. [in Russian]
- Lamarck, J.B.P.A. de M. de, 1819, Histoire Naturelle des Animaux sans Vertèbres: Paris, published by the author, v. 6, pt. 1, p. 1–343.
- Lamprell, K.L., and Healy, J., 1998, Bivalves of Australia: Leiden, Backhuys Publishers, v. 2, 288 p.
- Lamy, E., 1917, Révision des Mactridae vivants du Muséum d'Histoire Naturelle de Paris: Journal de Conchyliologie, v. 63, p. 197–270, 291–411.
- Le Renard, J., and Pacaud, J.M., 1995, Révision des Mollusques paléogènes du Bassin de Paris. 2.—Liste des références primaires des espèces: Cossmanniana, v. 3, p. 65–132.
- Lycett, J., 1848, Notes on the distribution of the fossil conchology of the Oolitic formations in the vicinity of Minchinhampton, Gloucestershire: Annals and Magazine of Natural History (series 2), v. 2, p. 248–259.
- Lynge, H., 1909, The Danish expedition to Siam 1899–1900. IV. Marine Lamellibranchiata: Det Kongelige Danske Videnskabernes Selskabs Skrifter, 7 Raekke, Naturvidenskab og Matematisk Afdeling: Geographical Journal, v. 3, p. 100–299.
- Manzoni, A., 1868, Saggio di Conchiologia Fossile Subappennina: Fauna delle Sabbie Gialle: Imola, Ignazio Galeati e Figlio, 74 p.
- Marcet, I Riba, J., 1955, Síntesis petrográfica y estratigráfica del Eoceno de la zona Palafrugell–Esclañá–Regencós (Bajo Ampurdán, provincia de Gerona): Annals de l'Institut d'Estudis Gironins, v. 10, p. 5–24.
- Marwick, J., 1943, Some Tertiary Mollusca from North Otago: Transactions of the Royal Society of New Zealand, v. 73, p. 181–192.
- Maxwell, P.A., 1978, Taxonomic and nomenclatural notes on some New Zealand Cenozoic Mollusca, with descriptions of new taxa: New Zealand Journal of Zoology, v. 5, p. 15–46.
- Meigen, J.W., 1830, Systematische Beschreibung der bekannten europäischen zweiflügeligen Insekten: Hamm: Schulz–Wundermann, v. 6, 401 p.
- Melville, J.C., and Standen, R., 1907, The Mollusca of the Persian Gulf, Gulf of Oman, and Arabian Sea, as evidenced mainly through the collections of Mr. F. W. Townsend, 1893–1906; with descriptions of new species. Part II. Pelecypoda: Proceedings of Zoological Society of London, v. 1906, p. 783–848.
- Nakagawa, T., 2009, Miocene tropical sandy and rocky bottom molluscan assemblages of the Shimo Formation, Uchiura Group, Fukui Prefecture, Central Japan: Bulletin of the Mizunami Fossil Museum, v. 35, p. 127–151.
- Newton, R.B., 1891, Systematic List of the Frederick E. Edwards Collection of British Oligocene and Eocene Mollusca in the British Museum (Natural History) with References to the Type Specimens from Similar Horizons Contained in Other Collections Belonging to the Geological Department of the Museum: London, Longmans and Co., 365 p.
- Nicklès, M., 1955, Scaphopodes et lamellibranches récoltés dans l'ouest Africain: Atlantide Report (Scientific results of the Danish expedition to the coasts of tropical west Africa (1945–1946)), v. 3, p. 93–237.
- Okamoto, K., and Terachi, M., 1974, Miocene molluscs from the Bihoku group at Miyauchi–Cho, Shobara City, southwest Japan: Transactions and Proceedings of the Palaeontological Society of Japan, n.s., v. 95, p. 341–352.
- Oliver, P.G., 1995, Bivalves (Bivalvia), in Dance, S.P., ed., Seashells of Eastern Arabia: Dubai, Abu Dhabi and London, Motivate Publishing, p. 194–281.
- Oostingh, C.H., 1934, Aanteekeningen over eenige bivalven uit het Neogeen van Java (mit Beschreibung einiger neuer Bivalven aus dem Neogen von Krawang, Java): De Ingenieur in Nederlands–Indie (IV), v. 4, p. 19–22.
- Orbigny, A. d', 1849–1852, Prodrome de Paléontologie Stratigraphique Universelle des Animaux and Mollusques et Rayonnés: Faisant Suite au Cours Élémentaire de Paléontologie et de Geologie Stratigraphiques: Paris, Victor Masson, v. 1, 394 p [1849], v. 2, 427 p. [1850], v. 3, 389 p. [1852].
- Otuka, Y., 1934, Tertiary structures of the northwestern end of the Kitakami mountainland, Iwate Prefecture: Japan, Bulletin of the Earthquake Research Institute, v. 12, p. 566–638.
- Paetel, F., 1875, Die bisher veröffentlichten Familien—und Gattungsnamen der Mollusken: Berlin, Gebrüder Paetel, 229 p.
- Paetel, F., 1883, Catalog der Conchylien–Sammlung: Berlin, Verlag von Gebrüder Paetel, 271 p.
- Pantanelli, D., 1892, Lamellibranchi Pliocenici. Enumerazione e sinonimia delle specie dell' Italia superiore e centrale: Bullettino della Societa Malacologica Italiana, v. 17, p. 1–295.
- Petit, R.E., 2009, George Brettingham Sowerby, I, II & III: their conchological publications and Molluscan taxa: Zootaxa, v. 2189, p. 1–218.
- Pictet, F.J., 1853–1857, Traité de Paléontologie, ou Histoire Naturelle des Animaux Fossiles Considérés dans leurs Rapports Zoologiques et Géologiques: Seconde ed. Paris, London, New York, J.–B. Baillière, v. 1, 584 p., v. 2, 727 p., v. 3, 654 p., v. 4, 768 p., atlas, 77 p.
- Roslim, A., Briguglio, A., Kocsis, L., Cori'c, S., and Razak, H., 2016, Late Miocene fossils from shallow marine sediments in Brunei Darussalam: systematics, palaeoenvironment and ecology: Geophysical Research Abstracts, v. 18, EGU2016–362, 2016 EGU General Assembly 2016.
- Ruhoff, F., 1980, Index to the Species of Mollusca Introduced from 1850 to 1870: Smithsonian Contributions to Zoology, v. 294, p. 1–640.
- Sacco, F., 1901, Parte 29. Donacidae, Psammobiidae, Solenidae, Mesodesmidae, Mactridae, Cardiidae, Myidae, Corbulidae, Glycymeridae, Gastrochaenidae, Pholadidae, Teredinidae, Cryptodontidae, Ungulinidae, Lucinidae, Tellinidae, Scrobulariidae, Cuspidariidae, Solenomyidae, Pandoridae, Verticordiidae, Lyonsiidae, Ceromyidae, Arcomyidae, Anatinidae, Poromyidae, Pholadomyidae e Clavagellidae, in Bellardi, L., and Sacco, F., 1872–1904, I Molluschi dei Terreni Terziari del Piemonte e della Liguria: Torino, C. Clausen, 31 pts., 216 p.
- Saul, L.R., 1973, Evidence for the origin of the Mactridae (Bivalvia) in the Cretaceous: University of California Publications in Geological Sciences, v. 97, p. 1–59.
- Savi, P., and Meneghini, G.G., 1850, Osservazioni stratigrafiche e paleontologiche concernanti la geologia della Toscana e dei paesi limitrofi (Appendix to Murchison, R.I.: Memoria sulla struttura geologica delle Alpi, degli Apennini e dei Carpati): Firenze, Stemparia granulea, 246 p.
- Schultz, O., 2003, Bivalvia neogenica (Lucinoidea–Mactroidea), in Piller, W.E., ed., Catalogus Fossilium Austriae: Wien, Österreichischen Akademie der Wissenschaften, 350 p.
- Sieber, R., 1958, Die Tortonfauna von Steinbrunn bei Drasenhofen (Bez. Mistelbach, N.–Ö): Verhandlungen der Geologischen Bundesanstalt, v. 1958, p. 142–155.
- Signorelli, J.H., and Carter, J.G., 2016, The Anatinellidae and Kymatoxinae: a reassessment of their affinities within the superfamily Mactroidea (Mollusca, Bivalvia): American Malacological Bulletin, v. 33, p. 204–211.
- Smith, E.A., 1906, Natural history notes from R.I.M.S. 'Investigator.'—Series III, No. 10. On Mollusca from the Bay of Bengal and the Arabian Sea: Annals and Magazine of Natural History, v. 18, p. 245–264.
- Sowerby, G.B. II, 1873, Monograph of the genus *Cardilia*, in Reeve, L.A., and Sowerby, G.B., II, eds., 1843–1878, Conchologia Iconica, or, Illustrations of the Shells of Molluscous Animals: London, L. Reeve and Co, v. 19, 1 pl. [unnumbered pages are explanations to plates]
- Stefanini, G., 1916, Fossili del Neogene Veneto: Memorie dell'Istituto Geologico della Regia Università di Padova, v. 4, p. 1–198.
- Studencka, B., Gontsharova, I., and Popov, S., 1998, The bivalve faunas as a basis for reconstruction of the middle Miocene history of the Paratethys: Acta Geologica Polonica, v. 48, p. 285–342.
- Taguchi, E., 2002, Stratigraphy, molluscan fauna and paleoenvironment of the Miocene Katsuta Group in Okayama Prefecture, Southwest Japan: Bulletin of the Mizunami Fossil Museum, v. 29, p. 95–133.
- Takayasu, K., 1981, Fossil Molluscs from the Miocene Masuda Group at Okuda, Masuda City, Molluscan Fossils from Various Localities in Shimane Prefecture, Part 2: Memoirs of the Faculty of Science of Shimane University, v. 15, p. 89–108.
- Taylor, J.D., Kennedy, W.J., and Hall, A., 1973, The shell structure and mineralogy of the Bivalvia. II. Lucinacea–Clavagellacea. Conclusions: Bulletin of the British Museum (Natural History), Zoology, v. 22, p. 253–294.
- Taylor, J.D., Williams, S.T., Glover, E.A., and Dyak, O., 2007, A molecular phylogeny of heterodont bivalves (Mollusca: Bivalvia: Heterodonta): new analyses of 18S and 28S rRNA genes: Zoologica Scripta, v. 36, p. 587–606.
- Tejkal, J., Ondrejčková, A., and Csepregy–Meznerics, I., 1967, Die Mollusken der Karpatischen Serie, in Cicha, I., Seneš, J., and Tejkal, J., eds., Chronostratigraphie und Neostatotypen M₃ (Karpatrien): Slowakische Akademie der Wissenschaften Bratislava, p. 149–212.
- Tennant, J., 1847, A stratigraphical list of British Fossils; arranged under the principal divisions of the British Museum strata, with a few

- elementary remarks on their character and localities: London, Reeve Brothers, 132 p.
- Tsuda, K., 1959, New Miocene molluscs from the Kurosedani Formation in Toyama Prefecture: Japan, Journal of the Faculty of Science, Niigata University series 2, v. 3, p. 67–110.
- Van Regteren, Altena, C.O. and Beets, C., 1945, Eine Neogene Molluskenfauna vom Tji Gugur (Priangan), W. Java. Verhandelingen van het Geologisch-Mijnbouwkundig Genootschap voor Nederland en Koloniën: Geologische serie, v. 14, p. 37–70. [the book carries the date 1944 but according to Winkler Prins, 1996 it appeared in 1945]
- Venzo, S., 1934, Il Neogene del Trentino, del Veronese e del Bresciano: Memorie del Museo di Storia Naturale della Venezia Tridentina, v. 2, p. 1–99.

Accepted 6 August 2017