

The balanced care model for global mental health

G. Thornicroft^{1*} and M. Tansella²

¹ Health Service and Population Research Department, King's College London, Institute of Psychiatry, London, UK

² Section of Psychiatry, Department of Public Health and Community Medicine, University of Verona, Italy

Background. For too long there have been heated debates between those who believe that mental health care should be largely or solely provided from hospitals and those who adhere to the view that community care should fully replace hospitals. The aim of this study was to propose a conceptual model relevant for mental health service development in low-, medium- and high-resource settings worldwide.

Method. We conducted a review of the relevant peer-reviewed evidence and a series of surveys including more than 170 individual experts with direct experience of mental health system change worldwide. We integrated data from these multiple sources to develop the balanced care model (BCM), framed in three sequential steps relevant to different resource settings.

Results. Low-resource settings need to focus on improving the recognition and treatment of people with mental illnesses in primary care. Medium-resource settings in addition can develop 'general adult mental health services', namely (i) out-patient clinics, (ii) community mental health teams (CMHTs), (iii) acute in-patient services, (iv) community residential care and (v) work/occupation. High-resource settings, in addition to primary care and general adult mental health services, can also provide specialized services in these same five categories.

Conclusions. The BCM refers both to a balance between hospital and community care and to a balance between all of the service components (e.g. clinical teams) that are present in any system, whether this is in low-, medium- or high-resource settings. The BCM therefore indicates that a comprehensive mental health system includes both community- and hospital-based components of care.

Received 11 January 2012; Revised 29 May 2012; Accepted 29 May 2012; First published online 11 July 2012

Key words: Community mental health, global mental health, health service research, needs assessment.

Introduction

Most people in the world who have mental illnesses receive no effective treatment (Thornicroft, 2007; Wang *et al.* 2007b; Kessler *et al.* 2009; Patel *et al.* 2010a). For example, of all adults affected by mental illnesses, the proportion who are treated ranges from 30.5% in the USA (Kessler *et al.* 2005b), and 27% across Europe (Wittchen & Jacobi, 2005; Alonso *et al.* 2007), to less than 1% in Nigeria (Demyttenaere *et al.* 2004; Kohn *et al.* 2004; Ormel *et al.* 2008; Wang *et al.* 2007a). This phenomenon, described by the World Health Organization (WHO) as the 'treatment gap' (Dua *et al.* 2011), is increasingly appreciated worldwide, and is seen as the difference between the true prevalence rate and the proportion who receive any kind of treatment (except for interventions by traditional, religious or similar

practitioners/healers) (Chisholm *et al.* 2007; Patel *et al.* 2007; Prince *et al.* 2007; Saxena *et al.* 2007) (see Table 1).

In 2008 the Department of Mental Health and Substance Abuse at the WHO recognized the importance of this challenge by launching as its centrepiece the Mental Health Global Action Programme (mhGAP). The first major product of this programme is the *mhGAP Intervention Guide* (mhGAP IG; WHO, 2010), which contains case findings and treatment guidelines for nine important categories of mental and neurological disorders that are particularly relevant in low-income settings, and that have a major global public health impact. The conditions included are: depression, psychoses, epilepsy/seizures, developmental disorders, behaviour disorders, dementia, alcohol use disorders, drug use disorders, and self-harm/suicide (Barbui *et al.* 2010).

In this paper we propose a 'Balanced Care Model' (BCM) to stimulate debate about how to conceptualize the planning and delivery of services to adults with mental disorders. For the purposes of this conceptual paper we refer to services for the range of mental disorders that affect adults, excluding drug and alcohol

* Address for correspondence: G. Thornicroft, Professor of Community Psychiatry, Health Service and Population Research Department, King's College London, Institute of Psychiatry, De Crespigny Park, London SE5 8AF, UK.
(Email: graham.thornicroft@kcl.ac.uk)

Table 1. Proportions (%) of people with key physical and mental disorders who are treated, by high- or low- and middle-income setting status (adapted from Ormel *et al.* 2008)

	High-income settings	Low- and middle-income settings
Physical disorders		
Diabetes	94	77
Heart disease	78	51
Asthma	65	44
Mental disorders		
Depression	29	8
Bipolar disorder	29	13
Panic disorder	33	9

misuse disorders, intellectual disabilities or any neurological disorders, such as epilepsy, that are commonly treated within mental health services in low-income countries, or services specifically for older adults. We have previously reviewed the evidence for hospital-based and community-based general adult mental health services, and we indicated that a comprehensive system of care should be based on a balance of both hospital and community components (Thornicroft & Tansella, 2004). Almost a decade later, is this still relevant?

Method

To develop this conceptual model we have integrated information from two sources: (i) a literature review of relevant peer-reviewed papers and other relevant published material and (ii) a consensus exercise conducted among experts in mental health services in many countries worldwide, supplemented by additional information from data sources in low- and middle-income countries (LMICs). The rationale for this hybrid approach is that our aim is to develop a high-level conceptual model that can be applied, for example, at country, regional or district levels, and that does not only include data from clinical studies, for example from randomized clinical trials.

For the first source of data, the published material, in 2011 we updated the literature review we initially conducted in 2004 (Thornicroft & Tansella, 2004). Primary data papers, along with review and discussion papers published in English, were examined from relevant journals and book chapters. The key search terms and combinations used were: (1) 'delivery of health care', (2) 'mental health', (3) 'mental OR psychiatric OR psychiatry', (4) 2 OR 3, (5) 4 AND 1. The only studies excluded were those directly

concerned with children/adolescents/young people, older adults, dementia, intellectual disability, alcohol and drug misuse, prison populations or forensic psychiatry, or homeless populations. Then the combined search categories were applied to Medline, EMBASE and PsycINFO (in relation to: title, abstract, subject headings, heading word, drug trade name, original title, device manufacturer, or drug manufacturer). This general adult version of the BCM can therefore be considered as a prototype model that may be suitable, in adapted or modified forms, to these other patients groups.

To identify any pertinent missing papers we cross-referenced the results of these searches with the peer-reviewed paper references contained in the relevant review papers by international experts who contributed chapters to an international textbook on community mental health (Thornicroft *et al.* 2011c). In addition, we accessed other electronic, non-indexed sources including the World Psychiatric Association (WPA) and country-specific Ministry of Health websites for relevant literature. Google was used to search for pdf documents that contained 'community mental health'. Other key texts were used to identify relevant papers and book chapters (Benegal *et al.* 2009; Mbuba & Newton, 2009; Patel & Thornicroft, 2009; Patel *et al.* 2009) along with sources from the 2007 special edition of *The Lancet* on Global Mental Health (Chisholm *et al.* 2007; Jacob *et al.* 2007; Patel *et al.* 2007; Prince *et al.* 2007; Saraceno *et al.* 2007; Saxena *et al.* 2007). WHO publications that provide information on community mental health services worldwide were also sourced (WHO, 2008).

For the second source of information used in developing this conceptual model, information from experts in many countries worldwide, we first contacted colleagues from across Africa, the Americas, Asia, Australasia and Europe whom we knew to be active in mental health service design in their respective countries. Using a semi-structured questionnaire we asked them to comment on 10 key challenges that we proposed in relation to mental health service development. A more detailed account of the method used has been published previously (Thornicroft & Tansella, 2009). To further broaden the scope of information gathered about international experience in developing mental health care, this review also drew upon the work of the WPA Task Force (2009–2010) on 'Steps, obstacles and mistakes to avoid in the implementation of community mental health care' (see Acknowledgements for membership). Task Force members were from seven world regions as defined by the WHO. Each member then contacted, in turn, colleagues with experience of developing mental health services within their own region of the world.

To further strengthen the information contributed from LMICs, we drew upon data gathered to assess how far mental health services in LMICs have been scaled up in recent years to reduce the treatment gap (Eaton *et al.* 2011). We also went further by including more specific data from the particular LMICs in sub-Saharan Africa. Within the work of the WPA Task Force, 21 experts completed a semi-structured questionnaire about their experience in implementing community mental health care in Cote d'Ivoire, Ethiopia, Kenya, Liberia, Malawi, Niger, Nigeria, South Africa, Sudan, Tanzania, Uganda and Zimbabwe (Hanlon *et al.* 2010; Thornicroft *et al.* 2011*b*). Using our pre-existing understanding of mental health systems (Thornicroft & Tansella, 1999), we integrated data from these multiple sources to develop the BCM, framed in three sequential steps relevant to low-, medium- or high-income settings and intended to be simple enough to be widely applicable, and flexible enough to be adapted to the complexities of local circumstances.

Results

In relation to the main literature search, the initial 55211 papers identified were reduced to 118 unique records when the eligibility criteria were applied. For the information gathered from experts worldwide, the first survey drew upon detailed responses from 27 colleagues in 25 countries worldwide (Thornicroft & Tansella, 2009), with the WPA Task Force comprising members from nine countries across seven world regions as defined by the WHO. In relation to the survey assessing progress made towards scaling up (Eaton *et al.* 2011), data were contributed by 142 experts from 59 countries worldwide, of which 19 (32%) were in the WHO Africa region, 16 (27%) in the Americas region, eight (14%) in the eastern Mediterranean region, six (10%) in the western Pacific, five (8%) in South East Asia, and five (8%) in Europe. In total, therefore, more than 170 individual experts worldwide actively contributed to this data collection process.

We present the BCM in relation to three 'levels of resources', using the World Bank classification (World Bank, 2010). In this system economies are divided according to 2009 Gross National Income (GNI) *per capita* calculated using the World Bank Atlas method. The groups are: low income (\leq US\$995), lower-middle income (US\$996–3945), upper-middle income (US\$3946–12195), and high income (\geq US\$12196). For the purposes of the BCM we have combined the lower-middle and upper-middle income setting groups.

The mental health resource disparities between low- and high-income settings are vast. In low-income

countries, for example, there are on average only 0.05 psychiatrists and 0.16 psychiatric nurses per 100 000 population, about 200 times less than in high-income settings (WHO, 2005). Many low-income countries in sub-Saharan Africa, for example, have on average less than one psychiatrist for every million people (e.g. Chad, Eritrea and Liberia), compared with 137 per million in the USA (Ndyanabangi *et al.* 2004; Miller, 2006). Furthermore, training programmes and facilities for mental health professionals in low-income settings are often grossly inadequate (WHO, 2005; Saxena *et al.* 2007). The BCM therefore organizes service components separately according to low-, middle- or high-resource settings as shown in Fig. 1 (Thornicroft & Tansella, unpublished observations).

Low-income settings

Most of the available provision in low-resource settings is by staff in primary health care and community settings (Ormel *et al.* 1994; Desjarlais *et al.* 1995; WHO, 2001; Seloilwe & Thupayagale-Tshweneagae, 2007; Deva, 2008). The roles of these staff include: case finding and assessment, brief talking and psychosocial treatments, and pharmacological treatments (Beaglehole & Bonita, 2008; Eaton, 2008). The very limited numbers of specialist mental health care staff (usually in the capital city and sometimes also in regional centres) are only able to provide: (i) training and supervision of primary care staff, (ii) consultation–liaison for complex cases, and (iii) out-patient and in-patient and assessment and treatment for cases that cannot be managed in primary care (Mubbashar, 1999; Alem, 2002; Njenga, 2002; Saxena & Maulik, 2003; Lund *et al.* 2011).

Medium-income settings

For medium-income settings it is important to appreciate that there is still a requirement for a strong primary care level of provision, so as to address the high levels of prevalence of common mental disorders in the general population (in many countries estimated at 20–30% annual period prevalence rate) (Kessler *et al.* 2005*a*; Wittchen *et al.* 2011). The literature from such middle-income settings, for example many of the countries of Eastern Europe and South America (Knapp *et al.* 2007; Semrau *et al.* 2011; Razzouk *et al.*, in press), indicates that modest levels of resource are usually allocated for mental health care compared with communicable and infectious diseases (Razali, 2004; Al-Krenawi, 2005; Furedi *et al.* 2006; Akiyama *et al.* 2008; Janse van Rensburg, 2009; Sharifi, 2009; Rodriguez, 2010). In addition, as resources allow, the BCM indicates that the five elements of




Fig. 1. Mental health service components relevant to low-, medium- and high-resource settings.

'general adult' mental health services are advisable, as discussed in the following sections.

Out-patient/ambulatory clinics

There is surprisingly little evidence on the effectiveness of out-patient, clinic or ambulatory care (Becker & Koesters, 2011) but there is a strong clinical consensus in many countries that they are a relatively efficient way to organize the provision of assessment and treatment, providing that the clinic sites are accessible to local populations. Nevertheless, these clinics are simply methods of arranging clinical contact between staff and patients, and so the key issue is the content of the clinical interventions, namely to deliver treatments that are effective (Roth & Fonagy, 1996; Nathan & Gorman, 2002; BMJ Books, 2003).

Community mental health teams (CMHTs)

CMHTs are the basic building block of community mental health services. The simplest model of provision of community care is for generic (non-specialized) CMHTs to provide the full range of interventions, staffed by multi-disciplinary personnel. These often prioritize adults with severe mental illness, for a local defined geographical catchment area (Tyrer *et al.* 1995, 1998, 2007; Sytema *et al.* 1997; Thornicroft *et al.* 1998, 1999; Burns, 2001; Simmonds

et al. 2001; Department of Health, 2002). The central issue here is that CMHTs can offer case management and continuity of care (Dieterich *et al.* 2010), in addition to mobility. In other words they can arrange appointments with patients at hospitals, clinics, community mental health centres, or at the patient's own homes. At the same time it needs to be recognized that, for patients not able or not willing to go to health facilities, this flexibility is necessary but not sufficient for proper care. Alongside the need for mobility is once again the requirement to deliver effective treatment when clinical encounters do take place (Malone *et al.* 2007).

Acute in-patient care

There continues to be relatively weak evidence about most aspects of in-patient care, and these studies are usually descriptive accounts (Holloway & Sederer, 2011). There are few systematic reviews in this field, one of which found that there were no differences in outcomes between routine admissions and planned shorter hospital stays (Johnstone & Zolse, 1999). More generally, although there is a consensus that acute in-patient services are necessary, the number of beds provided is highly contingent upon which other services exist locally, and upon local social, economic and cultural characteristics (Thornicroft & Tansella, 1999). Acute in-patient care commonly absorbs most

of the mental health budget (Knapp *et al.* 1997), therefore reducing the average length of stay may be an important system goal, especially if the resources released in this way can be used to pay for other service components (Lasalvia & Tansella, 2010; Lelliott & Bleksley, 2010; Sederer, 2010; Totman *et al.* 2010).

A related policy issue concerns how to provide acute beds in a humane and non-institutionalized way that is acceptable to patients, for example in general hospital units (Quirk & Lelliott, 2001; Tomov, 2001; Totman *et al.* 2010; The ITHACA Study Group, 2011). For example, descriptive research in England has identified 131 services that are alternative to traditional acute in-patient settings. Most were hospital based and situated in deprived areas, and about half were established after 2000. This suggests that such alternatives may represent an important, but previously undocumented and unevaluated, sector of the mental health economy (Johnson *et al.* 2009; Lloyd-Evans *et al.* 2009).

Long-term community-based residential care

It is important to know whether patients with severe and long-term disabilities should be cared for in larger, traditional institutions, or be transferred to long-term community-based residential care. Although there is no strong evidence on this question from low-income settings, the evidence from medium- and high-income settings is reasonably clear. When deinstitutionalization is carried out carefully, when patients who have previously received long-term in-patient care for many years are discharged to community care, then the outcomes are favourable for the majority (Shepherd & MacPherson, 2011). Nevertheless, the range and capacity of community residential long-term care that will be needed in any particular area is also highly dependent upon which other services are available locally, and upon social and cultural factors, such as the amount of family care available (van Wijngaarden *et al.* 2003).

Work and occupation

Rates of unemployment among people with mental disorders are usually much higher than in the general population (Warner, 2004). Traditional methods of occupation have not been shown to be effective in leading to open market employment (Shepherd, 1990; Marshall *et al.* 2001; Rosen & Barfoot, 2001). For settings with medium levels of resources it is reasonable at this stage to make pragmatic decisions about the provision of work and day care services, especially based upon the priorities and preferences of the patient/service user and carer/family members concerned (Cleary *et al.* 2006), where this is focusing

increasingly upon the importance of personal recovery (Slade, 2009; Slade *et al.* 2012). At the same time, it is reasonable to take into account the accumulating evidence for supported employment models (Marshall *et al.* 2001; Catty *et al.* 2007; Becker *et al.* 2011).

High-income settings

Superimposed upon a basic primary care system (Gask, 2005), and in addition to the provision of general adult mental health services, for high-income settings the application of the BCM implies that a series of specialized services can be provided, as resources allow (see Fig. 1). In fact, however, it is often the case that specialized services are developed in the absence of the first two layers of general services. This is often because advocates for a new team or service take a 'component view' of treatment rather than a public health orientation, using a 'system view' of the wider pattern of care and how the constituent parts contribute to the whole.

Such specialized services can be developed in the same five categories described earlier for medium-income settings.

Specialized out-patient/ambulatory clinics

Specialized out-patient facilities may provide services, for example, for those with eating disorders, treatment-resistant affective disorders, people with comorbid psychotic and substance misuse/dependence disorders, or for mentally ill mothers. Local decisions about whether to establish such specialist clinics will depend upon several factors, including their relative priority in relation to the other specialist services described below, identified services gaps, and the financial opportunities available (Becker & Koesters, 2011).

Specialized CMHTs

Although in-patient and out-patient care have rarely been evaluated, CMHTs continue to be subject to substantial research investigation.

Assertive community treatment (ACT) teams. These provide a form of specialized mobile outreach treatment for people with more disabling mental disorders (Deci *et al.* 1995; Teague *et al.* 1998; Killaspy & Rosen, 2011). There is now clear evidence that in high-income settings ACT can: reduce admissions to hospital and the use of acute beds; improve accommodation status and occupation; and increase service user satisfaction. ACT has not been shown to produce improvements in mental state or social behaviour. ACT can reduce the cost of in-patient services but does not change the

overall costs of care (Latimer, 1999; Marshall & Lockwood, 2000; Phillips *et al.* 2001).

Nevertheless, it is not known how far ACT is cross-culturally relevant, and indeed there is evidence that ACT may be less effective where usual services already offer high levels of continuity of care (Burns *et al.* 1999, 2001; Fiander *et al.* 2003). Indeed, data from outside the USA are somewhat equivocal about the benefits of ACT. A randomized controlled trial (RCT) in the UK (Killaspy *et al.* 2006, 2009; McCrone *et al.* 2009b) found no differences in any measure of in-patient service use, clinical or social functioning between ACT and standard community mental health team case management. However, patients treated by ACT were better engaged, less likely to drop out of contact and more satisfied with services. It may be that the lesser benefits from the UK studies are attributable to better functioning CMHTs in the control group than is the case in studies in the USA (Latimer, 2005; Burns *et al.* 2007; Ghosh & Killaspy, 2010; Killaspy & Rosen, 2011).

Early intervention (EI) teams. There is a growing body of research evidence on the short- to medium-term consequences of EI CMHTs in treating people with psychotic disorders (Power & McGorry, 2011), and also services to treat prodromal or ultra-high-risk groups (Addington *et al.* 2003; Raune *et al.* 2004; Preti & Cella, 2010). The more recent evidence gives conflicting views on whether EI services do deliver enduring patient benefit (Killackey, 2011; Larsen *et al.* 2011) or whether the advantages may only last while in contact with such relatively intense services (Bosanac *et al.* 2010; Gafoor *et al.* 2010). The most recent Cochrane systematic review (Marshall & Rathbone, 2011) concluded that 'There is emerging, but as yet inconclusive evidence, to suggest that people in the prodrome of psychosis can be helped by some interventions. There is some support for specialized early intervention services, but further trials would be desirable, and there is a question of whether gains are maintained'. This conclusion has been supported by some commentators (Castle, 2012) but challenged by others as being based upon too narrow a selection of the relevant literature (Power & McGorry, 2011; McGorry, 2012). It is therefore still too early to judge whether specialized EI teams should be seen as a priority in high-income settings (de Girolamo *et al.* 2012).

Alternatives to acute in-patient care

In recent years three main alternatives to acute in-patient care have been developed: acute day hospitals,

crisis houses, and home treatment/crisis resolution teams.

Acute day hospitals. These facilities offer programmes of day treatment for people with acute and severe psychiatric problems, as an alternative to admission to in-patient units. A recent systematic review found no clear difference between acute day hospital and out-patient care for the outcomes of: 'lost to follow-up', social functioning, or for mental state outcomes, a considerably less optimistic conclusion than previous reviews of acute day hospitals (Marshall *et al.* 2001, 2011).

Crisis houses. These are houses in community settings that are staffed by trained mental health professionals and offer admission for some patients who would otherwise be admitted acutely to hospital. A wide variety of respite houses, havens and refuges have been developed, but crisis house is used here to mean facilities that are alternatives to non-compulsory hospital admission. The limited available research evidence suggests that they are usually very acceptable to their residents (Davies *et al.* 1994), may be able to offer an alternative to hospital admission for about a quarter of otherwise admitted patients, and may be more cost-effective than hospital admission (Sledge *et al.* 1996a,b; Mosher, 1999). Furthermore, there is emerging evidence that female patients in particular prefer non-hospital alternatives (such as single-sex crisis houses) to acute in-patient treatment, and this may reflect the lack of perceived safety in those settings (Killaspy *et al.* 2000; Howard *et al.* 2008; Howard *et al.* 2009; Lawlor *et al.* 2010).

Home treatment/crisis resolution teams. These are mobile community mental health teams offering assessment for patients in psychiatric crises, and providing intensive treatment and care at home (Johnson *et al.* 2008, 2011). A recent Cochrane systematic review (Murphy *et al.* 2012) reported that 'care based on crisis intervention principles, with or without an ongoing home care package, appears to be a viable and acceptable way of treating people with serious mental illnesses'. Such studies have tended to find that crisis home care is more cost-effective than hospital care but data were often skewed (McCrone *et al.* 2009a). There were no data on staff satisfaction, carer input, compliance with medication, or the subsequent number of relapses. The authors concluded that home care crisis treatment, coupled with an ongoing home care package, is a viable and acceptable way of treating people with serious mental illnesses (Joy *et al.* 2006). A more widely based review (Johnson *et al.* 2008) has come to similar conclusions, namely that such teams can

reduce hospital admissions (especially voluntary admissions), may be associated with some reduction of overall treatment costs, and seem to be compatible with reasonable levels of staff morale (Johnson *et al.* 2005*a,b*; Glover *et al.* 2006; Jethwa *et al.* 2007; Keown *et al.* 2007; Johnson & Thornicroft, 2008; McCrone *et al.* 2009*c*; Nelson *et al.* 2009).

Alternative types of long-stay community residential care

These are usually replacements for long-stay wards in psychiatric institutions (Shepherd & MacPherson, 2011). Three main categories of such residential care can be identified: (i) 24-hour staffed residential care (high-staffed hostels, residential care homes or nursing homes, depending on whether the staff have professional qualifications; (ii) day-staffed residential facilities (hostels or residential homes that are staffed during the day); and (iii) lower supported accommodation (minimally supported hostels or residential homes with visiting staff). There is some evidence on the effectiveness and the cost-effectiveness of these types of residential care, but there are no systematic reviews (Thornicroft *et al.* 2005; Chilvers *et al.* 2006). It is therefore reasonable for policy makers to decide upon the need for such services with local stakeholders.

Specialized forms of work and occupation

Work represents an important goal for many people with severe mental illnesses (Lehman, 1995), but in high-income settings rates of unemployment among people with severe mental illness often exceed 90% (Thornicroft *et al.* 2004; Marwaha & Johnson, 2005). Furthermore, consumer and carer advocacy groups usually set work/occupation as one of their highest priorities, to enhance both functional status and quality of life (Becker *et al.* 1996; Thornicroft *et al.* 2002). Several RCTs, predominantly in the USA, have found that supported employment using the Individual Placement and Support (IPS) model can increase rates of competitive employment to 30–60% (Priebe *et al.* 1998; Drake *et al.* 1999; Crowther *et al.* 2001; Marshall *et al.* 2001; Lehman *et al.* 2002; Rinaldi & Perkins, 2007; Bond *et al.* 2008). This model is now also often referred to as ‘supported employment’. This model was elaborated in the 1990s and includes vocational rehabilitation as part of mental health treatment, rather than a separate entity. Its aim is to achieve rapid job placement into competitive employment (i.e. in the open labour market), followed by support and necessary training obtained while in the job. Supported employment is intended to be integrated within community mental health services and to be based on patients’ preferences.

RCTs in the USA have found this model of vocational rehabilitation to be more effective in gaining employment for people with severe mental illness compared with traditional vocational rehabilitation models. This has also been replicated outside the USA. One European study found favourable outcomes for IPS in countries with diverse labour markets and different welfare systems (Burns & Catty, 2008), but IPS was not effective in all the countries studied. A further UK RCT, the Supported Work and Needs (SWAN) Trial (Howard *et al.* 2010), found that the rate of employment at the 1-year follow-up was low for both the intervention group (13%) and the control group (7%), with no significant difference between them. At the 2-year follow-up, however, the difference was significant (22% *v.* 11% respectively) (Heslin *et al.* 2011). It is currently unclear, therefore, whether the IPS model is effective in populations with high rates of unemployment, and with relatively generous state benefits for unemployed people with severe mental illness (Heffernan & Pilkington, 2011). A potentially important variation of IPS is the employment of peer support workers, which shows some potential to assist patients in job finding and keeping (Nestor & Galletly, 2008; Robinson *et al.* 2010; Repper & Carter, 2011).

Discussion

In a previous paper on mental health service planning (Thornicroft & Tansella, 2004), we described a pragmatic balance of hospital and community care. In the current paper we have extended this concept to refer also to a balance between all of the service components (e.g. clinical teams) that are present in any system, whether this is in a low-, medium- or high-resource setting (see Fig. 1).

In low-resource settings, as an illustration, the crucial resource allocation decisions will be how to balance any investment in primary and community care sites against expenditure in psychiatric hospitals. In medium-resource settings the BCM approach proposes that services are provided in all of the five categories of care. If no provision for employment, or for community-based residential care, for example, is made, then in our view this is not a comprehensive and balanced system of care. In high-resource settings, these complex choices apply to an even greater extent, as there are even more mental health teams and agencies present, and so there are a greater number of possibilities for resource investment to achieve a balanced portfolio. Another important issue in high-resource settings is that a tendency to provide more specialized teams (e.g. CMHTs) leads to a situation in which there are many more interfaces to manage between such teams, in addition to the potential for more

points of overlap or dysfunction (e.g. poor referral pathways), and also a greater risk of patient experience of low continuity of care (Thornicroft *et al.* 1999).

There are several limitations to the method used in this paper. First, because the model has been developed to apply to higher-order levels of the health service, for example across a whole district or region, it cannot be derived from evaluations of individual service components. For this reason the BCM is not based on a systematic review of RCTs, but does draw upon a wider range of relevant publications and forms of expert experience and advice. There is therefore the risk that our selection of these sources of information, and their interpretation, may be biased. We have attempted to mitigate this risk by drawing upon multiple, independent sources of validation of the model, as described in the method. We propose the BCM to generate debate about whether this conceptual model is clear and practicable, and about how it can be modified as a whole, and adapted to differing local situations.

Furthermore, this paper sets out a somewhat oversimplified and linear view of how services can be developed in an additive and sequential way, building layer upon layer of increasingly differentiated services. In many places the reality on the ground is more piecemeal and more complex than this. Influential advocates may force the creation of particular services, for example because of their own clinical, research or commercial interests, without reference to wider public mental health needs.

This model also takes no account of whether the providers of services are state funded, are from the non-governmental sector, or are services run for profit. Indeed it is notable that publications about mental health service development or evaluation rarely mention the voluntary sector or the for-profit sector. There is a palpable lack of consideration in service planning about if and how for-profit and not-for-profit services can be mutually complementary within a wider system of care (Younes *et al.* 2005; Badrakalimuthu *et al.* 2009; Pollock, 2010; Wahlbeck *et al.* 2011).

In discussing the primary and community level of care, in this paper we have also not addressed very important questions regarding traditional, religious, alternative and complementary practitioners and healers (Ngoma *et al.* 2003; Shankar *et al.* 2006; Abbo *et al.* 2009). These issues are outside the scope of this review, but several considerations need to be taken into account: (i) such practitioners are relatively common across low-, medium- and high-resource settings; (ii) there is little published evidence on the outcomes of most of these interventions; and (iii) little is known about the population-level coverage provided in aggregate by these practitioners. For this latter reason it

is possible that this level of care may come to be considered as the first point of contact for people with mental illness in many settings (Raguram *et al.* 2002; Seloilwe & Thupayagale-Tshweneagae, 2007; Shibre *et al.* 2008; Ae-Ngibise *et al.* 2010; Campbell-Hall *et al.* 2010).

A further issue that we have not addressed in this paper is whether community mental health services provide worse or better physical health care than a hospital-based system (Roberts *et al.* 2007). It is now known that life expectancy for people with a wide range of mental illnesses is, even in relatively high-quality medical systems, up to 20 years less than for the general population (Amaddeo *et al.* 2007; Amaddeo, 2008; Grigoletti *et al.* 2009; Tiihonen *et al.* 2009; Amaddeo & Tansella, 2010). Yet at present there are no clear data that community-orientated services are associated with better physical health outcomes, or with a decrease in such mortality disparities, although two studies have shown a reduction in suicide rates (Pirkola *et al.* 2009; While *et al.* 2012).

A crucial structural issue in all settings is how far primary care staff see it as their responsibility to identify and treat cases of mental illness in their local communities. This approach has been assumed to be a vital initial stage of the care pathway since the Declaration of Alma-Ata (Passmore, 1979). Yet there is now strong evidence that primary care services in many countries provide little or no treatment to people with mental disorders (Greenhalgh, 2008; Lawn *et al.* 2008; Rohde *et al.* 2008; Wang *et al.* 2007a). In this case it may now be timely to consider new ways to provide sufficient staff capacity and training at the community level to identify and treat people with common mental disorders (Patel *et al.* 2010b), for example by using the mhGAP IG recently developed by the WHO (2010).

Conclusions

Does the BCM provide an approach to mental health service improvement that is relevant and, if so, how can it be implemented (Yamey, 2011)? In recent years an increasingly detailed appreciation has developed about the barriers that impede the implementation of evidence-based policies and practices (Saraceno *et al.* 2007), and about methods that can be used to successfully overcome these barriers (Institute of Medicine, 2001; Thornicroft *et al.* 2008, 2010, 2011a; Tansella & Thornicroft, 2009; Drake & Latimer, 2011; Semrau *et al.* 2011; Ito *et al.*, in press; McGeorge, in press; Razzouk *et al.*, in press; Thara & Padvatati, in press). In the future, therefore, it will be necessary to have available not only conceptual models that guide

planning but also pragmatic models that clearly guide implementation.

Acknowledgements

We thank all of the people who contributed to this paper, including A. A. Abdullah, T. Becker, C. K. Yoon, F. Crowley, C. C. Villares, N. Daumerie, I. De Coster, M. Freeman, N. Gionakis, P. G. Gökalp, S. Grozavu, L. Hansson, J. Harangozo, U. Junghan, Y. Kalakoutas, A. Latypov, B. Makenbaeva, G. Mellsop, R. Mezzina, P. Nawka, J. L. Roelandt, V. Švab, M. Taube, R. Teodorescu, R. Thom, C. Van Audenhove, J. van Weeghel, K. Wahlbeck, R. Warner and S. Weinmann, who have directly contributed to the review of experience gained in making mental health services changes. We also thank the members of the WPA Task Force (2009–2010) on 'Steps, obstacles and mistakes to avoid in the implementation of community mental health care', namely, A. Alem, R. A. Dos Santos, E. Barley, R. E. Drake, G. Gregorio, C. Hanlon, H. Ito, E. Latimer, A. Law, J. Mari, P. McGeorge, R. Padmavati, D. Razzouk, M. Semrau, Y. Setoya, R. Thara and D. Wondimagegn.

We also acknowledge the important contributions to this paper of E. Barley and M. Semrau. G.T. is funded by a National Institute for Health Research (NIHR) Applied Programme grant awarded to the South London and Maudsley National Health Service (NHS) Foundation Trust, and to the NIHR Specialist Mental Health Biomedical Research Centre at the Institute of Psychiatry, King's College London and the South London and Maudsley NHS Foundation Trust. All opinions expressed here are solely those of the authors.

Declaration of Interest

None.

References

- Abbo C, Ekblad S, Waako P, Okello E, Musisi S** (2009). The prevalence and severity of mental illnesses handled by traditional healers in two districts in Uganda. *African Health Sciences* 9 (Suppl. 1), S16–S22.
- Addington J, Coldham EL, Jones B, Ko T, Addington D** (2003). The first episode of psychosis: the experience of relatives. *Acta Psychiatrica Scandinavica* 108, 285–289.
- Ae-Ngibise K, Cooper S, Adiibokah E, Akpalu B, Lund C, Doku V; Mhapp Research Programme Consortium** (2010). 'Whether you like it or not people with mental problems are going to go to them': a qualitative exploration into the widespread use of traditional and faith

- healers in the provision of mental health care in Ghana. *International Review of Psychiatry* 22, 558–567.
- Akiyama T, Chandra N, Chen CN, Ganesan M, Koyama A, Kua EE, Lee MS, Lin CY, Ng C, Setoya Y, Takeshima T, Zou Y** (2008). Asian models of excellence in psychiatric care and rehabilitation [Review]. *International Review of Psychiatry* 20, 445–451.
- Alem A** (2002). Community-based vs. hospital-based mental health care: the case of Africa. *World Psychiatry* 1, 99–100.
- Al-Krenawi A** (2005). Mental health practice in Arab countries. *Current Opinion in Psychiatry* 18, 560–564.
- Alonso J, Codony M, Kovess V, Angermeyer MC, Katz SJ, Haro JM, De Girolamo G, De Graaf R, Demyttenaere K, Vilagut G, Almansa J, Lépine JP, Brugha TS** (2007). Population level of unmet need for mental healthcare in Europe. *British Journal of Psychiatry* 190, 299–306.
- Amaddeo F** (2008). Risk of death from natural causes increased in people with psychiatric admissions. *Evidence-Based Mental Health* 11, 106.
- Amaddeo F, Barbui C, Perini G, Biggeri A, Tansella M** (2007). Avoidable mortality of psychiatric patients in an area with a community-based system of mental health care. *Acta Psychiatrica Scandinavica* 115, 320–325.
- Amaddeo F, Tansella M** (2010). Mortality among people with mental disorders. *Epidemiologia e Psichiatria Sociale* 19, 1–3.
- Badrakalimuthu VR, Rangasamy S, Sathyavathy V** (2009). Mental health practice in private primary care in rural India: a survey of practitioners. *World Psychiatry* 8, 124–125.
- Barbui C, Dua T, van Ommeren M, Yasamy M, Fleischmann A, Clark N, Thornicroft G, Hill S, Saxena S** (2010). Challenges in developing evidence-based recommendations using the GRADE approach: the case of mental, neurological, and substance use disorders. *PLoS Medicine* 7, e1000322.
- Beaglehole R, Bonita R** (2008). Global public health: a scorecard. *Lancet* 372, 1988–1996.
- Becker DR, Bond GR, Drake RE** (2011). Individual placement and support: the evidence-based practice of supported employment. In *Oxford Textbook of Community Mental Health* (ed. G. Thornicroft, G. I. Szmukler, K. T. Mueser and R. E. Drake), pp. 204–217. Oxford University Press: Oxford.
- Becker DR, Drake RE, Farabaugh A, Bond GR** (1996). Job preferences of clients with severe psychiatric disorders participating in supported employment programs. *Psychiatric Services* 47, 1223–1226.
- Becker T, Koesters M** (2011). Psychiatric outpatient clinics. In *Oxford Textbook of Community Mental Health* (ed. G. Thornicroft, G. I. Szmukler, K. T. Mueser and R. E. Drake), pp. 179–191. Oxford University Press: Oxford.
- Benegal V, Chand PK, Obot IS** (2009). Packages of care for alcohol use disorders in low- and middle-income countries. *PLoS Medicine* 6, e1000170.
- BMJ Books** (2003). *Clinical Evidence*, Vol. 9. BMJ Books: London.
- Bond GR, Drake RE, Becker DR** (2008). An update on randomized controlled trials of evidence-based

- supported employment. *Psychiatric Rehabilitation Journal* 31, 280–290.
- Bosanac P, Patton GC, Castle DJ** (2010). Early intervention in psychotic disorders: faith before facts? *Psychological Medicine* 40, 353–358.
- Burns T** (2001). Generic versus specialist mental health teams. In *Textbook of Community Psychiatry* (ed. G. Thornicroft and G. Szumukler), pp. 231–241. Oxford University Press: Oxford.
- Burns T, Catty J** (2008). IPS in Europe: the EQOLISE trial. *Psychiatric Rehabilitation Journal* 31, 313–317.
- Burns T, Catty J, Dash M, Roberts C, Lockwood A, Marshall M** (2007). Use of intensive case management to reduce time in hospital in people with severe mental illness: systematic review and meta-regression. *British Medical Journal* 335, 336.
- Burns T, Creed F, Fahy T, Thompson S, Tyrer P, White I** (1999). Intensive versus standard case management for severe psychotic illness: a randomised trial. UK 700 Group. *Lancet* 353, 2185–2189.
- Burns T, Fioritti A, Holloway F, Malm U, Rossler W** (2001). Case management and assertive community treatment in Europe. *Psychiatric Services* 52, 631–636.
- Campbell-Hall V, Petersen I, Bhana A, Mjadu S, Hosegood V, Flisher AJ** (2010). Collaboration between traditional practitioners and primary health care staff in South Africa: developing a workable partnership for community mental health services. *Transcultural Psychiatry* 47, 610–628.
- Castle DJ** (2012). The truth, and nothing but the truth, about early intervention in psychosis. *Australian and New Zealand Journal of Psychiatry* 46, 10–13.
- Catty J, Burns T, Comas A** (2007). Day centres for severe mental illness. *Cochrane Database of Systematic Reviews* 1, CD001710.
- Chilvers R, Macdonald G, Hayes A** (2006). Supported housing for people with severe mental disorders. *Cochrane Database of Systematic Reviews* 4, CD000453.
- Chisholm D, Flisher AJ, Lund C, Patel V, Saxena S, Thornicroft G, Tomlinson M** (2007). Scale up services for mental disorders: a call for action. *Lancet* 370, 1241–1252.
- Cleary M, Freeman A, Walter G** (2006). Carer participation in mental health service delivery [Review]. *International Journal of Mental Health Nursing* 15, 189–194.
- Crowther RE, Marshall M, Bond GR, Huxley P** (2001). Helping people with severe mental illness to obtain work: systematic review. *British Medical Journal* 322, 204–208.
- Davies S, Presilla B, Strathdee G, Thornicroft G** (1994). Community beds: the future for mental health care? *Social Psychiatry and Psychiatric Epidemiology* 29, 241–243.
- Deci PA, Santos AB, Hiott DW, Schoenwald S, Dias JK** (1995). Dissemination of assertive community treatment programs. *Psychiatric Services* 46, 676–678.
- de Girolamo G, Dagani J, Purcell R, Cocchi A, McGorry P** (2012). Age of onset of mental disorders and use of mental health services: needs, opportunities and obstacles. *Epidemiology and Psychiatric Sciences* 21, 47–57.
- Demyttenaere K, Bruffaerts R, Posada-Villa J, Gasquet I, Kovess V, Lepine JP, Angermeyer MC, Bernert S, de Girolamo G, Morosini P, Polidori G, Kikkawa T, Kawakami N, Ono Y, Takeshima T, Uda H, Karam EG, Fayyad JA, Karam AN, Mneimneh ZN, Medina-Mora ME, Borges G, Lara C, de Graaf R, Ormel J, Gureje O, Shen Y, Huang Y, Zhang M, Alonso J, Haro JM, Vilagut G, Bromet EJ, Gluzman S, Webb C, Kessler RC, Merikangas KR, Anthony JC, Von Korff MR, Wang PS, Brugha TS, Aguilar-Gaxiola S, Lee S, Heeringa S, Pennell BE, Zaslavsky AM, Ustun TB, Chatterji S; WHO World Mental Health Survey Consortium** (2004). Prevalence, severity, and unmet need for treatment of mental disorders in the World Health Organization World Mental Health Surveys. *Journal of the American Medical Association* 291, 2581–2590.
- Department of Health** (2002). *Community Mental Health Teams, Policy Implementation Guidance*. Department of Health: London.
- Desjarlais R, Eisenberg L, Good B, Kleinman A** (1995). *World Mental Health. Problems and Priorities in Low-Income Countries*. Oxford University Press: Oxford.
- Deva MP** (2008). Bringing changes to Asian mental health. *International Review of Psychiatry* 20, 484–487.
- Dieterich M, Irving CB, Park B, Marshall M** (2010). Intensive case management for severe mental illness. *Cochrane Database of Systematic Reviews* 10, CD007906.
- Drake RE, Latimer E** (2012). Lessons learned in developing community mental health care in North America. *World Psychiatry* 11, 47–51.
- Drake RE, McHugo GJ, Bebout RR, Becker DR, Harris M, Bond GR, Quimby E** (1999). A randomized clinical trial of supported employment for inner-city patients with severe mental disorders. *Archives of General Psychiatry* 56, 627–633.
- Dua T, Barbui C, Clark N, Fleischmann A, van Ommeren M, Poznyak V, Yasamy MT, Thornicroft G, Saxena S** (2011). Evidence-based guidelines for mental, neurological and substance use disorders in low- and middle-income countries: summary of WHO recommendations. *PLoS Medicine* 8, 1–11.
- Eaton J** (2008). Ensuring access to psychotropic medication in sub-Saharan Africa. *African Journal of Psychiatry* 11, 179–181.
- Eaton J, McCay L, Semrau M, Chatterjee S, Baingana F, Araya R, Ntulo C, Thornicroft G, Saxena S** (2011). Scale up of services for mental health in low-income and middle-income countries. *Lancet* 378, 1592–1603.
- Fiander M, Burns T, McHugo GJ, Drake RE** (2003). Assertive community treatment across the Atlantic: comparison of model fidelity in the UK and USA. *British Journal of Psychiatry* 182, 248–254.
- Furedi J, Mohr P, Swingler D, Bitter I, Gheorghe MD, Hotujac L, Jarema M, Kocmur M, Koychev GI, Mosolov SN, Pecenek J, Rybakowski J, Svestka J, Sartorius N** (2006). Psychiatry in selected countries of Central and Eastern Europe: an overview of the current situation. *Acta Psychiatrica Scandinavica* 114, 223–231.
- Gafoor R, Nitsch D, McCrone P, Craig TK, Garety PA, Power P, McGuire P** (2010). Effect of early intervention

- on 5-year outcome in non-affective psychosis. *British Journal of Psychiatry* **196**, 372–376.
- Gask L** (2005). Overt and covert barriers to the integration of primary and specialist mental health care. *Social Science and Medicine* **61**, 1785–1794.
- Ghosh R, Killaspy H** (2010). A national survey of assertive community treatment services in England. *Journal of Mental Health* **19**, 500–508.
- Glover G, Arts G, Babu KS** (2006). Crisis resolution/home treatment teams and psychiatric admission rates in England. *British Journal of Psychiatry* **189**, 441–445.
- Greenhalgh T** (2008). Thirty years on from Alma-Ata: Where have we come from? Where are we going? *British Journal of General Practice* **58**, 798–804.
- Grigoletti L, Perini G, Rossi A, Biggeri A, Barbui C, Tansella M, Amaddeo F** (2009). Mortality and cause of death among psychiatric patients: a 20-year case-register study in an area with a community-based system of care. *Psychological Medicine* **39**, 1875–1884.
- Hanlon C, Wondimagegn D, Alem A** (2010). Lessons learned in developing community mental health care in Africa. *World Psychiatry* **9**, 185–189.
- Heffernan J, Pilkington P** (2011). Supported employment for persons with mental illness: systematic review of the effectiveness of individual placement and support in the UK. *Journal of Mental Health* **20**, 368–380.
- Heslin M, Howard L, Leese M, McCrone P, Rice C, Jarrett M, Spokes T, Huxley P, Thornicroft G** (2011). Randomized controlled trial of supported employment in England: 2 year follow-up of the Supported Work and Needs (SWAN) study. *World Psychiatry* **10**, 132–137.
- Holloway F, Sederer L** (2011). Inpatient treatment. In *Oxford Textbook of Community Mental Health* (ed. G. Thornicroft, G. I. Szmukler, K. T. Mueser and R. E. Drake), pp. 223–231. Oxford University Press: Oxford.
- Howard LM, Heslin M, Leese M, McCrone P, Rice C, Jarrett M, Spokes T, Huxley P, Thornicroft G** (2010). Supported employment: randomised controlled trial. *British Journal of Psychiatry* **196**, 404–411.
- Howard LM, Leese M, Byford S, Killaspy H, Cole L, Lawlor C, Johnson S** (2009). Methodological challenges in evaluating the effectiveness of women's crisis houses compared with psychiatric wards: findings from a pilot patient preference RCT. *Journal of Nervous and Mental Disease* **197**, 722–727.
- Howard LM, Rigon E, Cole L, Lawlor C, Johnson S** (2008). Admission to women's crisis houses or to psychiatric wards: women's pathways to admission. *Psychiatric Services* **59**, 1443–1449.
- Institute of Medicine** (2001). *Crossing the Quality Chasm: A New Health System for the 21st Century*. Institute of Medicine: Washington, DC.
- Ito H, Setoya Y, Suzuki Y** (in press). Lessons learned in developing community mental health care in East and Southeast Asia. *World Psychiatry*.
- Jacob KS, Sharan P, Mirza I, Garrido-Cumbrera M, Seedat S, Mari JJ, Sreenivas V, Saxena S** (2007). Mental health systems in countries: where are we now? *Lancet* **370**, 1061–1077.
- Janse van Rensburg AB** (2009). A changed climate for mental health care delivery in South Africa. *African Journal of Psychiatry* **12**, 157–165.
- Jethwa K, Galappathie N, Hewson P** (2007). Effects of a crisis resolution and home treatment team on in-patient admissions. *Psychiatric Bulletin* **31**, 170–172.
- Johnson S, Gilbert H, Lloyd-Evans B, Osborn DP, Boardman J, Leese M, Shepherd G, Thornicroft G, Slade M** (2009). In-patient and residential alternatives to standard acute psychiatric wards in England. *British Journal of Psychiatry* **194**, 456–463.
- Johnson S, Needle J, Bindman J, Thornicroft G** (2008). *Crisis Resolution and Home Treatment in Mental Health*. Cambridge University Press: Cambridge.
- Johnson S, Nolan F, Hoult J, White IR, Bebbington P, Sandor A, McKenzie N, Patel SN, Pilling S** (2005a). Outcomes of crises before and after introduction of a crisis resolution team. *British Journal of Psychiatry* **187**, 68–75.
- Johnson S, Nolan F, Pilling S, Sandor A, Hoult J, McKenzie N, White IR, Thompson M, Bebbington P** (2005b). Randomised controlled trial of acute mental health care by a crisis resolution team: the north Islington crisis study. *British Medical Journal* **331**, 599.
- Johnson S, Thornicroft G** (2008). The classic home treatment studies. In *Crisis Resolution and Home Treatment in Mental Health* (ed. S. Johnson, J. Needle, J. Bindman and G. Thornicroft), pp. 37–50. Cambridge University Press: Cambridge.
- Johnson S, Totman J, Hobbs L** (2011). Crisis and emergency services. In *Oxford Textbook of Community Mental Health* (ed. G. Thornicroft, G. Szmukler, K. Mueser and R. E. Drake), pp. 118–128. Oxford University Press: Oxford.
- Johnstone P, Zolese G** (1999). Systematic review of the effectiveness of planned short hospital stays for mental health care. *British Medical Journal* **318**, 1387–1390.
- Joy CB, Adams CE, Rice K** (2006). Crisis intervention for people with severe mental illnesses. *Cochrane Database of Systematic Reviews* **4**, CD001087.
- Keown P, Tacchi MJ, Niemiec S, Hughes J** (2007). Changes to mental healthcare for working age adults: impact of a crisis team and an assertive outreach team. *Psychiatric Bulletin* **31**, 288–292.
- Kessler RC, Aguilar-Gaxiola S, Alonso J, Chatterji S, Lee S, Ormel J, Ustun TB, Wang PS** (2009). The global burden of mental disorders: an update from the WHO World Mental Health (WMH) surveys. *Epidemiologia e Psichiatria Sociale* **18**, 23–33.
- Kessler RC, Chiu WT, Demler O, Merikangas KR, Walters EE** (2005a). Prevalence, severity, and comorbidity of 12-month DSM-IV disorders in the National Comorbidity Survey Replication. *Archives of General Psychiatry* **62**, 617–627.
- Kessler RC, Demler O, Frank RG, Olfson M, Pincus HA, Walters EE, Wang P, Wells KB, Zaslavsky AM** (2005b). Prevalence and treatment of mental disorders, 1990 to 2003. *New England Journal of Medicine* **352**, 2515–2523.
- Killackey E** (2011). Review: early intervention services can be clinically beneficial for people with early psychosis. *Evidence-Based Mental Health* **14**, 50.

- Killaspy H, Bebbington P, Blizard R, Johnson S, Nolan F, Pilling S, King M (2006). The REACT study: randomised evaluation of assertive community treatment in north London. *British Medical Journal* **332**, 815–820.
- Killaspy H, Dalton J, McNicholas S, Johnson S (2000). Drayton Park, an alternative to hospital admission for women in acute mental health crisis. *Psychiatric Bulletin* **24**, 101–104.
- Killaspy H, Kingett S, Bebbington P, Blizard R, Johnson S, Nolan F, Pilling S, King M (2009). Randomised evaluation of assertive community treatment: 3-year outcomes. *British Journal of Psychiatry* **195**, 81–82.
- Killaspy H, Rosen A (2011). Case management and assertive community treatment. In *Oxford Textbook of Community Mental Health* (ed. G. Thornicroft, G. I. Szmukler, K. T. Mueser and R. E. Drake), pp. 171–184. Oxford University Press: Oxford.
- Knapp M, Chisholm D, Astin J, Lelliott P, Audini B (1997). The cost consequences of changing the hospital-community balance: the mental health residential care study. *Psychological Medicine* **27**, 681–692.
- Knapp M, McDaid D, Mossialos E, Thornicroft G (eds) (2007). *Mental Health Policy and Practice Across Europe*. Open University Press: Maidenhead.
- Kohn R, Saxena S, Levav I, Saraceno B (2004). Treatment gap in mental health care. *Bulletin of the World Health Organization* **82**, 858–866.
- Larsen TK, Melle I, Auestad B, Haahr U, Joa I, Johannessen JO, Opjordsmoen S, Rund BR, Rossberg JI, Simonsen E, Vaglum P, Friis S, McGlashan T (2011). Early detection of psychosis: positive effects on 5-year outcome. *Psychological Medicine* **41**, 1461–1469.
- Lasalvia A, Tansella M (2010). Acute in-patient care in modern, community-based mental health services. Where and how? *Epidemiologia e Psichiatria Sociale* **19**, 275–281.
- Latimer E (2005). Economic considerations associated with assertive community treatment and supported employment for people with severe mental illness [Review]. *Journal of Psychiatry and Neuroscience* **30**, 355–359.
- Latimer EA (1999). Economic impacts of assertive community treatment: a review of the literature. *Canadian Journal of Psychiatry* **44**, 443–454.
- Lawlor C, Johnson S, Cole L, Howard LM (2010). Ethnic variations in pathways to acute care and compulsory detention for women experiencing a mental health crisis. *International Journal of Social Psychiatry* **58**, 3–15.
- Lawn JE, Rohde J, Rifkin S, Were M, Paul VK, Chopra M (2008). Alma-Ata 30 years on: revolutionary, relevant, and time to revitalise. *Lancet* **372**, 917–927.
- Lehman AF (1995). Vocational rehabilitation in schizophrenia. *Schizophrenia Bulletin* **21**, 645–656.
- Lehman AF, Goldberg R, Dixon LB, McNary S, Postrado L, Hackman A, McDonnell K (2002). Improving employment outcomes for persons with severe mental illnesses. *Archives of General Psychiatry* **59**, 165–172.
- Lelliott P, Bleksley S (2010). Improving the quality of acute inpatient care. *Epidemiologia e Psichiatria Sociale* **19**, 287–290.
- Lloyd-Evans B, Slade M, Jagielska D, Johnson S (2009). Residential alternatives to acute psychiatric hospital admission: systematic review. *British Journal of Psychiatry* **195**, 109–117.
- Lund C, De Silva M, Plagerson S, Cooper S, Chisholm D, Das J, Knapp M, Patel V (2011). Poverty and mental disorders: breaking the cycle in low-income and middle-income countries. *Lancet* **378**, 1502–1514.
- Malone D, Newron-Howes G, Simmonds S, Marriot S, Tyrer P (2007). Community mental health teams (CMHTs) for people with severe mental illnesses and disordered personality [Review]. *Cochrane Database of Systematic Reviews* **3**, CD000270.
- Marshall M, Crowther R, Almaraz-Serrano A, Creed F, Sledge W, Kluiters H, Roberts C, Hill E, Wiersma D, Bond GR, Huxley P, Tyrer P (2001). Systematic reviews of the effectiveness of day care for people with severe mental disorders: (1) acute day hospital versus admission; (2) vocational rehabilitation; (3) day hospital versus outpatient care. *Health Technology Assessment* **5**, 1–75.
- Marshall M, Crowther R, Sledge WH, Rathbone J, Soares-Weiser K (2011). Day hospital versus admission for acute psychiatric disorders. *Cochrane Database of Systematic Review* **12**, CD004026.
- Marshall M, Lockwood A (2000). Assertive community treatment for people with severe mental disorders. *Cochrane Database of Systematic Reviews* **2**, CD001089.
- Marshall M, Rathbone J (2011). Early intervention for psychosis. *Cochrane Database of Systematic Reviews* **6**, CD004718.
- Marwaha S, Johnson S (2005). Views and experiences of employment among people with psychosis: a qualitative descriptive study. *International Journal of Social Psychiatry* **51**, 302–316.
- Mbuba CK, Newton CR (2009). Packages of care for epilepsy in low- and middle-income countries. *PLoS Medicine* **6**, e1000162.
- McCrone P, Johnson S, Nolan F, Pilling S, Sandor A, Hoult J, McKenzie N, Thompson M, Bebbington P (2009a). Economic evaluation of a crisis resolution service: a randomised controlled trial. *Epidemiologia e Psichiatria Sociale* **18**, 54–58.
- McCrone P, Johnson S, Nolan F, Pilling S, Sandor A, Hoult J, McKenzie N, White IR, Bebbington P (2009c). Impact of a crisis resolution team on service costs in the UK. *Psychiatric Bulletin* **33**, 17–19.
- McCrone P, Killaspy H, Bebbington P, Johnson S, Nolan F, Pilling S, King M (2009b). The REACT study: cost-effectiveness analysis of assertive community treatment in north London. *Psychiatric Services* **60**, 908–913.
- McGeorge P (2012). Lessons learned in developing community mental health care in Australasia and the South Pacific. *World Psychiatry* **11**, 129–132.
- McGorry P (2012). At issue: Cochrane, early intervention, and mental health reform: analysis, paralysis, or evidence-informed progress? *Schizophrenia Bulletin* **38**, 221–224.
- Miller G (2006). Mental health in developing countries. The unseen: mental illness's global toll. *Science* **311**, 458–461.
- Mosher LR (1999). Soteria and other alternatives to acute psychiatric hospitalization: a personal and professional review [Review]. *Journal of Nervous and Mental Disease* **187**, 142–149.

- Mubbashar M** (1999). Mental health services in rural Pakistan. In *Common Mental Disorders in Primary Care* (ed. M. Tansella and G. Thornicroft), pp. 67–80. Routledge: London.
- Murphy S, Irving CB, Adams CE, Driver R** (2012). Crisis intervention for people with severe mental illnesses. *Cochrane Database of Systematic Reviews* 5, CD001087.
- Nathan P, Gorman J** (2002). *A Guide to Treatments that Work*, 2nd edn. Oxford University Press: Oxford.
- Ndyanabangi S, Basangwa D, Lutakome J, Mubiru C** (2004). Uganda mental health country profile. *International Review of Psychiatry* 16, 54–62.
- Nelson T, Johnson S, Bebbington P** (2009). Satisfaction and burnout among staff of crisis resolution, assertive outreach and community mental health teams. *Social Psychiatry and Psychiatric Epidemiology* 44, 541–549.
- Nestor P, Galletly C** (2008). The employment of consumers in mental health services: politically correct tokenism or genuinely useful? *Australasian Psychiatry* 16, 344–347.
- Ngoma MC, Prince M, Mann A** (2003). Common mental disorders among those attending primary health clinics and traditional healers in urban Tanzania. *British Journal of Psychiatry* 183, 349–355.
- Njenga F** (2002). Challenges of balanced care in Africa. *World Psychiatry* 1, 96–98.
- Ormel J, Petukhova M, Chatterji S, Aguilar-Gaxiola S, Alonso J, Angermeyer MC, Bromet EJ, Burger H, Demyttenaere K, de Girolamo G, Haro JM, Hwang I, Karam E, Kawakami N, Lépine JP, Medina-Mora ME, Posada-Villa J, Sampson N, Scott K, Ustün TB, Von Korff M, Williams DR, Zhang M, Kessler RC** (2008). Disability and treatment of specific mental and physical disorders across the world. *British Journal of Psychiatry* 192, 368–375.
- Ormel J, Von Korff M, Ustun B, Pini S, Korten A** (1994). Common mental disorders and disability across cultures. Results from the WHO Collaborative Study on Psychological Problems in General Health Care. *Journal of the American Medical Association* 272, 1741–1748.
- Passmore R** (1979). The declaration of Alma-Ata and the future of primary care. *Lancet* 2, 1005–1008.
- Patel V, Araya R, Chatterjee S, Chisholm D, Cohen A, De Silva M, Hosman C, McGuire H, Rojas G, van Ommeren M** (2007). Treatment and prevention of mental disorders in low-income and middle-income countries. *Lancet* 370, 991–1005.
- Patel V, Maj M, Flisher AJ, De Silva MJ, Koschorke M, Prince M** (2010a). Reducing the treatment gap for mental disorders: a WPA survey. *World Psychiatry* 9, 169–176.
- Patel V, Simon G, Chowdhary N, Kaaya S, Araya R** (2009). Packages of care for depression in low- and middle-income countries. *PLoS Medicine* 6, e1000159.
- Patel V, Thornicroft G** (2009). Packages of care for mental, neurological, and substance use disorders in low- and middle-income countries. *PLoS Medicine* 6, e1000160.
- Patel V, Weiss HA, Chowdhary N, Naik S, Pednekar S, Chatterjee S, De Silva MJ, Bhat B, Araya R, King M, Simon G, Verdelli H, Kirkwood BR** (2010b). Effectiveness of an intervention led by lay health counsellors for depressive and anxiety disorders in primary care in Goa, India (MANAS): a cluster randomised controlled trial. *Lancet* 376, 2086–2095.
- Phillips SD, Burns BJ, Edgar ER, Mueser KT, Linkins KW, Rosenheck RA, Drake RE, McDonel Herr EC** (2001). Moving assertive community treatment into standard practice. *Psychiatric Services* 52, 771–779.
- Pirkola S, Sund R, Sailas E, Wahlbeck K** (2009). Community mental-health services and suicide rate in Finland: a nationwide small-area analysis. *Lancet* 373, 147–153.
- Pollock AM** (2010). Should NHS mental health services fear the private sector? Yes. *British Medical Journal* 341, c5382.
- Power P, McGorry P** (2011). Early interventions for people with psychotic disorders. In *Oxford Textbook of Community Mental Health* (ed. G. Thornicroft, G. I. Szmukler, K. T. Mueser and R. E. Drake), pp. 159–160. Oxford University Press: Oxford.
- Preti A, Cella M** (2010). Randomized-controlled trials in people at ultra high risk of psychosis: a review of treatment effectiveness. *Schizophrenia Research* 123, 30–36.
- Priebe S, Warner R, Hubschmid T, Eckle I** (1998). Employment, attitudes toward work, and quality of life among people with schizophrenia in three countries. *Schizophrenia Bulletin* 24, 469–477.
- Prince M, Patel V, Saxena S, Maj M, Maselko J, Phillips MR, Rahman A** (2007). No health without mental health. *Lancet* 370, 859–877.
- Quirk A, Lelliott P** (2001). What do we know about life on acute psychiatric wards in the UK? A review of the research evidence. *Social Science and Medicine* 53, 1565–1574.
- Raguram R, Venkateswaran A, Ramakrishna J, Weiss MG** (2002). Traditional community resources for mental health: a report of temple healing from India. *British Medical Journal* 325, 38–40.
- Raune D, Kuipers E, Bebbington PE** (2004). Expressed emotion at first-episode psychosis: investigating a carer appraisal model. *British Journal of Psychiatry* 184, 321–326.
- Razali SM** (2004). Deinstitutionalization and community mental health services in Malaysia: an overview. *International Medical Journal* 11, 29–35.
- Razzouk D, Gregorio G, Antunes R, Mari J** (in press). Lessons learned in developing community mental health care in Latin American and Caribbean Countries. *World Psychiatry*.
- Repper J, Carter T** (2011). A review of the literature on peer support in mental health services. *Journal of Mental Health* 20, 392–411.
- Rinaldi M, Perkins R** (2007). Comparing employment outcomes for two vocational services: individual placement and support and non-integrated pre-vocational services in the UK. *Journal of Vocational Rehabilitation* 27, 21–27.
- Roberts L, Roalfe A, Wilson S, Lester H** (2007). Physical health care of patients with schizophrenia in primary care: a comparative study. *Family Practice* 24, 34–40.
- Robinson J, Bruxner A, Harrigan S, Bendall S, Killackey E, Tonin V, Monson K, Thurley M, Francey S, Yung AR** (2010). Study protocol: the development of a pilot study employing a randomised controlled design to investigate the feasibility and effects of a peer support program

- following discharge from a specialist first-episode psychosis treatment centre. *BMC Psychiatry* **10**, 37.
- Rodriguez JJ** (2010). Mental health care systems in Latin America and the Caribbean. *International Review of Psychiatry* **22**, 317–324.
- Rohde J, Cousens S, Chopra M, Tangcharoensathien V, Black R, Bhutta ZA, Lawn JE** (2008). 30 years after Alma-Ata: has primary health care worked in countries? *Lancet* **372**, 950–961.
- Rosen A, Barfoot K** (2001). Day care and occupation: structured rehabilitation and recovery programmes and work. In *Textbook of Community Psychiatry* (ed. G. Thornicroft and G. Szmukler), pp. 296–308. Oxford University Press: Oxford.
- Roth A, Fonagy P** (1996). *What Works for Whom? A Critical Review of Psychotherapy Research*. Guildford Press: New York.
- Saraceno B, van Ommeren M, Batniji R, Cohen A, Gureje O, Mahoney J, Sridhar D, Underhill C** (2007). Barriers to improvement of mental health services in low-income and middle-income countries. *Lancet* **370**, 1164–1174.
- Saxena S, Maulik P** (2003). Mental health services in low and middle income countries: an overview. *Current Opinion in Psychiatry* **16**, 437–442.
- Saxena S, Thornicroft G, Knapp M, Whiteford H** (2007). Resources for mental health: scarcity, inequity, and inefficiency. *Lancet* **370**, 878–889.
- Sederer LI** (2010). Inpatient psychiatry: why do we need it? *Epidemiologia e Psichiatria Sociale* **19**, 291–295.
- Selolilwe ES, Thupayagale-Tshweneagae G** (2007). Community mental health care in Botswana: approaches and opportunities. *International Nursing Review* **54**, 173–178.
- Semrau M, Barley E, Law A, Thornicroft G** (2011). Lessons learned in developing community mental health care in Europe. *World Psychiatry* **10**, 217–225.
- Shankar BR, Saravanan B, Jacob KS** (2006). Explanatory models of common mental disorders among traditional healers and their patients in rural south India. *International Journal of Social Psychiatry* **52**, 221–233.
- Sharifi V** (2009). Urban mental health in Iran: challenges and future directions. *Iranian Journal of Psychiatry and Behavioral Sciences* **3**, 9–14.
- Shepherd G** (1990). *Theory and Practice of Psychiatric Rehabilitation*. Wiley: Chichester.
- Shepherd G, MacPherson R** (2011). Residential care. In *Oxford Textbook of Community Psychiatry* (ed. G. Thornicroft, G. Szmukler, K. Mueser and R. E. Drake), pp. 178–187. Oxford University Press: Oxford.
- Shibre T, Spangeus A, Henriksson L, Negash A, Jacobsson L** (2008). Traditional treatment of mental disorders in rural Ethiopia. *Ethiopian Medical Journal* **46**, 87–91.
- Simmonds S, Coid J, Joseph P, Marriott S, Tyrer P** (2001). Community mental health team management in severe mental illness: a systematic review. *British Journal of Psychiatry* **178**, 497–502.
- Slade M** (2009). *Personal Recovery and Mental Illness. A Guide for Mental Health Professionals*. Cambridge University Press: Cambridge.
- Slade M, Leamy M, Bacon F, Janosik M, le Boutilier C, Williams J, Bird V** (2012). International differences in understanding recovery: systematic review. *Epidemiology and Psychiatric Sciences*. Published online: 15 March 2012. doi:http://dx.doi.org/10.1017/S2045796012000133.
- Sledge WH, Tebes J, Rakfeldt J, Davidson L, Lyons L, Druss B** (1996a). Day hospital/crisis respite care versus inpatient care, Part I: Clinical outcomes. *American Journal of Psychiatry* **153**, 1065–1073.
- Sledge WH, Tebes J, Wolff N, Helminiak TW** (1996b). Day hospital/crisis respite care versus inpatient care, Part II: Service utilization and costs. *American Journal of Psychiatry* **153**, 1074–1083.
- Sytema S, Micciolo R, Tansella M** (1997). Continuity of care for patients with schizophrenia and related disorders: a comparative South-Verona and Groningen case-register study. *Psychological Medicine* **27**, 1355–1362.
- Tansella M, Thornicroft G** (2009). Implementation science: understanding the translation of evidence into practice. *British Journal of Psychiatry* **195**, 283–285.
- Teague GB, Bond GR, Drake RE** (1998). Program fidelity in assertive community treatment: development and use of a measure. *American Journal of Orthopsychiatry* **68**, 216–232.
- Thara R, Padvamati R** (in press). Lessons learned in developing community mental health care in South Asia. *World Psychiatry*.
- The ITHACA Study Group** (2011). *The ITHACA Toolkit for Monitoring Human Rights and General Health Care in Mental Health and Social Care Institutions* (www.ithacastudy.eu). King's College London: London.
- Thornicroft G** (2007). Most people with mental illness are not treated. *Lancet* **370**, 807–808.
- Thornicroft G, Alem A, Antunes Dos Santos R, Barley E, Drake RE, Gregorio G, Hanlon C, Ito H, Latimer E, Law A, Mari J, McGeorge P, Padmavati R, Razzouk D, Semrau M, Setoya Y, Thara R, Wondimagegn D** (2010). WPA guidance on steps, obstacles and mistakes to avoid in the implementation of community mental health care. *World Psychiatry* **9**, 67–77.
- Thornicroft G, Bebbington P, Leff J** (2005). Outcomes for long-term patients one year after discharge from a psychiatric hospital. *Psychiatric Services* **56**, 1416–1422.
- Thornicroft G, Becker T, Holloway F, Johnson S, Leese M, McCrone P, Szmukler G, Taylor R, Wykes T** (1999). Community mental health teams: evidence or belief? *British Journal of Psychiatry* **175**, 508–513.
- Thornicroft G, Lempp H, Tansella M** (2011c). The place of implementation science in the translational medicine continuum. *Psychological Medicine* **41**, 2015–2021.
- Thornicroft G, Rose D, Huxley P, Dale G, Wykes T** (2002). What are the research priorities of mental health service users? *Journal of Mental Health* **11**, 1–5.
- Thornicroft G, Semrau M, Alem A, Drake RE, Ito H, Mari J, McGeorge P, Thara P** (2011a). *Global Mental Health: Putting Community Care into Practice*. Wiley-Blackwell: London.
- Thornicroft G, Szmukler G, Mueser K, Drake R** (2011b). *Oxford Textbook of Community Mental Health*. Oxford University Press: Oxford.
- Thornicroft G, Tansella M** (1999). *The Mental Health Matrix: A Manual to Improve Services*. Cambridge University Press: Cambridge.

- Thornicroft G, Tansella M (2004). Components of a modern mental health service: a pragmatic balance of community and hospital care: overview of systematic evidence. *British Journal of Psychiatry* **185**, 283–290.
- Thornicroft G, Tansella M (2009). *Better Mental Health Care*. Cambridge University Press: Cambridge.
- Thornicroft G, Tansella M, Becker T, Knapp M, Leese M, Schene A, Vazquez-Barquero JL (2004). The personal impact of schizophrenia in Europe. *Schizophrenia Research* **69**, 125–132.
- Thornicroft G, Tansella M, Law A (2008). Steps, challenges and lessons in developing community mental health care. *World Psychiatry* **7**, 87–92.
- Thornicroft G, Wykes T, Holloway F, Johnson S, Szukler G (1998). From efficacy to effectiveness in community mental health services. PRISM Psychosis Study. 10. *British Journal of Psychiatry* **173**, 423–427.
- Tiihonen J, Lonnqvist J, Wahlbeck K, Klaukka T, Niskanen L, Tanskanen A, Haukka J (2009). 11-year follow-up of mortality in patients with schizophrenia: a population-based cohort study (FIN11 study). *Lancet* **374**, 620–627.
- Tomov T (2001). Central and Eastern European countries. In *The Mental Health Matrix. A Manual to Improve Services* (ed. G. Thornicroft and M. Tansella), pp. 216–227. Cambridge University Press: Cambridge.
- Totman J, Mann F, Johnson S (2010). Is locating acute wards in the general hospital an essential element in psychiatric reform? The U.K. experience. *Epidemiologia e Psichiatria Sociale* **19**, 282–286.
- Tyrer P, Evans K, Gandhi N, Lamont A, Harrison-Read P, Johnson T (1998). Randomised controlled trial of two models of care for discharged psychiatric patients. *British Medical Journal* **316**, 106–109.
- Tyrer P, Morgan J, Van Horn E, Jayakody M, Evans K, Brummell R, White T, Baldwin D, Harrison-Read P, Johnson T (1995). A randomised controlled study of close monitoring of vulnerable psychiatric patients. *Lancet* **345**, 756–759.
- Tyrer S, Coid J, Simmonds S, Joseph P, Marriott S (2007). Community mental health teams (CMHTs) for people with severe mental illnesses and disordered personality. *Cochrane Database of Systematic Reviews* **3**, CD000270.
- van Wijngaarden GK, Schene A, Koeter M, Becker T, Knudsen HC, Tansella M, Thornicroft G, Vazquez-Barquero JL; EPSILON Study Group (2003). People with schizophrenia in five European countries: conceptual similarities and intercultural differences in family caregiving. *Schizophrenia Bulletin* **29**, 573–586.
- Wahlbeck K, Westman J, Nordentoft M, Gissler M, Laursen TM (2011). Outcomes of Nordic mental health systems: life expectancy of patients with mental disorders. *British Journal of Psychiatry* **199**, 453–458.
- Wang PS, Angermeyer M, Borges G, Bruffaerts R, Tat CW, de Girolamo G, Fayyad J, Gureje O, Haro JM, Huang Y, Kessler RC, Kovess V, Levinson D, Nakane Y, Oakley Brown MA, Ormel JH, Posada-Villa J, Aguilar-Gaxiola S, Alonso J, Lee S, Heeringa S, Pennell BE, Chatterji S, Ustün TB (2007a). Delay and failure in treatment seeking after first onset of mental disorders in the World Health Organization's World Mental Health Survey Initiative. *World Psychiatry* **6**, 177–185.
- Wang PS, Aguilar-Gaxiola S, Alonso J, Angermeyer MC, Borges G, Bromet EJ, Bruffaerts R, de Girolamo G, de Graaf R, Gureje O, Haro JM, Karam EG, Kessler RC, Kovess V, Lane MC, Lee S, Levinson D, Ono Y, Petukhova M, Posada-Villa J, Seedat S, Wells JE (2007b). Use of mental health services for anxiety, mood, and substance disorders in 17 countries in the WHO world mental health surveys. *Lancet* **370**, 841–850.
- Warner R (2004). *Recovery from Schizophrenia: Psychiatry and Political Economy*. Brunner-Routledge: Hove.
- While D, Bickley H, Roscoe A, Windfuhr K, Rahman S, Shaw J, Appleby L, Kapur N (2012). Implementation of mental health service recommendations in England and Wales and suicide rates, 1997–2006: a cross-sectional and before-and-after observational study. *Lancet* **379**, 1005–1012.
- WHO (2001). *World Health Report 2001. Mental Health: New Understanding, New Hope*. World Health Organization: Geneva.
- WHO (2005). *Mental Health Atlas: Revised Edition*. World Health Organization: Geneva.
- WHO (2008). *mhGAP. Mental Health Gap Action Programme: Scaling Up Care for Mental, Neurological, and Substance Use Disorders*. World Health Organization: Geneva.
- WHO (2010). *mhGAP Intervention Guide for Mental, Neurological and Substance Use Disorders in Non-Specialized Health Settings*. World Health Organization: Geneva.
- Wittchen HU, Jacobi F (2005). Size and burden of mental disorders in Europe – a critical review and appraisal of 27 studies. *European Neuropsychopharmacology* **15**, 357–376.
- Wittchen HU, Jacobi F, Rehm J, Gustavsson A, Svensson M, Jönsson B, Olesen J, Allgulander C, Alonso J, Faravelli C, Fratiglioni L, Jennum P, Lieb R, Maercker A, van Os J, Preisig M, Salvador-Carulla L, Simon R, Steinhausen HC (2011). The size and burden of mental disorders and other disorders of the brain in Europe 2010. *European Neuropsychopharmacology* **21**, 655–679.
- World Bank (2010). *World Bank List of Economies* (<http://data.worldbank.org/about/country-classifications>). World Bank: Washington, DC.
- Yamey G (2011). Scaling up global health interventions: a proposed framework for success. *PLoS Medicine* **8**, e1001049.
- Younes N, Hardy-Bayle MC, Falissard B, Kovess V, Chaillet MP, Gasquet I (2005). Differing mental health practice among general practitioners, private psychiatrists and public psychiatrists. *BMC Public Health* **5**, 104.